CLAIMS

1. Use of N-[(1S)-2,2,5,7-tetrafluoro-2,3-dihydro-1H-inden-1-yl]sulfamide



(Ia)


or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the treatment of epilepsy, which is to be administered in combination with an AMPA-type glutamate receptor antagonist.

2. Use according to claim 1, wherein N-[(1S)-2,2,5,7-tetrafluoro-2,3-dihydro-1H-inden-1-yl]sulfamide or a pharmaceutically acceptable salt thereof and the AMPA-type glutamate receptor antagonist are to be administered simultaneously or separately.

3. Use according to claim 1, wherein N-[(1S)-2,2,5,7-tetrafluoro-2,3-dihydro-1H-inden-1-yl]sulfamide or a pharmaceutically acceptable salt thereof and the AMPA-type glutamate receptor antagonist are to be administered simultaneously.


4. Use according to claim 1 or 2, wherein the AMPA-type glutamate receptor antagonist is a compound selected from the following group consisting of:

3-(2-cyanophenyl)-5-(2-pyridyl)-1-phenyl-1,2-dihydropyridin-2-one,




(II)

9-(2-chlorophenyl)-7-(pyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(III)

2-fluoro-6-(7-(5-methoxypyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(IV)

2-fluoro-6-(7-(6-methylpyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(V)

9-(2-chloro-3-fluorophenyl)-7-(6-methylpyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(VI)

2-fluoro-6-(7-(2-methoxypyrimidin-5-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(VII)

7-(pyridin-3-yl)-9-(2,3,5,6-tetrafluorophenyl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(VIII)

3-(8-oxo-7-(thiophen-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(IX)

3-(8-oxo-7-(thiophen-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)pyrazine-2-carbonitrile,




(X)

9-(2-fluorophenyl)-7-phenyl-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(XI)

2-(7-(4-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XII)

3-(7-(4-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XIII)

3-(7-(2-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XIV)

3-(3-fluoro-8-oxo-7-phenyl-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XV)

2-fluoro-6-(3-fluoro-8-oxo-7-(pyridin-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)ben
zonitrile,



(XVI)

2-fluoro-6-(7-(5-fluoropyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)ben
zonitrile,



(XVII)


2-fluoro-6-(10-fluoro-3-oxo-4-(pyridin-3-yl)-4,5-dihydro-3H-chromeno[3,4-b]pyridin-2-yl)benzonitrile,



(XVIII)


9-(2-chloro-3-fluorophenyl)-7-(5-fluoropyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one

,




(XIX)

2-fluoro-6-(8-oxo-7-(pyrimidin-5-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XX)

3,6-difluoro-2-(8-oxo-7-(pyridin-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XXI)

2-(7-(5-chloropyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)-6-fluoroben zonitrile,



(XXII)

2-fluoro-6-(7-(2-methylpyrimidin-5-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl) benzonitrile, and



(XXIII)


9-(3-fluoro-2-methylphenyl)-7-(pyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one



(XXIV)

or a pharmaceutically acceptable salt thereof.

5. Use according to claim 1 or 2, wherein the AMPA-type glutamate receptor antagonist is 3-(2-cyanophenyl)-5-(2-pyridyl)-1-phenyl-1,2-dihydropyridin-2-one




(II)

or a pharmaceutically acceptable salt thereof.


6. Use according to claim 1 or 2, wherein the AMPA-type glutamate receptor antagonist is a compound selected from the following group consisting of:

9-(2-chlorophenyl)-7-(pyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,



(III)


2-fluoro-6-(7-(5-methoxypyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,



(IV)


2-fluoro-6-(7-(6-methylpyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)ben

zonitrile,




(V)

9-(2-chloro-3-fluorophenyl)-7-(6-methylpyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-on
e,




(VI)

2-fluoro-6-(7-(2-methoxypyrimidin-5-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-y
l)benzonitrile,




(VII)

7-(pyridin-3-yl)-9-(2,3,5,6-tetrafluorophenyl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(VIII)

3-(8-oxo-7-(thiophen-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(IX)

3-(8-oxo-7-(thiophen-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)pyrazine-2-carbonitrile,




(X)

9-(2-fluorophenyl)-7-phenyl-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(XI)

2-(7-(4-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,



(XII)

3-(7-(4-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XIII)

3-(7-(2-fluorophenyl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XIV)

3-(3-fluoro-8-oxo-7-phenyl-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)picolinonitrile,




(XV)

2-fluoro-6-(3-fluoro-8-oxo-7-(pyridin-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,



(XVI)

2-fluoro-6-(7-(5-fluoropyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XVII)

2-fluoro-6-(10-fluoro-3-oxo-4-(pyridin-3-yl)-4,5-dihydro-3H-chromeno[3,4-b]pyridin-2-yl)benzonitrile,




(XVIII)

9-(2-chloro-3-fluorophenyl)-7-(5-fluoropyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one,




(XIX)

2-fluoro-6-(8-oxo-7-(pyrimidin-5-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XX)

3,6-difluoro-2-(8-oxo-7-(pyridin-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,




(XXI)

2-(7-(5-chloropyridin-3-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)-6-fluorobenzonitrile,




(XXII)

2-fluoro-6-(7-(2-methylpyrimidin-5-yl)-8-oxo-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile,



(XXIII)


and 9-(3-fluoro-2-methylphenyl)-7-(pyridin-3-yl)-6H-pyrano[3,2-b:5,4-b']dipyridine-8(7H)-one



(XXIV)

or a pharmaceutically acceptable salt thereof.

7. Use according to claim 1 or 2, wherein the AMPA-type glutamate receptor antagonist is 2-fluoro-6-(3-fluoro-8-oxo-7-(pyridin-3-yl)-7,8-dihydro-6H-pyrano[3,2-b:5,4-b']dipyridin-9-yl)benzonitrile



(XVI)

or a pharmaceutically acceptable salt thereof.