


## COMBINATION THERAPY FOR CANCER TREATMENT


[0001] The present disclosure generally relates to combination therapies for cancer treatment, including administration of a pentaaza macrocyclic ring complex, and active agent that is at least one of a thioredoxin reductase inhibitor and a glutathione-depleting agent.

[0002] Transition metal-containing pentaaza macrocyclic ring complexes having the macrocyclic ring system corresponding to Formula A have been shown to be effective in a number of animal and cell models of human disease, as well as in treatment of conditions afflicting human patients.





FORMULA A

For example, in a rodent model of colitis, one such compound, GC4403, has been reported to very significantly reduce the injury to the colon of rats subjected to an experimental model of colitis (see Cuzzocrea et al., *Europ. J. Pharmacol.*, 432, 79-89 (2001)).



GC4403 has also been reported to attenuate the radiation damage arising both in a clinically relevant hamster model of acute, radiation-induced oral mucositis (Murphy et al., *Clin. Can. Res.*, 14(13), 4292 (2008)), and lethal total body irradiation of adult mice (Thompson et al., *Free Radical Res.*, 44(5), 529-40 (2010)). Similarly, another such compound, GC4419, has been shown to attenuate VEGFr inhibitor-induced pulmonary disease in a rat model (Tuder, et al., , *Am. J. Respir. Cell Mol. Biol.*, 29, 88-97 (2003)).

Additionally, another such compound, GC4401 has been shown to provide protective effects in animal models of septic shock (S. Cuzzocrea, et.al., *Crit. Care Med.*, 32(1), 157 (2004) and pancreatitis (S. Cuzzocrea, et.al., *Shock*, 22(3), 254-61 (2004)).



- 5            [0003] Certain of these compounds have also been shown to possess potent anti-inflammatory activity and prevent oxidative damage *in vivo*. For example, GC4403 has been reported to inhibit inflammation in a rat model of inflammation (Salvemini, et.al., *Science*, 286, 304 (1999)), and prevent joint disease in a rat model of collagen-induced arthritis (Salvemini et al., *Arthritis & Rheumatism*, 44(12), 2009-2021 (2001)).
- 10 Yet others of these compounds, MdPAM and MnBAM, have shown *in vivo* activity in the inhibition of colonic tissue injury and neutrophil accumulation into colonic tissue (Weiss et al., *The Journal of Biological Chemistry*, 271(42), 26149-26156 (1996)). In addition, these compounds have been reported to possess analgesic activity and to reduce inflammation and edema in the rat-paw carrageenan hyperalgesia model, see, e.g.,
- 15 U.S. Pat. No. 6,180,620. Furthermore, certain of these compounds may variously be considered anti-oxidants, or to reduce oxidative stress.


- [0004] Compounds of this class have also been shown to be safe and effective in the prevention and treatment of disease in human subjects. For example, GC4419 has been shown to reduce oral mucositis in head-and-neck cancer patients
- 20 undergoing chemoradiation therapy (Anderson, C., *Phase 1 Trial of Superoxide Dismutase (SOD) Mimetic GC4419 to Reduce Chemoradiotherapy (CRT)-Induced Mucositis (OM) in Patients (pts) with Mouth or Oropharyngeal Carcinoma (OCC)*, Oral Mucositis Research Workshop, MASCC/ISOO Annual Meeting on Supportive Care in Cancer, Copenhagen, Denmark (June 25, 2015)).

- 25            [0005] In addition, transition metal-containing pentaaza macrocyclic ring complexes corresponding to this class have shown efficacy in the treatment of various cancers. For example, certain compounds corresponding to this class have been

provided in combination with agents such as paclitaxel and gemcitabine to enhance cancer therapies, such as in the treatment of colorectal cancer and lung cancer (non-small cell lung cancer) (see, e.g., *U.S. Patent No. 9,998,893*). The GC4403 compound above has also been used for treatment in *in vivo* models of Meth A spindle cell spindle cell ascites tumor and renal carcinoma (Samlowski et al., *Madame Curie Bioscience Database (Internet)*, 230-249 (2006)), and has also been used for treatment in *in vivo* models of spindle-cell squamous carcinoma (Samlowski et al., *Nature Medicine*, 9(6), 750-755 (2003)).

[0006] However, a need remains for enhanced methods for cancer treatment that provide improved efficacy in the killing of cancer cells, while also providing good selectivity in the killing of cancer cells as compared to normal cells. There also remains a need for enhanced methods of treatment to supplement cancer treatments such as radiation therapy and chemotherapy, to improve outcomes for patients receiving these treatments.

[0007] Among the various aspects of the present disclosure, therefore, is method of treating a cancer in a mammalian subject afflicted therewith. The method includes administering to the subject at least one active agent including one or more of a thioredoxin reductase inhibitor and a glutathione depleting agent, and administering to the subject a pentaaza macrocyclic ring complex corresponding to formula (I) below:



wherein

M is  $\text{Mn}^{2+}$  or  $\text{Mn}^{3+}$ ;

$R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$ , and  $R_{10}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and  $R_1$  and  $R_{10}$  attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;


n is an integer from 0 to 3; and

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[0008] Among the various aspects of the present disclosure, therefore, is a method of treating a cancer in a mammalian subject afflicted therewith, the method including administering to the subject at least one active agent including one or more of

a thioredoxin reductase inhibitor and a glutathione-depleting agent, administering to the subject a pentaaza macrocyclic ring complex, and further administering to the subject a cancer therapy that is at least one of radiation therapy and chemotherapy. The pentaaza macrocyclic ring complex can correspond to the formula (I) below:

5



wherein

M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

10  $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9,$  and  $R_{10}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}, -NR_{11}R_{12}, -COR_{11}, -CO_2R_{11}, -CONR_{11}R_{12}, -SR_{11}, -SOR_{11}, -SO_2R_{11}, -SO_2NR_{11}R_{12}, -N(OR_{11})(R_{12}), -P(O)(OR_{11})(OR_{12}), -P(O)(OR_{11})(R_{12}),$  and  $-OP(O)(OR_{11})(OR_{12}),$  wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

15

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

20 V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted

or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R<sub>1</sub> and R<sub>10</sub> attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

Z is a counterion;

n is an integer from 0 to 3; and

the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[0009] Other objects and features will be in part apparent and in part pointed out hereinafter.

#### Brief Description of the Drawings

[0010] FIG. 1 shows the normalized mean fluorescence intensity (MFI) of 5-(and 6) carboxy-2',7' dichlorodihydrofluorescein diacetate (DCFH2) in cancer cells MB231 and H1299, as compared to normal HMEC or HBEpC cells, using the DCFH2 fluorescent probe as a non-specific oxidation sensitive probe.

[0011] FIG. 2 shows the normalized mean fluorescence intensity (MFI) for PeroxyOrange-1 (PO-1) oxidation as a measure of steady-state H<sub>2</sub>O<sub>2</sub> in H292 cells treated with GC4419 alone, as well as for H292 cells treated with combinations of buthionine sulfoximine (BSO) and 3-aminotriazole (AT), and combinations of BSO and AT with GC4419, as compared to a control.

[0012] FIG. 3 shows normalized levels of peroxide flux, measured as the 3-aminotriazole inactivation of catalase, in head and neck cancer cells (Cal27), lung cancer cells (H1299) and breast cancer cells (MB231) treated with GC4419, compared to a control untreated with GC4419.

[0013] FIG. 4 shows the surviving fraction of MB231 and H1299 cancer cells as compared to HMEC and HBEpC normal epithelial cells, following treatment with (i) GC4419, (ii) GC4419 and BSO, (iii) GC4419 and auranofin (Au), (iv) BSO and Au, and (v) GC4419, BSO and Au, as compared to a control.

5 [0014] FIG. 5 shows the surviving fraction of MB231 and H1299 cancer cells, following treatment with (i) GC4419, (ii) BSO and Au, and (iii) GC4419, BSO and Au, both in the presence and in the absence of the thiol antioxidant N-acetylcysteine (NAC).

[0015] FIG. 6 shows the normalized surviving fraction of H1299 cancer cells, following treatment with (i) Au, (ii) ionizing radiation (IR), and (iii) Au in combination with  
10 IR, as compared to a control.

[0016] FIG. 7 shows total glutathione levels as measured using a DTNB recycling spectrophotometric assay, for breast cancer cells (MB231) and lung cancer cells (H292) treated with BSO.

[0017] FIG. 8 shows tumor volumes over time post treatment of MB231 breast  
15 cancer xenografts in mice, where treatment included exposure to (i) GC4419 alone, (ii) combination of the Au and BSO, and (iii) the combination of GC4419, Au and BSO.

[0018] FIG. 9 shows average tumor volumes, tumor volumes for individual animals, and the percentage of tumors with volumes less than 1000 mm<sup>3</sup> for a non-small cell lung cancer tumor xenograft model, over time post treatment (post-final IR  
20 exposure), with treatment with GC4419, treatment with IR (18 Gy X1 dose or 2 GY X 16 doses), and treatment with a combination of IR and GC4419, as compared to control.

[0019] FIG. 10 shows average tumor volumes over time post treatment, in animal models of squamous cell carcinoma of the head and neck (HNSCC), for  
25 treatment with (i) GC4419, (ii) IR (12 Gy x 1), and (iii) IR and GC4419, in comparison to control.

[0020] FIGS. 11A-11B are schematic diagrams showing mechanisms by which intracellular levels of H<sub>2</sub>O<sub>2</sub> can be increased with pentaaza macrocyclic ring complexes, thioredoxin reductase inhibitors and/or glutathione depleting agents.

[0021] FIG. 12 shows percentage growth of leukemia cell lines incubated at  
30 different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to

provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0022] FIG. 13 shows percentage growth of non-small cell lung cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0023] FIG. 14 shows percentage growth of colon cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0024] FIG. 15 shows percentage growth of CNS cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0025] FIG. 16 shows percentage growth of melanoma cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0026] FIG. 17 shows percentage growth of ovarian cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0027] FIG. 18 shows percentage growth of renal cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0028] FIG. 19 shows percentage growth of prostate cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0029] FIG. 20 shows percentage growth of breast cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40401 (GC4401), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

5 [0030] FIG. 21 shows percentage growth of leukemia cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

10 [0031] FIG. 22 shows percentage growth of non-small cell lung cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

15 [0032] FIG. 23 shows percentage growth of colon cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

20 [0033] FIG. 24 shows percentage growth of CNS cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0034] FIG. 25 shows percentage growth of melanoma cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

25 [0035] FIG. 26 shows percentage growth of ovarian cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

30 [0036] FIG. 27 shows percentage growth of renal cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403),

to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0037] FIG. 28 shows percentage growth of prostate cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0038] FIG. 29 shows percentage growth of breast cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40403 (GC4403), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0039] FIG. 30 shows percentage growth of leukemia cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0040] FIG. 31 shows percentage growth of non-small cell lung cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0041] FIG. 32 shows percentage growth of colon cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0042] FIG. 33 shows percentage growth of CNS cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0043] FIG. 34 shows percentage growth of melanoma cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

[0044] FIG. 35 shows percentage growth of ovarian cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

5 [0045] FIG. 36 shows percentage growth of renal cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

10 [0046] FIG. 37 shows percentage growth of prostate cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

15 [0047] FIG. 38 shows percentage growth of breast cancer cell lines incubated at different concentrations of the pentaaza macrocyclic ring complex M40432 (GC4432), to provide *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex in the cancerous cell lines.

#### Abbreviations and Definitions

20 [0048] The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art.

25 [0049] "AcyI" means a -COR moiety where R is alkyl, haloalkyl, optionally substituted aryl, or optionally substituted heteroaryl as defined herein, e.g., acetyl, trifluoroacetyl, benzoyl, and the like.

[0050] "Acyloxy" means a -OCOR moiety where R is alkyl, haloalkyl, optionally substituted aryl, or optionally substituted heteroaryl as defined herein, e.g., acetyl, trifluoroacetyl, benzoyl, and the like.

30 [0051] "Alkoxy" means a -OR moiety where R is alkyl as defined above, e.g., methoxy, ethoxy, propoxy, or 2-propoxy, n-, iso-, or tert-butoxy, and the like.

[0052] "Alkyl" means a linear saturated monovalent hydrocarbon moiety such as of one to six carbon atoms, or a branched saturated monovalent hydrocarbon moiety, such as of three to six carbon atoms, e.g., C<sub>1</sub>-C<sub>6</sub> alkyl groups such as methyl, ethyl, propyl, 2-propyl, butyl (including all isomeric forms), pentyl (including all isomeric forms), and the like.

[0053] Moreover, unless otherwise indicated, the term "alkyl" as used herein is intended to include both "unsubstituted alkyls" and "substituted alkyls," the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Indeed, unless otherwise indicated, all groups recited herein are intended to include both substituted and unsubstituted options.

[0054] The term "C<sub>x-y</sub>" when used in conjunction with a chemical moiety, such as alkyl and aralkyl, is meant to include groups that contain from x to y carbons in the chain. For example, the term C<sub>x-y</sub> alkyl refers to substituted or unsubstituted saturated hydrocarbon groups, including straight chain alkyl and branched chain alkyl groups that contain from x to y carbon atoms in the chain.

[0055] "Alkylene" means a linear saturated divalent hydrocarbon moiety, such as of one to six carbon atoms, or a branched saturated divalent hydrocarbon moiety, such as of three to six carbon atoms, unless otherwise stated, e.g., methylene, ethylene, propylene, 1-methylpropylene, 2-methylpropylene, butylene, pentylene, and the like.

[0056] "Alkenyl" a linear unsaturated monovalent hydrocarbon moiety, such as of two to six carbon atoms, or a branched saturated monovalent hydrocarbon moiety, such as of three to six carbon atoms, e.g., ethenyl (vinyl), propenyl, 2-propenyl, butenyl (including all isomeric forms), pentenyl (including all isomeric forms), and the like.

[0057] "Alkaryl" means a monovalent moiety derived from an aryl moiety by replacing one or more hydrogen atoms with an alkyl group.

[0058] "Alkenylcycloalkenyl" means a monovalent moiety derived from an alkenyl moiety by replacing one or more hydrogen atoms with a cycloalkenyl group.

[0059] "Alkenylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with an alkenyl group.

[0060] "Alkylcycloalkenyl" means a monovalent moiety derived from a cycloalkenyl moiety by replacing one or more hydrogen atoms with an alkyl group.

[0061] "Alkylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with an alkyl group.

5 [0062] "Alkynyl" means a linear unsaturated monovalent hydrocarbon moiety, such of two to six carbon atoms, or a branched saturated monovalent hydrocarbon moiety, such as of three to six carbon atoms, e.g., ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like.

10 [0063] "Alkoxy" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a hydroxy group.

[0064] "Amino" means a  $-NR^aR^b$  group where  $R^a$  and  $R^b$  are independently hydrogen, alkyl or aryl.

15 [0065] "Aralkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an aryl group. The term "arylalkyl" may also be used equivalently herein to refer to "aralkyl."

[0066] "Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon moiety of 6 to 10 ring atoms e.g., phenyl or naphthyl.

[0067] "Cycle" means a carbocyclic saturated monovalent hydrocarbon moiety of three to ten carbon atoms.

20 [0068] "Cycloalkyl" means a cyclic saturated monovalent hydrocarbon moiety of three to ten carbon atoms, e.g., cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, and the like.

25 [0069] "Cycloalkylalkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a cycloalkyl group, e.g., cyclopropylmethyl, cyclobutylmethyl, cyclopentylethyl, or cyclohexylethyl, and the like.

[0070] "Cycloalkylcycloalkyl" means a monovalent moiety derived from a cycloalkyl moiety by replacing one or more hydrogen atoms with a cycloalkyl group.

30 [0071] "Cycloalkenyl" means a cyclic monounsaturated monovalent hydrocarbon moiety of three to ten carbon atoms, e.g., cyclopropenyl, cyclobutenyl, cyclopentenyl, or cyclohexenyl, and the like.

[0072] "Cycloalkenylalkyl" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with a cycloalkenyl group, e.g., cyclopropenylmethyl, cyclobutenylmethyl, cyclopentenylethyl, or cyclohexenylethyl, and the like.

5 [0073] "Ether" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an alkoxy group.

[0074] "Halo" means fluoro, chloro, bromo, or iodo, preferably fluoro or chloro.

[0075] "Heterocycle" or "heterocyclyl" means a saturated or unsaturated monovalent monocyclic group of 4 to 8 ring atoms in which one or two ring atoms are  
10 heteroatom selected from N, O, or S(O)<sub>n</sub>, where n is an integer from 0 to 2, the remaining ring atoms being C. The heterocyclyl ring is optionally fused to a (one) aryl or heteroaryl ring as defined herein provided the aryl and heteroaryl rings are monocyclic. The heterocyclyl ring fused to monocyclic aryl or heteroaryl ring is also referred to in this Application as "bicyclic heterocyclyl" ring. Additionally, one or two ring carbon atoms in  
15 the heterocyclyl ring can optionally be replaced by a –CO– group. More specifically the term heterocyclyl includes, but is not limited to, pyrrolidino, piperidino, homopiperidino, 2-oxopyrrolidinyl, 2-oxopiperidinyl, morpholino, piperazino, tetrahydropyranyl, thiomorpholino, and the like. When the heterocyclyl ring is unsaturated it can contain one or two ring double bonds provided that the ring is not aromatic. When the  
20 heterocyclyl group is a saturated ring and is not fused to aryl or heteroaryl ring as stated above, it is also referred to herein as saturated monocyclic heterocyclyl.

[0076] "Heteroaryl" means a monovalent monocyclic or bicyclic aromatic moiety of 5 to 10 ring atoms where one or more, preferably one, two, or three, ring atoms are heteroatom selected from N, O, or S, the remaining ring atoms being carbon.  
25 Representative examples include, but are not limited to, pyrrolyl, pyrazolyl, thienyl, thiazolyl, imidazolyl, furanyl, indolyl, isoindolyl, oxazolyl, isoxazolyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, quinoliny, isoquinoliny, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl, tetrazolyl, and the like.

[0077] "Nitro" means –NO<sub>2</sub>.

30 [0078] "Organosulfur" means a monovalent moiety a –SR group where R is hydrogen, alkyl or aryl.

[0079] "Substituted alkyl," "substituted cycle," "substituted phenyl," "substituted aryl," "substituted heterocycle," and "substituted nitrogen heterocycles" means an alkyl, cycle, aryl, phenyl, heterocycle or nitrogen-containing heterocycle, respectively, optionally substituted with one, two, or three substituents, such as those independently selected from alkyl, alkoxy, alkoxyalkyl, halo, hydroxy, hydroxyalkyl, or organosulfur.

[0080] "Thioether" means a monovalent moiety derived from an alkyl moiety by replacing one or more hydrogen atoms with an –SR group wherein R is alkyl.

[0081] As used herein, (i) the compound referred to herein and in the Figures as compound 401, 4401 or GC4401 is a reference to the same compound, (ii) the compound referred to herein and in the Figures as compound 403, 4403 or GC4403 is a reference to the same compound, (iii) the compound referred to herein and in the Figures as compound 419, 4419 or GC4419 is a reference to the same compound, and (iv) the compound referred to herein and in the Figures as compound 444, 4444 or GC4444 is a reference to the same compound.

## DETAILED DESCRIPTION

[0082] Aspects of the present disclosure are directed to the treatment of cancer by administration of a pentaaza macrocyclic ring complex according to Formula (I) with at least one active agent that is at least one of a thioredoxin reductase inhibitor and a glutathione depleting agent. Embodiments of the treatment can provide for the enhanced killing of cancerous cells in patients in need thereof, as well as improved selectivity in the killing of cancer cells versus normal cells. The combinations of these compounds may also be administered as a supplement to another cancer therapy, such as a radiation therapy and/or chemotherapy, to improve the efficacy thereof.

[0083] Without being limited to any particular theory, it is believed that the combination of the pentaaza macrocyclic ring complex of Formula (I) with the active agent that is at least one of a thioredoxin reductase inhibitor and a glutathione depleting agent may result in an increase in intracellular H<sub>2</sub>O<sub>2</sub> levels in cancer cells, thereby leading to increased cytotoxicity. In particular, differences in mitochondrial oxidative metabolism between cancer and normal cells lead to increased levels of reactive oxygen species (ROS) such as O<sub>2</sub><sup>•-</sup> (superoxide anion) and H<sub>2</sub>O<sub>2</sub>. Glutathione and thioredoxin pathways are thus upregulated in cancer cells to compensate for this

difference, by providing reducing agents to remove the reactive oxygen species. (Zhou et al., *Adv. Cancer Res.* 122, 1-67 (2014); Aykin-Burns et al., *Biochem J.*, 418 (1), 29-37 (2009); Aykin-Burns et al., *J. Biol. Chem.*, 280(6), 4254-4263 (2005); Spitz et al., *Ann. NY Acad. Sci.*, 899, 349-362 (2000); Fath et al. *Clin Cancer Res.*, 17(19), 6206-6217 (2011); Hadzic et al., *Free Radic. Biol. Med.*, 48(8), 1024-1033 (2010); Scarbrough et al., *Free Radic Biol Med.*, 52 (2), 436-443 (2012)).

[0084] Accordingly, embodiments of the invention are directed to exploiting this difference in oxygen metabolism between cancer and normal cells to provide enhanced cancer treatment, with a combination of compounds that unexpectedly exhibits synergistic effects. In particular, embodiments of the invention provide for the administration of an active agent that reduces that ability of cells to regulate  $H_2O_2$ , which may selectively impact cancer cells that exhibit increased levels of reactive oxygen species. Synergistic effects in cancer cell cytotoxicity are then unexpectedly achieved via administration of a compound that removes one reactive oxygen species, superoxide anion, and converts it into another,  $H_2O_2$ , with two superoxide equivalents being removed for each  $H_2O_2$  equivalent created. The combined administration of the compounds both attacks the mechanism by which  $H_2O_2$  is eliminated from cancer cells, and also promotes the formation of new  $H_2O_2$ , resulting in a synergistic increase in  $H_2O_2$  and cytotoxicity for the cancer cells.


[0085] In one embodiment, to inhibit one or more of the glutathione and/or thioredoxin-dependent metabolism of  $H_2O_2$ , at least one of a thioredoxin reductase inhibitor and a glutathione-depleting agent can be provided. Furthermore, the compound to promote  $H_2O_2$  formation is a pentaaza macrocyclic ring complex according to Formula (I), which may increase the production of  $H_2O_2$  from superoxide in cells. (Buettner et al., *Free Radic Biol. Med.*, 41(8), 1338-1350 (2006). Embodiments of a mechanism by which intracellular levels or fluxes of  $H_2O_2$  may be increased by the combined administration of the compounds are shown in FIGS. 11A-11B, where GC4419 is used as the pentaaza macrocyclic ring complex, and auranofin is provided as a thioredoxin reductase inhibitor. The mechanism as shown thus results in an overall increase in the  $H_2O_2$  levels in cancer cells by promotion of  $H_2O_2$  formation from superoxide by the GC4419, and simultaneous inhibition of thioredoxin reductase by auranofin to reduce metabolism of the  $H_2O_2$ . Similarly, a glutathione depleting agent can be provided to inhibit glutathione-dependent  $H_2O_2$  metabolism pathways. An

embodiment of a mechanism by which glutathione depletion can contribute to increased intracellular levels of  $H_2O_2$ , with BSO as a glutathione depleting agent provided as a part of a combination therapy, is also shown in FIGS. 11A-11B.

[0086] Accordingly, by providing a combination of the pentaaza macrocyclic ring complex according to Formula (1) and the active agent that is at least one of a thioredoxin reductase inhibitor and a glutathione-depleting agent, it has been unexpectedly discovered that synergistic effects in the killing of cancer cells can be provided. The combination can thus be administered for the treatment of cancer, as well as to supplement conventional cancer treatment therapies, with improved treatment efficacy.

### Transition Metal Pentaaza Macrocyclic Ring Complex

[0087] In one embodiment, the pentaaza macrocyclic ring complex corresponds to the complex of Formula (I):



wherein

[0088] M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

[0089]  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

[0090] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

5 [0091] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[0092] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused  
10 heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R<sub>1</sub> and R<sub>10</sub> attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

[0093] X and Y represent suitable ligands which are derived from any  
15 monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[0094] Z is a counterion;

[0095] n is an integer from 0 to 3; and

[0096] the dashed lines represent coordinating bonds between the nitrogen  
20 atoms of the macrocycle and the transition metal, manganese.

[0097] As noted above in connection with the pentaaza macrocyclic ring complex of Formula (I), M is Mn<sup>2+</sup> or Mn<sup>3+</sup>. In one particular embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I), M is Mn<sup>2+</sup>. In another particular embodiment in which the pentaaza macrocyclic ring complex corresponds to  
25 Formula (I), M is Mn<sup>3+</sup>.

[0098] In the embodiments in which one or more of R<sub>1</sub>, R<sub>2</sub>, R'<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R'<sub>5</sub>, R<sub>6</sub>, R'<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R'<sub>9</sub>, and R<sub>10</sub> are hydrocarbyl, for example, suitable hydrocarbyl moieties include, but are not limited to alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, and aralkyl. In one embodiment,  
30 R<sub>1</sub>, R<sub>2</sub>, R'<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R'<sub>5</sub>, R<sub>6</sub>, R'<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R'<sub>9</sub>, and R<sub>10</sub> are independently hydrogen,

hydrocarbyl, substituted hydrocarbyl, or heterocyclyl. More preferably in this embodiment,  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen or lower alkyl (e.g.,  $C_1$ - $C_6$  alkyl, more typically  $C_1$ - $C_4$  alkyl).

Thus, for example,  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  may be independently hydrogen, methyl, ethyl, propyl, or butyl (straight, branched, or cyclic). In one preferred embodiment,  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen or methyl.

[0099] In one preferred embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I),  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are each hydrogen and one of  $R_6$  and  $R'_6$  is hydrogen and the other of  $R_6$  and  $R'_6$  is methyl. In this embodiment, for example,  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  may each be hydrogen while  $R'_6$  is methyl. Alternatively, for example,  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  may each be hydrogen while  $R_6$  is methyl. In another preferred embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I),  $R_1$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ , and  $R_{10}$  are each hydrogen, one of  $R_2$  and  $R'_2$  is hydrogen and the other of  $R_2$  and  $R'_2$  is methyl, and one of  $R_9$  and  $R'_9$  is hydrogen and the other of  $R_9$  and  $R'_9$  is methyl. In this embodiment, for example,  $R_1$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_7$ ,  $R_8$ ,  $R_9$ , and  $R_{10}$  may each be hydrogen while  $R_2$  and  $R'_9$  are methyl. Alternatively, for example,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_7$ ,  $R_8$ ,  $R'_9$ , and  $R_{10}$  may each be hydrogen while  $R'_2$  and  $R_9$  are methyl. In another embodiment in which the pentaaza macrocyclic ring complex corresponds to Formula (I),  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are each hydrogen.

[00100] In certain embodiments the U and V moieties are independently substituted or unsubstituted fused cycloalkyl moieties having 3 to 20 ring carbon atoms, more preferably 4 to 10 ring carbon atoms. In a particular embodiment, the U and V moieties are each trans-cyclohexanyl fused rings.

[00101] In certain embodiments the W moiety is a substituted or unsubstituted fused heteroaromatic moiety. In a particular embodiment, the W moiety is a substituted or unsubstituted fused pyridino moiety. Where W is a substituted fused pyridino moiety, for example, the W moiety is typically substituted with a hydrocarbyl or substituted hydrocarbyl moiety (e.g., alkyl, substituted alkyl) at the ring carbon atom positioned para

to the nitrogen atom of the heterocycle. In a one preferred embodiment, the W moiety is an unsubstituted fused pyridino moiety.

[00102] As noted above, X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the  
5 corresponding anion thereof (for example benzoic acid or benzoate anion, phenol or phenoxide anion, alcohol or alkoxide anion). For example, X and Y may be selected from the group consisting of halo, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxy, hydroperoxy, alkylperoxy, arylperoxy, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl  
10 hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol  
15 thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl  
20 phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate, thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate,  
25 aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions  
30 of ion exchange resins, or the corresponding anions thereof, among other possibilities. In one embodiment, X and Y if present, are independently selected from the group consisting of halo, nitrate, and bicarbonate ligands. For example, in this embodiment, X and Y, if present, are halo ligands, such as chloro ligands.

[00103] Furthermore, in one embodiment X and Y correspond to  $-O-C(O)-X_1$ , where each  $X_1$  is  $-C(X_2)(X_3)(X_4)$ , and each  $X_1$  is independently substituted or unsubstituted phenyl or  $-C(-X_2)(-X_3)(-X_4)$ ;

[00104] each  $X_2$  is independently substituted or unsubstituted phenyl, methyl,  
5 ethyl or propyl;

[00105] each  $X_3$  is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino,  $-X_5C(=O)R_{13}$  where  $X_5$  is NH or O, and  $R_{13}$  is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or  $-OR_{14}$ , where  $R_{14}$  is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with  $X_4$  is  $(=O)$ ; and

10 [00106] each  $X_4$  is independently hydrogen or together with  $X_3$  is  $(=O)$ .

[00107] In yet another embodiment, X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof; or X and Y are independently attached to one or more of  $R_1$ ,  
15  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$ .

[00108] In the pentaaza macrocyclic ring complex corresponding to Formula (I), Z is a counterion (e.g., a charge-neutralizing anion), wherein n is an integer from 0 to 3. In general, Z may correspond to counterions of the moieties recited above in connection for X and Y.

20 [00109] In combination, among certain preferred embodiments are pentaaza macrocyclic ring complexes corresponding to Formula (I) wherein

[00110] M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

[00111]  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen or lower alkyl;

25 [00112] U and V are each trans-cyclohexanyl fused rings;

[00113] W is a substituted or unsubstituted fused pyridino moiety;


[00114] X and Y are ligands; and

[00115] Z, if present, is a charge-neutralizing anion.

[00116] More preferably in these embodiments, M is  $Mn^{2+}$ ;  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  
30  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen or methyl; U and V

are each trans-cyclohexanyl fused rings; W is an unsubstituted fused pyridino moiety; and X and Y are independently halo ligands (e.g., fluoro, chloro, bromo, iodo). Z, if present, may be a halide anion (e.g., fluoride, chloride, bromide, iodide).

[00117] In yet another embodiment, the pentaaza macrocyclic ring complex is represented by formula (II) below:




(II)

5 [00118] wherein


[00119] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

[00120]  $R_A$ ,  $R_B$ ,  $R_C$ , and  $R_D$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl.

15 [00121] Furthermore, in one embodiment, the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):



(III)



(IV)







[00122] wherein




X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and


[00123]  $R_A$ ,  $R_B$ ,  $R_C$ , and  $R_D$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl.


[00124] Certain particularly preferred pentaaza macrocyclic ring complexes for use in the methods and compositions described herein include those corresponding to Formulae (V)-(XVI):








5



10




wherein X and Y in each of Formulae are independently ligands. For example, according to one embodiment, the pentaaza macrocyclic ring complex for use in the methods and compositions described herein include those corresponding to the

Formulae above, with X and Y in each of these formulae being halo, such as chloro. Alternatively, X and Y may be ligands other than chloro.

[00126] In one embodiment, X and Y in any of the formulas herein are independently selected from the group consisting of fluoro, chloro, bromo and iodo anions. In yet another embodiment, X and Y in any of the formulas herein are independently selected from the group consisting of alkyl carboxylates, aryl carboxylates and arylalkyl carboxylates. In yet another embodiment, X and Y in any of the formulas herein are independently amino acids.

[00127] In one embodiment, the pentaaza macrocyclic ring complex has the following Formula (IA):



wherein

[00128] M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

[00129]  $R_{1A}$ ,  $R_{1B}$ ,  $R_2$ ,  $R_3$ ,  $R_{4A}$ ,  $R_{4B}$ ,  $R_5$ ,  $R_6$ ,  $R_{7A}$ ,  $R_{7B}$ ,  $R_8$ ,  $R_9$ ,  $R_{10A}$ , and  $R_{10B}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety independently selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-C(=O)NR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(=O)(OR_{11})(OR_{12})$ ,  $-P(=O)(OR_{11})(R_{12})$ , and  $-OP(=O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

[00130] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

5 [00131] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [00132] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R<sub>5</sub> and R<sub>6</sub> attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent; wherein

15 [00133] each X<sub>1</sub> is independently substituted or unsubstituted phenyl or -C(-X<sub>2</sub>)(-X<sub>3</sub>)(-X<sub>4</sub>);

[00134] each X<sub>2</sub> is independently substituted or unsubstituted phenyl or alkyl;

20 [00135] each X<sub>3</sub> is independently hydrogen, hydroxyl, alkyl, amino, -X<sub>5</sub>C(=O)R<sub>13</sub> where X<sub>5</sub> is NH or O, and R<sub>13</sub> is C<sub>1</sub>-C<sub>18</sub> alkyl, substituted or unsubstituted aryl or C<sub>1</sub>-C<sub>18</sub> aralkyl, or -OR<sub>14</sub>, where R<sub>14</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl, substituted or unsubstituted aryl or C<sub>1</sub>-C<sub>18</sub> aralkyl, or together with X<sub>4</sub> is (=O);

[00136] each X<sub>4</sub> is independently hydrogen or together with X<sub>3</sub> is (=O); and

[00137] the bonds between the transition metal M and the macrocyclic nitrogen atoms and the bonds between the transition metal M and the oxygen atoms of the axial ligands -OC(=O)X<sub>1</sub> are coordinate covalent bonds.

25 [00138] In one embodiment, within Formula (IA), and groups contained therein, in one group of compounds X<sub>1</sub> is -C(-X<sub>2</sub>)(-X<sub>3</sub>)(-X<sub>4</sub>) and each X<sub>2</sub>, X<sub>3</sub>, and X<sub>4</sub>, in combination, corresponds to any of the combinations identified in the following table:







| Combination | X <sub>2</sub>  | X <sub>3</sub>                                           | X <sub>4</sub> |
|-------------|-----------------|----------------------------------------------------------|----------------|
| 1           | Ph              | H                                                        | H              |
| 2           | Ph              | OH                                                       | H              |
| 3           | Ph              | NH <sub>2</sub>                                          | H              |
| 4           | Ph              | =O<br>(X <sub>3</sub> and X <sub>4</sub> in combination) |                |
| 5           | Ph              | CH <sub>3</sub>                                          | H              |
| 6           | CH <sub>3</sub> | H                                                        | H              |
| 7           | CH <sub>3</sub> | OH                                                       | H              |
| 8           | CH <sub>3</sub> | NH <sub>2</sub>                                          | H              |
| 9           | CH <sub>3</sub> | =O<br>(X <sub>3</sub> and X <sub>4</sub> in combination) |                |

[00139] Furthermore, within embodiment (IA), and groups contained therein, in one group of compounds X<sub>1</sub> is C(-X<sub>2</sub>)(-X<sub>3</sub>)(-X<sub>4</sub>), and X<sub>3</sub> is -X<sub>5</sub>C(=O)R<sub>13</sub>, such that the combinations of X<sub>2</sub>, X<sub>3</sub> and X<sub>4</sub> include any of the combinations identified in the following table:

| Combination | X <sub>2</sub>  | X <sub>3</sub>         | X <sub>4</sub> |
|-------------|-----------------|------------------------|----------------|
| 1           | Ph              | NHC(=O)R <sub>13</sub> | H              |
| 2           | Ph              | OC(=O)R <sub>13</sub>  | H              |
| 3           | CH <sub>3</sub> | NHC(=O)R <sub>13</sub> | H              |
| 4           | CH <sub>3</sub> | OC(=O)R <sub>13</sub>  | H              |

[00140] where R<sub>13</sub> is C<sub>1</sub>-C<sub>18</sub> alkyl, substituted or unsubstituted aryl or C<sub>1</sub>-C<sub>18</sub> aralkyl, or -OR<sub>14</sub>, where R<sub>14</sub> is C<sub>1</sub>-C<sub>18</sub> alkyl, substituted or unsubstituted aryl or C<sub>1</sub>-C<sub>18</sub> aralkyl.

[00141] In one embodiment, the pentaaza macrocyclic ring complex corresponding to Formula (IA) is one of the complexes Formula (IE), such as (IE<sub>R1</sub>), (IE<sub>S1</sub>), (IE<sub>R2</sub>), (IE<sub>S2</sub>), (IE<sub>R3</sub>), or (IE<sub>S3</sub>):



5

wherein

M is Mn<sup>+2</sup> or Mn<sup>+3</sup>;

each  $X_1$  is independently substituted or unsubstituted phenyl or  $-C(X_2)(X_3)(X_4)$ ;

each  $X_2$  is independently substituted or unsubstituted phenyl, methyl, ethyl, or propyl;

each  $X_3$  is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino, or  
5 together with  $X_4$  is  $=O$ ;


each  $X_4$  is independently hydrogen or together with  $X_3$  is  $=O$ ; and

the bonds between the manganese and the macrocyclic nitrogen atoms and the bonds between the manganese and the oxygen atoms of the axial ligands  $-OC(O)X_1$  are coordinate covalent bonds.


10 [00142] In one embodiment, each  $X_1$  is  $-C(X_2)(X_3)(X_4)$  and each  $-C(X_2)(X_3)(X_4)$  corresponds to any of combinations 1 to 9 appearing in the table for Formula (IA) above.

[00143] In yet another embodiment, the X and Y in pentaaza macrocyclic ring complex of Formula (I) correspond to the ligands in Formulas (IA) or (IE). For example,  
15 X and Y in the complex of Formula (I) may correspond to  $-O-C(O)-X_1$ , where  $X_1$  is as defined for the complex of Formula (IA) and (IE) above.

[00144] In one embodiment, pentaaza macrocyclic ring complexes corresponding to Formula (I) (e.g., of Formula (I) or any of the subsets of Formula (I) corresponding to Formula (II)-(XIV), (IA and (IE) and others described herein), can  
20 comprise any of the following structures:




[00145]




[00146]

(4403)



4401


[00147]




[00148]

4444

[00149]




4702




[00150]

4711

[00151] In a particularly preferred embodiment, the pentaaza macrocyclic complex corresponds to Formula (XVII) or Formula (XVIII):




(XVII)




(XVIII)

[00152] For example, the pentaaza macrocyclic ring complex may correspond to at least one of the complexes below:





(4419)




(4403)

[00153] The chemical structures of 4403 (dichloro complex form of XVII described, for example, in Riley, D.P., Schall, O.F., 2007, *Advances in Inorganic Chemistry*, 59: 233-263) and of 4419 herein (such as the dichloro complex form of XVIII), are identical except that they possess mirror image chirality; that is, the enantiomeric structures are non-superimposable.

[00154] In yet another embodiment, the pentaaza macrocyclic ring complex may correspond to at least one of the complexes below, and/or an enantiomer thereof:



[00155] In yet another embodiment, the pentaaza macrocyclic ring complex corresponds to Formula (XIX):



wherein

5 M is a transition metal (e.g.,  $Mn^{2+}$ ,  $Mn^{3+}$ );

$R_{1A}$ ,  $R_{1B}$ ,  $R_{2A}$ ,  $R_{2B}$ ,  $R_{3A}$ ,  $R_{3B}$ ,  $R_{4A}$ ,  $R_{4B}$ ,  $R_{5A}$ ,  $R_{5B}$ ,  $R_{6A}$ ,  $R_{6B}$ ,  $R_{7A}$ ,  $R_{7B}$ ,  $R_{8A}$ ,  $R_{8B}$ ,  $R_{9A}$ ,  $R_{9B}$ ,  $R_{10A}$ , and  $R_{10B}$  are independently:

(i) hydrogen;

(ii) a moiety independently selected from the group consisting of  
 10 alkenyl, alkenylcycloalkenyl, alkenylcycloalkyl, alkyl, alkylcycloalkenyl, alkylcycloalkyl, alkynyl, aralkyl, aryl, cycloalkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylcycloalkyl, cycloalkenylalkyl, heterocyclyl, and aralkyl radicals and radicals attached to the  $\alpha$ -carbon of amino acids (i.e.,  $\alpha$ -amino acids); or

(iii) a moiety independently selected from the group consisting of  
 15  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ ,  $-OP(O)(OR_{11})(OR_{12})$ , and substituents attached to the  $\alpha$ -carbon of amino acids (i.e.,  $\alpha$ -amino acids), wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

(iv) a member of a substituted or unsubstituted, saturated, partially  
 20 saturated, or unsaturated cycle or heterocycle containing 3 to 20 carbon ring atoms comprising

(a)  $R_{1A}$  or  $R_{1B}$  and  $R_{2A}$  or  $R_{2B}$ ;  $R_{3A}$  or  $R_{3B}$  and  $R_{4A}$  or  $R_{4B}$ ;  $R_{5A}$  or  $R_{5B}$  and  $R_{6A}$  or  $R_{6B}$ ;  $R_{7A}$  or  $R_{7B}$  and  $R_{8A}$  or  $R_{8B}$ ;  $R_{9A}$  or  $R_{9B}$  and  $R_{10A}$  or  $R_{10B}$  together with the carbon atoms to which they are respectively attached;

(b)  $R_{10A}$  or  $R_{10B}$  and  $R_{1A}$  or  $R_{1B}$ ;  $R_{2A}$  or  $R_{2B}$  and  $R_{3A}$  or  $R_{3B}$ ;  $R_{4A}$  or  $R_{4B}$  and  $R_{5A}$  or  $R_{5B}$ ;  $R_{6A}$  or  $R_{6B}$  and  $R_{7A}$  or  $R_{7B}$ ; or  $R_{8A}$  or  $R_{8B}$  and  $R_{9A}$  or  $R_{9B}$  together with the carbon atoms to which they are respectively attached; or

- 5 (c)  $R_{1A}$  and  $R_{1B}$ ;  $R_{2A}$  and  $R_{2B}$ ;  $R_{3A}$  and  $R_{3B}$ ;  $R_{4A}$  and  $R_{4B}$ ;  $R_{5A}$  and  $R_{5B}$ ;  $R_{6A}$  and  $R_{6B}$ ;  $R_{7A}$  and  $R_{7B}$ ;  $R_{8A}$  and  $R_{8B}$ ;  $R_{9A}$  and  $R_{9B}$ ; or  $R_{10A}$  and  $R_{10B}$  together with the carbon atoms to which they are respectively attached; or  
(v) a combination of any of (i) through (iv) above;

each  $X_1$  is independently substituted or unsubstituted phenyl or

10  $-C(-X_2)(-X_3)(-X_4)$ ;

each  $X_2$  is independently substituted or unsubstituted phenyl or substituted or unsubstituted alkyl;

each  $X_3$  is independently hydrogen, hydroxyl, alkyl, amino, or together with  $X_4$  is =O;

15 each  $X_4$  is independently hydrogen or together with  $X_3$  is =O; and

the bonds between the transition metal, manganese, and the macrocyclic nitrogen atoms and the bonds between the transition metal, manganese, and the oxygen atoms of the axial ligands  $-OC(O)X_1$  are coordinate covalent bonds.

[00156] In one embodiment, the pentaaza macrocyclic ring complex  
20 corresponding to Formula (XIX) is one of the complexes Formulae (IE<sub>R1</sub>), (IE<sub>S1</sub>), (IE<sub>R2</sub>), (IE<sub>S2</sub>), (IE<sub>R3</sub>), or (IE<sub>S3</sub>) described above.

[00157] In one embodiment, each  $X_1$  is  $-C(X_2)(X_3)(X_4)$  and each  $-C(X_2)(X_3)(X_4)$  corresponds to any of combinations 1 to 7 appearing in the following table:

| Combination | $X_2$           | $X_3$                                   | $X_4$ |
|-------------|-----------------|-----------------------------------------|-------|
| 1           | Ph              | H                                       | H     |
| 2           | Ph              | OH                                      | H     |
| 3           | Ph              | NH <sub>2</sub>                         | H     |
| 4           | Ph              | =O<br>( $X_3$ and $X_4$ in combination) |       |
| 5           | Ph              | CH <sub>3</sub>                         | H     |
| 6           | CH <sub>3</sub> | H                                       | H     |
| 7           | CH <sub>3</sub> | OH                                      | H     |

[00158] In yet another embodiment, the X and Y in pentaaza macrocyclic ring complex of formula (1) correspond to the ligands in formula (XIX). For example, X and Y in the complex of formula (1) may correspond to -O-C(O)-X<sub>1</sub>, where X<sub>1</sub> is as defined for the complex of Formula (XIX) above.

[00159] In one embodiment, the enantiomeric purity of the pentaaza macrocyclic ring complex is greater than 95%, more preferably greater than 98%, more preferably greater than 99%, and most preferably greater than 99.5%. As used herein, the term "enantiomeric purity" refers to the amount of a compound having the depicted absolute stereochemistry, expressed as a percentage of the total amount of the depicted compound and its enantiomer. In one embodiment, the diastereomeric purity of the pentaaza macrocyclic ring complex is greater than 98%, more preferably greater than 99%, and most preferably greater than 99.5%. As used herein, the term "diastereomeric purity" refers to the amount of a compound having the depicted absolute stereochemistry, expressed as a percentage of the total amount of the depicted compound and its diastereomers. Methods for determining diastereomeric and enantiomeric purity are well-known in the art. Diastereomeric purity can be determined by any analytical method capable of quantitatively distinguishing between a compound and its diastereomers, such as high performance liquid chromatography (HPLC). Similarly, enantiomeric purity can be determined by any analytical method capable of quantitatively distinguishing between a compound and its enantiomer. Examples of suitable analytical methods for determining enantiomeric purity include, without limitation, optical rotation of plane-polarized light using a polarimeter, and HPLC using a chiral column packing material.

[00160] In one embodiment, a therapeutically effective amount of the pentaaza macrocyclic ring complex may be an amount sufficient to provide a peak plasma concentration of at least 0.1  $\mu$ M when administered to a patient. For example, in one embodiment, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration of at least 1  $\mu$ M when administered to a patient. In yet another embodiment, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration of at least 10  $\mu$ M when administered to a patient. Generally, the pentaaza macrocyclic ring

complex will not be administered in an amount that would provide a peak plasma concentration greater than 40  $\mu\text{M}$  when administered to a patient. For example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 0.1  $\mu\text{M}$  to 40  $\mu\text{M}$  in a patient.

5 As another example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 0.5  $\mu\text{M}$  to 20  $\mu\text{M}$  in a patient. As another example, the pentaaza macrocyclic ring complex may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 1  $\mu\text{M}$  to 10  $\mu\text{M}$  in a patient.

10 [00161] In yet another embodiment, a dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be at least 0.1 mg/kg, such as at least 0.2 mg/kg. For example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be at least 0.5 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is  
15 administered per kg body weight of the patient may be at least 1 mg/kg. In another example, the pentaaza macrocyclic ring complex that is administered per kg body weight may be at least 2 mg/kg, such as at least 3 mg/kg, and even at least about 15 mg/kg, such as at least 24 mg/kg and even at least 40 mg/kg. Generally, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the  
20 patient will not exceed 1000 mg/kg. For example the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 0.1 to 1000 mg/kg, such as from 0.2 mg/kg to 40 mg/kg, such as 0.2 mg/kg to 24 mg/kg, and even 0.2 mg/kg to 10 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight may be in a  
25 range of from 1 mg/kg to 1000 mg/kg, such as from 3 mg/kg to 1000 mg/kg, and even from 5 mg/kg to 1000 mg/kg, such as 10 mg/kg to 1000 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight may be in a range of from 2 mg/kg to 15 mg/kg. As yet another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight  
30 may be in a range of from 3 mg/kg to 10 mg/kg. As another example, the dose of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 0.5 to 5 mg/kg. As yet a further example, the dose

of the pentaaza macrocyclic ring complex that is administered per kg body weight of the patient may be in the range of from 1 to 5 mg/kg.

[00162] In one embodiment, the dosages and/or plasma concentrations discussed above may be particularly suitable for the pentaaza macrocyclic ring complex corresponding to GC4419, although they may also be suitable for other pentaaza macrocyclic ring complexes. In addition, one of ordinary skill in the art would recognize how to adjust the dosages and/or plasma concentrations based on factors such as the molecular weight and/or activity of the particular compound being used. For example, for a pentaaza macrocyclic ring complex having an activity twice that of GC4419, the dosage and/or plasma concentration may be halved, or for a pentaaza macrocyclic ring complex having a higher molecular weight than GC4419, a correspondingly higher dosage may be used.

[00163] The dosing schedule of the pentaaza macrocyclic ring complex can similarly be selected according to the intended treatment. For example, in one embodiment, a suitable dosing schedule can comprise dosing a patient at least once per week, such as at least 2, 3, 4, 5, 6 or 7 days per week (e.g., daily), during a course of treatment. As another example, in one embodiment, the dosing may be at least once a day (qd), or even at least twice a day (bid). In one embodiment, the course of treatment with the pentaaza macrocyclic ring complex may last at least as long as a course of treatment with one or more other agents, such as least one of the thioredoxin reductase inhibitor and glutathione-depleting agent, and may even exceed the duration during which the at least one other agent is provided. The course of therapy with the pentaaza macrocyclic ring complex may also start on the same date as treatment with the other agent, such as on a same date as treatment with at least one of the thioredoxin reductase inhibitor and glutathione-depleting agent, or may start sometime after initial dosing with at least one of the other agents.

### **Thioredoxin Reductase Inhibitor**

[00164] In one embodiment, the thioredoxin reductase inhibitor is a compound that inhibits thioredoxin reductase, the enzyme that catalyzes the reduction of thioredoxin. Thioredoxin can act as a reducing agent to reduce levels of reactive oxygen species, such as H<sub>2</sub>O<sub>2</sub>. Accordingly, the inhibition of thioredoxin reductase

maintains thioredoxin in its reduced state, thereby decreasing the ability of thioredoxin to remove reactive oxygen species such as  $H_2O_2$ . In one embodiment, a thioredoxin reductase inhibitor selected for combination with the pentaaza macrocyclic ring complex is a compound that exhibits the thioredoxin reductase inhibition effect while also being therapeutically acceptable to the patient receiving the compound. For example, the thioredoxin reductase inhibitor may be at least one of auranofin, auro-thio-glucose, chloro(triethylphosphine)gold(I) (TEPAu), aurothiomalate, gold sodium thiomalate, sodium aurothiosulfate, gold acetate, 1,2,5-selenadiazole and derivatives thereof (e.g., as described in Liang et al, *Eur. J. Med. Chem.*, 84, 335-342 (2014)), metal complexes with 2-acetylpyridine-N(4)-ortho-chlorophenylthiosemicarbazone, such as palladium (II), platinum (II), bismuth (III), antimony (III) and gold (III) metal complexes (e.g., as described in Parilla et al, *Eur. J. Med. Chem.*, 84, 537-544 (2014)), and/or a pharmaceutically acceptable salt thereof. By way of further example, the thioredoxin reductase inhibitor may be at least one of auranofin (gold(+1)cation; 3,4,5-triacetyloxy-6-(acetyloxymethyl)oxane-2-thiolate; triethylphosphonium) and auro-thio-glucose (gold(1) (2S,3S,4R,5S)-3,4,5-trihydroxy-6-(hydroxymethyl)-oxane-2-thiolate).

[00165] In one embodiment, a therapeutically effective amount of the thioredoxin reductase inhibitor, such as auranofin, may be an amount sufficient to provide a peak plasma concentration of at least 0.5  $\mu M$  when administered to a patient. For example, in one embodiment, the thioredoxin reductase inhibitor, such as auranofin, may be administered in an amount sufficient to provide a peak plasma concentration of at least 0.8  $\mu M$  when administered to a patient. In yet another embodiment, the thioredoxin reductase inhibitor, such as auranofin, may be administered in an amount sufficient to provide a peak plasma concentration of at least 1  $\mu M$  when administered to a patient. Generally, the thioredoxin reductase inhibitor, such as auranofin, will not be administered in an amount that would provide a peak plasma concentration greater than 8  $\mu M$  when administered to a patient, and may even provide a peak plasma concentration of less than 5  $\mu M$ , and even less than 2  $\mu M$ . For example, the thioredoxin reductase inhibitor, such as auranofin, may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 0.5  $\mu M$  to 5  $\mu M$  in a patient, such as in a range of from 0.5  $\mu M$  to 2  $\mu M$ . In yet another embodiment, a thioredoxin reductase inhibitor, such as auranofin, may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 2.5  $\mu M$  to 3.5  $\mu M$ .

[00166] In yet another embodiment, a dose of the thioredoxin reductase inhibitor, such as auranofin, that is administered per kg body weight of the patient may be at least 0.05 mg/kg, such as at least 0.1 mg/kg, and even at least 0.2 mg/kg. For example, the dose of the thioredoxin reductase inhibitor, such as auranofin, that is administered per kg body weight of the patient may be at least 0.5 mg/kg. As another example, the dose of the thioredoxin reductase inhibitor, such as auranofin, that is administered per kg body weight of the patient may be at least 1 mg/kg. In another example, the thioredoxin reductase inhibitor, such as auranofin, that is administered per kg body weight may be at least 1.5 mg/kg, such as at least 3 mg/kg, and even at least about 5 mg/kg. Generally, the dose of the thioredoxin reductase inhibitor, such as auranofin that is administered per kg body weight of the patient will not exceed 300 mg/kg. For example the dose of the thioredoxin reductase inhibitor, such as auranofin, that is administered per kg body weight of the patient may be in the range of from 0.1 to 300 mg/kg, such as from 0.2 mg/kg to 40 mg/kg, and even from 1 mg/kg to 5 mg/kg.








[00167] Furthermore, it is noted that the dosage of the thioredoxin reductase inhibitor may vary according to the particular inhibitor selected, as well as according to the route of administration, etc. For example, for thioredoxin reductase inhibitors having high potency, a lower dosage may be used to provide the same treatment effect, as compared to thioredoxin reductase inhibitors having less potency. Accordingly, while exemplary dosage ranges for the particular thioredoxin reductase inhibitor auranofin are being provided herein, the exemplary ranges are not limiting, and auranofin and/or other thioredoxin reductase inhibitors may be provided in different dosages, administration routes, or administration frequencies, other than those specifically exemplified herein, according to the particular thioredoxin reductase inhibitor to be administered and the therapeutic effect to be achieved.


[00168] The dosing schedule of the thioredoxin reductase inhibitor, such as auranofin, can similarly be selected according to the intended treatment. For example, in one embodiment, a suitable dosing schedule can comprise dosing a patient at least once per week, such as at least 2, 3, 4, 5, 6 or 7 days per week (e.g., daily), during a course of treatment. As another example, in one embodiment, the dosing may be at least once a day (qd), or even at least twice a day (bid). In one embodiment, the course of treatment with the thioredoxin reductase inhibitor, such as auranofin, may last at least as long as a course of treatment with one or more other agents, such as least one of the

pentaaza macrocyclic ring complex and glutathione-depleting agent, and may even exceed the duration during which the at least one other agent is provided. The course of therapy with the thioredoxin reductase inhibitor, such as auranofin, may also start on the same date as treatment with the other agent, such as on a same date as treatment with at least one of the pentaaza macrocyclic ring complex and glutathione-depleting agent, or may start sometime after initial dosing with at least one of the other agents.


### **Glutathione Depleting Agent**

[00169] In one embodiment, the glutathione depleting agent is an agent that decreases levels of glutathione in the cancerous cells. The glutathione depleting agent may be a compound that acts to deplete glutathione by any of a number of different mechanisms. For example, in one embodiment, the glutathione depleting agent is a glutathione synthesis inhibitor, such as buthionine sulfoximine. In another embodiment, the glutathione depleting agent is an inhibitor of  $x_c^-$  cysteine/glutamate antiporter, such as sulfasalazine. In yet another embodiment, the glutathione depleting agent is a glutathione reductase inhibitor, such as 2-acetylamino-3-[4-(2-acetylamino-2-carboxyethylsulfanyltiocarbonylamino)phenylthiocarbamoylsulfanyl] propionic acid (2-AAPA). In one embodiment, the glutathione depleting agent can comprise at least one of buthionine sulfoximine, sulfasalazine, piperlongumine, N-ethylmaleimide, N-pyrenylmaleimide, 2-AAPA, erastin, sorafenib, 1S,3R-RSL3, DPI19, DPI18, DPI17, DPI13, DPI12, DPI10 (ML210), DPI7(ML162) (Cao et al., *Cell Mol. Life Sci.*, (2016)), and altretamine, and/or pharmaceutically acceptable salts thereof. Structures of some of these suitable glutathione depleting agents are as follows:






DPI13 = N-[2-(allyloxy)benzyl]-N-1,3-benzodioxol-5-yl-2-chloroacetamide




DPI17 = 6-(chloromethyl)-N-(2-naphthyl)-1,3,5-triazine-2,4-diamine

5



DPI18 = 6-(chloromethyl)-N-(4-ethoxyphenyl)-1,3,5-triazine-2,4-diamine

10



DPI19 = 2-chloro-N-heptyl-N-m-tolyl-acetamide

15

[00170] In one embodiment, the glutathione depleting agent comprises at least one of buthionine sulfoximine (BSO) and sulfasalazine, and/or a pharmaceutically acceptable salt thereof.

20

[00171] In one embodiment, a therapeutically effective amount of the glutathione-depleting agent, such as sulfasalazine, may be an amount sufficient to provide a peak plasma concentration of at least 200  $\mu\text{M}$  when administered to a patient. For example, in one embodiment, the glutathione-depleting agent, such as sulfasalazine, may be administered in an amount sufficient to provide a peak plasma concentration of at least 250  $\mu\text{M}$  when administered to a patient. In yet another embodiment, the glutathione-depleting agent, sulfasalazine, may be administered in an

amount sufficient to provide a peak plasma concentration of at least 300  $\mu\text{M}$  when administered to a patient. Generally, the glutathione-depleting agent, such as sulfasalazine, will not be administered in an amount that would provide a peak plasma concentration greater than 600  $\mu\text{M}$  when administered to a patient. For example, the glutathione-depleting agent, such as sulfasalazine, may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 200  $\mu\text{M}$  to 600  $\mu\text{M}$  in a patient.

[00172] In yet another embodiment, a dose of the glutathione-depleting agent, such as sulfasalazine, that is administered per kg body weight of the patient (e.g., administered intravenously) may be at least 3 mg/kg, such as at least 5 mg/kg, and even at least 8 mg/kg. Generally, the dose of the glutathione-depleting agent, such as sulfasalazine, that is administered per kg body weight of the patient will not exceed 40 mg/kg. For example the dose of the glutathione-depleting agent, such as sulfasalazine, that is administered per kg body weight of the patient may be in the range of from 3 mg/kg to 40 mg/kg, such as from 4 mg/kg to 10 mg/kg.

[00173] In one embodiment, a therapeutically effective amount of the glutathione-depleting agent, such as BSO, may be an amount sufficient to provide a peak plasma concentration of at least 100  $\mu\text{M}$  when administered to a patient. For example, in one embodiment, the glutathione-depleting agent, such as BSO, may be administered in an amount sufficient to provide a peak plasma concentration of at least 200  $\mu\text{M}$  when administered to a patient. In yet another embodiment, the glutathione-depleting agent, such as BSO, may be administered in an amount sufficient to provide a peak plasma concentration of at least 250  $\mu\text{M}$  when administered to a patient. Generally, the glutathione-depleting agent, such as BSO, will not be administered in an amount that would provide a peak plasma concentration greater than 9000  $\mu\text{M}$  when administered to a patient, and may even provide a peak plasma concentration equal to or less than 1000  $\mu\text{M}$  in a patient. For example, the glutathione-depleting agent, such as BSO, may be administered in an amount sufficient to provide a peak plasma concentration in the range of from 100  $\mu\text{M}$  to 1000  $\mu\text{M}$  in a patient.

[00174] In yet another embodiment, a dose of the glutathione-depleting agent, such as BSO, that is administered per  $\text{m}^2$  of the patient (e.g., administered intravenously) may be at least 8  $\text{g}/\text{m}^2$ , such as at least 10  $\text{g}/\text{m}^2$ , and even at least 13

g/m<sup>2</sup>. Generally, the dose of the glutathione-depleting agent, such as BSO, that is administered per m<sup>2</sup> of the patient will not exceed 40 g/m<sup>2</sup>. For example the dose of the glutathione-depleting agent, such as BSO, that is administered per m<sup>2</sup> of the patient may be in the range of from 8 g/m<sup>2</sup> to 40 g/m<sup>2</sup>, such as from 10 g/m<sup>2</sup> to 17 g/m<sup>2</sup>.

5           [00175] Furthermore, it is noted that the dosage of the glutathione-depleting agent may vary according to the particular agent selected, as well as according to the route of administration, etc. For example, for glutathione-depleting agents having high potency, a lower dosage may be used to provide the same treatment effect, as compared to glutathione-depleting agents having less potency. Accordingly, while  
10           exemplary dosage ranges for the particular glutathione-depleting agents BSO and sulfasalazine are being provided herein, the exemplary ranges are not limiting, and BSO, sulfasalazine and/or other glutathione-depleting agents may be provided in different dosages, administration routes, or administration frequencies, other than those specifically exemplified herein, according to the particular glutathione-depleting agent to  
15           be administered and the therapeutic effect to be achieved.

          [00176] The dosing schedule of the glutathione-depleting agent, such as BSO or sulfasalazine, can similarly be selected according to the intended treatment. For example, in one embodiment, a suitable dosing schedule can comprise dosing a patient at least once per week, such as at least 2, 3, 4, 5, 6 or 7 days per week (e.g., daily),  
20           during a course of treatment. As another example, in one embodiment, the dosing may be at least once a day (qd), or even at least twice a day (bid). In one embodiment, the course of treatment with the glutathione-depleting agent, such as BSO or sulfasalazine, may last at least as long as a course of treatment with one or more other agents, such as least one of the pentaaza macrocyclic ring complex and thioredoxin reductase  
25           inhibitor agent, and may even exceed the duration during which the at least one other agent is provided. The course of therapy with the glutathione-depleting agent, such as BSO or sulfasalazine, may also start on the same date as treatment with the other agent, such as on a same date as treatment with at least one of the pentaaza macrocyclic ring complex and thioredoxin reductase inhibitor, or may start sometime  
30           after initial dosing with at least one of the other agents.

## METHODS OF ADMINISTRATION

[00177] According to one embodiment, the at least one active agent that is at least one of a thioredoxin reductase inhibitor and a glutathione depleting agent, is administered as a co-therapy or combination therapy with the pentaaza macrocyclic ring complex. Co-therapy or combination therapy according to the methods described herein is intended to embrace administration of each compound in a sequential manner in a regimen that will provide beneficial effects of the drug combination, and is intended as well to embrace co-administration of these agents in a substantially simultaneous manner, such as in a single capsule having a fixed ratio of these active agents or in multiple, separate capsules for each agent, or single or multiple parenteral administrations, or other routes of administration and dosage forms. When administered in combination, therefore, the therapeutic agents (i.e., the pentaaza macrocyclic ring complex, the thioredoxin reductase inhibitor, and/or the glutathione-depleting agent) can be formulated as separate compositions that are administered at the same time or sequentially at different times, or the therapeutic agents can be given as a single composition. Pharmaceutical compositions and formulations are discussed elsewhere herein. Furthermore, while the at least one "active agent" is referred to herein as being at least one of the thioredoxin reductase inhibitor and glutathione depleting agent, it is noted that all combinations of these are also explicitly included herein, including administration of thioredoxin reductase inhibitor without a glutathione depleting agent, administration of a glutathione-depleting agent without a thioredoxin reductase inhibitor, and administration of both a thioredoxin reductase inhibitor and a glutathione-depleting agent.

[00178] It is not necessary that the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione-depleting agent, be administered simultaneously or essentially simultaneously; the agents and compounds may be administered in sequence. The advantage of a simultaneous or essentially simultaneous administration, or sequential administration, is well within the determination of the skilled clinician. For instance, while a pharmaceutical composition or formulation comprising a pentaaza macrocyclic ring complex may be advantageous for administering first in the combination for one particular treatment, prior administration of the thioredoxin reductase inhibitor and/or glutathione depleting agent(s) (or prior administration of the pentaaza macrocyclic ring complex) may be

advantageous in another treatment. It is also understood that the instant combination of pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and glutathione depleting agent may be used in conjunction with other methods of treating cancer (typically cancerous tumors) including, but not limited to, radiation therapy and surgery, or other chemotherapy. It is further understood that a cytostatic or quiescent agent, or antiemetic agent, or hematopoietic growth factor, if any, may be administered sequentially or simultaneously with any or all of the other synergistic therapies.

[00179] Thus, embodiment of the therapeutic method include wherein a pentaaza macrocyclic ring complex and at least one active agent selected from thioredoxin reductase inhibitor, glutathione-depleting agent, and combinations thereof, are administered simultaneously or sequentially. For instance, the present disclosure encompasses a method for the treatment of cancer wherein a pentaaza macrocyclic ring complex and a thioredoxin reductase inhibitor are administered simultaneously or sequentially. By way of another example, the present disclosure encompasses a method for the treatment of cancer wherein a pentaaza macrocyclic ring complex and a glutathione depleting agent are administered simultaneously or sequentially. Further, the present disclosure encompasses a method for the treatment of cancer wherein a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and a glutathione inhibiting agent are administered simultaneously or sequentially.

[00180] As noted above, if the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and glutathione depleting agent are not administered simultaneously or essentially simultaneously, then the initial order of administration of the components may be varied.

[00181] Thus, for example, a pentaaza macrocyclic ring complex may be administered first, followed by the administration of a thioredoxin reductase inhibitor; or a thioredoxin reductase may be administered first, followed by the administration of a pentaaza macrocyclic ring complex. Similarly, a pentaaza macrocyclic ring complex may be administered first, followed by the administration of a glutathione depleting agent; or a glutathione depleting agent may be administered first, followed by the administration of a pentaaza macrocyclic ring complex. Where a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and a glutathione-depleting agent are administered in sequence, the sequence may vary accordingly. This alternate administration may be

repeated during a single treatment protocol. The determination of the order of administration, and the number of repetitions of administration of each therapeutic agent during a treatment protocol, is well within the knowledge of the skilled physician after evaluation of the disease being treated and the condition of the patient. By way of another example, at least one of the thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered initially (e.g., to decrease hydrogen peroxide metabolism). The treatment is then continued with the administration of the pentaaza macrocyclic ring complex (e.g., to produce new to hydrogen peroxide from superoxide), until the treatment protocol is complete. Other sequences of administration to exploit the effects described herein are contemplated.

[00182] In one embodiment, the subject is pre-treated with the pentaaza macrocyclic ring complex (i.e., the pentaaza macrocyclic ring complex is pre-administered), followed by administration of at least one of the other active agents (i.e., the thioredoxin reductase inhibitor and/or the glutathione depleting agent), or vice versa. In accordance with such embodiments, the other active agents is/are is preferably administered at least 1 hour, but no more than 3 days, after administration of the pentaaza macrocyclic ring complex, or vice versa. For example, in one embodiment, the other active agents is/are administered between 1 hour and 2 days after administration of the pentaaza macrocyclic ring complex, or vice versa. In another embodiment, for example, the other active agents is/are administered between 1 hour and 1 day after administration of the pentaaza macrocyclic ring complex, or vice versa. For example, the other active agent may be administered within 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 12 hours, 18 hours, 24 hours, 36 hours, or 48 hours after administration of the pentaaza macrocyclic ring complex, or vice versa. In one particular embodiment, for example, the active agents is/are administered within 24 hours after administration of the pentaaza macrocyclic ring complex, or vice versa. In these and other embodiments, the pentaaza macrocyclic ring complex may be administered in multiple doses leading up to administration of the other active agent.

[00183] Alternatively, the subject may be pre-treated with the active agents (i.e., the thioredoxin reductase inhibitor(s) and/or the glutathione depleting agent(s)), followed by administration of the pentaaza macrocyclic ring complex, or vice versa. In accordance with such embodiments, the pentaaza macrocyclic ring complex is preferably administered within at least 1 plasma half-life of the other active agents, but

no more than 4 plasma half-lives of the other active agents, or vice versa. For example, the pentaaza macrocyclic ring complex may be administered within 1, 2, or 3 plasma half-lives of the other active agents, or vice versa.

[00184] In other alternative embodiments, the subject may be pre-treated with the pentaaza macrocyclic ring complex, followed by administration of at least one of the other active agent, which is further followed by an additional administration of the pentaaza macrocyclic ring complex. In accordance with this embodiment, for example, the standard pentaaza macrocyclic ring complex dose may be separated into two (or more) portions, one portion of which is administered prior to administration of the other active agent, and the second portion of which is administered after administration of the other active agent. This staggered therapy regime could also be employed where the active agents is/are administered first. In addition, the subject could be pre-treated with a partial or full dose of pentaaza macrocyclic ring complex, followed by administration of a first active agent (e.g., one of the thioredoxin reductase inhibitor), which is then followed by the administration of additional (or partial) dose of pentaaza macrocyclic ring complex, which may be further followed by administration of a second active agent (e.g., the other of the thioredoxin reductase inhibitor and glutathione depleting agent).

[00185] As described in further detail below, the combinations of the disclosure may also be co-administered with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. Combinations may alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a multiple combination formulation is inappropriate.

[00186] The pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and glutathione depleting agent are generally administered according to therapeutic protocols well known in the art. It will be apparent to those skilled in the art that the administration of the various components can be varied depending on the disease being treated and the known effects of pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione depleting agent on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic protocols (e.g., dosage amounts and times of administration) can be varied in view of the observed effects of the administered therapeutic agents (i.e., pentaaza macrocyclic ring complex,

thioredoxin reductase inhibitor, and glutathione depleting agent) on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

[00187] Also, in general, pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione depleting agent do not have to be administered in the same pharmaceutical composition, and may, because of different physical and chemical characteristics, have to be administered by different routes. For example, the pentaaza macrocyclic ring complex may be administered orally to generate and maintain good blood levels thereof, while the thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered intravenously, or vice versa. The determination of the mode of administration and the advisability of administration, where possible, in the same pharmaceutical composition, or in separate pharmaceutical compositions (e.g., two or three separate compositions) is well within the knowledge of the skilled clinician. The initial administration can be made according to established protocols known in the art, and then, based upon the observed effects, the dosage, modes of administration and times of administration can be modified by the skilled clinician.

[00188] The particular choice of pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and glutathione depleting agent (each of which are described in detail herein), and other related therapies (such as chemotherapy or radiation), will depend upon the diagnosis of the attending physicians and their judgment of the condition of the patient and the appropriate treatment protocol.

[00189] Thus, in accordance with experience and knowledge, the practicing physician can modify each protocol for the administration of a component (pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione depleting agent) of the treatment according to the individual patient's needs, as the treatment proceeds.

[00190] The attending clinician, in judging whether treatment is effective at the dosage administered, will consider the general well-being of the patient as well as more definite signs such as relief of disease-related symptoms, inhibition of tumor growth, actual shrinkage of the tumor, or inhibition of metastasis. Size of the tumor can be measured by standard methods such as radiological studies, e.g., CAT or MRI scan, and successive measurements can be used to judge whether or not growth of the tumor

has been retarded or even reversed. Relief of disease-related symptoms such as pain, and improvement in overall condition can also be used to help judge effectiveness of treatment.

[00191] The products of which the combination are composed may be  
5 administered simultaneously, separately or spaced out over a period of time so as to obtain the maximum efficacy of the combination; it being possible for each administration to vary in its duration from a rapid administration to a relatively continuous perfusion of either component (in separate formulations or in a single formulation). As a result, for the purposes of the present disclosure, the combinations  
10 are not exclusively limited to those which are obtained by physical association of the constituents, but also to those which permit a separate administration, which can be simultaneous or spaced out over a period of time.

[00192] Accordingly, administration of the components described herein can occur as a single event or over a time course of treatment. For example, one or more  
15 of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione depleting agent can be administered (simultaneously or in sequence) hourly (e.g., every hour, every two hours, every three hours, every four hours, every five hours, every six hours, and so on), daily, weekly, bi-weekly, or monthly. For treatment of acute conditions, the time course of treatment may be at least several hours or days.  
20 Certain conditions could extend treatment from several days to several weeks. For example, treatment could extend over one week, two weeks, or three weeks. For more chronic conditions, treatment could extend from several weeks to several months, a year or more, or the lifetime of the patient in need of such treatment. Alternatively, the compounds and agents can be administered hourly, daily, weekly, bi-weekly, or  
25 monthly, for a period of several weeks, months, years, or over the lifetime of the patient as a prophylactic measure.

[00193] The dose or amount of pharmaceutical compositions including the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor, and/or glutathione depleting agent administered to the patient should be an effective amount for the intended purpose, i.e., treatment or prophylaxis of one or more of the diseases, pathological disorders, and medical conditions discussed herein, particularly cancer. Generally speaking, the effective amount of the composition administered can vary according to a variety of factors such as, for example, the age, weight, sex, diet, route of administration, and the medical condition of the patient in need of the treatment. Specifically preferred doses are discussed more fully below. It will be understood, however, that the total daily usage of the compositions described herein will be decided by the attending physician or veterinarian within the scope of sound medical judgment.

[00194] As noted above, the combinations can be co-administered (via a co-formulated dosage form or in separate dosage forms administered at about the same time). The combinations can also be administered separately, at different times, with each agent in a separate unit dosage form. Numerous approaches for administering anti-cancer drugs and pentaaza macrocyclic ring complex are known in the art, and can readily be adapted for use in the present disclosure. The pharmaceutical compositions may be delivered orally, e.g., in a tablet or capsule unit dosage form, or parenterally, e.g., in an injectable unit dosage form, or by some other route. For systemic administration, for example, the drugs can be administered by, for example, intravenous infusion (continuous or bolus). The compositions can be used for any therapeutic or prophylactic treatment where the patient benefits from treatment with the combination.

[00195] The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound(s) employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the route of administration; the rate of excretion of the specific compound(s) employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound(s) employed and like factors well known in the medical and/or veterinary arts. For example, it is well within the skill of the art to start doses of the compound(s) at

levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. If desired, the effective daily doses may be divided into multiple doses for purposes of administration. Consequently, single dose compositions may contain such amounts or submultiples to make up the daily dose.

[00196] Suitable or preferred doses for each of the components employed in the methods or included in the compositions described herein are generally known in the art. Preferred dosages for the pentaaza macrocyclic ring complex, for instance, may be within the range of 10 to 500 mg per patient per day. A typical dose of auranofin, a thioredoxin reductase inhibitor, can be a dose that results in blood levels of 0.5  $\mu$ M to 5  $\mu$ M, such as 0.5  $\mu$ M to 2  $\mu$ M. For buthionine sulfoximine, a glutathione-depleting agent, a suitable dose range may be one that provides a blood level of 100  $\mu$ M to 9000  $\mu$ M, such as from 100  $\mu$ M to 1000  $\mu$ M. For sulfasalazine, a glutathione-depleting agent, a suitable dose range may be one that provides a blood level of 200  $\mu$ M to 600  $\mu$ M. However, the dosage may vary depending on the dosing schedule, which can be adjusted as necessary to achieve the desired therapeutic effect. It should be noted that the ranges of effective doses provided herein are not intended to limit the disclosure and represent exemplary dose ranges. The most preferred dosage will be tailored to the individual subject, taking into account, among other things, the particular combinations employed, and the patient's age, sex, weight, physical condition, diet, etc., as is understood and determinable by one of ordinary skill in the art without undue experimentation.

[00197] Treatment of cancer, or cancer therapies, described herein includes achieving a therapeutic benefit and/or a prophylactic benefit. Therapeutic benefits generally refer to at least a partial eradication or amelioration of the underlying disorder being treated. For example, in a cancer patient, therapeutic benefit includes (partial or complete) eradication or amelioration of the underlying cancer. Also, a therapeutic benefit is achieved with at least partial, or complete, eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the patient, notwithstanding the fact that the patient may still be afflicted with the underlying disorder. For prophylactic benefit, a method of the disclosure may be performed on, or a composition of the invention administered to, a patient at risk of developing cancer, or to a patient reporting one or more of the

physiological symptoms of such conditions, even though a diagnosis of the condition may not have been made.

### **Other Cancer Therapies**

[00198] In one embodiment, the treatment provided herein can further  
5 comprise treatment with another therapy other than those specifically described above, such as for example a radiation therapy, a chemotherapy, or other treatment. For example, in one embodiment, one or more of radiation therapy and chemotherapy is administered to the subject prior to, concomitantly with, or after administration of one or more of at least one of the thioredoxin reductase inhibitor and glutathione-depleting  
10 agent, and the pentaaza macrocyclic ring complex. Further detailed description of radiation therapies and chemotherapies suitable for the treatment of cancer are provided below.

[00199] In one embodiment, one or more of radiation therapy and chemotherapy can be administered concomitantly with administration of at least one of  
15 the thioredoxin reductase inhibitor and glutathione-depleting agent, and pentaaza macrocyclic ring complex. For example, one or more of the agent(s) and pentaaza macrocyclic ring complexes may be administered during a course of radiation therapy and/or chemotherapy, such as in between, before or after, or on the same day as dosing with radiation and/or chemotherapy.

[00200] In yet another embodiment, the combination therapy of the pentaaza macrocyclic ring complex and agents(s) (e.g. thioredoxin reductase inhibitor, glutathione-depleting agent), can be administered in the absence of any other cancer treatment. As demonstrated further in the examples below, it has been unexpectedly  
20 discovered that the pentaaza macrocyclic ring complexes exhibit synergistic results when combined with at least one of the thioredoxin reductase inhibitor and glutathione-depleting agent, even when administered without radiation therapy or chemotherapy. Accordingly, in one embodiment, the cancer treatment provided to the subject may consist essentially of the pentaaza macrocyclic ring complex and one or more of the agents, without the administration of a chemotherapeutic agent or radiation exposure  
25 (i.e. without administering a radiation dose or dose fraction). For example, the combination of the pentaaza macrocyclic ring complex and at least one agent may be administered to a subject that is not receiving radiation therapy, and/or that is not  
30

receiving chemotherapy. In one embodiment, the treatment comprises administering the pentaaza macrocyclic ring complex to a subject that is not receiving radiation therapy. In yet another embodiment, the treatment comprises administering both (i) at least one of the agents, and (ii) the pentaaza macrocyclic ring complex, to a subject that is not receiving radiation therapy. In yet another embodiment, where a course of therapy comprises administration of the pentaaza macrocyclic ring complex and at least one of the agents (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent), they are administered to a subject that does not receive radiation therapy during the course of therapy.

[00201] In one embodiment, the subject receiving the combination of pentaaza macrocyclic ring complex and the at least one agent (e.g. thioredoxin reductase inhibitor, glutathione-depleting agent), may be one that has not been exposed to radiation (i.e., received a dose or dose fraction of radiation) and/or has not received a dose of chemotherapeutic agent for at least one day, such as at least one week, and even at least one month, and even at least 6 months, and/or that has not ever received such treatment at all before initial treatment with one or more of the pentaaza macrocyclic ring complex and agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent). In yet another embodiment, any radiation therapy and/or chemotherapy that is administered to the subject after the combination treatment with the pentaaza macrocyclic ring complex and agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent) is delayed by at least one day, such as at least one week, and even at least one month, such as at least 6 months, after a final dose of one or more of the pentaaza macrocyclic ring complex and agent(s) provided during the course of the combination therapy treatment. That is, the combination therapy of the pentaaza macrocyclic ring complex and agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent) can be administered to a subject that has never before received radiation therapy and/or chemotherapy, or that has received such therapy only in the distant past. Furthermore, the combination therapy of the pentaaza macrocyclic ring complex and agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent) can be administered to provide a course of treatment that does not include any exposure to radiation or doses of chemotherapeutic agent. As yet a further embodiment, the combination therapy of the pentaaza macrocyclic ring complex and agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent) can be

provided to form a course of treatment substantially without performing any radiation therapy or chemotherapy after the course of treatment, or with such radiation or chemotherapeutic treatment being performed only after a significant period of time has elapsed after the course of combination treatment has ended. In one embodiment, the treatment comprises administering one or more of the pentaaza macrocyclic ring complex agent(s) (e.g., thioredoxin reductase inhibitor, glutathione-depleting agent) to the subject on a day other than a day that the subject is receiving radiation therapy.

### **Cancer Treatment Methods**

[00202] In general, any subject having, or suspected of having, a cancer or other proliferative disorder may be treated using the compositions and methods of the present disclosure. Subjects receiving treatment according to the methods described herein are mammalian subjects, and typically human patients. Other mammals that may be treated according to the present disclosure include companion animals such as dogs and cats, farm animals such as cows, horses, and swine, as well as birds and more exotic animals (e.g., those found in zoos or nature preserves). In one embodiment of the disclosure, a method is provided for the treatment of cancerous tumors, particularly solid tumors. Advantageously, the methods described herein may reduce the development of tumors, reduce tumor burden, or produce tumor regression in a mammalian host. Cancer patients and individuals desiring cancer prophylaxis can be treated with the combinations described herein.

[00203] Cancer and tumors generally refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. By means of the pharmaceutical combinations, co-formulations, and combination therapies of the present disclosure, various tumors can be treated such as tumors of the breast, heart, lung, small intestine, colon, spleen, kidney, bladder, head and neck, ovary, prostate, brain, pancreas, skin, bone, bone marrow, blood, thymus, uterus, testicles, cervix, and liver.

[00204] In one embodiment, the tumor or cancer is chosen from adenoma, angio-sarcoma, astrocytoma, epithelial carcinoma, germinoma, glioblastoma, glioma, hamartoma, hemangioendothelioma, hemangiosarcoma, hematoma, hepatoblastoma, leukemia, lymphoma, medulloblastoma, melanoma, neuroblastoma, osteosarcoma, retinoblastoma, rhabdomyosarcoma, sarcoma, and teratoma. The

tumor can be chosen from acral lentiginous melanoma, actinic keratoses, adenocarcinoma, adenoid cystic carcinoma, adenomas, adenosarcoma, adenosquamous carcinoma, astrocytic tumors, Bartholin gland carcinoma, basal cell carcinoma, bronchial gland carcinomas, capillary, carcinoids, carcinoma, 5 carcinosarcoma, cavernous, cholangio-carcinoma, chondrosarcoma, choroid plexus papilloma/carcinoma, clear cell carcinoma, cystadenoma, endodermal sinus tumor, endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, ependymal, epitheloid, Ewing's sarcoma, fibrolamellar, focal nodular hyperplasia, gastrinoma, germ cell tumors, glioblastoma, glucagonoma, 10 hemangioblastomas, hemangioendothelioma, hemangiomas, hepatic adenoma, hepatic adenomatosis, hepatocellular carcinoma, insulinoma, intraepithelial neoplasia, interepithelial squamous cell neoplasia, invasive squamous cell carcinoma, large cell carcinoma, leiomyosarcoma, lentigo maligna melanomas, malignant melanoma, malignant mesothelial tumors, medulloblastoma, medulloepithelioma, melanoma, 15 meningeal, mesothelial, metastatic carcinoma, mucoepidermoid carcinoma, neuroblastoma, neuroepithelial adenocarcinoma nodular melanoma, oat cell carcinoma, oligodendroglial, osteosarcoma, pancreatic, papillary serous adeno-carcinoma, pineal cell, pituitary tumors, plasmacytoma, pseudo-sarcoma, pulmonary blastoma, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, small cell 20 carcinoma, soft tissue carcinomas, somatostatin-secreting tumor, squamous carcinoma, squamous cell carcinoma, submesothelial, superficial spreading melanoma, undifferentiated carcinoma, uveal melanoma, verrucous carcinoma, vipoma, well differentiated carcinoma, and Wilm's tumor.

[00205] Thus, for example, the present disclosure provides methods for the 25 treatment of a variety of cancers, including, but not limited to, the following: carcinoma including that of the bladder (including accelerated and metastatic bladder cancer), breast, colon (including colorectal cancer), kidney, liver, lung (including small and non-small cell lung cancer and lung adenocarcinoma), ovary, prostate, testes, genitourinary tract, lymphatic system, rectum, larynx, pancreas (including exocrine pancreatic 30 carcinoma), esophagus, stomach, gall bladder, cervix, thyroid, and skin (including squamous cell carcinoma); hematopoietic tumors of lymphoid lineage including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma,

histiocytic lymphoma, and Burkett's lymphoma; hematopoietic tumors of myeloid lineage including acute and chronic myelogenous leukemias, myelodysplastic syndrome, myeloid leukemia, and promyelocytic leukemia; tumors of the central and peripheral nervous system including astrocytoma, neuroblastoma, glioma, and schwannomas; tumors of mesenchymal origin including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors including melanoma, xeroderma pigmentosum, keratoactanthoma, seminoma, thyroid follicular cancer, and teratocarcinoma.

[00206] For example, particular leukemias that can be treated with the combinations and methods described herein include, but are not limited to, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, acute granulocytic leukemia, chronic granulocytic leukemia, acute promyelocytic leukemia, adult T-cell leukemia, aleukemic leukemia, a leukocythemic leukemia, basophilic leukemia, blast cell leukemia, bovine leukemia, chronic myelocytic leukemia, leukemia cutis, embryonal leukemia, eosinophilic leukemia, Gross' leukemia, hairy-cell leukemia, hemoblastic leukemia, hemocytoblastic leukemia, histiocytic leukemia, stem cell leukemia, acute monocytic leukemia, leukopenic leukemia, lymphatic leukemia, lymphoblastic leukemia, lymphocytic leukemia, lymphogenous leukemia, lymphoid leukemia, lymphosarcoma cell leukemia, mast cell leukemia, megakaryocytic leukemia, micromyeloblastic leukemia, monocytic leukemia, myeloblastic leukemia, myelocytic leukemia, myeloid granulocytic leukemia, myelomonocytic leukemia, Naegeli leukemia, plasma cell leukemia, plasmacytic leukemia, promyelocytic leukemia, Rieder cell leukemia, Schilling's leukemia, stem cell leukemia, subleukemic leukemia, and undifferentiated cell leukemia.

[00207] Lymphomas can also be treated with the combinations and methods described herein. Lymphomas are generally neoplastic transformations of cells that reside primarily in lymphoid tissue. Lymphomas are tumors of the immune system and generally are present as both T cell- and as B cell-associated disease. Among lymphomas, there are two major distinct groups: non-Hodgkin's lymphoma (NHL) and Hodgkin's disease. Bone marrow, lymph nodes, spleen and circulating cells, among others, may be involved. Treatment protocols include removal of bone marrow from the patient and purging it of tumor cells, often using antibodies directed against antigens present on the tumor cell type, followed by storage. The patient is then given a toxic

dose of radiation or chemotherapy and the purged bone marrow is then re-infused in order to repopulate the patient's hematopoietic system.

[00208] Other hematological malignancies that can be treated with the combinations and methods described herein include myelodysplastic syndromes (MDS), myeloproliferative syndromes (MPS) and myelomas, such as solitary myeloma and multiple myeloma. Multiple myeloma (also called plasma cell myeloma) involves the skeletal system and is characterized by multiple tumorous masses of neoplastic plasma cells scattered throughout that system. It may also spread to lymph nodes and other sites such as the skin. Solitary myeloma involves solitary lesions that tend to occur in the same locations as multiple myeloma.

[00209] In one embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of breast cancer, melanoma, oral squamous cell carcinoma, lung cancer including non-small cell lung cancer, renal cell carcinoma, colorectal cancer, prostate cancer, brain cancer, spindle cell carcinoma, urothelial cancer, bladder cancer, colorectal cancer, head and neck cancers such as squamous cell carcinoma, and pancreatic cancer. In one embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of breast cancer, lung cancer including non-small cell lung cancer, renal cell carcinoma, spindle cell carcinoma, colorectal cancer, head and neck squamous cell carcinoma, pancreatic cancer. In yet another embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of head and neck cancer and lung cancer. In yet another embodiment, the methods and pharmaceutical compositions described herein are used to treat a cancer that is any of breast cancer and lung cancer.

## **Pharmaceutical Formulations**

[00210] Another aspect of the present disclosure relates to the pharmaceutical compositions comprising the combinations described herein, together with a pharmaceutically acceptable excipient. The pharmaceutical compositions include the pentaaza macrocyclic ring complex (e.g., those corresponding to Formula (I)), and at least one active agent selected from a thioredoxin reductase inhibitor, a glutathione depleting agent, and combinations thereof, as discussed above, typically formulated as a pharmaceutical dosage form, optionally in combination with a pharmaceutically

acceptable carrier, additive or excipient. In one embodiment, for example, the pharmaceutical composition comprises a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and a pharmaceutically acceptable excipient. In another embodiment, the pharmaceutical composition comprises a pentaaza macrocyclic ring complex, a glutathione depleting agent, and a pharmaceutically acceptable excipient. In yet another embodiment, the pharmaceutical composition comprises a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, a glutathione depleting agent, and a pharmaceutically acceptable excipient. Pharmaceutical compositions according to the present disclosure may be used in the treatment of cancer.

[00211] The pharmaceutical compositions described herein can comprise products that result from the mixing or combining of more than one active ingredients, and includes both fixed and non-fixed combinations of the active ingredients. Fixed combinations are those in which the active ingredients, e.g., a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent described herein, are both administered to a patient simultaneously in the form of a single entity or dosage. Non-fixed combinations are those in which the active ingredients, e.g., a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent described herein, are administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific intervening time limits, wherein such administration provides effective levels of the two compounds in the body of the patient. The latter also applies to cocktail therapy, e.g., the administration of three or more active ingredients.

[00212] The above-described a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent may be dispersed in a pharmaceutically acceptable carrier prior to administration to the mammal; i.e., the components described herein are preferably co-formulated. The carrier, also known in the art as an excipient, vehicle, auxiliary, adjuvant, or diluent, is typically a substance which is pharmaceutically inert, confers a suitable consistency or form to the composition, and does not diminish the efficacy of the compound. The carrier is generally considered to be "pharmaceutically or pharmacologically acceptable" if it does not produce an unacceptably adverse, allergic or other untoward reaction when administered to a mammal, especially a human.

[00213] The selection of a pharmaceutically acceptable carrier will also, in part, be a function of the route of administration. In general, the compositions of the described herein can be formulated for any route of administration so long as the blood circulation system is available via that route, and in accordance with the conventional route of administration of the component (e.g., a pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent). For example, suitable routes of administration include, but are not limited to, oral, parenteral (e.g., intravenous, intraarterial, subcutaneous, rectal, subcutaneous, intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, or intrasternal), topical (nasal, transdermal, intraocular), intravesical, intrathecal, enteral, pulmonary, intralymphatic, intracavitary, vaginal, transurethral, intradermal, aural, intramammary, buccal, orthotopic, intratracheal, intralesional, percutaneous, endoscopic, transmucosal, sublingual and intestinal administration.

[00214] Pharmaceutically acceptable carriers for use in combination with the compositions of the present disclosure are well known to those of ordinary skill in the art and are selected based upon a number of factors: the particular compound(s) and agent(s) used, and its/their concentration, stability and intended bioavailability; the subject, its age, size and general condition; and the route of administration. Suitable nonaqueous, pharmaceutically-acceptable polar solvents include, but are not limited to, alcohols (e.g., 1,2-glycerol formal, 6-glycerol formal, 1,3-butyleneglycol, aliphatic or aromatic alcohols having 2 to 30 carbon atoms such as methanol, ethanol, propanol, isopropanol, butanol, t-butanol, hexanol, octanol, amylene hydrate, benzyl alcohol, glycerin (glycerol), glycol, hexylene glycol, tetrahydrofurfuryl alcohol, lauryl alcohol, cetyl alcohol, or stearyl alcohol, fatty acid esters of fatty alcohols such as polyalkylene glycols (e.g., polypropylene glycol, polyethylene glycol), sorbitan, sucrose and cholesterol); amides (e.g., dimethylacetamide (DMA), benzyl benzoate DMA, dimethylformamide, N-(6-hydroxyethyl)-lactamide, N,N-dimethylacetamide amides, 2-pyrrolidinone, 1-methyl-2-pyrrolidinone, or polyvinylpyrrolidone); esters (e.g., 1-methyl-2-pyrrolidinone, 2-pyrrolidinone, acetate esters such as monoacetin, diacetin, and triacetin, aliphatic or aromatic esters such as ethyl caprylate or octanoate, alkyl oleate, benzyl benzoate, benzyl acetate, dimethylsulfoxide (DMSO), esters of glycerin such as mono, di-, or tri-glyceryl citrates or tartrates, ethyl benzoate, ethyl acetate, ethyl carbonate, ethyl lactate, ethyl oleate, fatty acid esters of sorbitan, fatty acid derived PEG esters, glyceryl

monostearate, glyceride esters such as mono, di-, or tri-glycerides, fatty acid esters such as isopropyl myristate, fatty acid derived PEG esters such as PEG-hydroxyoleate and PEG-hydroxystearate, N-methyl pyrrolidinone, pluronic 60, polyoxyethylene sorbitol oleic polyester, polyoxyethylene sorbitan esters such as polyoxyethylene-sorbitan monooleate, polyoxyethylene-sorbitan monopalmitate, polyoxyethylene-sorbitan monolaurate, polyoxyethylene-sorbitan monostearate, and Polysorbate® 20, 40, 60 or 80 from ICI Americas, Wilmington, DE, polyvinylpyrrolidone, alkyleneoxy modified fatty acid esters such as polyoxyl 40 hydrogenated castor oil and polyoxyethylated castor oils (e.g., Cremophor® EL solution or Cremophor® RH 40 solution), saccharide fatty acid esters (i.e., the condensation product of a monosaccharide (e.g., pentoses such as ribose, ribulose, arabinose, xylose, lyxose and xylulose, hexoses such as glucose, fructose, galactose, mannose and sorbose, trioses, tetroses, heptoses, and octoses), disaccharide (e.g., sucrose, maltose, lactose and trehalose) or oligosaccharide or mixture thereof with a C<sub>4</sub> to C<sub>22</sub> fatty acid(s) (e.g., saturated fatty acids such as caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid and stearic acid, and unsaturated fatty acids such as palmitoleic acid, oleic acid, elaidic acid, erucic acid and linoleic acid)), or steroidal esters); alkyl, aryl, or cyclic ethers having 2 to 30 carbon atoms (e.g., diethyl ether, tetrahydrofuran, dimethyl isosorbide, diethylene glycol monoethyl ether); glycofurool (tetrahydrofurfuryl alcohol polyethylene glycol ether); ketones having 3 to 30 carbon atoms (e.g., acetone, methyl ethyl ketone, methyl isobutyl ketone); aliphatic, cycloaliphatic or aromatic hydrocarbons having 4 to 30 carbon atoms (e.g., benzene, cyclohexane, dichloromethane, dioxolanes, hexane, n-decane, n-dodecane, n-hexane, sulfolane, tetramethylenesulfon, tetramethylenesulfoxide, toluene, dimethylsulfoxide (DMSO), or tetramethylenesulfoxide); oils of mineral, vegetable, animal, essential or synthetic origin (e.g., mineral oils such as aliphatic or wax-based hydrocarbons, aromatic hydrocarbons, mixed aliphatic and aromatic based hydrocarbons, and refined paraffin oil, vegetable oils such as linseed, tung, safflower, soybean, castor, cottonseed, groundnut, rapeseed, coconut, palm, olive, corn, corn germ, sesame, persic and peanut oil and glycerides such as mono-, di- or triglycerides, animal oils such as fish, marine, sperm, cod-liver, haliver, squalene, squalane, and shark liver oil, oleic oils, and polyoxyethylated castor oil); alkyl or aryl halides having 1 to 30 carbon atoms and optionally more than one halogen substituent; methylene chloride; monoethanolamine; petroleum benzin; triethylamine; omega-3 polyunsaturated fatty acids

(e.g., alpha-linolenic acid, eicosapentaenoic acid, docosapentaenoic acid, or docosahexaenoic acid); polyglycol ester of 12-hydroxystearic acid and polyethylene glycol (Solutol® HS-15, from BASF, Ludwigshafen, Germany); polyoxyethylene glycerol; sodium laurate; sodium oleate; or sorbitan monooleate.

5           [00215] In some embodiments, oils or non-aqueous solvents may be employed in the formulations, e.g., to bring one or more of the compounds into solution, due to, for example, the presence of large lipophilic moieties. Alternatively, emulsions, suspensions, or other preparations, for example, liposomal preparations, may be used. With respect to liposomal preparations, for example, any known  
10 methods for preparing liposomes may be used. See, for example, Bangham *et al.*, *J. Mol. Biol.*, 23: 238-252 (1965) and Szoka *et al.*, *Proc. Natl Acad. Sci* 75: 4194-4198 (1978), incorporated herein by reference. Thus, in one embodiment, one or more of the compounds are administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles.  
15 Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. Ligands may also be attached to the liposomes, for instance, to direct these compositions to particular sites of action.

          [00216] Other pharmaceutically acceptable solvents for use in the pharmaceutical compositions described herein are well known to those of ordinary skill  
20 in the art, and are identified in The Chemotherapy Source Book (Williams & Wilkins Publishing), The Handbook of Pharmaceutical Excipients, (American Pharmaceutical Association, Washington, D.C., and The Pharmaceutical Society of Great Britain, London, England, 1968), Modern Pharmaceutics, (G. Banker *et al.*, eds., 3d ed.) (Marcel Dekker, Inc., New York, New York, 1995), The Pharmacological Basis of  
25 Therapeutics, (Goodman & Gilman, McGraw Hill Publishing), Pharmaceutical Dosage Forms, (H. Lieberman *et al.*, eds.) (Marcel Dekker, Inc., New York, New York, 1980), Remington's Pharmaceutical Sciences (A. Gennaro, ed., 19th ed.) (Mack Publishing, Easton, PA, 1995), The United States Pharmacopeia 24, The National Formulary 19, (National Publishing, Philadelphia, PA, 2000), and A.J. Spiegel *et al.*, Use of  
30 Nonaqueous Solvents in Parenteral Products, *Journal of Pharmaceutical Sciences*, Vol. 52, No. 10, pp. 917-927 (1963).

[00217] Formulations containing the pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent may take the form of solid, semi-solid, lyophilized powder, or liquid dosage forms such as, for instance, aerosols, capsules, creams, emulsions, foams, gels/jellies, lotions, ointments, pastes, powders, soaps, solutions, sprays, suppositories, suspensions, sustained-release formulations, tablets, tinctures, transdermal patches, and the like, preferably in unit dosage forms suitable for simple administration of precise dosages. If formulated as a fixed dose, such pharmaceutical compositions or formulation products employ the pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent within accepted dosage ranges.

[00218] Formulations for certain pentaaza macrocyclic ring complexes are also described in, for example, in U.S. Patent Nos. 5,610,293, 5,637,578, 5,874,421, 5,976,498, 6,084,093, 6,180,620, 6,204,259, 6,214,817, 6,245,758, 6,395,725, and 6,525,041 (each of which is hereby incorporated herein by reference in its entirety).

[00219] It is contemplated that co-formulations of the pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent may employ conventional formulation techniques for these components individually, or alternative formulation routes, subject to compatibility and efficacy of the various components, in combination.

[00220] The above-described pharmaceutical compositions including the pentaaza macrocyclic ring complex, a thioredoxin reductase inhibitor, and/or a glutathione depleting agent may additionally include one or more pharmaceutically active components. Suitable pharmaceutically active agents that may be included in the compositions of the present invention include, for instance, antiemetics, anesthetics, antihypertensives, antianxiety agents, anticlotting agents, anticonvulsants, blood glucose-lowering agents, decongestants, antihistamines, antitussives, antineoplastics, beta blockers, anti-inflammatory agents, antipsychotic agents, cognitive enhancers, cholesterol-reducing agents, antiobesity agents, autoimmune disorder agents, anti-impotence agents, antibacterial and antifungal agents, hypnotic agents, anti-Parkinsonism agents, anti-Alzheimer's Disease agents, antibiotics, anti-depressants, and antiviral agents. The individual components of such combinations

may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.

### **Combination Treatment with Cancer Therapy**

[00221] In one embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent can be administered in combination with another cancer therapy, to provide therapeutic treatment. For example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered as a part of at least one of a chemotherapy treatment and radiation therapy.

[00222] In general, the temporal aspects of the administration of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may depend for example, on the particular compound, radiation therapy, or chemotherapy that is selected, or the type, nature, and/or duration of the radiation exposure. Other considerations may include the disease or disorder being treated and the severity of the disease or disorder; activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors. For example, the compounds may be administered in various embodiments before, during, and/or after the administration of the cancer therapy (e.g., radiation therapy or chemotherapy, or before, during or after a course of radiation therapy or chemotherapy comprising multiple exposures and/or doses). By way of another example, the compound may be administered in various embodiments before, during, and/or after an exposure to radiation.

[00223] If desired, the effective dose can be divided into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the dose.

[00224] In one embodiment, for example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient prior to or simultaneous with the cancer therapy corresponding to at least one of radiation therapy and chemotherapy, such as prior to or

simultaneous with a dose or dose fraction of such treatment, or prior to or simultaneous with a course of such treatment comprising multiple doses. In another embodiment, for example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient prior to, but not after, the cancer therapy, such as before but not after a cancer therapy dose or dose fraction or prior to but not after a course of cancer therapy comprising multiple doses or dose fractions. In yet another embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient at least 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, prior to an initial dose or dose fraction of cancer therapy corresponding to at least one of radiation therapy and chemotherapy. In still other embodiments, for example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient after a dose or dose fraction of the cancer therapy; thus, for example, the compound may be administered up to 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, or 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, after the cancer treatment, such as after a single dose or dose fraction and/or final dose or dose fraction in a course of cancer treatment corresponding to one or more of radiation therapy and chemotherapy.

[00225] In another embodiment, for example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient prior to or simultaneous with the radiation exposure. In another embodiment, for example, the compounds are administered to the patient prior to, but not after, the radiation exposure. In yet another embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient at least 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, prior to the radiation exposure, such as an initial radiation exposure in a course of radiation treatment, or

prior to another dose or dose fraction of radiation that is one of the doses or dose fractions of radiation in the course of treatment. In still other embodiments, for example, pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered to the patient after the radiation exposure; thus, for  
5 example, the compound may be administered up to 15 minutes, 30 minutes, 45 minutes, 60 minutes, 90 minutes, or 180 minutes, 0.5 days, 1 day, 3 days, 5 days, one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks, ten weeks, eleven weeks, twelve weeks, or longer, after the radiation exposure, which may be a dose or dose fraction of radiation in a multi-dose  
10 course of radiation therapy, or may be the single or final dose or dose fraction of radiation in the radiation therapy.

[00226] In one embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered as a part of a course of therapy that includes the radiation therapy. In radiation therapy, a  
15 patient receives a dose of ionizing radiation to kill or control the growth of cancerous cells. The dose of radiation may be directed at a specific part of the body, and the beam of radiation may also be shaped according to a predetermined treatment regimen, to reduce deleterious effects on parts of the body not afflicted with cancer. A typical course of radiation therapy may include one or a plurality of doses of radiation, which  
20 can be administered over the course of days, weeks and even months. A total “dose” of radiation given during a course of radiation therapy typically refers to the amount of radiation a patient receives during the entire course of radiation therapy, which doses may be administered as dose “fractions” corresponding to multiple radiation exposures in the case where the total dose is administered over several sessions, with the sum of  
25 the fractions administered corresponding to the overall dose. As is discussed in more detail in the Examples section below, the administration of pentaaza macrocyclic ring complex with at least one of thioredoxin reductase inhibitor and glutathione depleting agent demonstrates unexpected synergistic effects in sensitizing cancer cells to radiation therapy, thereby improving the efficacy of radiation treatment.

30 [00227] In one embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin compound and glutathione depleting agent are administered within a predetermined time period before or after a radiation dose is administered. For example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or

glutathione depleting agent may be administered within 1 week, 48 hours, 24 hours, 12 hours, 6, hours, 2 hours, 1 hour or even within 30 minutes of the patient receiving the radiation exposure, such as the dose or dose fraction (either before or after the radiation exposure corresponding to the radiation dose or dose fraction). Other durations

5 between the radiation dose and administration of the compound that result in the enhanced the killing of cancer cells may also be suitable. In one embodiment, one or more of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered before the radiation exposure (such as before a dose or dose fraction), and the remaining one or more of the pentaaza  
10 macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent can be administered after the exposure (such as after a dose or dose fraction). One or more of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may also be administered both before and after administration of a radiation exposure.

15 [00228] In one embodiment, a course of radiation therapy includes a plurality of radiation doses or dose fractions given over a predetermined period of time, such as over the course of hours, weeks, days and even months, with the plural doses or dose fractions being either of the same magnitude or varying. That is, course of radiation therapy can comprise the administration of a series of multiple doses or dose fractions  
20 of radiation. In one embodiment, the active agent and pentaaza macrocyclic ring complex can be administered before one or more radiation doses or dose fractions in the series, such as before each radiation dose, or before some fraction of the radiation doses. Furthermore, the administration of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent during the course of  
25 radiation therapy can be selected to enhance the cancer treating effects of the radiation therapy, such as by sensitizing cancer cells to the radiation therapy. In one embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered within a predetermined duration before or after each dose or dose fraction, such as the predetermined duration  
30 discussed above. In another embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered within a predetermined duration of time before or after only select doses or dose fractions. In yet another embodiment, at least one of the pentaaza macrocyclic ring

complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered within a predetermined duration of time before the doses or dose fractions, while another of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered within a predetermined duration of time after the doses or dose fractions. In a further embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered only within the predetermined duration before or after select doses or dose fractions, while another of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered only within the predetermined duration before or after doses or dose fractions other than the select doses or dose fractions.

[00229] A suitable overall dose to provide during a course of therapy can be determined according to the type of treatment to be provided, the physical characteristics of the patient and other factors, and the dose fractions that are to be provided can be similarly determined. In one embodiment, a dose fraction of radiation that is administered to a patient may be at least 1.8 Gy, such as at least 2 Gy, and even at least 3 Gy, such as at least 5 Gy, and even at least 6 Gy. In yet another embodiment, a dose fraction of radiation that is administered to a patient may be at least 10 Gy, such as at least 12 Gy, and even at least 15 Gy, such as at least 18 Gy, and even at least 20 Gy, such as at least 24 Gy. In general, a dose fraction of radiation administered to a patient will not exceed 54 Gy. Furthermore, it should be noted that, in one embodiment, a dose fraction delivered to a subject may refer to an amount delivered to a specific target region of a subject, such as a target region of a tumor, whereas other regions of the tumor or surrounding tissue may be exposed to more or less radiation than that specified by the nominal dose fraction amount.

[00230] In yet another embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered as a part of a course of therapy that includes chemotherapy. In chemotherapy, chemotherapeutic agents are administered to a patient to kill or control the growth of cancerous cells. A typical course of chemotherapy may include one or a plurality of doses of one or more chemotherapeutic agents, which can be administered over the course of days, weeks and even months. Chemotherapeutic agents can include at least one of: alkylating antineoplastic agents such as nitrogen mustards (e.g.

cyclophosphamide, chlorambucil), nitrosoureas (e.g. n-nitroso-n-methylurea, carmustine, semustine), tetrazines (e.g. dacarbazine, mitozolimide), aziridines (e.g. thiotepa, mytomyacin), cisplatins (e.g. cisplatin, carboplatin, oxaliplatin); anti-metabolites such as anti-folates (e.g. methotrexate and pemetrexed), fluoropyrimidines (e.g.,  
5 fluorouracil, capecitabine), deoxynucleoside analogs (e.g. cytarabine, gemcitabine, decitabine) and thiopurines (e.g., thioguanine, mercaptopurine); anti microtubule agents such as taxanes (e.g. paclitaxel, docetaxel); and topoisomerase inhibitors (e.g. etoposide, doxorubicin, mitoxantrone, teniposide). For example, the chemotherapeutic agent may be selected from the group consisting of all-trans retinoic acid, azacitidine,  
10 azathioprine, bleomycin, carboplatin, capecitabine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, gemcitabine, hydroxyurea, idarubicin, imatinib, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, teniposide, tioguanine, valrubicin, vinblastine, vincristine,  
15 vindesine, and vinorelbine. The administration of many of the chemotherapeutic agents is described in the "Physicians' Desk Reference" (PDR), e.g., 1996 edition (Medical Economics Company, Montvale, N.J. 07645-1742, USA).

[00231] In one embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin compound and glutathione depleting agent are administered within  
20 a predetermined time period before or after a dose of a chemotherapeutic agent is administered. For example, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered within 1 week, 48 hours, 24 hours, 12 hours, 6, hours, 2 hours, 1 hour or even within 30 minutes of the patient receiving the dose of chemotherapeutic agent (either before or after the  
25 dose of chemotherapeutic agent). Other durations between the chemotherapeutic agent dose and administration of the compound that result in the enhanced the killing of cancer cells may also be suitable. In one embodiment, one or more of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may be administered before the dose of the chemotherapeutic agent, and the  
30 remaining one or more of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent can be administered after the dose of the chemotherapeutic agent. One or more of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent may also be

administered both before and after administration of the dose of chemotherapeutic agent.

[00232] In one embodiment, a course of chemotherapy includes a plurality of doses of a chemotherapeutic agent given over a predetermined period of time, such as over the course of hours, weeks, days and even months. The plural doses may be either of the same magnitude or varying, and can include doses of the same or different chemotherapeutic agents and/or a combination of chemotherapeutic agents. The administration of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent during the course of chemotherapy can be selected to enhance the cancer treating effects of the chemotherapy, such as by increasing intracellular levels of hydrogen peroxide in the cancer cells. In one embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered within a predetermined duration before or after each dose, such as the predetermined duration discussed above. In another embodiment, the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent are administered within a predetermined duration of time before or after only select doses. In yet another embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered within a predetermined duration of time before the doses, while another of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered within a predetermined duration of time after the doses. In a further embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered only within the predetermined duration before or after select doses, while another of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered only within the predetermined duration before or after doses other than the select doses.

[00233] In yet another embodiment, at least one of the pentaaza macrocyclic ring complex, thioredoxin reductase inhibitor and/or glutathione depleting agent is administered in combination with both a radiation therapy and chemotherapy.

## Examples

[00234] The following non-limiting examples are provided to further illustrate the present invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent approaches the inventors have found function well in the practice of the invention, and thus can be considered to constitute examples of modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

### **Example 1**

[00235] Steady-state levels of pro-oxidants in human breast (MB231) and lung cancer (H1299) cells were investigated versus normal breast (HMEC) and lung cancer epithelial cells (HBEpC), to determine whether the cancerous cells exhibited a higher level of the pro-oxidants. The fluorescent probe 5-(and 6) carboxy-2',7' dichlorodihydrofluorescein diacetate (DCFH2) was used as a non-specific oxidation sensitive fluorescence probe. Cells were grown in culture for 48 hours, and stained with oxidation sensitive dyes at 4% O<sub>2</sub>, to approximate physiologically relevant O<sub>2</sub> concentrations. The fluorescence intensity of 10,000 cells/sample were analyzed using a Becton Dickinson FACScan system. The results are shown in FIG. 1, and show that the cancer cells MB231 and H1299 have approximately 3-5 times more DCFH2 oxidation as compared to the normal HMEC or HBEpC cells. The results demonstrated that the cancer cells have higher steady-state levels of pro-oxidants at baseline than normal cells, and thus show a mechanism that can be exploited for cancer-cell selective treatment.

### **Example 2**

[00236] The effects on H<sub>2</sub>O<sub>2</sub> fluxes of a pentaaza macrocyclic ring complex (GC4419) provided in combination with inhibitors of H<sub>2</sub>O<sub>2</sub> metabolism were tested in human lung cancer cells (H292). The H292 cells were treated with 20 μM GC4419 for 3 and 24 hours, and PeroxyOrange-1 (PO-1) oxidation was monitored as a measure of

steady-state  $\text{H}_2\text{O}_2$ . The results are shown in FIG. 2. The results demonstrate that a 1.5-fold increase in PO-1 oxidation is exhibited at both 3 hours (4419 3) and 24 hours (4419 24) after exposure to the 20  $\mu\text{M}$  GC4419, as compared to the control. The results also show that depletion of glutathione by treating the cells with 100  $\mu\text{M}$  buthionine sulfoximine (BSO) while inhibiting catalase using 3-aminotriazole (AT) (a compound that irreversibly inhibits catalase enzyme and thus inhibits the metabolism of  $\text{H}_2\text{O}_2$  independently of glutathione) increases PO-1 oxidation 2-fold. Also, the combination of GC4419 + BSO + AT for 3 hours (B-AT 4419 3) and 24 hours (B-AT 4419 24) further increased PO-1 oxidation to 3-fold greater than a control. The PO-1 oxidation was also monitored in dosing the cells three times with 100  $\mu\text{M}$  hydrogen peroxide ( $\text{H}_2\text{O}_2$ ). Accordingly, the results show that the pentaaza macrocyclic ring complex GC4419 could increase intracellular steady-state levels of  $\text{H}_2\text{O}_2$ , with the increase being further enhanced by compounds that inhibit  $\text{H}_2\text{O}_2$  metabolism pathways.

### Example 3

[00237] The effect of the pentaaza macrocyclic ring complex GC4419 in increasing levels of  $\text{H}_2\text{O}_2$  in cancer cells was further tested by measuring the 3-aminotriazole mediated inactivation of catalase, using a method as described in Ahmad et al, *J. Biol Chem*, 280(6), 4254-4263 (2005). Briefly, head and neck cancer cells (Cal27), lung cancer cells (H1299) and breast cancer cells (MB231) were grown in 60 mm dishes, transduced with 50 MOI adCat to genetically increase catalase expression, and allowed 48 hours to recover so that the activity of catalase could increase to maximal levels. Afterwards, 50 mM AT was added for 5 minutes, which irreversibly binds to and inhibits the catalase intermediate formed in the presence of  $\text{H}_2\text{O}_2$ . 20  $\mu\text{M}$  GC4419 was added, and cells were collected at 10 minute intervals for 60 minutes, and the inhibition of catalase was determined. A Lowry's protein assay determined the total protein concentration of individual samples. This method provides a stoichiometry of one catalase molecule being inhibited for each molecule of  $\text{H}_2\text{O}_2$ , thereby providing for precise measurements of changes in  $\text{H}_2\text{O}_2$  flux. The results are shown in FIG. 3 (left hand side is Cal27, middle is H1299, and right hand side is MB231). The results demonstrated significant increases in steady-state levels of intracellular  $\text{H}_2\text{O}_2$  flux in both the lung and breast cancer cell lines when treated with GC4419, as normalized to

the control. Thus, GC4419 has been demonstrated to significantly increase the H<sub>2</sub>O<sub>2</sub> flux in cancer cells.

#### **Example 4**

5            [00238] The effect of the pentaaza macrocyclic ring complex GC4419 in combination with inhibitors of hydrogen peroxide metabolism was tested on breast cancer cells (MB231) and non-small cell lung cancer cells (H1299) versus normal cells (HMEC normal mammary epithelial cells and HBEpC normal bronchial epithelial cells). The cells were plated and grown exponentially for 2 days. Either or both of 20 µM of  
10   the CG4419 compound and 100 µM BSO were added at time 0 of exposure, and 500 nM Auranofin (a thioredoxin reductase inhibitor) was added 48 hours after, for 15 minutes prior to a clonogenic assay. The results are shown in FIG. 4. The results showed that the combination of GC4419 with auranofin (Au) and/or BSO produced synergistic effects in the selective killing of cancer cells as compared to any of the  
15   compounds alone. In particular, the combination of GC4419 with auranofin provides significantly improved selective killing of cancer cells as compared to GC4419 alone. The combination of all of GC4419, BSO and AU provided the greatest killing of cancer cells, and the greatest selectivity in killing of cancer cells versus normal cells. Thus, the results demonstrate the effectiveness of inhibiting hydrogen peroxide detoxification  
20   pathways in the selective killing of cancer cells using GC4419.

#### **Example 5**

             [00239] To determine whether the synergistic effects of the pentaaza macrocyclic ring complex (GC4419) with hydrogen peroxide metabolism inhibitors (BSO  
25   and/or Au) was related to oxidative stress, tests on cancer cell clonogenic killing was performed in the presence and absence of the thiol antioxidant N-acetylcysteine (NAC). Breast cancer cells (MB231) and lung cancer cells (H1299) were grown in culture to 75% confluence and treated with either 20 µM GC4419, or 100 µM BSO with 500 nM Au added after 48 hours, or all of GC4419, BSO and Au. The cancer cells were also  
30   treated with the same sets of compounds, but with the further addition of 10 mM of NAC. The results are shown in FIG. 5. Significantly, the addition of NAC nearly completely inhibited the cell killing caused by GC4419 + Au and/or BSO. Thus, the

results support that pentaaza macrocyclic ring complexes such as GC4419 combined with inhibitors of hydrogen peroxide metabolism can selectively kill cancer cells versus normal cells via oxidative stress correlated with increase  $H_2O_2$  fluxes.

## 5            **Example 6**

[00240] The effects of an inhibitor of hydrogen peroxide metabolism, auranofin (a thioredoxin reductase inhibitor), on cancer cells was tested in a clonogenic cell survival assay to determine the effects of the compound alone and in combination with ionizing radiation. Non-small cell lung cancer cells (H1299) were exponentially grown and treated with auranofin (Au) either by itself for 15 minutes, or 15 minutes prior to the first of two ionizing radiation (IR) doses (2 Gy each) spaced 24 hours apart. After the last ionizing radiation dose, the cells were trypsinized, counted, replated at varying lower densities and incubated for 12 days. Plates were stained and counted for clonogenic survival assays. The results are shown in FIG. 6. The results show that treatment with 250 nM of Au (Auranofin 250 nM) resulted in significant cell killing, as compared to a control. Furthermore, treatment with auranofin sensitized the cancer cells to radiation, thereby increasing the efficiency of the ionizing radiation in cell killing (Auranofin + IR) as compared to administration of ionizing radiation alone (IR 4Gy). Accordingly, the results show that auranofin significantly increases the response of cancer cells to ionizing radiation.

[00241] Furthermore, comparing the results in FIG. 6 with those of FIG. 4 above, it can be seen that the combination of the thioredoxin reductase inhibitor auranofin with the pentaaza macrocyclic ring complex GC4419 (GC4419 + Au in FIG. 4) provide synergistic effects in the selective killing of cancer cells, as compared to GC4419 alone (GC4419 in FIG. 4) or auranofin alone (Auranofin 250 nM in FIG. 6). In particular, while 20  $\mu$ M GC4419 in FIG. 4 results in a normalized surviving cell fraction (on a log scale) of H1299 cancer cells of about 1 (with the control also being about 1), and 250 nM of auranofin in FIG. 6 results in a normalized surviving cell fraction of H1299 cancer cells of a little over 0.5, the combination of 20  $\mu$ M GC4419 with 500 nM of auranofin in FIG. 6 results in a normalized survival fraction (on a log scale) of H1299 cancer cells of less than 0.2, meaning that the combination of GC4419 and auranofin

provides a greater than additive effect in the selective killing of cancer cells over either the GC4419 or auranofin alone.

### **Example 7**

5            [00242] The effect of buthionine sulfoximine (BSO) on reduced glutathione (GSH) levels in breast cancer cells (MB231) and lung cancer cells (H292) was examined. The cells were plated and grown to 75% confluency, then treated with either 50  $\mu$ M BSO or 100  $\mu$ M BSO for another 24 hours before harvesting for analysis for total glutathione levels using a DTNB recycling spectrophotometric assay. The results are  
10 shown in FIG. 7, and demonstrate that BSO significantly decreases glutathione levels in the cancer cells.

### **Example 8**

             [00243] The effects of the pentaaza macrocyclic ring complex GC4419 alone,  
15 and in combination with the thioredoxin reductase inhibitor auranofin (Au) and the glutathione depleting agent BSO, were tested in an *in vivo* model of breast cancer. MB231 breast cancer xenografts were grown in flanks of nude mice, and once tumors reached 2-4 mm diameter, the mice were treated with either BSO at 225 mg/kg i.p and Au at 1.5 mg/kg i.p. daily (Au + BSO), GC4419 at 9 mg/kg i.p. daily (GC4419), or  
20 GC4419 given daily in combination with BSO and Au (GC4419 + AU + BSO). Tumor growth averages were measured. The results are shown in FIG. 8. The results demonstrate that the combined therapy with GC4419 and Au + BSO resulted in a dramatic decrease in tumor volume as compared to GC4419 alone, or Au + BSO alone. The results thus show the synergistic effects of the combination of the pentaaza  
25 macrocyclic ring complex GC4419 with the thioredoxin reductase inhibitor auranofin and the glutathione-depleting agent BSO in an *in vivo* therapy regimen.

### **Example 9**

             [00244] The effect of the pentaaza macrocyclic ring complex GC4419 was  
30 tested *in vivo* on non-small cell lung cancer (NSCLC) in a tumor xenograft model. Subcutaneous H1299 tumors were subjected to biologically equivalent fractionation

schemes of 18 Gy X 1 dose or 2 Gy X 16 doses (delivered on consecutive days), 1 hour following an initial injection of 24 mg/kg GC4419. GC4419 was also administered on the next 4 consecutive days after the first injection (given, in the case of the 2 Gy fraction dose experiment, 1 hour prior to each of those 4 days' radiation treatments).

- 5 The results are shown in FIG. 9. The data demonstrate that GC4419 exhibits an anti-cancer effect in reduced cancer size as well as improved survival as compared to the control. The data also shows that GC4419 increases the response of NSCLC tumors to radiation, significantly enhancing radiation induced growth delay, and survival.


#### 10 **Example 10**

- [00245] The effect of GC4419 was tested *in vivo* on animal models of squamous cell carcinoma of the head and neck (HNSCC). GC4419 was delivered as a 24 mg/kg daily dose delivered as an ip bolus on 5 consecutive days starting 1 hour prior to irradiation. Radiation was delivered as a single dose of 12 Gy using a 10 mm  
15 collimated beam aimed at the center of the tumor. The results are shown in FIG. 10. The data shows that GC4419 treatment induces a significant growth delay to HN5 (HNSCC) tumors *in vivo*. Not only does the administration of GC4419 result in reduced tumor volumes as compared to a control, but the combination of GC4419 with radiation provides even greater growth delays, indicating that the GC4419 increases the  
20 response of the cancerous cells to the ionizing radiation.

#### **Example 11**

- [00246] In this example, *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex M40401 (GC4401) were conducted. Cancerous cell  
25 lines were incubated with M40401 (GC4401) for 6 days, at 5 different dosages from  $1 \times 10^{-8}$  to  $1 \times 10^{-4}$  M. Anti-proliferative effects were observed.

[00247] The structure of M40401 (GC4401) is below:




[00248] The compound was tested *in vitro* against cancers including leukemia (including lines CCRF-CEM, HB-60(TB), K-562, MOLT-4, RPMI-8226, SR), non-small cell lung cancer (including lines A549/ATCC, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H322M, NCI-H460, NCI-H522), colon cancer (including lines COLO 205, HCC-2998, HCT-116, HCT-15, HJ29, KM12, SW-620), CNS cancer (including lines SF 268, SF-295, SF-539, SNB-19, SNB-75, U251), melanoma (including lines MALME-3M, M14, SK-MEL-2, SK-MEL-28, SK-MEL-5, UACC-257, ACC-62), ovarian cancer (IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3), renal cancer (including lines 786-0, A498, ACHN, CAKI-J, RXF 393, SN12C, TK 10, UO-31), prostate cancer (including lines PC-3, DU-145) and breast cancer (including lines MCF7, NCVADR-RES, MDA-M8-231/ATCC, HS 578T, MDA-MB-435, MDA-N, BT-549, T-47D). FIGS. 12-20 show the anti-proliferative effects of the compound in the various different cell lines.

### Example 12

[00249] In this example, *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex M40403 (G4403) were conducted. Cancerous cell lines were incubated with M40403 (GC4403) for 6 days, at 5 different dosages from  $1 \times 10^{-8}$  to  $1 \times 10^{-4}$  M. Anti-proliferative effects were observed.

[00250] The structure of M40403 (GC4403) is below:




[00251] The compound was tested *in vitro* against cancers including leukemia (including lines CCRF-CEM, HB-60(TB), K-562, MOLT-4, RPMI-8226, SR), non-small cell lung cancer (including lines A549/ATCC, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H322M, NCI-H460, NCI-H522), colon cancer (including lines COLO 205, HCC-2998, HCT-116, HCT-15, HJ29, KM12, SW-620), CNS cancer (including lines SF 268, SF-295, SF-539, SNB-19, SNB-75, U251), melanoma (including lines MALME-3M, M14, SK-MEL-2, SK-MEL-28, SK-MEL-5, UACC-257, ACC-62), ovarian cancer (IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3), renal cancer (including lines 786-0, A498, ACHN, CAKI-J, RXF 393, SN12C, TK 10, UO-31), prostate cancer (including lines PC-3, DU-145) and breast cancer (including lines MCF7, NCVADR-RES, MDA-M8-231/ATCC, HS 578T, MDA-MB-435, MDA-N, BT-549, T-47D). FIGS. 21-29 show the anti-proliferative effects of the compound in the various different cell lines.

### Example 13

[00252] In this example, *in vitro* screening for the anti-proliferative effects of the pentaaza macrocyclic ring complex M40432 (GC4432) were conducted. Cancerous cell lines were incubated with M40432 (G4432) for 6 days, at 5 different dosages from  $1 \times 10^{-8}$  to  $1 \times 10^{-4}$  M. Anti-proliferative effects were observed.

[00253] The structure of M40432 (GC4432) is below:

[00254] GC4432:




[00255] The compound was tested *in vitro* against cancers including leukemia (including lines CCRF-CEM, HB-60(TB), K-562, MOLT-4, RPMI-8226, SR), non-small cell lung cancer (including lines A549/ATCC, EKVX, HOP-62, HOP-92, NCI-H226, NCI-H23, NCI-H322M, NCI-H460, NCI-H522), colon cancer (including lines COLO 205, HCC-2998, HCT-116, HCT-15, HJ29, KM12, SW-620), CNS cancer (including lines SF 268, SF-295, SF-539, SNB-19, SNB-75, U251), melanoma (including lines MALME-3M, M14, SK-MEL-2, SK-MEL-28, SK-MEL-5, UACC-257, ACC-62), ovarian cancer (IGROV1, OVCAR-3, OVCAR-4, OVCAR-5, OVCAR-8, SK-OV-3), renal cancer (including lines 786-0, A498, ACHN, CAKI-J, RXF 393, SN12C, TK 10, UO-31), prostate cancer (including lines PC-3, DU-145) and breast cancer (including lines MCF7, NCVADR-RES, MDA-M8-231/ATCC, HS 578T, MDA-MB-435, MDA-N, BT-549, T-47D). FIGS. 30-38 show the anti-proliferative effects of the compound in the various different cell lines.

[00256] Embodiments according to aspects of the disclosure are provided below, although the disclosure is not limited thereto.

[00257] Embodiment 1. A method of treating a cancer in a mammalian subject afflicted with the cancer, the method comprising:

[00258] administering to the subject at least one active agent comprising one or more of a thioredoxin reductase inhibitor and a glutathione depleting agent; and

[00259] administering to the subject a pentaaza macrocyclic ring complex corresponding to formula (I) below:



[00260] (I)

[00261] wherein

[00262] M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

[00263]  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are

5 independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

10 [00264] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

[00265] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or  
15 heterocycle having 3 to 20 ring carbon atoms;

[00266] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic  
20 heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and  $R_1$  and  $R_{10}$  attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

[00267] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00268] Z is a counterion;

5 [00269] n is an integer from 0 to 3; and


[00270] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

[00271] Embodiment 2. A method in accordance with Embodiment 1, wherein  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are each hydrogen.


10 [00272] Embodiment 3. A method in accordance with any preceding Embodiment, wherein W is an unsubstituted pyridine moiety.

[00273] Embodiment 4. A method in accordance with any preceding Embodiment, wherein U and V are trans-cyclohexanyl fused rings.


15 [00274] Embodiment 5. A method in accordance with any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is a complex selected from the group consisting of the complexes of formula (2)-(7) below:




2 (4459)




3 (4444)




4 (4462)



5 (4401)



6 (4419)



7 (4403)

[00275] Embodiment 6. A method in accordance with any preceding Embodiment, wherein X and Y, are independently selected from the group consisting of fluoro, chloro, bromo and iodo anions.


- 5 [00276] Embodiment 7. A method in accordance with any preceding Embodiment, wherein X and Y correspond to  $-O-C(O)-X_1$ , where each  $X_1$  is  $-C(X_2)(X_3)(X_4)$ , and each  $-C(X_2)(X_3)(X_4)$  corresponds to any of combinations 1 to 7 appearing in the following table:

| Combination | $X_2$           | $X_3$                                                                                        | $X_4$ |
|-------------|-----------------|----------------------------------------------------------------------------------------------|-------|
| 1           | Ph              | H                                                                                            | H     |
| 2           | Ph              | OH                                                                                           | H     |
| 3           | Ph              | NH <sub>2</sub>                                                                              | H     |
| 4           | Ph              | $\begin{array}{c} =O \\ (X_3 \text{ and } X_4 \text{ in} \\ \text{combination}) \end{array}$ |       |
| 5           | Ph              | CH <sub>3</sub>                                                                              | H     |
| 6           | CH <sub>3</sub> | H                                                                                            | H     |
| 7           | CH <sub>3</sub> | OH                                                                                           | H     |


10

[00277] Embodiment 8. A method in accordance with any preceding Embodiment, wherein the pentaaza macrocyclic ring complex is at least one of the following:

[00278]




(4403)



(4419)


5



(4432)

10

15



(4409)

20

[00286] Embodiment 9. A method in accordance with any preceding Embodiment, wherein the at least one active agent is one that increases intracellular

H<sub>2</sub>O<sub>2</sub> levels in cancerous cells, when provided to such cancerous cells *in vitro* in combination with the pentaaza macrocyclic ring complex.

[00287] Embodiment 10. A method in accordance with any preceding Embodiment, wherein the combination of the at least one active agent and pentaaza macrocyclic ring complex selectively kills cancer cells over normal cells when administered for the cancer treatment.

[00288] Embodiment 11. A method in accordance with any preceding Embodiment, wherein the thioredoxin reductase inhibitor is at least one of auranofin, auro-thio-glucose, chloro(triethylphosphine)gold(I), aurothiomalate, gold sodium thiomalate, sodium aurothiosulfate, gold acetate, 1,2,5-selenadiazole and derivatives thereof, metal complexes with 2-acetylpyridine-N(4)-orthochlorophenylthiosemicarbazone and/or a pharmaceutically acceptable salt thereof.

[00289] Embodiment 12. A method in accordance with Embodiment 10, wherein the thioredoxin reductase inhibitor is auranofin.

[00290] Embodiment 13. A method in accordance with any preceding Embodiment, wherein the glutathione depleting agent is at least one of a glutathione synthesis inhibitor, an inhibitor of x<sub>c</sub><sup>-</sup> cysteine/glutamate antiporter, and a glutathione reductase inhibitor,

[00291] Embodiment 14. A method in accordance with Embodiment 13, wherein the glutathione-depleting agent is at least one of buthionine sulfoximine (BSO), sulfasalazine, piperlongumine, N-ethylmaleimide, N-pyrenylmaleimide, 2-AAPA, erastin, sorafenib, 1S,3R-RSL3, DPI19, DPI18, DPI17, DPI13, DPI12, DPI10 (ML210), DPI7(ML162), and altretamine, and/or pharmaceutically acceptable salts thereof

[00292] Embodiment 15. A method in accordance with Embodiment 14, wherein the glutathione-depleting agent is at least one of buthionine sulfoximine and sulfasalazine.

[00293] Embodiment 16. A method in accordance with any preceding Embodiment, comprising administering both a thioredoxin reductase inhibitor and a glutathione depleting agent in combination with the pentaaza macrocyclic ring complex.

[00294] Embodiment 17. A method in accordance with any preceding Embodiment, wherein the mammal is a human patient.

[00295] Embodiment 18. A method in accordance with any preceding Embodiment, wherein the cancer is selected from the group consisting of cancer of the breast, heart, lung, small intestine, colon, spleen, kidney, bladder, head and neck, ovary, prostate, brain, pancreas, skin, bone, bone marrow, blood, thymus, uterus, testicles, cervix, and liver.

[00296] Embodiment 19. A method in accordance with Embodiment 18, wherein the subject is afflicted with cancer selected from the group consisting of breast cancer, lung cancer, renal cell carcinoma, spindle cell carcinoma, colorectal cancer, and head and neck cancer.

[00297] Embodiment 20. A method in accordance with Embodiment 19, wherein the subject is afflicted with cancer selected from the group consisting of breast cancer and lung cancer.

[00298] Embodiment 21. A method in accordance with any preceding Embodiment, wherein the active agent is administered substantially simultaneously with the pentaaza macrocyclic ring complex.

[00299] Embodiment 22. A method in accordance with any preceding Embodiment, wherein the active agent is administered before the pentaaza macrocyclic ring complex.

[00300] Embodiment 23. A method in accordance with any preceding Embodiment, wherein the active agent is administered after the pentaaza macrocyclic ring complex.

[00301] Embodiment 24. A method in accordance with any preceding Embodiment, wherein the active agent and pentaaza macrocyclic ring complex are administered within 24 hours of each other.

[00302] Embodiment 25. A method in accordance with any preceding Embodiment, wherein the active agent and pentaaza macrocyclic ring complex are administered within 1 hour of each other.

[00303] Embodiment 26. A method in accordance with any preceding Embodiment, wherein the active agent and pentaaza macrocyclic ring complex are administered by an administration route selected from the group consisting of oral, parenteral, intravenous, intraarterial, subcutaneous, rectal, subcutaneous,

intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, intrasternal, topical, nasal, transdermal, intraocular, intravesical, intrathecal, enteral, pulmonary, intralymphatic, intracavitary, vaginal, transurethral, intradermal, aural, intramammary, buccal, orthotopic, intratracheal, intralesional, percutaneous, endoscopic, transmucosal, sublingual and intestinal administration.

[00304] Embodiment 27. A method in accordance with any preceding Embodiment, wherein the method further comprises administering a cancer therapy comprising at least one of a radiation therapy and a chemotherapeutic therapy to the subject.

[00305] Embodiment 28. A method in accordance with Embodiment 27, wherein the active agent and pentaaza macrocyclic ring complex are administered prior to the cancer therapy.

[00306] Embodiment 29. A method in accordance with any of Embodiments 27-28, wherein the active agent and pentaaza macrocyclic ring complex are administered within 24 hours of the cancer therapy.

[00307] Embodiment 30. A method in accordance with any of Embodiments 27-29, wherein the active agent and pentaaza macrocyclic ring complex are administered within 1 hour of the cancer therapy.

[00308] Embodiment 31. A method in accordance with Embodiment 27, wherein at least one of the active agent and pentaaza macrocyclic ring complex are administered after the cancer therapy.

[00309] Embodiment 32. A method in accordance with any of Embodiments 27-30, wherein the cancer therapy comprises the administration of a series of multiple doses of radiation, and wherein the active agent and pentaaza macrocyclic ring complex are administered before one or more radiation dose in the series.

[00310] Embodiment 33. A method in accordance with any of Embodiments 27-30, wherein the chemotherapy dose comprises administration of a chemotherapeutic agent selected from the group consisting of all-trans retinoic acid, azacitidine, azathioprine, bleomycin, carboplatin, capecitabine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, gemcitabine, hydroxyurea, idarubicin,

imatinib, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, teniposide, tiguane, valrubicin, vinblastine, vincristine, vindesine, and vinorelbine.


[00311] Embodiment 34. A method of treating a cancer in a mammalian  
5 subject afflicted with the cancer, the method comprising:

[00312] administering to the subject at least one active agent comprising at least one of a thioredoxin reductase inhibitor and a glutathione-depleting agent;

[00313] administering to the subject a pentaaza macrocyclic ring complex; and

[00314] administering to the subject a cancer therapy comprising at least one  
10 of radiation therapy and chemotherapy;

[00315] wherein the pentaaza macrocyclic ring complex corresponds to the formula (I) below:



[00316]

[00317] wherein

15 [00318] M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

[00319]  $R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ ,  
20 wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl;

[00320] U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

5 [00321] V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms;

10 [00322] W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and  $R_1$  and  $R_{10}$  attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent;

15 [00323] X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof;

[00324] Z is a counterion;


[00325] n is an integer from 0 to 3; and

20 [00326] the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

WHAT IS CLAIMED IS:

1. Use of at least one active agent comprising a thioredoxin reductase inhibitor, and a pentaaza macrocyclic ring complex in the manufacture of a medicament for treating a cancer in a mammalian subject afflicted with the cancer,

5 wherein the pentaaza macrocyclic ring complex corresponds to formula (I) below:



wherein

10 M is  $Mn^{2+}$  or  $Mn^{3+}$ ;

$R_1$ ,  $R_2$ ,  $R'_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R'_5$ ,  $R_6$ ,  $R'_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R'_9$ , and  $R_{10}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ ,  
15 wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or alkyl,

U, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms,

20 V, together with the adjacent carbon atoms of the macrocycle, forms a fused substituted or unsubstituted, saturated, partially saturated or unsaturated, cycle or heterocycle having 3 to 20 ring carbon atoms,

W, together with the nitrogen of the macrocycle and the carbon atoms of the macrocycle to which it is attached, forms an aromatic or alicyclic, substituted or

unsubstituted, saturated, partially saturated or unsaturated nitrogen-containing fused heterocycle having 2 to 20 ring carbon atoms, provided that when W is a fused aromatic heterocycle the hydrogen attached to the nitrogen which is both part of the heterocycle and the macrocycle and R<sub>1</sub> and R<sub>10</sub> attached to the carbon atoms which are both part of the heterocycle and the macrocycle are absent,

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof,

Z is a counterion,

n is an integer from 0 to 3, and


the dashed lines represent coordinating bonds between the nitrogen atoms of the macrocycle and the transition metal, manganese.

2. The use of claim 1, wherein R<sub>1</sub>, R<sub>2</sub>, R'<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R'<sub>5</sub>, R<sub>6</sub>, R'<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R'<sub>9</sub>, and R<sub>10</sub> are each hydrogen.

3. The use according to any preceding claim, wherein W is an unsubstituted pyridine moiety.

4. The use according to any preceding claim, wherein U and V are trans-cyclohexanyl fused rings.

5. The use according to any preceding claim, wherein the pentaaza macrocyclic ring complex is represented by formula (II):




5 wherein

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and


10  $R_A$ ,  $R_B$ ,  $R_C$ , and  $R_D$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or

15 alkyl.

6. The use according to any preceding claim, wherein the pentaaza macrocyclic ring complex is represented by formula (III) or formula (IV):



(III)



(IV)


5 wherein

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and


10  $R_A$ ,  $R_B$ ,  $R_C$ , and  $R_D$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting of  $-OR_{11}$ ,  $-NR_{11}R_{12}$ ,  $-COR_{11}$ ,  $-CO_2R_{11}$ ,  $-CONR_{11}R_{12}$ ,  $-SR_{11}$ ,  $-SOR_{11}$ ,  $-SO_2R_{11}$ ,  $-SO_2NR_{11}R_{12}$ ,  $-N(OR_{11})(R_{12})$ ,  $-P(O)(OR_{11})(OR_{12})$ ,  $-P(O)(OR_{11})(R_{12})$ , and  $-OP(O)(OR_{11})(OR_{12})$ , wherein  $R_{11}$  and  $R_{12}$  are independently hydrogen or

15 alkyl.


7. The use according to any preceding claim, wherein the pentaaza macrocyclic ring complex is a compound represented by a formula selected from the group consisting of the complexes of formula (V)-(XVI) below:




(V)




(VI)




(VII)







(VIII)




(IX)




(X)





(XV)



(XVI)

8. The use according to any preceding claim, wherein X and Y are independently selected from substituted or unsubstituted moieties of the group consisting of halide, oxo, aquo, hydroxo, alcohol, phenol, dioxygen, peroxo, hydroperoxo, alkylperoxo, arylperoxo, ammonia, alkylamino, arylamino, heterocycloalkyl amino, heterocycloaryl amino, amine oxides, hydrazine, alkyl hydrazine, aryl hydrazine, nitric oxide, cyanide, cyanate, thiocyanate, isocyanate, isothiocyanate, alkyl nitrile, aryl nitrile, alkyl isonitrile, aryl isonitrile, nitrate, nitrite, azido, alkyl sulfonic acid, aryl sulfonic acid, alkyl sulfoxide, aryl sulfoxide, alkyl aryl sulfoxide, alkyl sulfenic acid, aryl sulfenic acid, alkyl sulfinic acid, aryl sulfinic acid, alkyl thiol carboxylic acid, aryl thiol carboxylic acid, alkyl thiol thiocarboxylic acid, aryl thiol thiocarboxylic acid, alkyl carboxylic acid, aryl carboxylic acid, urea, alkyl urea, aryl urea, alkyl aryl urea, thiourea, alkyl thiourea, aryl thiourea, alkyl aryl thiourea, sulfate, sulfite, bisulfate, bisulfite, thiosulfate, thiosulfite, hydrosulfite, alkyl phosphine, aryl phosphine, alkyl phosphine oxide, aryl phosphine oxide, alkyl aryl phosphine oxide, alkyl phosphine sulfide, aryl phosphine sulfide, alkyl aryl phosphine sulfide, alkyl phosphonic acid, aryl phosphonic acid, alkyl phosphinic acid, aryl phosphinic acid, alkyl phosphinous acid, aryl phosphinous acid, phosphate,

thiophosphate, phosphite, pyrophosphite, triphosphate, hydrogen phosphate, dihydrogen phosphate, alkyl guanidino, aryl guanidino, alkyl aryl guanidino, alkyl carbamate, aryl carbamate, alkyl aryl carbamate, alkyl thiocarbamate, aryl thiocarbamate, alkylaryl thiocarbamate, alkyl dithiocarbamate, aryl dithiocarbamate, alkylaryl dithiocarbamate, bicarbonate, carbonate, perchlorate, chlorate, chlorite, hypochlorite, perbromate, bromate, bromite, hypobromite, tetrahalomanganate, tetrafluoroborate, hexafluoroantimonate, hypophosphite, iodate, periodate, metaborate, tetraaryl borate, tetra alkyl borate, tartrate, salicylate, succinate, citrate, ascorbate, saccharinate, amino acid, hydroxamic acid, thiotosylate, and anions of ion exchange resins, or the corresponding anions thereof;

or X and Y correspond to  $-O-C(O)-X_1$ , where each  $X_1$  is  $-C(X_2)(X_3)(X_4)$ , and each  $X_1$  is independently substituted or unsubstituted phenyl or  $-C(-X_2)(-X_3)(-X_4)$ ;

each  $X_2$  is independently substituted or unsubstituted phenyl, methyl, ethyl or propyl;

each  $X_3$  is independently hydrogen, hydroxyl, methyl, ethyl, propyl, amino,  $-X_5C(=O)R_{13}$  where  $X_5$  is NH or O, and  $R_{13}$  is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or  $-OR_{14}$ , where  $R_{14}$  is C1-C18 alkyl, substituted or unsubstituted aryl or C1-C18 aralkyl, or together with  $X_4$  is  $(=O)$ ; and

each  $X_4$  is independently hydrogen or together with  $X_3$  is  $(=O)$ ;

or X and Y are independently selected from the group consisting of charge-neutralizing anions which are derived from any monodentate or polydentate coordinating ligand and a ligand system and the corresponding anion thereof;

or X and Y are independently attached to one or more of  $R_1, R_2, R'_2, R_3, R_4, R_5, R'_5, R_6, R'_6, R_7, R_8, R_9, R'_9$ , and  $R_{10}$ .

9. The use according to any preceding claim, wherein X and Y are independently selected from the group consisting of fluoro, chloro, bromo, and iodo anions.


10. The use according to any one of claims 1-8, wherein X and Y are independently selected from the group consisting of alkyl carboxylates, aryl

carboxylates and arylalkyl carboxylates.

11. The use according to any one of claims 1-8, wherein X and Y are independently amino acids.


5

12. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




10

13. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




15


14. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is a compound represented by the formula:




5 15. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



16. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



17. The use according to any one of claims 1-8, wherein the pentaaza macrocyclic ring complex is represented by the formula:



4711

18. The use according to any preceding claim, wherein the at least one active agent is one that increases intracellular H<sub>2</sub>O<sub>2</sub> levels in cancerous cells, when provided to such cancerous cells *in vitro* in combination with the pentaaza macrocyclic ring complex.

19. The use according to any preceding claim, wherein the combination of the at least one active agent and pentaaza macrocyclic ring complex selectively kills cancer cells over normal cells when administered for the cancer treatment.

20. The use according to any preceding claim, wherein the thioredoxin reductase inhibitor is at least one of auranofin, auro-thio-glucose, chloro(triethylphosphine)gold(I), aurothiomalate, gold sodium thiomalate, sodium aurothiosulfate, gold acetate, 1,2,5-selenadiazole and derivatives thereof, metal complexes with 2-acetylpyridine-N(4)-orthochlorophenylthiosemicarbazone and/or a pharmaceutically acceptable salt thereof.

21. The use of claim 20, wherein the thioredoxin reductase inhibitor is auranofin.

22. The use according to claim 25, wherein the glutathione depleting agent is at least one of a glutathione synthesis inhibitor, an inhibitor of  $\gamma$ -cysteine/glutamate antiporter, and a glutathione reductase inhibitor.

5

23. The use of claim 22, wherein the glutathione-depleting agent is at least one of buthionine sulfoximine (BSO), sulfasalazine, piperlongumine, N-ethylmaleimide, N-pyrenylmaleimide, 2-AAPA, erastin, sorafenib, 1S,3R-RSL3, DPI19, DPI18, DPI17, DPI13, DPI12, DPI10 (ML210), DPI7(ML162), and altretamine, and/or pharmaceutically acceptable salts thereof.

10

24. The use of claim 23, wherein the glutathione-depleting agent is at least one of buthionine sulfoximine and sulfasalazine.

15

25. The use according to any preceding claim, comprising administering both a thioredoxin reductase inhibitor and a glutathione depleting agent in combination with the pentaaza macrocyclic ring complex.

20

26. The use according to any preceding claim, wherein the mammal is a human patient.

25

27. The use according to any preceding claim, wherein the cancer is selected from the group consisting of cancer of the breast, heart, lung, small intestine, colon, spleen, kidney, bladder, head and neck, ovary, prostate, brain, pancreas, skin, bone, bone marrow, blood, thymus, uterus, testicles, cervix, and liver.

30

28. The use of claim 27, wherein the subject is afflicted with cancer selected from the group consisting of breast cancer, lung cancer, renal cell carcinoma, spindle cell carcinoma, colorectal cancer, and head and neck cancer.

29. The use of claim 28, wherein the subject is afflicted with cancer selected from the group consisting of breast cancer and lung cancer.

30. The use according to any preceding claim, wherein the active agent is

administered substantially simultaneously with the pentaaza macrocyclic ring complex.

31. The use according to any of claims 1-29, wherein the active agent is administered before the pentaaza macrocyclic ring complex.

5

32. The use according to any of claims 1-29, wherein the active agent is administered after the pentaaza macrocyclic ring complex.

33. The use according to any preceding claim, wherein the active agent and pentaaza macrocyclic ring complex are administered within 24 hours of each other.

10

34. The use according to any preceding claim, wherein the active agent and pentaaza macrocyclic ring complex are administered within 1 hour of each other.

15

35. The use according to any preceding claim, wherein the active agent and pentaaza macrocyclic ring complex are administered by an administration route selected from the group consisting of oral, parenteral, intravenous, intraarterial, subcutaneous, rectal, subcutaneous, intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, intrasternal, topical, nasal, transdermal, intraocular, intravesical, intrathecal, enteral, pulmonary, intralymphatic, intracavitary, vaginal, transurethral, intradermal, aural, intramammary, buccal, orthotopic, intratracheal, intralesional, percutaneous, endoscopic, transmucosal, sublingual and intestinal administration.

20

25

36. The use according to any preceding claim, wherein the use further comprises use of at least one of a radiation therapy and a chemotherapeutic therapy in the manufacture of a medicament for a cancer therapy to the subject.

30

37. The use of claim 36, wherein the active agent and pentaaza macrocyclic ring complex are administered prior to the cancer therapy.

38. The use according to any of claims 36-37, wherein the active agent and pentaaza macrocyclic ring complex are administered within 24 hours of the cancer therapy.

39. The use according to any of claims 36-38, wherein the active agent and pentaaza macrocyclic ring complex are administered within 1 hour of the cancer therapy.

5


40. The use of claim 36, wherein at least one of the active agent and pentaaza macrocyclic ring complex are administered after the cancer therapy.

41. The use according to any of claims 36-40, wherein the cancer therapy comprises the administration of a series of multiple doses of radiation, and wherein the active agent and pentaaza macrocyclic ring complex are administered before one or more radiation dose in the series.

42. The use according to any of claims 36-40, wherein the chemotherapy dose comprises administration of a chemotherapeutic agent selected from the group consisting of all-trans retinoic acid, azacitidine, azathioprine, bleomycin, carboplatin, capecitabine, cisplatin, chlorambucil, cyclophosphamide, cytarabine, daunorubicin, docetaxel, doxifluridine, doxorubicin, epirubicin, epothilone, etoposide, fluorouracil, gemcitabine, hydroxyurea, idarubicin, imatinib, mechlorethamine, mercaptopurine, methotrexate, mitoxantrone, oxaliplatin, paclitaxel, pemetrexed, teniposide, tiguane, valrubicin, vinblastine, vincristine, vindesine, and vinorelbine.

43. The use according to any of claims 1-42, wherein the pentaaza macrocyclic ring complex is a compound which can be represented by the following formulas:

a) formula (II):




wherein

X and Y represent suitable ligands which are derived from any monodentate or  
 5 polydentate coordinating ligand or ligand system or the corresponding anion  
 thereof; and


R<sub>A</sub>, R<sub>B</sub>, R<sub>C</sub>, and R<sub>D</sub> are independently hydrogen, hydrocarbyl, substituted  
 hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected  
 from the group consisting

10 of -OR<sub>11</sub>, -NR<sub>11</sub>R<sub>12</sub>, -COR<sub>11</sub>, -CO<sub>2</sub>R<sub>11</sub>, -CONR<sub>11</sub>R<sub>12</sub>, -SR<sub>11</sub>, -SOR<sub>11</sub>, -SO<sub>2</sub>R<sub>11</sub>, -SO  
 2NR<sub>11</sub>R<sub>12</sub>, -N(OR<sub>11</sub>)(R<sub>12</sub>), -P(O)(OR<sub>11</sub>)(OR<sub>12</sub>), -P(O)(OR<sub>11</sub>)(R<sub>12</sub>),  
 and -OP(O)(OR<sub>11</sub>)(OR<sub>12</sub>), wherein R<sub>11</sub> and R<sub>12</sub> are independently hydrogen or  
 alkyl; or

15 b) formula (III) or formula (IV):



(III)



(IV)

wherein

X and Y represent suitable ligands which are derived from any monodentate or polydentate coordinating ligand or ligand system or the corresponding anion thereof; and

RA, RB, RC, and RD are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, heterocyclyl, an amino acid side chain moiety, or a moiety selected from the group consisting

of -OR<sub>11</sub>, -NR<sub>11</sub>R<sub>12</sub>, -COR<sub>11</sub>, -CO<sub>2</sub>R<sub>11</sub>, -CONR<sub>11</sub>R<sub>12</sub>, -SR<sub>11</sub>, -SOR<sub>11</sub>, -SO<sub>2</sub>R<sub>11</sub>, -SO<sub>2</sub>NR<sub>11</sub>R<sub>12</sub>, -N(OR<sub>11</sub>)(R<sub>12</sub>), -P(O)(OR<sub>11</sub>)(OR<sub>12</sub>), -P(O)(OR<sub>11</sub>)(R<sub>12</sub>), and -OP(O)(OR<sub>11</sub>)(OR<sub>12</sub>), wherein R<sub>11</sub> and R<sub>12</sub> are independently hydrogen or alkyl.