
The claims defining the invention are as follows

1. Use of a compound of Formula (I), or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment of a wound in a subject:




2. The use according to claim 1, wherein the wound is an incision or an excision.

3. The use according to claim 1 or claim 2, wherein the medicament is adapted to be administered topically to the site of the wound.

4. The use according to any one of claims 1 to 3, wherein the medicament is in the form of a balm, a cream, a gel, a liniment, a lotion, an ointment, a paste, a rub or a salve.

5. A pharmaceutical composition for use in treating a wound in a subject, comprising from 0.1 to 2% (w/w) of a compound of Formula (I), or a pharmaceutically acceptable salt thereof, and from 99.9 to 98% (w/w) of a carrier:



6. The pharmaceutical composition according to claim 5, wherein the compound, or a pharmaceutically acceptable salt thereof is present in the range of 0.1 to 1% (w/w), and the carrier is present in the range 99.9 to 99% (w/w).

7. The pharmaceutical composition according to claim 6, wherein the compound, or a pharmaceutically acceptable salt thereof is present at about 0.5% (w/w), and the carrier is present at about 99.5% (w/w).

8. The pharmaceutical composition according to any one of claims 5 to 7, wherein the composition comprises 0.1 to 2% (w/w) of a compound of Formula (I), or a pharmaceutically acceptable salt thereof; one or more emollients at about 5 to about 10% (w/w); one or more surfactants at about 10 to about 15% (w/w); one or more humectants at about 1 to about 10% (w/w); one or more preservatives at about 1 to about 5% (w/w); one or more antioxidants at about 0.1 to about 2% (w/w); and a solvent making the remaining balance up to 100% (w/w).

9. The pharmaceutical composition according to any one of claims 8, wherein the one or more humectants is glycerine.

10. The pharmaceutical composition according to any one of claims 8 and 9, wherein the one or more preservatives is selected from the group consisting of phenoxy ethanol, potassium sorbate and sodium benzoate.

11. The pharmaceutical composition according to any one of claims 8 to 10, wherein the one or more antioxidants is vitamin E.

12. The pharmaceutical composition according to claim 5 wherein the composition comprises 0.5% (w/w) of a compound of Formula (I), or a pharmaceutically acceptable salt thereof; heavy liquid paraffin - 2.5% (w/w); glyceryl mono stearate - 1.85% (w/w);stearic acid - 2.0% (w/w); kokum butter - 2.0% (w/w); polyethylene glycol - 5.0% (w/w); glycerine - 5.0% (w/w); cetostearyl alcohol - 2.5% (w/w); cetyl alcohol - 2.0% (w/w); ethylene glycol monostearate - 1.0% (w/w); phenoxy ethanol - 0.5% (w/w); potassium sorbate - 0.5% (w/w); sodium benzoate - 0.5% (w/w); vitamin E - 0.5% (w/w); and purified water - 73.65% (w/w).