

4-Aryl-N-phenyl-1,3,5-triazin-2-amines containing a sulfoximine group

The present invention relates to 4-aryl-N-phenyl-1,3,5-triazin-2-amines containing a sulfoximine group of general formula (I) or (Ia) as described and defined herein, and methods for their preparation, their use for the treatment and/or prophylaxis of disorders, in particular of hyper-proliferative disorders and/or virally induced infectious diseases and/or of cardiovascular diseases. The invention further relates to intermediate compounds useful in the preparation of said compounds of general formula (I) or (Ia).

The family of cyclin-dependent kinase (CDK) proteins consists of members that are key regulators of the cell division cycle (cell cycle CDK's), that are involved in regulation of gene transcription (transcriptional CDK's), and of members with other functions. CDKs require for activation the association with a regulatory cyclin subunit. The cell cycle CDKs CDK1/cyclin B, CDK2/cyclin A, CDK2/cyclinE, CDK4/cyclinD, and CDK6/cyclinD get activated in a sequential order to drive a cell into and through the cell division cycle. The transcriptional CDKs CDK9/cyclin T and CDK7/cyclin H regulate the activity of RNAPolymerase II via phosphorylation of the carboxy-terminal domain (CTD). Positive transcription factor b (P-TEFb) is a heterodimer of CDK9 and one of four cyclin partners, cyclin T1, cyclin K, cyclin T2a or T2b.

Whereas CDK9 (NCBI GenBank Gene ID 1025) is exclusively involved in transcriptional regulation, CDK7 in addition participates in cell cycle regulation as CDK-activating kinase (CAK).

Transcription of genes by RNA polymerase II is initiated by assembly of the pre-initiation complex at the promoter region and phosphorylation of Ser 5 and Ser 7 of the CTD by CDK7/cyclin H. For a major fraction of genes RNA polymerase II stops mRNA transcription after it moved 20-40 nucleotides along the DNA template. This promoter-proximal pausing of RNA polymerase II is mediated by negative elongation factors and is recognized as a major control mechanism to regulate expression of rapidly induced genes in response to a variety of stimuli (Cho et al., Cell Cycle **2010**, 9, 1697). P-TEFb is crucially involved in overcoming promoter-proximal pausing of RNA polymerase II and transition into a productive elongation state by phosphorylation of Ser 2 of the CTD as well as by phosphorylation and inactivation of negative elongation factors.

Activity of P-TEFb itself is regulated by several mechanisms. About half of cellular P-TEFb exists in an inactive complex with 7SK small nuclear RNA (7SK snRNA), La-related protein 7 (LARP7/PIP7S) and hexamethylene bis-acetamide inducible proteins 1/2 (HEXIM1/2, He et al., Mol. Cell **2008**, 29, 588). The remaining half of P-TEFb exists in an active complex containing the bromodomain protein Brd4 (Yang et al., Mol. Cell **2005**, 19, 535). Brd4 recruits P-TEFb through interaction with acetylated histones to chromatin areas primed for gene transcription. Through alternately interacting with its positive and negative regulators, P-TEFb is maintained in a functional equilibrium: P-TEFb bound to the 7SK snRNA complex represents a reservoir from which active P-TEFb can be released on demand of cellular transcription and cell proliferation (Zhou & Yik, Microbiol. Mol. Biol. Rev. **2006**, 70, 646).

Furthermore, the activity of P-TEFb is regulated by posttranslational modifications including phosphorylation/de-phosphorylation, ubiquitination, and acetylation (reviewed in Cho et al., *Cell Cycle* **2010**, 9, 1697).

- 5 Deregulated activity of CDK9 kinase activity of the P-TEFb heterodimer is associated with a variety of human pathological settings such as hyper-proliferative diseases (e.g. cancer), virally induced infectious diseases or cardiovascular diseases.

10 Cancer is regarded as a hyper-proliferative disorder mediated by a disbalance of proliferation and cell death (apoptosis). High levels of anti-apoptotic Bcl-2-family proteins are found in various human tumors and account for prolonged survival of tumor cells and therapy resistance. Inhibition of P-TEFb kinase activity was shown to reduce transcriptional activity of RNA polymerase II leading to a decline of short-lived anti-apoptotic proteins, especially Mcl-1 and XIAP, reinstalling the ability of tumor cells to undergo apoptosis. A number of other proteins associated with the transformed tumor phenotype (such as
15 Myc, NF-kB responsive gene transcripts, mitotic kinases) are either short-lived proteins or are encoded by short-lived transcripts which are sensitive to reduced RNA polymerase II activity mediated by P-TEFb inhibition (reviewed in Wang & Fischer, *Trends Pharmacol. Sci.* **2008**, 29, 302).

20 Many viruses rely on the transcriptional machinery of the host cell for the transcription of their own genome. In case of HIV-1 RNA polymerase II gets recruited to the promoter region within the viral LTR's. The viral transcription activator (Tat) protein binds to nascent viral transcripts and overcomes promoter-proximal RNA polymerase II pausing by recruitment of P-TEFb which in turn promotes transcriptional elongation. Furthermore, the Tat protein increases the fraction of active P-TEFb by replacement of the P-TEFb inhibitory proteins HEXIM1/2 within the 7SK snRNA complex. Recent data
25 have shown that inhibition of the kinase activity of P-TEFb is sufficient to block HIV-1 replication at kinase inhibitor concentrations that are not cytotoxic to the host cells (reviewed in Wang & Fischer, *Trends Pharmacol. Sci.* **2008**, 29, 302). Similarly, recruitment of P-TEFb by viral proteins has been reported for other viruses such as B-cell cancer-associated Epstein-Barr virus, where the nuclear antigen EBNA2 protein interacts with P-TEFb (Bark-Jones et al., *Oncogene* **2006**, 25, 1775), and the human T-
30 lymphotropic virus type 1 (HTLV-1), where the transcriptional activator Tax recruits P-TEFb (Zhou et al., *J. Virol.* **2006**, 80, 4781).

Cardiac hypertrophy, the heart's adaptive response to mechanical overload and pressure (hemodynamic stress e.g. hypertension, myocardial infarction), can lead, on a long term, to heart failure and death.
35 Cardiac hypertrophy was shown to be associated with increased transcriptional activity and RNA polymerase II CTD phosphorylation in cardiac muscle cells. P-TEFb was found to be activated by dissociation from the inactive 7SK snRNA/HEXIM1/2 complex. These findings suggest

pharmacological inhibition of P-TEFb kinase activity as a therapeutic approach to treat cardiac hypertrophy (reviewed in Dey et al., Cell Cycle **2007**, 6, 1856).

5 In summary, multiple lines of evidence suggest that selective inhibition of the CDK9 kinase activity of the P-TEFb heterodimer (= CDK9 and one of four cyclin partners, cyclin T1, cyclin K, cyclin T2a or T2b) represents an innovative approach for the treatment of diseases such as cancer, viral diseases, and/or diseases of the heart. CDK9 belongs to a family of at least 13 closely related kinases of which the subgroup of the cell cycle CDK's fulfills multiple roles in regulation of cell proliferation. Thus, co-inhibition of cell cycle CDK's (e.g. CDK1/cyclin B, CDK2/cyclin A, CDK2/cyclinE, CDK4/cyclinD,
10 CDK6/cyclinD) and of CDK9 is expected to impact normal proliferating tissues such as intestinal mucosa, lymphatic and hematopoietic organs, and reproductive organs. To maximize the therapeutic margin of CDK9 kinase inhibitors, molecules with high selectivity towards CDK9 are therefore required.

CDK inhibitors in general as well as CDK9 inhibitors are described in a number of different publications:

15 WO2008129070 and WO2008129071 both describe 2,4 disubstituted aminopyrimidines as CDK inhibitors in general. It is also asserted that some of these compounds may act as selective CDK9 inhibitors (WO2008129070) and as CDK5 inhibitors (WO2008129071), respectively, but no specific CDK9 IC50 (WO2008129070) or CDK5 IC50 (WO200812971) data is presented.

WO2008129080 discloses 4,6 disubstituted aminopyrimidines and demonstrates that these compounds show
20 an inhibitory effect on the protein kinase activity of various protein kinases, such as CDK1, CDK2, CDK4, CDK5, CDK6 and CDK9, with a preference for CDK9 inhibition (example 80).

EP1218360 B1 describes triazin derivatives as kinase inhibitors, but does not disclose potent or selective CDK9 inhibitors.

WO2008079933 discloses aminopyridine and aminopyrimidine derivatives and their use as CDK1, CDK2,
25 CDK3, CDK4, CDK5, CDK6, CDK7, CDK8 or CDK9 inhibitors.

WO2011012661 describes aminopyridine derivatives useful as CDK inhibitors.

Wang et al. (Chemistry & Biology **2010**, 17, 1111-1121) describe 2-anilino-4-(thiazol-5-yl)pyrimidine
transcriptional CDK inhibitors, which show anticancer activity in animal models.

30

WO2004009562 discloses substituted triazine kinase inhibitors. For selected compounds CDK1 and CDK 4 test data, but no CDK9 data is presented.

WO2004072063 describes heteroaryl (pyrimidine, triazine) substituted pyrroles as inhibitors of protein kinases such as ERK2, GSK3, PKA or CDK2.

WO2010009155 discloses triazine and pyrimidine derivatives as inhibitors of histone deacetylase and/or cyclin dependent kinases (CDKs). For selected compounds CDK2 test data is described.

WO2003037346 (corresponding to US7618968B2, US7291616B2, US2008064700A1, US2003153570A1) relates to aryl triazines and uses thereof, including to inhibit lysophosphatidic acid acyltransferase beta (LPAAT-beta) activity and/or proliferation of cells such as tumor cells.

WO2008025556 describes carbamoyl sulfoximides having a pyrimidine core, which are useful as kinase inhibitors. No CDK9 data is presented.

WO2002066481 describes pyrimidine derivatives as cyclin dependent kinase inhibitors. CDK9 is not mentioned and no CDK9 data is presented.

WO2008109943 concerns phenyl aminopyri(mi)dine compounds and their use as kinase inhibitors, in particular as JAK2 kinase inhibitors. The specific examples focus on compounds having a pyrimidine core.

WO2009032861 describes substituted pyrimidinyl amines as JNK kinase inhibitors. The specific examples focus on compounds having a pyrimidine core.

WO2011046970 concerns amino-pyrimidine compounds as inhibitors of TBKL and/or IKK epsilon. The specific examples focus on compounds having a pyrimidine core.

Despite the fact that various inhibitors of CDK's are known, there remains a need for selective CDK9 inhibitors to be used for the treatment of diseases such as hyper-proliferative diseases, viral diseases, and/or diseases of the heart, which offer one or more advantages over the compounds known from prior art, such as :

- improved activity and / or efficacy,
- beneficial kinase selectivity profile according to the respective therapeutic need,
- improved side effect profile, such as fewer undesired side effects,
- lower intensity of side effects, or reduced (cyto)toxicity,
- improved physicochemical properties, such as solubility in water and body fluids,
- improved pharmacokinetic properties, allowing e.g. for dose reduction or an easier dosing scheme.

A particular object of the invention is to provide CDK9 kinase inhibitors which, compared to the compounds known from prior art, show an increased solubility in water.

A particular object of the invention is to provide CDK9 kinase inhibitors which, compared to the compounds known from prior art, show an increased selectivity for CDK9/Cyclin T1 as compared to CDK2/Cyclin E.


Another object of the invention is to provide CDK9 kinase inhibitors which show an increased potency to inhibit CDK9 activity (demonstrated by a lower IC50 value for CDK9/Cyc T1) compared to the compounds known from prior art.

Another object of the invention is to provide CDK9 kinase inhibitors, which show an improved anti-proliferative activity in one or more of certain tumor cell lines such as for example HeLa or DU145 compared to the compounds known from prior art.


Another object of the invention is to provide CDK9 kinase inhibitors, which show an improved side effect profile such as a reduced carbonic anhydrase-1 and -2 inhibition compared to the compounds known from prior art.

Further, it is also an object of the present invention to provide CDK9 kinase inhibitors, which, compared to the compounds known from prior art, show an increased solubility in water and/or which are highly selective for CDK9/Cyclin T1 as compared to CDK2/Cyclin E, and/or which show an increased potency to inhibit CDK9 activity (demonstrated by a lower IC₅₀ value for CDK9/Cyc T1) and/or which show an improved anti-proliferative activity in one or more of certain tumor cell lines such as HeLa or DU145.

The present invention relates to compounds of general formula (I)



or of general formula (Ia)




, wherein

R¹ represents a group selected from C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocyclyl-, phenyl, heteroraryl, phenyl-C₁-C₃-alkyl- or heteroaryl-C₁-C₃-alkyl-,

wherein said group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of hydroxy, cyano, halogen, halo-C₁-C₃-alkyl-, C₁-C₆-alkoxy-, C₁-C₃-fluoroalkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines;

R² represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, C_1 - C_6 -alkyl-, C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl, heteroaryl

wherein said C_1 - C_6 -alkyl, C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, amino, alkylamino-, dialkylamino-, acetilamino-, N-methyl-N-acetyl-amino-, cyclic amines, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;

R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen,

hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-, C₂-C₃-alkenyl-, C₂-C₃-alkynyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl, heteroaryl, wherein said C₃-C₇-cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, cyano, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-;

b) a C₃-C₇-cycloalkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-, C₂-C₃-alkenyl-, C₂-C₃-alkynyl-;

c) a heterocycyl-group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-, C₂-C₃-alkenyl-, C₂-C₃-alkynyl-;

d) a phenyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

e) a heteroaryl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

f) a phenyl-C₁-C₃-alkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

g) a heteroaryl-C₁-C₃-alkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

R⁹ represents a group selected from C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl, benzyl or heteroaryl wherein said group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-;

R¹⁰, R¹¹ represent, independently from each other, a group selected from hydrogen, C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl or heteroaryl wherein said C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-;

or their salts, solvates or salts of solvates.

Compounds according to the invention are the compounds of the formula (I) or (Ia) or (Ib) and the salts, solvates and solvates of the salts thereof, the compounds of the hereinafter recited formula which are encompassed by formula (I) or (Ia) or (Ib) and the salts, solvates and solvates of the salts thereof, and the compounds which are encompassed by formula (I) or (Ia) or (Ib) and which are mentioned hereinafter as exemplary embodiments and the salts, solvates and solvates of the salts thereof, where the compounds which are encompassed by formula (I) or (Ia) or (Ib) and are mentioned hereinafter are not already salts, solvates and solvates of the salts.

The compounds according to the invention may, depending on their structure, exist in stereoisomeric forms (enantiomers, diastereomers). The invention therefore relates to the enantiomers or diastereomers and respective mixtures thereof. The stereoisomerically pure constituents can be isolated in a known manner from such mixtures of enantiomers and/or diastereomers.

If the compounds according to the invention can be in tautomeric forms, the present invention encompasses all tautomeric forms.

Further, the compounds of the present invention can exist in free form, e.g. as a free base, or as a free acid, or as a zwitterion, or can exist in the form of a salt. Said salt may be any salt, either an organic or inorganic addition salt, particularly any physiologically acceptable organic or inorganic addition salt, customarily used in pharmacy.

Salts which are preferred for the purposes of the present invention are physiologically acceptable salts of the compounds according to the invention. However, salts which are not suitable for pharmaceutical applications per se, but which, for example, can be used for the isolation or purification of the compounds according to the invention, are also comprised.

The term "physiologically acceptable salt" refers to a relatively non-toxic, inorganic or organic acid addition salt of a compound of the present invention, for example, see S. M. Berge, et al. "Pharmaceutical Salts," J. Pharm. Sci. **1977**, 66, 1-19.

5 Physiologically acceptable salts of the compounds according to the invention encompass acid addition salts of mineral acids, carboxylic acids and sulfonic acids, for example salts of hydrochloric acid, hydrobromic acid, hydroiodic, sulfuric acid, bisulfuric acid, phosphoric acid, nitric acid or with an organic acid, such as formic, acetic, acetoacetic, pyruvic, trifluoroacetic, propionic, butyric, hexanoic, heptanoic, undecanoic, lauric, benzoic, salicylic, 2-(4-hydroxybenzoyl)-benzoic, camphoric, cinnamic, cyclopentanepropionic, digluconic, 3-hydroxy-2-naphthoic, nicotinic, pamoic, pectinic, persulfuric, 3-phenylpropionic, picric, pivalic, 2-hydroxyethanesulfonate, itaconic, sulfamic, trifluoromethanesulfonic, dodecylsulfuric, ethansulfonic, benzenesulfonic, para-toluenesulfonic, methansulfonic, 2-naphthalenesulfonic, naphthalenedisulfonic, camphorsulfonic acid, citric, tartaric, stearic, lactic, oxalic, malonic, succinic, malic, adipic, alginic, maleic, fumaric, D-gluconic, mandelic, ascorbic, glucoheptanoic, glycerophosphoric, aspartic, sulfosalicylic, hemisulfuric, or thiocyanic acid, for example.

Physiologically acceptable salts of the compounds according to the invention also comprise salts of conventional bases, such as, by way of example and by preference, alkali metal salts (for example sodium and potassium salts), alkaline earth metal salts (for example calcium and magnesium salts) and ammonium salts derived from ammonia or organic amines with 1 to 16 C atoms, such as, by way of example and by preference, ethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, dimethylaminoethanol, procaine, dibenzylamine, N-methylmorpholine, arginine, lysine, ethylenediamine, N-methylpiperidine, N-methylglucamine, dimethylglucamine, ethylglucamine, 1,6-hexadamine, glucosamine, sarcosine, serinol, tris(hydroxymethyl)aminomethane, aminopropanediol, Sovak base, and 1-amino-2,3,4-butanetriol. Additionally, basic nitrogen containing groups may be quaternised with such agents as lower alkylhalides such as methyl-, ethyl-, propyl-, and butylchlorides, -bromides and -iodides ; dialkylsulfates like dimethyl-, diethyl-, dibutyl- and diamylsulfates, long chain halides such as decyl-, lauryl-, myristyl- and stearylchlorides, -bromides and -iodides, aralkylhalides like benzyl- and phenethylbromides and others.

30 The present invention includes all possible salts of the compounds of the present invention as single salts, or as any mixture of said salts, in any ratio.

Solvates is the term used for the purposes of the invention for those forms of the compounds according to the invention which form a complex with solvent molecules by coordination in the solid or liquid state. Hydrates are a special form of solvates in which the coordination takes place with water. Hydrates are preferred as solvates within the scope of the present invention.

The invention also includes all suitable isotopic variations of a compound of the invention. An isotopic variation of a compound of the invention is defined as one in which at least one atom is replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually or predominantly found in nature. Examples of isotopes that can be incorporated into a compound of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulfur, fluorine, chlorine, bromine and iodine, such as ^2H (deuterium), ^3H (tritium), ^{13}C , ^{14}C , ^{15}N , ^{17}O , ^{18}O , ^{32}P , ^{33}P , ^{33}S , ^{34}S , ^{35}S , ^{36}S , ^{18}F , ^{36}Cl , ^{82}Br , ^{123}I , ^{124}I , ^{129}I and ^{131}I , respectively. Certain isotopic variations of a compound of the invention, for example, those in which one or more radioactive isotopes such as ^3H or ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution studies. Tritiated and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased in vivo half-life or reduced dosage requirements and hence may be preferred in some circumstances. Isotopic variations of a compound of the invention can generally be prepared by conventional procedures known by a person skilled in the art such as by the illustrative methods or by the preparations described in the examples hereafter using appropriate isotopic variations of suitable reagents.

In addition, the present invention also encompasses prodrugs of the compounds according to the invention. The term "prodrugs" encompasses compounds which themselves may be biologically active or inactive, but are converted (for example by metabolism or hydrolysis) to compounds according to the invention during their residence time in the body.

For the purposes of the present invention, the substituents have the following meaning, unless otherwise specified:

The term "halogen", "halogen atom" or "halo" represents fluorine, chlorine, bromine and iodine, particularly chlorine or fluorine, preferably fluorine atoms.

The term "alkyl" represents a linear or branched alkyl radical having the number of carbon atoms specifically indicated, e.g. C_1 - C_{10} one, two, three, four, five, six, seven, eight, nine or ten carbon atoms, e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, hexyl, heptyl, octyl, nonyl-, decyl-, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, 1,2-dimethylpropyl, neo-pentyl, 1,1-dimethylpropyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 2-ethylbutyl, 1-ethylbutyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 2,3-dimethylbutyl, 1,3-dimethylbutyl, or 1,2-dimethylbutyl. If the number of carbon atoms is not specifically indicated the term "alkyl" represents a linear or branched alkyl radical having, as a rule, 1 to 9, particularly 1 to 6, preferably 1 to 4 carbon atoms. Particularly, the alkyl group has 1, 2, 3, 4, 5 or 6 carbon atoms ("C₁-C₆-alkyl"), e.g. methyl, ethyl, n-propyl-, isopropyl, n-butyl, tert-butyl, pentyl, isopentyl, hexyl, 2-methylbutyl, 1-methylbutyl, 1-ethylpropyl, 1,2-dimethylpropyl, neo-pentyl, 1,1-dimethylpropyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 2-ethylbutyl, 1-ethylbutyl, 3,3-

dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 2,3-dimethylbutyl, 1,3-dimethylbutyl, or 1,2-dimethylbutyl. Preferably, the alkyl group has 1, 2 or 3 carbon atoms ("C₁-C₃-alkyl"), methyl, ethyl, n-propyl or isopropyl.

5 The term "C₂-C₃-alkenyl" is to be understood as preferably meaning a linear or branched, monovalent hydrocarbon group, which contains one double bond, and which has 2 or 3 carbon atoms ("C₂-C₃-alkenyl"). Said alkenyl group is, for example, a vinyl, allyl, (E)-2-methylvinyl, (Z)-2-methylvinyl or isopropenyl group.

10 The term "C₂-C₃-alkynyl" is to be understood as preferably meaning a linear, monovalent hydrocarbon group which contains one triple bond, and which contains 2 or 3 carbon atoms. Said C₂-C₃-alkynyl group is, for example, ethynyl, prop-1-ynyl or prop-2-ynyl group.

15 The term "C₃-C₇-cycloalkyl" is to be understood as preferably meaning a saturated, monovalent, monocyclic hydrocarbon ring which contains 3, 4, 5, 6 or 7 carbon atoms. Said C₃-C₇-cycloalkyl group is for example, a monocyclic hydrocarbon ring, e.g. a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl group. Said cycloalkyl ring can optionally contain one or more double bonds e.g. cycloalkenyl, such as a cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl or cycloheptenyl group, wherein the bond between said ring with the rest of the molecule may be to any carbon atom of said ring, be it saturated or unsaturated. Preferably the "C₃-C₇-cycloalkyl" is a cyclopropyl group.

20 The term "heterocyclyl" is to be understood as meaning a saturated or partially unsaturated, monovalent, mono- or bicyclic hydrocarbon ring which contains 3, 4, 5, 6, 7, 8 or 9 carbon atoms and further containing 1, 2 or 3 heteroatom-containing groups selected from oxygen, sulfur, nitrogen. Particularly, the term "heterocyclyl" is to be understood as meaning a "4- to 10-membered heterocyclic ring".

25 The term a "4- to 10-membered heterocyclic ring" is to be understood as meaning a saturated or partially unsaturated, monovalent, mono- or bicyclic hydrocarbon ring which contains 3, 4, 5, 6, 7, 8 or 9 carbon atoms, and further containing 1, 2 or 3 heteroatom-containing groups selected from oxygen, sulfur, nitrogen. Said heterocyclic ring is for example, a monocyclic heterocyclic ring such as an oxetanyl, azetidiny, tetrahydrofuranyl, pyrrolidiny, 1,3-dioxolanyl, imidazolidiny, pyrazolidiny, oxazolidiny, isoxazolidiny, 1,4-dioxanyl, pyrroliny, tetrahydropyranyl, piperidiny, morpholiny, 1,3-dithianyl, thiomorpholiny, piperaziny, or chinuclidiny group. Optionally, said heterocyclic ring can contain one or more double bonds, e.g. 4H-pyranyl, 2H-pyranyl, 2,5-dihydro-1H-pyrrolyl, 1,3-dioxolyl, 4H-1,3,4-thiadiaziny, 2,5-dihydrofuranyl, 2,3-dihydrofuranyl, 2,5-dihydrothienyl, 2,3-dihydrothienyl, 4,5-dihydrooxazolyl, 4,5-dihydroisoxazolyl, or 4H-1,4-thiaziny group, or, it may be benzo fused.

30

35 Particularly, the term "heterocyclyl" is to be understood as being a heterocyclic ring which contains 3, 4 or 5 carbon atoms, and 1, 2 or 3 of the above-mentioned heteroatom-containing groups (a "4- to 7-membered heterocyclic ring"), more particularly said ring can contain 4 or 5 carbon atoms, and 1, 2 or 3 of the above-mentioned heteroatom-containing groups (a "5- to 7-membered heterocyclic ring"), more

particularly said heterocyclic ring is a "6-membered heterocyclic ring", which is to be understood as containing 4 carbon atoms and 2 of the above-mentioned heteroatom-containing groups or 5 carbon atoms and one of the above-mentioned heteroatom-containing groups, preferably 4 carbon atoms and 2 of the above-mentioned heteroatom-containing groups.

5 The term "C₁-C₆-alkoxy-" is to be understood as preferably meaning a linear or branched, saturated, monovalent, hydrocarbon group of formula -O-alkyl, in which the term "alkyl" is defined supra, e.g. a methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, tert-butoxy, sec-butoxy, pentyloxy, iso-pentyloxy, n-hexyloxy group, or an isomer thereof. Particularly, the "C₁-C₆-alkoxy-" group is a "C₁-C₄-alkoxy-", a "C₁-C₃-alkoxy-", a methoxy, ethoxy, or propoxy group, preferably a methoxy, ethoxy or
10 propoxy group.

The term „C₁-C₃-fluoroalkoxy-“ is to be understood as preferably meaning a linear or branched, saturated, monovalent, C₁-C₃-alkoxy- group, as defined supra, in which one or more of the hydrogen atoms is replaced, identically or differently, by one or more fluoro atoms. Said C₁-C₃-fluoroalkoxy-
15 group is, for example a 1,1-difluoromethoxy-, a 1,1,1-trifluoromethoxy-, a 2-fluoroethoxy-, a 3-fluoropropoxy-, a 2,2,2-trifluoroethoxy-, a 3,3,3-trifluoropropoxy- particularly a "C₁-C₂-fluoroalkoxy-" group.

The term „alkylamino-“ is to be understood as preferably meaning an alkylamino group with a linear or branched alkyl group as defined supra. (C₁-C₃)-alkylamino- for example means a monoalkylamino group with 1, 2 oder 3 carbon atoms, (C₁-C₆)-alkylamino- with 1, 2, 3, 4, 5 or 6 carbon atoms. The term
20 "alkylamino-" comprises for example methylamino-, ethylamino-, n-propylamino-, isopropylamino-, tert.-butylamino-, n-pentylamino- or n-hexylamino-.

The term „dialkylamino-“ is to be understood as preferably meaning an alkylamino group having two linear or branched alkyl groups as defined supra, which are independent from each other. (C₁-C₃)-dialkylamino- for example represents a dialkylamino group with two alkyl groups each of them having 1
25 to 3 carbon atoms per alkyl group. The term "dialkylamino-" comprises for example: N,N-Dimethylamino-, N,N-Diethylamino-, N-Ethyl-N-methylamino-, N-Methyl-N-n-propylamino-, N-Isopropyl-N-n-propylamino-, N-t-Butyl-N-methylamino-, N-Ethyl-N-n-pentylamino- und N-n-Hexyl-N-methylamino-.

The term "cyclic amine" is to be understood as preferably meaning a cyclic amine group. Suitable cyclic
30 amines are especially azetidine, pyrrolidine, piperidine, piperazine, 1-methylpiperazine, morpholine, thiomorpholine, which could be optionally substituted by one or two methyl groups.

The term "halo-C₁-C₃-alkyl-" is to be understood as preferably meaning a linear or branched, saturated, monovalent hydrocarbon group in which the term "C₁-C₃-alkyl" is defined supra, and in which one or more hydrogen atoms is replaced by a halogen atom, identically or differently, i.e. one halogen atom
35 being independent from another. Particularly, said halogen atom is fluorine. Preferred is fluoro-C₁-C₃-

alkyl such as for example- CF_3 , $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_2\text{CF}_3$, or $-\text{CH}_2\text{CF}_3$. Said halo- $\text{C}_1\text{-C}_3$ -alkyl- group is, for example, a halo- $\text{C}_1\text{-C}_2$ -alkyl- group, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_2\text{CF}_3$, or $-\text{CH}_2\text{CF}_3$, preferably it is $-\text{CF}_3$.

The term “phenyl- $\text{C}_1\text{-C}_3$ -alkyl-” is to be understood as preferably meaning a phenyl group, in which one of the hydrogen atoms is replaced by a $\text{C}_1\text{-C}_3$ -alkyl group, as defined supra, that links the phenyl- $\text{C}_1\text{-C}_3$ -alkyl- group to the molecule. Particularly, the “phenyl- $\text{C}_1\text{-C}_3$ -alkyl-” is a phenyl- $\text{C}_1\text{-C}_2$ -alkyl-, preferably a benzyl- group.

The term “heteroaryl” is to be understood as preferably meaning a monovalent, aromatic, mono- or bicyclic aromatic ring system having 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 ring atoms (a “5- to 14-membered heteroaryl” group), particularly 5 (a “5-membered heteroaryl”) or 6 (a “6-membered heteroaryl”) or 9 (a “9-membered heteroaryl”) or 10 ring atoms (a “10-membered heteroaryl”), and which contains at least one heteroatom which may be identical or different, said heteroatom being such as oxygen, nitrogen or sulfur, and can be monocyclic, bicyclic, or tricyclic, and in addition in each case can be benzo-condensed. Particularly, heteroaryl is selected from thienyl, furanyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, tetrazolyl etc., and benzo derivatives thereof, such as, for example, benzofuranyl, benzothienyl, benzoxazolyl, benzisoxazolyl, benzimidazolyl, benzotriazolyl, indazolyl, indolyl, isoindolyl, etc.; or pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, etc., and benzo derivatives thereof, such as, for example, quinoliny, quinazoliny, isoquinoliny, etc.; or azociny, indoliziny, puriny, etc., and benzo derivatives thereof; or cinnoliny, phthalazinyl, quinazoliny, quinoxaliny, naphthyridiny, pteridiny, carbazolyl, acridiny, phenazinyl, phenothiazinyl, phenoxazinyl, xanthenyl, or oxepiny, 1,4-benzodioxanyl etc. Preferably, heteroaryl is selected from monocyclic heteroaryl, 5- membered heteroaryl or 6- membered heteroaryl.

The term “5-membered heteroaryl” is understood as preferably meaning a monovalent, aromatic, mono-aromatic ring system having 5 ring atoms and which contains at least one heteroatom which may be identical or different, said heteroatom being such as oxygen, nitrogen or sulfur. Particularly, “5 membered heteroaryl” is selected from thienyl, furanyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, tetrazolyl.

The term “6-membered heteroaryl” is understood as preferably meaning a monovalent, aromatic, mono-aromatic ring system having 6 ring atoms and which contains at least one heteroatom which may be identical or different, said heteroatom being such as oxygen, nitrogen or sulfur. Particularly, “6 membered heteroaryl” is selected from pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl.

The term “heteroaryl- $\text{C}_1\text{-C}_3$ -alkyl-” is to be understood as preferably meaning a heteroaryl, a 5-membered heteroaryl or a 6- membered heteroaryl group, each as defined supra, in which one of the hydrogen atoms is replaced by a $\text{C}_1\text{-C}_3$ -alkyl group, as defined supra, that links the heteroaryl- $\text{C}_1\text{-C}_3$ -alkyl- group to the molecule. Particularly, the “heteroaryl- $\text{C}_1\text{-C}_3$ -alkyl-” is a heteroaryl- $\text{C}_1\text{-C}_2$ -alkyl-, a pyridinyl- $\text{C}_1\text{-C}_3$ -alkyl-, a pyridinylmethyl-, a pyridinylethyl-, a pyridinylpropyl-, a pyrimidinyl- $\text{C}_1\text{-C}_3$ -


alkyl-, a pyrimidinylmethyl-, a pyrimidinylethyl-, a pyrimidinylpropyl-, preferably a pyridinylmethyl- or a pyridinylethyl- or a pyrimidinylethyl- or a pyrimidinylpropyl- group.

- The term “C₁-C₁₀”, as used throughout this text, e.g. in the context of the definition of “C₁-C₁₀-alkyl” is to be understood as meaning an alkyl group having a finite number of carbon atoms of 1 to 10, i.e. 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms. It is to be understood further that said term “C₁-C₁₀” is to be interpreted as any sub-range comprised therein, e.g. C₁-C₁₀, C₁-C₉, C₁-C₈, C₁-C₇, C₁-C₆, C₁-C₅, C₁-C₄, C₁-C₃, C₁-C₂, C₂-C₁₀, C₂-C₉, C₂-C₈, C₂-C₇, C₂-C₆, C₂-C₅, C₂-C₄, C₂-C₃, C₃-C₁₀, C₃-C₉, C₃-C₈, C₃-C₇, C₃-C₆, C₃-C₅, C₃-C₄, C₄-C₁₀, C₄-C₉, C₄-C₈, C₄-C₇, C₄-C₆, C₄-C₅, C₅-C₁₀, C₅-C₉, C₅-C₈, C₅-C₇, C₅-C₆, C₆-C₁₀, C₆-C₉, C₆-C₈, C₆-C₇, C₇-C₁₀, C₇-C₉, C₇-C₈, C₈-C₁₀, C₈-C₉, C₉-C₁₀.
- 10 Similarly, as used herein, the term “C₁-C₆”, as used throughout this text, e.g. in the context of the definition of “C₁-C₆-alkyl”, “C₁-C₆-alkoxy” is to be understood as meaning an alkyl group having a finite number of carbon atoms of 1 to 6, i.e. 1, 2, 3, 4, 5 or 6 carbon atoms. It is to be understood further that said term “C₁-C₆” is to be interpreted as any sub-range comprised therein, e.g. C₁-C₆, C₁-C₅, C₁-C₄, C₁-C₃, C₁-C₂, C₂-C₆, C₂-C₅, C₂-C₄, C₂-C₃, C₃-C₆, C₃-C₅, C₃-C₄, C₄-C₆, C₄-C₅, C₅-C₆.
- 15 Similarly, as used herein, the term “C₁-C₃”, as used throughout this text, e.g. in the context of the definition of “C₁-C₃-alkyl”, “C₁-C₃-alkoxy” or “C₁-C₃-fluoroalkoxy” is to be understood as meaning an alkyl group having a finite number of carbon atoms of 1 to 3, i.e. 1, 2 or 3 carbon atoms. It is to be understood further that said term “C₁-C₃” is to be interpreted as any sub-range comprised therein, e.g. C₁-C₃, C₁-C₂, C₂-C₃.
- 20 Further, as used herein, the term “C₃-C₇”, as used throughout this text, e.g. in the context of the definition of “C₃-C₇-cycloalkyl”, is to be understood as meaning a cycloalkyl group having a finite number of carbon atoms of 3 to 7, i.e. 3, 4, 5, 6 or 7 carbon atoms, particularly 3, 4, 5 or 6 carbon atoms. It is to be understood further that said term “C₃-C₇” is to be interpreted as any sub-range comprised therein, e.g. C₃-C₇, C₃-C₆, C₃-C₅, C₃-C₄, C₄-C₇, C₄-C₆, C₄-C₅, C₅-C₇, C₅-C₆, C₆-C₇.
- 25 A symbol  at a bond denotes the linkage site in the molecule.
- As used herein, the term “one or more times”, e.g. in the definition of the substituents of the compounds of the general formulae of the present invention, is understood as meaning one, two, three, four or five times, particularly one, two, three or four times, more particularly one, two or three times, even more particularly one or two times.
- 30 Where the plural form of the word compounds, salts, hydrates, solvates and the like, is used herein, this is taken to mean also a single compound, salt, isomer, hydrate, solvate or the like.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a group selected from C_1 - C_6 -alkyl-, C_3 - C_7 -cycloalkyl- or heterocyclyl-;

R^2 represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom or C_1 - C_3 -alkyl-;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, C_1 - C_6 -alkyl-;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

10 R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two or three substituents, selected from the group consisting of halogen, hydroxy, C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-, C_3 - C_7 -cycloalkyl-, heterocyclyl-, phenyl, heteroaryl,

wherein said C_3 - C_7 -cycloalkyl-, heterocyclyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-;

b) a phenyl group;

c) a phenyl- C_1 - C_3 -alkyl- group, which phenyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-;

20

d) a heteroaryl-C₁-C₃-alkyl- group, which heteroaryl group is optionally substituted with one substituent selected from the group consisting of halogen;

R⁹ represents a C₁-C₆-alkyl group;


R¹⁰, R¹¹ represent, independently from each other, a group selected from hydrogen or C₁-C₆-alkyl-;

5 or their salts, solvates or salts of solvates.

In a preferred embodiment the present invention relates to compounds of general formula (I) or (Ia), wherein

R¹ represents a C₁-C₆-alkyl- or C₃-C₇-cycloalkyl- group;

R² represents a group selected from



10 R³, R⁴ represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, C₁-C₃-alkyl-;

R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)R⁹, -C(O)OR⁹, -C(O)NR¹⁰R¹¹;

R⁶, R⁷ represent, independently from each other, a hydrogen atom or fluoro atom;

15 R⁸ represents a group selected from

a) a C₁-C₁₀-alkyl group, which is optionally substituted with one or two substituents, selected from the group consisting of hydroxy, C₂-C₃-alkenyl-, C₂-C₃-alkynyl-, C₃-C₇-cycloalkyl-, phenyl, heteroaryl, wherein said C₃-C₇-cycloalkyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, cyano, halo-C₁-C₃-alkyl-;

b) a phenyl-C₁-C₃-alkyl- group, which phenyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, cyano, halo-C₁-C₃-alkyl-;

c) a heteroaryl-C₁-C₃-alkyl- group, which heteroaryl group is optionally substituted with one substituent selected from the group consisting of halogen;

25


R^9 represents a C_1 - C_6 -alkyl group;

R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen or C_1 - C_6 -alkyl-; or their salts, solvates or salts of solvates.

In another preferred embodiment the present invention relates to compounds of general formula (I) or (Ia),
5 wherein

R^1 represents a C_1 - C_6 -alkyl- group;

R^2 represents a group selected from



R^3 represents a hydrogen atom;

R^4 represents a hydrogen atom;

10 R^5 represents a group selected from a hydrogen atom, cyano group or $-C(O)NR^{10}R^{11}$;


R^6 , R^7 represent, independently from each other, a hydrogen atom or fluoro atom;

R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one substituent selected from the group consisting of C_2 - C_3 -alkynyl;

15 R^{10} , R^{11} represent, independently from each other, a hydrogen atom or a C_1 - C_6 -alkyl- group; or their salts, solvates or salts of solvates.

In another preferred embodiment the present invention relates to compounds of general formula (Ib)




(Ib), wherein

- 5 R^1 represents a C_1 - C_6 -alkyl- group;
- R^3 represents a hydrogen or fluoro or chloro atom;
- R^4 represents a hydrogen or fluoro atom;
- R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$,
 10 $C(O)NR^{10}R^{11}$;
- R^8 represents a group selected from
- a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two substituents,
 selected from the group consisting of C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-, phenyl, heteroaryl,
 wherein said phenyl or heteroaryl group is optionally substituted with one, two or three
 15 substituents, identically or differently, selected from halogen, cyano, halo- C_1 - C_3 -alkyl-;
- b) a phenyl group;
- c) a phenyl- C_1 - C_3 -alkyl- group, which phenyl group is optionally substituted with one or
 two or three substituents, identically or differently, selected from the group consisting of
 halogen, cyano, halo- C_1 - C_3 -alkyl-;
- 20 d) a heteroaryl- C_1 - C_3 -alkyl- group, which heteroaryl group is optionally substituted with
 one substituent selected from the group consisting of halogen;
- R^9 represents a C_1 - C_6 -alkyl group;
- R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen or C_1 - C_6 -alkyl-;
 or their salts, solvates or salts of solvates.

25 The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a group selected from C_1 - C_6 -alkyl- or heterocyclyl-;

R^2 represents a group selected from



R^3 , R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

5 R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, C_1 - C_6 -alkyl-;

R^6 , R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

R^8 represents a group selected from

10 a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one substituent, selected from the group consisting of C_3 - C_7 -cycloalkyl-, heterocyclyl-, phenyl, wherein said C_3 - C_7 -cycloalkyl-, heterocyclyl- or phenyl group is optionally substituted with one, or two substituents, identically or differently, selected from halogen;

b) a phenyl group;

15 c) a phenyl- C_1 - C_3 -alkyl- group, which is optionally substituted with one or two substituents, identically or differently, selected from the group consisting of halogen;

R^9 represents a C_1 - C_6 -alkyl group;


R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen or C_1 - C_6 -alkyl-;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a C_1 - C_6 -alkyl- group;

R^2 represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom;

10 R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one substituent selected from the group consisting of C_3 - C_7 -cycloalkyl-, phenyl, wherein said C_3 - C_7 -cycloalkyl- or phenyl group is optionally substituted with one halogen atom;

b) a phenyl- C_1 - C_3 -alkyl- group, which is optionally substituted with one halogen atom;


15 R^9 represents a C_1 - C_6 -alkyl group;

R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen, C_1 - C_6 -alkyl-; or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a C_1 - C_6 -alkyl- group;

20 R^2 represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

5 R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^{10}R^{11}$;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom;

10 R^8 represents a group selected from
 a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one phenyl group, wherein said phenyl group is optionally substituted with one halogen atom;

b) a phenyl- C_1 - C_3 -alkyl- group, which is optionally substituted with one halogen atom;


R^9 represents a C_1 - C_6 -alkyl group;

R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen, C_1 - C_6 -alkyl-; or their salts, solvates or salts of solvates.

15 The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a C_1 - C_6 -alkyl- group;

R^2 represents a group selected from



R^3, R^4 represent a hydrogen atom;

R^5 represents a group selected from a hydrogen atom or $-C(O)NR^{10}R^{11}$;

R^6, R^7 represent, independently from each other, a hydrogen or a fluoro atom;

5 R^8 represents a C_1 - C_{10} -alkyl group;


R^{10}, R^{11} represent, independently from each other, a hydrogen atom or C_1 - C_6 -alkyl- group;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a C_1 - C_6 -alkyl-group;

10 R^2 represents a group selected from



R^3, R^4 represent a hydrogen atom;

R^5 represents a group selected from a hydrogen atom or $-C(O)NR^{10}R^{11}$;

R^6, R^7 represent, independently from each other, a hydrogen atom or a fluoro atom;

R^8 represents a C_1 - C_{10} -alkyl group;


15 R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen, C_1 - C_6 -alkyl-;

or their salts, solvates or salts of solvates.

In another embodiment the present invention concerns compounds of general formula (I) or (Ia), wherein

R¹ represents C₁-C₆-alkyl-,

R² represents a group selected from



R³, R⁴ represent a hydrogen atom;

5 R⁵ represents a group selected from a hydrogen atom or -C(O)OR⁹;

R⁶, R⁷ represent, independently from each other, a group selected from a hydrogen atom or a fluoro atom;

R⁸ represents a group selected from

a) a C₁-C₁₀-alkyl- group or

10 b) a phenyl-C₁-C₃-alkyl- group;


R⁹ represents a C₁-C₆-alkyl- group;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R¹ represents a group selected from methyl, ethyl, cyclopropyl, tetrahydro-2H-pyranyl;

15 R² represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom or methyl;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$ or C_1 - C_6 -alkyl-;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

R^8 represents a group selected from

10

a) a methyl, ethyl propyl or butyl group, which group is optionally substituted with one or two or three substituents, selected from the group consisting of halogen, hydroxy, ethenyl, propenyl, ethynyl, propynyl, cyclopentyl, cyclohexyl, tetrahydro-2H-pyranyl, phenyl, pyridinyl, thiazolyl, oxazolyl, wherein said phenyl or pyridinyl group is optionally substituted with one, two or three substituents, identically or differently, selected from a fluoro or chloro atom, cyano, methyl, or trifluoromethyl;

15

b) a (2H_2)methyl group substituted with a (2H_5)phenyl group;

c) a phenyl group;

d) a benzyl group, which phenyl ring is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of a fluoro atom, chloro atom, cyano, methyl or trifluoromethyl-;

20

e) a pyridin-2-ylmethyl- group, which pyridine is optionally substituted with one fluoro atom;

f) a pyridin-3-ylmethyl- group, which pyridine is optionally substituted with one fluoro atom;

g) a pyridin-4-ylmethyl- group, which pyridine is optionally substituted with one fluoro atom;

5 h) a thiazolylmethyl- group;

i) an oxazolylmethyl- group

R^9 represents a methyl or ethyl group;

R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen, methyl;


or their salts, solvates or salts of solvates.

10

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a group selected from methyl, tetrahydro-2H-pyranyl;

R^2 represents a group selected from



15 R^3 , R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, methyl;

R^6 , R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

20 R^8 represents a group selected from

a) a methyl group, which is optionally substituted with one substituent, selected from the group consisting of cyclohexyl, tetrahydro-2H-pyranyl, phenyl, wherein said phenyl group is optionally substituted with one, or two substituents, identically or differently, selected from a fluoro or chloro atom;

5 b) a ($^2\text{H}_2$)methyl group substituted with a ($^2\text{H}_5$)phenyl group;

c) an ethyl group, which group is optionally substituted with one substituent selected from the group consisting of tetrahydro-2H-pyranyl, cyclopentyl;

d) a phenyl group;

10 e) a phenyl- C_1 - C_3 -alkyl- group, which is optionally substituted with one or two substituents, identically or differently, selected from a fluoro or chloro atom;

R^9 represents a methyl or ethyl group;


R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen, methyl;

or their salts, solvates or salts of solvates.

15 The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a methyl or ethyl or cyclopropyl group;

R^2 represents a group selected from



R^3 , represents a group selected from a hydrogen atom, fluoro atom, chloro atom or methyl;

R^4 represents a group selected from a hydrogen atom or fluoro atom

20 R^5 represents a group selected from a hydrogen atom, cyano, $-\text{C}(\text{O})\text{R}^9$, $-\text{C}(\text{O})\text{OR}^9$, $-\text{C}(\text{O})\text{NR}^{10}\text{R}^{11}$;

R^6 , R^7 represent, independently from each other, a group selected from a hydrogen atom or fluoro atom;

R^8 represents a group selected from

methyl, ethyl, prop-2-en-1-yl, 2-methylprop-2-en-1-yl, 2-(hydroxymethyl)prop-2-en-1-yl, (2Z)-but-2-en-1-yl, prop-2-yn-1-yl, but-2-yn-1-yl, cyclohexylmethyl, benzyl, 3-cyanobenzyl, 3-fluorobenzyl, 3-chlorobenzyl, 4-fluorobenzyl, 4-chlorobenzyl, 3-fluoro-5-(trifluoromethyl)benzyl, 3-chloro-5-fluorobenzyl, pyridin-4-yl, 2-fluoropyridin-4-yl, 2,3,5-trifluorobenzyl, 3,4,5-trifluorobenzyl;

R^9 represents a methyl or ethyl group;


R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen, C_1 - C_6 -alkyl-;

or their salts, solvates or salts of solvates.

10 The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a group selected from methyl;

R^2 represents a group selected from



R^3 represents a hydrogen atom;

R^4 represents a hydrogen atom;

15 R^5 represents a hydrogen atom, a cyano group or $-C(O)NR^{10}R^{11}$;

R^6 , R^7 represent, independently from each other, a hydrogen atom or fluoro atom;

R^8 represents a group selected from

a) a methyl group, which is optionally substituted with one substituent selected from the group consisting of ethynyl, propynyl;

20 b) an ethyl group;

R^{10} represents a hydrogen atom;


R^{11} represents a methyl group;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a methyl group;

R^2 represents a group selected from



R^3, R^4 represent, independently from each other, a hydrogen, fluoro or chloro atom;

5 R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom;

R^8 represents a group selected from

10 a) methyl group, which is optionally substituted with one substituent, selected from the group consisting of cyclohexyl and phenyl, wherein said phenyl group is optionally substituted with one fluoro atom;

b) a benzyl group, which is optionally substituted with one fluoro atom;

R^9 represents a methyl or ethyl group;


15 R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen or methyl;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

R^1 represents a methyl group;

R^2 represents a group selected from




20 R^3, R^4 represent, independently from each other, a hydrogen, fluoro or chloro atom;

- R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^{10}R^{11}$;
- R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom;
- 5 R^8 represents a group selected from
- a) a methyl group, which is optionally substituted with one phenyl group, wherein said phenyl group is optionally substituted with one fluoro atom;
- b) a benzyl group, which is optionally substituted with one fluoro atom;
- R^9 represents a methyl or ethyl group;
- 10 R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen or methyl;
- or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein


- R^1 represents a methyl group;
- R^2 represents a group selected from



- 15 R^3, R^4 each represent a hydrogen atom;
- R^5 represents a group selected from a hydrogen atom, $-C(O)NR^{10}R^{11}$;
- R^6, R^7 represent, independently from each other, a hydrogen or a fluoro atom;
- R^8 represents a methyl group;
- R^{10}, R^{11} represent, independently from each other, a group selected from hydrogen, methyl;
- 20 or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (I) or (Ia), wherein

- R^1 represents a methyl group;
- R^2 represents a group selected from



R^3, R^4 represent a hydrogen atom;

R^5 represents a hydrogen atom or a $-C(O)NR^{10}R^{11}$ group;

R^6, R^7 represent, independently from each other, a hydrogen or a fluoro atom;

5 R^8 represents a methyl group;


R^{10}, R^{11} represent, independently from each other, a hydrogen atom or a methyl group;

or their salts, solvates or salts of solvates.

In another embodiment the present invention concerns compounds of general formula (I) or (Ia), wherein

10 R^1 represents methyl,

R^2 represents a group selected from



R^3, R^4 represent a hydrogen atom;

R^5 represents a group selected from a hydrogen atom or $-C(O)OR^9$;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom or fluoro atom;

15 R^8 represents a group selected from methyl or benzyl;


R^9 represents ethyl;

or their salts, solvates or salts of solvates.

In another embodiment the present invention concerns compounds of general formula (I) or (Ia), wherein

R¹ represents methyl;

R² represents a group selected from



R³ represents a fluoro atom;

5 R⁴ represents a hydrogen atom;

R⁵ represents a group selected from a hydrogen atom or cyano;


R⁶, R⁷ represent, independently from each other, a group selected from a hydrogen atom or fluoro atom;

R⁸ represents a group selected from methyl or benzyl;

or their salts, solvates or salts of solvates.

10

The present invention relates to compounds of general formula (Ia)



, wherein

R¹ represents a methyl group;

15 R² represents a group selected from 2-methoxyphenyl-, 4-fluoro-2-methoxyphenyl-, 5-fluoro-2-methoxyphenyl-, 2-(benzyloxy)-4-fluorophenyl-, 4,5-difluoro-2-methoxyphenyl-, 4-chloro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-, 2-(cyclohexylmethoxy)-4-fluorophenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-, 2-[(3,4-dichlorobenzyl)oxy]phenyl-, 2-(1-cyclopentylethoxy)-4-fluorophenyl-, 3-chloro-2-methoxyphenyl-, 2-phenoxyphenyl-;

R³ represents a group selected from a hydrogen atom, fluoro atom, chloro atom;


20 R⁴ represents a group selected from a hydrogen atom or chloro atom;

R^5 represents a group selected from a hydrogen atom, $-C(O)R^9$, $-C(O)OR^9$, methyl;

R^9 represents methyl or ethyl;

or their salts, solvates or salts of solvates.

5 The present invention relates to compounds of general formula (Ia)



(Ia)

, wherein

R^1 represents a methyl group;

R^2 represents a group selected from 2-methoxyphenyl-, 4-fluoro-2-methoxyphenyl-,
5-fluoro-2-methoxyphenyl-, 2-(benzyloxy)-4-fluorophenyl-, 4,5-difluoro-2-methoxyphenyl-,
10 4-chloro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-, 2-(cyclohexylmethoxy)-4-fluorophenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-, 2-[(3,4-dichlorobenzyl)oxy]phenyl-, 2-(1-cyclopentylethoxy)-4-fluorophenyl-;


R^3 represents a group selected from a hydrogen atom, fluoro atom, chloro atom;

15 R^4 represents a group selected from a hydrogen atom, chloro atom;

R^5 represents a hydrogen atom;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (Ia)



(Ia)

, wherein

R^1 represents a methyl group;

R^2 represents a group selected from 2-methoxyphenyl-, 4-fluoro-2-methoxyphenyl-,
2-(benzyl)oxy-4-fluorophenyl-, 4,5-difluoro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-,
20 2-(cyclohexylmethoxy)-4-fluorophenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-;

R^3 represents a group selected from a hydrogen atom, fluoro atom, chloro atom;


R^4 represents a group selected from a hydrogen atom, chloro atom;

R^5 represents a group selected from a hydrogen atom;

or their salts, solvates or salts of solvates.

5

The present invention relates to compounds of general formula (Ia)



(Ia)

, wherein

R^1 represents a group selected from methyl;

R^2 represents a group selected from 4-fluoro-2-methoxyphenyl-, 2-(benzyloxy)-4-fluorophenyl-,
4,5-difluoro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-,
4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-;

R^3 represents a group selected from a hydrogen atom, fluoro atom, chloro atom;


R^4 represents a group selected from a hydrogen atom, chloro atom;

R^5 represents a group selected from a hydrogen atom, $-C(O)R^9$, $-C(O)OR^9$;

R^9 represents methyl or ethyl;

or their salts, solvates or salts of solvates.

The present invention relates to compounds of general formula (Ia)



(Ia)

, wherein

R^1 represents a group selected from methyl;

R^2 represents a group selected from 4-fluoro-2-methoxyphenyl-, 5-fluoro-2-methoxyphenyl-,
4,5-difluoro-2-methoxyphenyl-;

R^3 , R^4 represent, independently from each other, a hydrogen atom;


R^5 represents a hydrogen atom;

or their salts, solvates or salts of solvates.

20

25

The present invention relates to compounds of general formula (Ia)



, wherein

R¹ represents a group selected from methyl;


R² represents a group selected from 4-fluoro-2-methoxyphenyl-, 4,5-difluoro-2-methoxyphenyl-;

5 R³, R⁴ represent, independently from each other, a hydrogen atom;

R⁵ represents a hydrogen atom;

or their salts, solvates or salts of solvates.

In another preferred embodiment the present invention relates to compounds of general formula (Ib)



(Ib), wherein

R¹ represents a methyl group;

R³ represents a group selected from a hydrogen atom or fluoro atom or chloro atom;

R⁴ represents a group selected from a hydrogen atom or fluoro atom;

15 R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)CH₃ -C(O)OCH₂CH₃-,
-C(O)NHCH₃;

R⁸ represents a group selected from methyl, ethyl, prop-2-en-1-yl, 2-methylprop-2-en-1-yl,
(2Z)-but-2-en-1-yl, prop-2-yn-1-yl, but-2-yn-1-yl, 2-(hydroxymethyl)prop-2-en-1-yl, phenyl,
benzyl, 3-cyanobenzyl, 3-fluorobenzyl, 3-chlorobenzyl, 4-fluorobenzyl, 4-chlorobenzyl,
20 3-fluoro-5-(trifluoromethyl)benzyl, 3-chloro-5-fluorobenzyl, pyridin-4-yl, 2-fluoropyridin-4-yl,
2,3,5-trifluorobenzyl, 3,4,5-trifluorobenzyl;

or their salts, solvates or salts of solvates.

25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹
represents a C₁-C₃-alkyl-, a C₃-C₅-cycloalkyl- or a heterocyclyl group.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹
represents a C₁-C₃-alkyl- or a heterocyclyl group.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹ represents a C₁-C₃-alkyl- group.

- 5 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹ represents a methyl group.


In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹ represents a methyl group.

10


In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹ represents a C₃-C₅-cycloalkyl-group, preferably a cycopropyl group.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents


15



In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents




In a preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents



20

In another preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents



- In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents a group selected from 2-methoxyphenyl-, 4-fluoro-2-methoxyphenyl-, 5-fluoro-2-methoxyphenyl-,
 5 2-(benzyloxy)-4-fluorophenyl-, 4,5-difluoro-2-methoxyphenyl-, 4-chloro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-, 2-(cyclohexylmethoxy)-4-fluorophenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-, 2-[(3,4-dichlorobenzyl)oxy]phenyl-, 2-(1-cyclopentylethoxy)-4-fluorophenyl-, 3-chloro-2-methoxyphenyl-, 2-phenoxyphenyl-.
- 10 In a preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents a group selected from 4-fluoro-2-methoxyphenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-, 4-fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl-, 2-(but-2-yn-1-yloxy)-4-fluorophenyl-, 4-fluoro-2-(prop-2-yn-1-yloxy)phenyl-.
- 15 In another, particularly preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents a group selected from 4-fluoro-2-methoxyphenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl-, 4-fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl-, 4-fluoro-2-(prop-2-yn-1-yloxy)phenyl-.
- 20 In another, particularly preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R² represents a group selected from 4-fluoro-2-methoxyphenyl-.
- In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R³ represents a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C₁-C₂-alkyl-, C₁-C₂-alkoxy-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-.
- 25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R³ represents a hydrogen atom or a fluoro atom or a chloro atom.
- In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R³ represents a hydrogen atom or a fluoro atom.
- 30 In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R³ represents a hydrogen atom.

In another preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R³ represents a fluoro atom.

5 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R³ represents a fluoro atom and in which R³ is in meta position of the N-phenyl substituent in 2-position of the 1, 3, 5-triazine core.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R³ represents a fluoro atom and in which R³ is in meta position of the N-phenyl substituent in 2-position of the 1, 3, 5-triazine core, and in which R⁴ represents a hydrogen atom.

10 In another embodiment the invention relates to compounds of formula (I), in which R³ represents a fluoro atom and in which R³ is in para position of the N-phenyl substituent in 2-position of the 1, 3, 5-triazine core.

In another embodiment the invention relates to compounds of formula (I), in which R³ represents a fluoro atom and in which R³ is in para position of the N-phenyl substituent in 2-position of the 1, 3, 5-triazine core, and in which R⁴ represents a hydrogen atom.

15 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁴ represents a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C₁-C₂-alkyl-, C₁-C₂-alkoxy-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁴ represents a group selected from a hydrogen, a fluoro or a chloro atom.

20 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁴ represents a group selected from a hydrogen atom or fluoro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁴ represents a hydrogen atom.

25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁴ represents a chloro atom.

In a preferred embodiment the invention relates to compounds of formula (Ia) or (Ib), in which R³ represents a fluoro atom and R⁴ represents a hydrogen atom.

In another embodiment the invention relates to compounds of formula (Ia) or (Ib), in which R³ represents a hydrogen atom and R⁴ represents a fluoro atom.

30 In another preferred embodiment the invention relates to compounds of formula (Ia) or (Ib), in which R³ represents a hydrogen atom and R⁴ represents a hydrogen atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)R⁹, -C(O)OR⁹, -S(O)₂R⁹, -C(O)NR¹⁰R¹¹, C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl, phenyl, heteroaryl, wherein said C₁-C₆-alkyl, C₃-C₇-cycloalkyl-, heterocycyl, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, cyano, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)R⁹, -C(O)OR⁹, -S(O)₂R⁹, -C(O)NR¹⁰R¹¹, methyl.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from cyano, -C(O)R⁹, -C(O)OR⁹, -S(O)₂R⁹, -C(O)NR¹⁰R¹¹, methyl.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)OR⁹, C(O)NR¹⁰R¹¹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from cyano, -C(O)OR⁹, C(O)NR¹⁰R¹¹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom, cyano or -C(O)OR⁹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from cyano or -C(O)OR⁹.

In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom or a cyano group.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from cyano, -S(O)₂R⁹, -C(O)NR¹⁰R¹¹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a group selected from a hydrogen atom or -C(O)OR⁹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a hydrogen atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents -C(O)OR⁹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents -C(O)NR¹⁰R¹¹.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents -S(O)₂R⁹.

5 In another preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a cyano group.

In another preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁵ represents a hydrogen atom.

10 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ represents a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C₁-C₂-alkyl-, C₁-C₂-alkoxy-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ represents a group selected from a hydrogen atom or fluoro atom or chloro atom.

15 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ represents a group selected from a hydrogen atom or fluoro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ represents a hydrogen atom.

20 In another preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and represents a fluoro atom.

In another preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which

R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and represents a fluoro atom;

R⁷ represents a hydrogen atom.

25 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ represents a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C₁-C₂-alkyl-, C₁-C₂-alkoxy-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ represents a group selected from a hydrogen atom or fluoro or chloro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ represents a group selected from a hydrogen atom or fluoro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ represents a fluoro atom.

- 5 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ represents a chloro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁷ is a hydrogen atom.

- 10 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and R⁶ represents a fluoro atom and in which R⁷ represents a group selected from a hydrogen atom, fluoro atom, chloro atom, C₁-C₂-alkyl-, C₁-C₂-alkoxy-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-.

- 15 In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and R⁶ represents a fluoro atom and in which R⁷ represents a group selected from a hydrogen atom or fluoro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and R⁶ represents a fluoro atom and in which R⁷ represents a hydrogen atom.

- 20 In a preferred embodiment the invention relates to compounds of formula (I) or (Ia), in which R⁶ is in para position of the phenyl substituent in 4-position of the 1, 3, 5- triazine core and R⁶ represents a fluoro atom and in which R⁷ represents a hydrogen atom.

- 25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a C₁-C₂-alkyl group, which is optionally substituted with one substituent, selected from the group consisting of C₃-C₇-cycloalkyl-, heterocycyl-, phenyl, wherein said C₃-C₇-cycloalkyl-, heterocycyl- or phenyl group is optionally substituted with one, or two substituents, identically or differently, selected from halogen.

- 30 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a methyl group, which is optionally substituted with one substituent selected from the group consisting of phenyl, wherein said phenyl group is optionally substituted with one or two substituents, identically or differently, selected from a fluoro or chloro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a methyl group, which is optionally substituted with one substituent selected from the group

consisting of heteroaryl, wherein said heteroaryl group is optionally substituted with one or two substituents, identically or differently, selected from a fluoro or chloro atom.

In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a methyl group.

- 5 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a group selected from -CH₂-CH₂CF₃, -CH₂CH₂CF₂CF₃.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a methyl or a (²H₃)methyl group.

- 10 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a phenyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₂-alkyl-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-, C₁-C₂-alkoxy-.

- 15 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a heteroaryl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₂-alkyl-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-, C₁-C₂-alkoxy-.

- 20 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a phenyl-C₁-C₃-alkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of fluoro atom, chloro atom, cyano, methyl-, trifluoromethyl-.

- 25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a phenyl-C₁-C₃-alkyl- group, which is optionally substituted with one or two substituents, identically or differently, selected from the group consisting of halogen.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a benzyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of fluoro atom, chloro atom, cyano, methyl-, trifluoromethyl-.

- 30 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a benzyl group, which is optionally substituted with one or two substituents, identically or differently, selected from the group consisting of a fluoro or chloro atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a benzyl group, which is optionally substituted with one fluoro atom.

5 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a heteroaryl-C₁-C₂-alkyl-, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, halo-C₁-C₂-alkyl- and C₁-C₂-alkoxy-.

10 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a pyridyl-C₁-C₂-alkyl- group, wherein said pyridyl is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, NH₂, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₂-alkyl-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-, C₁-C₂-alkoxy-.

15 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a pyridyl-C₁-C₂-alkyl- group, wherein said pyridyl is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, halo-C₁-C₂-alkyl- and C₁-C₂-alkoxy-.

20 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a oxazolyl-C₁-C₂-alkyl- or thiazolyl-C₁-C₂-alkyl- group, wherein said oxazolyl or thiazolyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, NH₂, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₂-alkyl-, halo-C₁-C₂-alkyl-, C₁-C₂-fluoroalkoxy-, C₁-C₂-alkoxy-.

25 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a oxazolyl-C₁-C₂-alkyl- or thiazolyl-C₁-C₂-alkyl- group, wherein said oxazolyl or thiazolyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, halo-C₁-C₂-alkyl-, C₁-C₂-alkoxy-.

30 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a group selected from methyl, ethyl, prop-2-en-1-yl-, 2-methylprop-2-en-1yl-, 2-(hydroxymethyl)prop-2-en-1-yl-, (2Z)-but-2-en-1-yl-, prop-2-yn-1yl-, but-2-yn-1yl-, cyclohexylmethyl-, benzyl-, 3-cyanobenzyl-, 4-cyanobenzyl-, 3-fluorobenzyl-, 3-chlorobenzyl-, 4-fluorobenzyl-, 4-chlorobenzyl-, 3-fluoro-5-(trifluoromethyl)benzyl-, 3-chloro-5-fluorobenzyl-, pyridin-4-yl-, 2-fluoropyridin-4-yl-, 2,3,5-trifluorobenzyl-, 3,4,5-trifluorobenzyl-.

In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a group selected from methyl, ethyl, prop-2-en-1-yl-, 2-methylprop-2-en-1yl-, 2-(hydroxymethyl)prop-2-en-1-yl-, (2Z)-but-2-en-1-yl-, prop-2-yn-1yl-, but-2-yn-1yl-, phenyl, 3-

cyanobenzyl-, 3-fluorobenzyl-, 3-chlorobenzyl-, 4-fluorobenzyl-, 4-chlorobenzyl-, 3-fluoro-5-(trifluoromethyl)benzyl-, 3-chloro-5-fluorobenzyl-, pyridin-4-yl-, 2-fluoropyridin-4-yl-, 2,3,5-trifluorobenzyl-, 3,4,5-trifluorobenzyl-.

5 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a group selected from C₁-C₃-alkyl or benzyl.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents a group selected from methyl or benzyl.

In a preferred embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents methyl.

10 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁸ represents benzyl.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R⁹ represents a C₁-C₆-alkyl-group.

15 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹⁰ represents a group selected from a hydrogen atom or C₁-C₆-alkyl-.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹⁰ represents a hydrogen atom.

In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹¹ represents a group selected from a hydrogen atom or C₁-C₆-alkyl-.

20 In another embodiment the invention relates to compounds of formula (I) or (Ia) or (Ib), in which R¹¹ represents methyl.

It is to be understood that the present invention relates to any sub-combination within any embodiment of the present invention of compounds of formula (I) or (Ia) or (Ib), supra.

25 More particularly still, the present invention covers compounds of formula (I) or (Ia) or (Ib) which are disclosed in the Example section of this text, infra.

Very specially preferred are combinations of two or more of the abovementioned preferred ranges.

In particular, further preferred subjects of the present invention are the compounds selected from:

(rac)-Ethyl [(3- {[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
 (rac)-4-(4-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
 4-(4-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,
 4-(4-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,
 (rac)-Ethyl {[3-({4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl} amino)benzyl)(methyl)oxido- λ^6 -sulfanylidene} carbamate,
 (rac)-4-[2-(Benzyloxy)-4-fluorophenyl]-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
 4-[2-(Benzyloxy)-4-fluorophenyl]-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,
 4-[2-(Benzyloxy)-4-fluorophenyl]-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,
 (rac)-Ethyl [(3- {[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
 (rac)- 4-(4,5-Difluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
 (rac)-Ethyl [(3- {[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
 (rac) 4-(4-Chloro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
 4-(4-Chloro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,
 4-(4-Chloro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,
 (rac)-1-[(3- {[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea,
 1-[(3- {[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 1,
 1-[(3- {[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 2

(rac)-Ethyl [(3- {[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,

(rac)-Ethyl [(3- {[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(5-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,

4-(5-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,

4-(5-Fluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,

(rac)-N-[(3- {[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]acetamide,

(rac)-Ethyl [(3- {[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(2-Methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,

4-(2-Methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,

4-(2-Methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,

(rac)-Ethyl [(3- {[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,

4-(3,4-Dihydro-2H-chromen-8-yl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 1,

4-(3,4-Dihydro-2H-chromen-8-yl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine; enantiomer 2,

(rac)-Ethyl [(3- {[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,

4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,

4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,

(rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,

4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,

(rac)-N-{3-[(N,S-Dimethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,

(rac)-Ethyl [(3-{[4-(2-{[4-(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl})(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-{2-{[4-(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-N-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]methanesulfonamide,

(rac)-Ethyl [(3-{[4-(3-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-Ethyl [(3-{[4-{5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-Ethyl [methyl(oxido)(3-{[4-(2-phenoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)- λ^6 -sulfanylidene]carbamate,

(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,

[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 1,

[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2,

(rac)-Ethyl [(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, enantiomer 1,
4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, enantiomer 2,
(rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-{4-Fluoro-2-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-(3-{[S-(tetrahydro-2H-pyran-4-yl)sulfonimidoyl]methyl}-phenyl)-1,3,5-triazin-2-amine,
(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
(rac)-Ethyl [{3-[(4-{2-[(3,4-dichlorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl}(methyl)-oxido- λ^6 -sulfanylidene]carbamate,
(rac)-4-{2-[(3,4-Dichlorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-(4-Fluoro-2-{[($^2\text{H}_5$)phenyl($^2\text{H}_2$)methyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
4-[2-(1-cyclopentylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(3,3,3-trifluoropropoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(pyridin-2-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

4-{4-Fluoro-2-[1-(4-fluorophenyl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of 4 stereoisomers,
(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ 6-sulfanylidene)cyanamide],
[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ 6-sulfanylidene)cyanamide; enantiomer 1,
[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ 6-sulfanylidene)cyanamide; enantiomer 2,
(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
(rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-{[4-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-{[3-fluoro-5-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-3-({5-Fluoro-2-[4-(3-[(S-methylsulfonimidoyl)methyl]phenyl)amino]-1,3,5-triazin-2-yl}phenoxy)methylbenzonitrile,

(rac)-4-{4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4- {4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]-phenyl} -1,3,5-triazin-2-amine,

(rac)-4- {4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]-phenyl} -1,3,5-triazin-2-amine,

(rac)-[(2,3-Difluoro-5- {4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,

(rac)-N- {3,4-Difluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,

(rac)-[Ethyl(3- {4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)oxido- λ^6 -sulfanylidene]cyanamide,

(rac)-N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine
 N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine;
 enantiomer 1,

N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine;
 enantiomer 2,

(rac)-[(3- {4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} -5-methylbenzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,

(rac)-2-({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl} amino)-1,3,5-triazin-2-yl]phenoxy} methyl)prop-2-en-1-ol,


(rac)-[Cyclopropyl(3- {4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)oxido- λ^6 -sulfanylidene]cyanamide,

or their salts, solvates or salts of solvates.


The above-mentioned definitions of radicals which have been detailed in general terms or in preferred ranges also apply to the end products of the formula (I) or (Ia) or (Ib) and, analogously, to the starting materials or intermediates required in each case for the preparation.

5 Scheme 1:

The invention furthermore relates to a process for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the invention, in which N-unprotected sulfoximines of formula (6) are reacted to give N-functionalized sulfoximines of formula (I) or (Ia) or (Ib).



The present invention therefore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib), in which R⁵ is not a hydrogen atom, according to the invention, in which method the nitrogen of the sulfoximine group of a compound of formula (6)



in which R¹, R², R³ and R⁴ are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention, is functionalized according to methods known in the art, thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R⁵ is not hydrogen, and the resulting compounds are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.


There are multiple methods for the preparation of N-functionalized sulfoximines by functionalization of the nitrogen of the sulfoximine group:

- 15 - Alkylation: see for example: a) U. Lücking et al, US 2007/0232632; b) C.R. Johnson, J. Org. Chem. **1993**, 58, 1922; c) C. Bolm et al, Synthesis **2009**, 10, 1601.
- Acylation: see for example: a) C. Bolm et al, Chem. Europ. J. **2004**, 10, 2942; b) C. Bolm et al, Synthesis **2002**, 7, 879; c) C. Bolm et al, Chem. Europ. J. **2001**, 7, 1118.
- Arylation: see for example: a) C. Bolm et al, Tet. Lett. **1998**, 39, 5731; b) C. Bolm et al., J. Org. Chem. **2000**, 65, 169; c) C. Bolm et al, Synthesis **2000**, 7, 911; d) C. Bolm et al, J. Org. Chem. **2005**, 70, 2346; e) U. Lücking et al, WO2007/71455.
- Reaction with isocyanates: see for example: a) V.J. Bauer et al, J. Org. Chem. **1966**, 31, 3440; b) C. R. Johnson et al, J. Am. Chem. Soc. **1970**, 92, 6594; c) S. Allenmark et al, Acta Chem. Scand. Ser. B **1983**, 325; d) U. Lücking et al, US2007/0191393.
- 25 - Reaction with sulfonylchlorides: see for example: a) D.J. Cram et al, J. Am. Chem. Soc. **1970**, 92, 7369; b) C.R. Johnson et al, J. Org. Chem. **1978**, 43, 4136; c) A.C. Barnes, J. Med. Chem. **1979**, 22, 418;


d) D. Craig et al, Tet. **1995**, 51, 6071; e) U. Lücking et al, US2007/191393.

- Reaction with chloroformates: see for example: a) P.B. Kirby et al, DE2129678; b) D.J. Cram et al, J. Am. Chem. Soc. **1974**, 96, 2183; c) P. Stoss et al, Chem. Ber. **1978**, 111, 1453; d) U. Lücking et al, WO2005/37800.

- 5 N-unprotected sulfoximines of formula (6) can be prepared by deprotection of N-protected sulfoximines of formula (5). Preferred it the use of sodium ethanolate in ethanol at 60°C (see for example: U. Lücking et al, WO2005/37800).




- 10 The invention therefore furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the present invention, in which R⁵ is a hydrogen atom (identical to the N-unprotected sulfoximines of formula (6) shown above), according to the invention, in which method the –C(O)O-Ethyl group of an N-protected compound of formula (5)




- 15 in which R¹, R², R³ and R⁴ are as defined for the compound of general formula (I) or (Ia) or (Ib), is deprotected according to methods known in the art, thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R⁵ is a hydrogen atom, and the resulting compounds (the N-unprotected sulfoximines of formula (6) shown above) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.
- 20

The invention furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the present invention, in which R⁵ is –C(O)O-Ethyl (identical to the N-protected sulfoximines of formula (5) shown above), in which method an intermediate compound of formula (3),



in which R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib), is reacted with a compound of formula (4)



in which R^2 is as defined for the compound of general formula (I) or (Ia) or (Ib), and R represent, independently from each other, a hydrogen atom, or a C_1 - C_{10} -alkyl group or, alternatively, both R together form a R-R group, which is $-C(CH_3)_2-C(CH_3)_2-$,

thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R^5 is $-C(O)O$ -Ethyl, and

the resulting compounds (see N-protected sulfoximines of formula (5) shown above) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.

Compounds of general formula (4) can be prepared analogously to known processes (review: D.G. Hall, Boronic Acids, **2005** WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, ISBN 3-527-30991-8 and references cited herein). Further, a wide variety of compounds of general formula (4) are commercially available.

The coupling reaction of compounds of formula (3) with compounds of formula (4) is catalyzed by Pd catalysts, e.g. by Pd(0) catalysts or by Pd(II) catalysts. Examples for Pd(0) catalysts are tetrakis(triphenylphosphine)palladium(0) $[Pd(PPh_3)_4]$ or tris(dibenzylideneacetone)di-palladium(0) $[Pd_2(dba)_3]$, examples for Pd(II) catalysts dichlorobis(triphenylphosphine)-palladium(II) $[Pd(PPh_3)_2Cl_2]$, palladium(II) acetate and triphenylphosphine or $[1,1'$ -bis(diphenylphosphino)ferrocene]palladium dichloride $[Pd(dppf)Cl_2]$ (review: D.G. Hall, Boronic Acids, **2005** WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, ISBN 3-527-30991-8 and references cited therein).

This reaction is preferably carried out in aprotic or protic solvents, preferably in a mixture of aprotic and protic solvents, more preferably in solvents like, for example, 1,2-dimethoxyethane, dioxane,


dimethylformamid, tetrahydrofuran, or isopropanol with water (review: D.G. Hall, Boronic Acids, **2005** WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, ISBN 3-527-30991-8 and references cited therein).

Preferably the reaction is carried out in the presence of a suitable base, such as for example aqueous potassium carbonate, aqueous sodium bicarbonate or aqueous potassium phosphate (review: D.G. Hall, Boronic Acids, **2005** WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, ISBN 3-527-30991-8 and references cited therein).

The reaction is performed at temperatures ranging from room temperature ($=20^{\circ}\text{C}$) to the boiling point of the solvent. Further on, the reaction can be performed at temperatures above the boiling point using pressure tubes and a microwave oven. (review: D.G. Hall, Boronic Acids, **2005** WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, ISBN 3-527-30991-8 and references cited therein).

The reaction is preferably completed after 1 to 36 hours of reaction time.

Compounds of general formula (3) can be obtained as follows:




2,4-dichloro-1,3,5-triazine (1), which is known [CAS-Registry Number: 2831-66-5] or can be prepared analogously to known processes, is reacted with suitable anilines of formula (2) to give the corresponding 4-chloro-N-phenyl-1,3,5-triazin-2-amines of formula (3).

This reaction can be carried out with one equivalent of the aniline of formula (2) in an inert solvent like, for example, 1,2-dimethoxyethane, dioxane, dimethylformamid, tetrahydrofuran, or an alcohol like, for example, isopropanol, or mixtures of such solvents. Preferably, the reaction is carried out at a temperature below 0°C in such a way that the reaction mixture is kept homogenous. Preferred conditions use an additional base like for example triethylamine or N,N-diisopropylethylamine. The reaction is preferably completed after 1 to 6 hours.

Anilines of formula (2) can be prepared by the following processes:

Reaction of suitable benzylchlorides or -bromides of formula (7) with suitable thiols of formula (8) under basic conditions yields the corresponding thioethers of formula (9) (see for example: Sammond et al, Bioorg. Med. Chem. Lett. **2005**, 15, 3519).



LG = Cl, Br


Oxidation of thioethers of formula (9) gives the corresponding sulfoxides of formula (10). The oxidation can be performed analogously to known processes (see for example: (a) M.H. Ali et al, *Synthesis* **1997**, 764; (b) M.C. Carreno, *Chem. Rev.* **1995**, 95, 1717; (c) I. Patel et al, *Org. Proc. Res. Dev.* **2002**, 6, 225; (d) N. Khair et al, *Chem. Rev.* **2003**, 103, 3651). Preferred is the herein described use of periodic acid and iron(III)chloride.



Rhodium-catalyzed imination of the sulfoxides of formula (10) followed by deprotection gives the corresponding N-unprotected sulfoximines of formula (11) (see for example: Bolm et al, *Org. Lett.* **2004**, 6, 1305).




Introduction of a suitable protecting group leads to N-protected sulfoximines of formula (12) (see for example: Lücking et al, WO 2005/037800).



Reduction of the nitro group finally gives the desired anilines of formula (2). The reduction can be prepared analogously to known processes (see for example: (a) Sammond et al, *Bioorg. Med. Chem. Lett.* **2005**, 15, 3519; (b) R.C. Larock, *Comprehensive Organic Transformations*, VCH, New York, **1989**, 411-415).




The preparation of the compounds of general formula (I) or (Ia) or (Ib) according to the invention can be illustrated by the following synthesis scheme (Scheme 1):



Compounds of general formula (Ia), (Ib), (2a), (3a), (5a), (6a) can be prepared analogously.

Scheme 2:

The invention furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the present invention, in which R⁵ is a hydrogen atom (identical to the N-unprotected sulfoximines of formula (6) shown above), in which method a compound of formula (15)



15

in which R¹, R², R³ and R⁴ are as defined for the compound of formula (I) or (Ia) or (Ib) according to the present invention,

is reacted with sodium azide in the presence of an acid thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R⁵ is a hydrogen atom, and, the resulting compounds (the N-unprotected sulfoximines of formula (6) shown above) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.


This reaction is preferably carried out in the presence of sulfuric acid or polyphosphoric acid. It is preferably carried out in trichloromethane in the presence of sulfuric acid and at temperatures ranging from 40°C to 50°C, preferably from 43°C to 47°C, most preferred at about 45°C. This reaction is preferably completed after 12 hours to 120 hours of reaction time (see for example: a) H. R. Bentley et al, J. Chem. Soc. **1952**, 1572; b) C. R. Johnson et al, J. Am. Chem. Soc. **1970**, 92, 6594; c) Satzinger et al, Angew. Chem. **1971**, 83, 83).

In another embodiment of the present invention the method of the present invention is performed with sodium azide in polyphosphoric acid at temperatures ranging from 40°C to 70°C, preferably from 55°C to 65°C, most preferred at about 60°C (see for example: a) M.D. Sindkhedkar et al, WO 2007/023507).


The preparation of compounds of general formula (15) is described below in context of the description of synthesis Scheme 2. Compounds of general formula (Ia) or (Ib) can be prepared analogously.

Scheme 3

The invention furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the present invention, in which R⁵ is a cyano group (identical to the N-cyanosulfoximines of formula (20)),



in which method a compound of formula (19)



in which R^1 , R^2 , R^3 and R^4 are as defined for the compound of formula (I) or (Ia) or (Ib) according to the present invention,

is oxidized according to methods known in the art, thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R^5 is a cyano group, and the resulting compounds (the N-cyanosulfoximines of formula (20) as shown above) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.

There are multiple methods for the oxidation of N-cyanosulfilimines of formula (19) to N-cyanosulfoximines of formula (20):


a) C. Bolm et al, Org. Lett. **2007**, 9, 3809

b) J.E.G. Kemp et al, Tet. Lett. **1979**, 39, 3785


c) M.R. Loso et al, US patent publication US2007/0203191.

The preparation of compounds of general formula (19) is described below in context of the description of synthesis Scheme 3. Compounds of general formula (Ia) or (Ib) can be prepared analogously.

The invention furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib), in which R^5 is a hydrogen atom (identical to the sulfoximines of formula (6)),

**6 (formula (I), R⁵ = H)**

in which method the N-cyano group of the compound of formula (20)

**20**


in which R¹, R², R³ and R⁴ are as defined for the compound of formula (I) or (Ia) or (Ib),

is converted upon treatment with TFAA to the corresponding N-trifluoroacetylsulfoximine, which N-trifluoroacetylsulfoximine is converted by methanolysis into the compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R⁵ is a hydrogen group, and the resulting compounds (compounds of general formula (6)) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.

The preparation of compounds of general formula (6) from compounds of general formula (20) is described in more detail below in context of the description of synthesis Scheme 3.

15 Scheme 4:

The invention furthermore relates to a method for the preparation of the compounds of formula (I) or (Ia) or (Ib) according to the present invention, in which R⁵ is a hydrogen atom (identical to the N-unprotected sulfoximines of formula (6) shown above), in which method a compound of formula (22)

**22**

in which R^1 , R^3 , R^4 , R^6 and R^7 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention,

is solubilized in an alcohol R^8 -OH,

5 in which R^8 is as defined for the compounds of general formula (I) or (Ia) or (Ib) according to the invention

and is reacted with at least two equivalents of alkali hydride,


thus providing a compound of general formula (I) or (Ia) or (Ib) according to the invention, in which R^5 is a hydrogen atom, and

10 the resulting compounds (the N-protected sulfoximines of formula (6)) are optionally, if appropriate, reacted with the corresponding (i) solvents and/or (ii) bases or acids to the solvates, salts and/or solvates of the salts thereof.


The reaction can be performed with potassium hydride or sodium hydride. Preferably, it is performed with sodium hydride. The reaction is performed at temperatures ranging from 40°C to the boiling point of the alcohol R^8 -OH, preferably at 50°C to 70°C. The reaction is preferably completed after 10 to 100 hours of reaction time.

The preparation of compounds of general formula (22) is described below in context of the description of synthesis Scheme 4. Compounds of general formula (Ia) and (Ib) can be prepared analogously.

In another embodiment the present invention concerns intermediate compounds of general formula (3)




or of general formula (3a)




, wherein

R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

In another embodiment the present invention concerns compounds of general formula (5)



(5) or of general formula (5a)




5a

, wherein


R^1 , R^2 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

In another embodiment the present invention concerns compounds of general formula (2)



2

or of general formula (2a)





2a

, wherein



R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

In another embodiment the present invention concerns compounds of general formula (11)




R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

In another embodiment the present invention concerns compounds of general formula (12)




10 R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

In another embodiment the present invention concerns compounds of general formula (20)



or of general formula (20a)



R^1 , R^2 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) or (Ib) according to the invention.

The compounds according to the invention show a valuable pharmacological and pharmacokinetic spectrum of action which could not have been predicted.

They are therefore suitable for use as medicaments for the treatment and/or prophylaxis of disorders in humans and animals.

Within the scope of the present invention, the term “treatment” includes prophylaxis.

The pharmaceutical activity of the compounds according to the invention can be explained by their action as inhibitors of CDK9. Thus, the compounds according to the general formula (I) or (Ia) or (Ib) as well as pharmaceutically acceptable salts thereof are used as inhibitors for CDK9.

Furthermore, the compounds according to the invention show a particularly high potency (demonstrated by a low IC_{50} value in the CDK9/CycT1 assay) for inhibiting CDK9 activity.

In context of the present invention, the IC_{50} value with respect to CDK9 can be determined by the methods described in the method section below. Preferably, it is determined according to Method 1. (“CDK9/CycT1 kinase assay”) described in the Materials and Method section below.

Surprisingly it turned out that the compounds according to the general formula (I) or (Ia) or (Ib) as well as pharmaceutically acceptable salts thereof selectively inhibit CDK9 in comparison to other cyclin-dependent protein kinases, preferably in comparison to CDK2. Thus, the compounds according to the

general formula (I) or (Ia) or (Ib) as well as pharmaceutically acceptable salts thereof are preferably used as selective inhibitors for CDK9.

Compounds of the present invention according to general formula (I) or (Ia) or (Ib) show a significantly stronger CDK9 than CDK2 inhibition. Preferred compounds of the present invention show a CDK2 IC₅₀ / CDK9 IC₅₀ ratio of more than 40, preferably of more than 55 and even more preferably of more than 70. The CDK9 IC₅₀ is determined according to Method 1., the CDK2 IC₅₀ according to Method 2, both described in more detail in the Materials and Method section below.

In context of the present invention, the IC₅₀ value with respect to CDK2 can be determined by the methods described in the method section below. Preferably, it is determined according to Method 2. (“CDK2/CycE kinase assay”) described in the Materials and Method section below.

Further, compounds of the present invention according to formula (I) or (Ia) or (Ib) mediate a surprisingly strong anti-proliferative activity in tumor cell lines such as HeLa and/or DU145. In context of the present invention, the IC₅₀ values of the compounds with respect to these cell lines is preferably determined according to Method 3. (“Proliferation Assay”) described in the Materials and Method section below.

Further, preferred compounds of the present invention according to formula (I) or (Ia) or (Ib) surprisingly show an increased solubility in water at pH 6.5 compared to the compounds described in the prior art.

In context of the present invention the solubility in water at pH 6.5 is preferably determined according to Method 4. (“Equilibrium Shake Flask Solubility Assay”) described in the Materials and Method section below.

Further, compounds of the present invention according to formula (I) or (Ia) or (Ib) show no significant inhibition of carbonic anhydrase-1 or -2 (IC₅₀ values of more than 10 μM) and therefore show an improved side effect profile as compared to those CDK inhibitors described in the prior art containing a sulfonamide group, which inhibit carbonic anhydrase-1 or -2. In context of the present invention, the carbonic anhydrase-1 and -2 inhibition is preferably determined according to Method 5. (“Carbonic anhydrase Assay”) described in the Materials and Method section below.

A further subject matter of the present invention is the use of the compounds of general formula (I) or (Ia) or (Ib) according to the invention for the treatment and/or prophylaxis of disorders, preferably of disorders relating to or mediated by CDK9 activity, in particular of hyper-proliferative disorders, virally induced infectious diseases and/or of cardiovascular diseases, more preferably of hyper-proliferative disorders.

The compounds of the present invention may be used to inhibit the activity or expression of CDK9. Therefore, the compounds of formula (I) or (Ia) or (Ib) are expected to be valuable as therapeutic agents.

Accordingly, in another embodiment, the present invention provides a method of treating disorders relating to or mediated by CDK9 activity in a patient in need of such treatment, comprising administering to the patient an effective amount of a compound of formula (I) or (Ia) or (Ib) as defined above. In certain embodiments, the disorders relating to CDK9 activity are hyper-proliferative disorders, virally induced infectious diseases and/or of cardiovascular diseases, more preferably hyper-proliferative disorders, particularly cancer.

The term "treating" or "treatment" as stated throughout this document is used conventionally, *e.g.*, the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of a disease or disorder, such as a carcinoma.

The term "subject" or "patient" includes organisms which are capable of suffering from a cell proliferative disorder or a disorder associated with reduced or insufficient programmed cell death (apoptosis) or who could otherwise benefit from the administration of a compound of the invention, such as human and non-human animals. Preferred humans include human patients suffering from or prone to suffering from a cell proliferative disorder or associated state, as described herein. The term "non-human animals" includes vertebrates, *e.g.*, mammals, such as non-human primates, sheep, cow, dog, cat and rodents, *e.g.*, mice, and non-mammals, such as chickens, amphibians, reptiles, etc.

The term "disorders relating to or mediated by CDK9" shall include diseases associated with or implicating CDK9 activity, for example the hyperactivity of CDK9, and conditions that accompany with these diseases. Examples of "disorders relating to or mediated by CDK9" include disorders resulting from increased CDK9 activity due to mutations in genes regulating CDK9 activity such as LARP7, HEXIM1/2 or 7sk snRNA, or disorders resulting from increased CDK9 activity due to activation of the CDK9/cyclinT/RNAPolymerase II complex by viral proteins such as HIV-TAT or HTLV-TAX or disorders resulting from increased CDK9 activity due to activation of mitogenic signaling pathways.

The term "hyperactivity of CDK9" refers to increased enzymatic activity of CDK9 as compared to normal non-diseased cells, or it refers to increased CDK9 activity leading to unwanted cell proliferation, or to reduced or insufficient programmed cell death (apoptosis), or mutations leading to constitutive activation of CDK9.

The term "hyper-proliferative disorder" includes disorders involving the undesired or uncontrolled proliferation of a cell and it includes disorders involving reduced or insufficient programmed cell death (apoptosis). The compounds of the present invention can be utilized to prevent, inhibit, block, reduce, decrease, control, etc., cell proliferation and/or cell division, and/or produce apoptosis. This method comprises administering to a subject in need thereof, including a mammal, including a human, an amount of a compound of this invention, or a pharmaceutically acceptable salt, hydrate or solvate thereof which is effective to treat or prevent the disorder.

Hyper-proliferative disorders in the context of this invention include, but are not limited to, *e.g.*, psoriasis, keloids and other hyperplasias affecting the skin, endometriosis, skeletal disorders, angiogenic or blood vessel proliferative disorders, pulmonary hypertension, fibrotic disorders, mesangial cell proliferative disorders, colonic polyps, polycystic kidney disease, benign prostate hyperplasia (BPH), and solid tumors, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid, and their distant metastases. Those disorders also include lymphomas, sarcomas and leukemias.

Examples of breast cancer include, but are not limited to invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma *in situ*, and lobular carcinoma *in situ*. Canine or feline mammary carcinoma.

Examples of cancers of the respiratory tract include, but are not limited to small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma, pleuropulmonary blastoma, and mesothelioma.

Examples of brain cancers include, but are not limited to brain stem and hypophthalmic glioma, cerebellar and cerebral astrocytoma, glioblastoma, medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumor.

Tumors of the male reproductive organs include, but are not limited to prostate and testicular cancer. Tumors of the female reproductive organs include, but are not limited to endometrial, cervical, ovarian, vaginal and vulvar cancer, as well as sarcoma of the uterus.

Tumors of the digestive tract include, but are not limited to anal, colon, colorectal, esophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers. Anal gland adenocarcinomas, mast cell tumors.

Tumors of the urinary tract include, but are not limited to bladder, penile, kidney, renal pelvis, ureter, urethral, and hereditary and sporadic papillary renal cancers.

Eye cancers include, but are not limited to intraocular melanoma and retinoblastoma.

Examples of liver cancers include, but are not limited to hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed hepatocellular cholangiocarcinoma.

Skin cancers include, but are not limited to squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer. Mast cell tumors.

Head-and-neck cancers include, but are not limited to laryngeal, hypopharyngeal, nasopharyngeal, oropharyngeal cancer, lip and oral cavity cancer, and squamous cell cancer. Oral melanoma.

Lymphomas include, but are not limited to AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Burkitt lymphoma, Hodgkin's disease, and lymphoma of the central nervous system.

Sarcomas include, but are not limited to sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma. Malignant histiocytosis, fibrosarcoma, hemangiosarcoma, hemangiopericytoma, leiomyosarcoma.

Leukemias include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia.

Fibrotic proliferative disorders, *i.e.* the abnormal formation of extracellular matrices, that may be treated with the compounds and methods of the present invention include lung fibrosis, atherosclerosis, restenosis, hepatic cirrhosis, and mesangial cell proliferative disorders, including renal diseases such as glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, transplant rejection, and glomerulopathies.

Other conditions in humans or other mammals that may be treated by administering a compound of the present invention include tumor growth, retinopathy, including diabetic retinopathy, ischemic retinal-vein occlusion, retinopathy of prematurity and age-related macular degeneration, rheumatoid arthritis, psoriasis, and bullous disorders associated with subepidermal blister formation, including bullous pemphigoid, erythema multiforme and dermatitis herpetiformis.

The compounds of the present invention may also be used to prevent and treat diseases of the airways and the lung, diseases of the gastrointestinal tract as well as diseases of the bladder and bile duct.

The disorders mentioned above have been well characterized in humans, but also exist with a similar etiology in other animals, including mammals, and can be treated by administering pharmaceutical compositions of the present invention.

In a further aspect of the present invention, the compounds according to the invention are used in a method for preventing and/or treating infectious diseases, in particular virally induced infectious diseases. The virally induced infectious diseases, including opportunistic diseases, are caused by retroviruses, hepadnaviruses, herpesviruses, flaviviridae, and/or adenoviruses. In a further preferred embodiment of this method, the retroviruses are selected from lentiviruses or oncoretroviruses, wherein the lentivirus is selected from the group comprising: HIV-1, HIV-2, FIV, BIV, SIVs, SHIV, CAEV, VMV or EIAV, preferably HIV-1 or HIV-2 and wherein the oncoretrovirus is selected from the group consisting of: HTLV-I, HTLV-II or BLV. In a further preferred embodiment of this method, the hepadnavirus is selected from HBV, GSHV or WHV, preferably HBV, the herpesvirus is selected from the group comprising: HSV I, HSV II, EBV, VZV, HCMV or HHV 8, preferably HCMV and the flaviviridae is selected from HCV, West Nile or Yellow Fever.

The compounds according to general formula (I) or (Ia) or (Ib) are also useful for prophylaxis and/or treatment of cardiovascular diseases such as cardiac hypertrophy, adult congenital heart disease, aneurysm, stable angina, unstable angina, angina pectoris, angioneurotic edema, aortic valve stenosis, aortic aneurysm, arrhythmia, arrhythmogenic right ventricular dysplasia, arteriosclerosis, arteriovenous malformations, atrial fibrillation, Behcet syndrome, bradycardia, cardiac tamponade, cardiomegaly, congestive cardiomyopathy, hypertrophic cardiomyopathy, restrictive cardiomyopathy, cardiovascular disease prevention, carotid stenosis, cerebral hemorrhage, Churg-Strauss syndrome, diabetes, Ebstein's Anomaly, Eisenmenger complex, cholesterol embolism, bacterial endocarditis, fibromuscular dysplasia, congenital heart defects, heart diseases, congestive heart failure, heart valve diseases, heart attack,

epidural hematoma, hematoma, subdural, Hippel-Lindau disease, hyperemia, hypertension, pulmonary hypertension, hypertrophic growth, left ventricular hypertrophy, right ventricular hypertrophy, hypoplastic left heart syndrome, hypotension, intermittent claudication, ischemic heart disease, Klippel-Trenaunay-Weber syndrome, lateral medullary syndrome, long QT syndrome mitral valve prolapse, moyamoya disease, mucocutaneous lymph node syndrome, myocardial infarction, myocardial ischemia, myocarditis, pericarditis, peripheral vascular diseases, phlebitis, polyarteritis nodosa, pulmonary atresia, Raynaud disease, restenosis, Sneddon syndrome, stenosis, superior vena cava syndrome, syndrome X, tachycardia, Takayasu's arteritis, hereditary hemorrhagic telangiectasia, telangiectasis, temporal arteritis, tetralogy of fallot, thromboangiitis obliterans, thrombosis, thromboembolism, tricuspid atresia, varicose veins, vascular diseases, vasculitis, vasospasm, ventricular fibrillation, Williams syndrome, peripheral vascular disease, varicose veins and leg ulcers, deep vein thrombosis, Wolff-Parkinson-White syndrome.

Preferred are cardiac hypertrophy, adult congenital heart disease, aneurysms, angina, angina pectoris, arrhythmias, cardiovascular disease prevention, cardiomyopathies, congestive heart failure, myocardial infarction, pulmonary hypertension, hypertrophic growth, restenosis, stenosis, thrombosis and arteriosclerosis.

A further subject matter of the present invention is the use of the compounds of general formula (I) or (Ia) or (Ib) according to the invention for the treatment and/or prophylaxis of disorders, in particular of the disorders mentioned above.

A further subject matter of the present invention are the compounds according to the invention for use in a method for the treatment and/or prophylaxis of the disorders mentioned above.

A further subject matter of the present invention is the use of the compounds according to the invention in the manufacture of a medicament for the treatment and/or prophylaxis of disorders, in particular the disorders mentioned above.

A further subject matter of the present invention is a method for the treatment and/or prophylaxis of disorders, in particular the disorders mentioned above, using an effective amount of the compounds according to the invention.

Another aspect of the present invention relates to pharmaceutical combinations comprising a compound of general formula (I) or (Ia) or (Ib) according to the invention in combination with at least one or more further active ingredients.

As used herein the term "pharmaceutical combination" refers to a combination of at least one compound of general formula (I) or (Ia) or (Ib) according to the invention as active ingredient together with at least one other active ingredient with or without further ingredients, carrier, diluents and/or solvents.

Another aspect of the present invention relates to pharmaceutical compositions comprising a compound of general formula (I) or (Ia) or (Ib) according to the invention in combination with an inert, nontoxic, pharmaceutically suitable adjuvant.

5 As used herein the term “pharmaceutical composition” refers to a galenic formulation of at least one pharmaceutically active agent together with at least one further ingredient, carrier, diluent and/or solvent.

Another aspect of the present invention relates to the use of the pharmaceutical combinations and/or the pharmaceutical compositions according to the invention for the treatment and/or prophylaxis of disorders, in particular of the disorders mentioned above.

10 Compounds of formula (I) or (Ia) or (Ib) may be administered as the sole pharmaceutical agent or in combination with one or more additional therapeutic agents where the combination causes no unacceptable adverse effects. This pharmaceutical combination includes administration of a single pharmaceutical dosage formulation which contains a compound of formula (I) or (Ia) or (Ib) and one or more additional therapeutic agents, as well as administration of the compound of formula (I) or (Ia) or (Ib) and each additional therapeutic agent in its own separate pharmaceutical dosage formulation. For
15 example, a compound of formula (I) or (Ia) or (Ib) and a therapeutic agent may be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent may be administered in separate dosage formulations.

Where separate dosage formulations are used, the compound of formula (I) or (Ia) or (Ib) and one or more additional therapeutic agents may be administered at essentially the same time (*e.g.*, concurrently)
20 or at separately staggered times (*e.g.*, sequentially).

In particular, the compounds of the present invention may be used in fixed or separate combination with other anti-tumor agents such as alkylating agents, anti-metabolites, plant-derived anti-tumor agents, hormonal therapy agents, topoisomerase inhibitors, camptothecin derivatives, kinase inhibitors, targeted drugs, antibodies, interferons and/or biological response modifiers, anti-angiogenic compounds, and
25 other anti-tumor drugs. In this regard, the following is a non-limiting list of examples of secondary agents that may be used in combination with the compounds of the present invention:

- Alkylating agents include, but are not limited to, nitrogen mustard *N*-oxide, cyclophosphamide, ifosfamide, thiotepa, ranimustine, nimustine, temozolomide, altretamine, apaziquone, brostallicin, bendamustine, carmustine, estramustine, fotemustine, glufosfamide, mafosfamide, bendamustin, and
30 mitolactol; platinum-coordinated alkylating compounds include, but are not limited to, cisplatin, carboplatin, eptaplatin, lobaplatin, nedaplatin, oxaliplatin, and satraplatin;
- Anti-metabolites include, but are not limited to, methotrexate, 6-mercaptopurine riboside, mercaptopurine, 5-fluorouracil alone or in combination with leucovorin, tegafur, doxifluridine, carmofur, cytarabine, cytarabine ocfosfate, enocitabine, gemcitabine, fludarabin, 5-azacitidine,
35 capecitabine, cladribine, clofarabine, decitabine, eflornithine, ethynylcytidine, cytosine arabinoside,

hydroxyurea, melphalan, nelarabine, nolatrexed, ocfosfite, disodium premetrexed, pentostatin, pelitrexol, raltitrexed, triapine, trimetrexate, vidarabine, vincristine, and vinorelbine;

- 5 • Hormonal therapy agents include, but are not limited to, exemestane, Lupron, anastrozole, doxercalciferol, fadrozole, formestane, 11-beta hydroxysteroid dehydrogenase 1 inhibitors, 17-alpha hydroxylase/17,20 lyase inhibitors such as abiraterone acetate, 5-alpha reductase inhibitors such as finasteride and epristeride, anti-estrogens such as tamoxifen citrate and fulvestrant, Trelstar, toremifene, raloxifene, lasofoxifene, letrozole, anti-androgens such as bicalutamide, flutamide, mifepristone, nilutamide, Casodex, and anti-progesterones and combinations thereof;
- 10 • Plant-derived anti-tumor substances include, *e.g.*, those selected from mitotic inhibitors, for example epothilones such as sagopilone, ixabepilone and epothilone B, vinblastine, vinflunine, docetaxel, and paclitaxel;
- 15 • Cytotoxic topoisomerase inhibiting agents include, but are not limited to, aclarubicin, doxorubicin, amonafide, belotecan, camptothecin, 10-hydroxycamptothecin, 9-aminocamptothecin, diflomotecan, irinotecan, topotecan, edotecarin, epimbicin, etoposide, exatecan, gimatecan, lurtotecan, mitoxantrone, pirambicin, pixantrone, rubitecan, sobuzoxane, tafluposide, and combinations thereof;
- 20 • Immunologicals include interferons such as interferon alpha, interferon alpha-2a, interferon alpha-2b, interferon beta, interferon gamma-1a and interferon gamma-n1, and other immune enhancing agents such as L19-IL2 and other IL2 derivatives, filgrastim, lentinan, sizofilan, TheraCys, ubenimex, aldesleukin, alemtuzumab, BAM-002, dacarbazine, daclizumab, denileukin, gentuzumab, ozogamicin, ibritumomab, imiquimod, lenograstim, lentinan, melanoma vaccine (Corixa), molgramostim, sargramostim, tasonermin, teceleukin, thymalasin, tositumomab, Vimlizin, epratuzumab, mitumomab, oregovomab, pentumomab, and Provenge; Merial melanoma vaccine
- 25 • Biological response modifiers are agents that modify defense mechanisms of living organisms or biological responses such as survival, growth or differentiation of tissue cells to direct them to have anti-tumor activity; such agents include, *e.g.*, krestin, lentinan, sizofiran, picibanil, ProMune, and ubenimex;
- 30 • Anti-angiogenic compounds include, but are not limited to, acitretin, aflibercept, angiostatin, aplidine, asentar, axitinib, recentin, bevacizumab, brivanib alaninat, cilengtide, combretastatin, DAST, endostatin, fenretinide, halofuginone, pazopanib, ranibizumab, rebimastat, removab, revlimid, sorafenib, vatalanib, squalamine, sunitinib, telatinib, thalidomide, ukrain, and vitaxin;
- Antibodies include, but are not limited to, trastuzumab, cetuximab, bevacizumab, rituximab, ticilimumab, ipilimumab, lumiliximab, catumaxomab, atacicept, oregovomab, and alemtuzumab;

- VEGF inhibitors such as, *e.g.*, sorafenib, DAST, bevacizumab, sunitinib, recentin, axitinib, aflibercept, telatinib, brivanib alaninate, vatalanib, pazopanib, and ranibizumab; Palladia
- EGFR (HER1) inhibitors such as, *e.g.*, cetuximab, panitumumab, vectibix, gefitinib, erlotinib, and Zactima;
- 5 • HER2 inhibitors such as, *e.g.*, lapatinib, tratuzumab, and pertuzumab;
- mTOR inhibitors such as, *e.g.*, temsirolimus, sirolimus/Rapamycin, and everolimus;
- c-Met inhibitors;
- PI3K and AKT inhibitors;
- CDK inhibitors such as roscovitine and flavopiridol;
- 10 • Spindle assembly checkpoints inhibitors and targeted anti-mitotic agents such as PLK inhibitors, Aurora inhibitors (*e.g.* Hesperadin), checkpoint kinase inhibitors, and KSP inhibitors;
- HDAC inhibitors such as, *e.g.*, panobinostat, vorinostat, MS275, belinostat, and LBH589;
- HSP90 and HSP70 inhibitors;
- Proteasome inhibitors such as bortezomib and carfilzomib;
- 15 • Serine/threonine kinase inhibitors including MEK inhibitors (such as *e.g.* RDEA 119) and Raf inhibitors such as sorafenib;
- Farnesyl transferase inhibitors such as, *e.g.*, tipifarnib;
- Tyrosine kinase inhibitors including, *e.g.*, dasatinib, nilotinib, DAST, bosutinib, sorafenib, bevacizumab, sunitinib, AZD2171, axitinib, aflibercept, telatinib, imatinib mesylate, brivanib alaninate, pazopanib, ranibizumab, vatalanib, cetuximab, panitumumab, vectibix, gefitinib, erlotinib, lapatinib, tratuzumab, pertuzumab, and c-Kit inhibitors; Palladia, masitinib
- 20 • Vitamin D receptor agonists;
- Bcl-2 protein inhibitors such as obatoclax, oblimersen sodium, and gossypol;
- Cluster of differentiation 20 receptor antagonists such as, *e.g.*, rituximab;
- 25 • Ribonucleotide reductase inhibitors such as, *e.g.*, gemcitabine;

- Tumor necrosis apoptosis inducing ligand receptor 1 agonists such as, *e.g.*, mapatumumab;
- 5-Hydroxytryptamine receptor antagonists such as, *e.g.*, rEV598, xaliprode, palonosetron hydrochloride, granisetron, Zindol, and AB-1001;
- Integrin inhibitors including alpha5-beta1 integrin inhibitors such as, *e.g.*, E7820, JSM 6425, volociximab, and endostatin;
- Androgen receptor antagonists including, *e.g.*, nandrolone decanoate, fluoxymesterone, Android, Prost-aid, andromustine, bicalutamide, flutamide, apo-cyproterone, apo-flutamide, chlormadinone acetate, Androcur, Tabi, cyproterone acetate, and nilutamide;
- Aromatase inhibitors such as, *e.g.*, anastrozole, letrozole, testolactone, exemestane, aminoglutethimide, and formestane;
- Matrix metalloproteinase inhibitors;
- Other anti-cancer agents including, *e.g.*, alitretinoin, ampligen, atrasentan bexarotene, bortezomib, bosentan, calcitriol, exisulind, fotemustine, ibandronic acid, miltefosine, mitoxantrone, I-asparaginase, procarbazine, dacarbazine, hydroxycarbamide, pegaspargase, pentostatin, tazaroten, velcade, gallium nitrate, canfosfamide, darinaparsin, and tretinoin.

The compounds of the present invention may also be employed in cancer treatment in conjunction with radiation therapy and/or surgical intervention.

In a further embodiment of the present invention the compounds of the present invention may be used in fixed or separate combination with one or more other active ingredients such as:

- 131I-chTNT, abarelix, abiraterone, aclarubicin, aldesleukin, alemtuzumab, alitretinoin, altretamine, aminoglutethimide, amrubicin, amsacrine, anastrozole, arglabin, arsenic trioxide, asparaginase, azacitidine, basiliximab, BAY 80-6946, BAY 1000394, BAY 86-9766 (RDEA 119), belotecan, bendamustine, bevacizumab, bexarotene, bicalutamide, bisantrene, bleomycin, bortezomib, buserelin, busulfan, cabazitaxel, calcium folinate, calcium levofolinate, capecitabine, carboplatin, carmofur, carmustine, catumaxomab, celecoxib, celmoleukin, cetuximab, chlorambucil, chlormadinone, chlormethine, cisplatin, cladribine, clodronic acid, clofarabine, crisantaspase, cyclophosphamide, cyproterone, cytarabine, dacarbazine, dactinomycin, darbepoetin alfa, dasatinib, daunorubicin, decitabine, degarelix, denileukin diftotox, denosumab, deslorelin, dibrospidium chloride, docetaxel, doxifluridine, doxorubicin, doxorubicin + estrone, eculizumab, edrecolomab, elliptinium acetate, eltrombopag, endostatin, enocitabine, epirubicin, epitiostanol, epoetin alfa, epoetin beta, eptaplatin, eribulin, erlotinib, estradiol, estramustine, etoposide, everolimus, exemestane, fadrozole, filgrastim, fludarabine, fluorouracil, flutamide, formestane, fotemustine, fulvestrant, gallium nitrate, ganirelix,

gefitinib, gemcitabine, gemtuzumab, glutoxim, goserelin, histamine dihydrochloride, histrelin, hydroxycarbamide, I-125 seeds, ibandronic acid, ibritumomab tiuxetan, idarubicin, ifosfamide, imatinib, imiquimod, improsulfan, interferon alfa, interferon beta, interferon gamma, ipilimumab, irinotecan, ixabepilone, lanreotide, lapatinib, lenalidomide, lenograstim, lentinan, letrozole, leuporelin, levamisole, 5 lisuride, lobaplatin, lomustine, lonidamine, masoprocol, medroxyprogesterone, megestrol, melphalan, mepitiostane, mercaptopurine, methotrexate, methoxsalen, Methyl aminolevulinate, methyltestosterone, mifamurtide, miltefosine, miriplatin, mitobronitol, mitoguazone, mitolactol, mitomycin, mitotane, mitoxantrone, nedaplatin, nelarabine, nilotinib, nilutamide, nimotuzumab, nimustine, nitracrine, ofatumumab, omeprazole, oprelvekin, oxaliplatin, p53 gene therapy, paclitaxel, palifermin, palladium-10 103 seed, pamidronic acid, panitumumab, pazopanib, pegaspargase, PEG-epoetin beta (methoxy PEG-epoetin beta), pegfilgrastim, peginterferon alfa-2b, pemetrexed, pentazocine, pentostatin, peplomycin, perfosfamide, picibanil, pirarubicin, plerixafor, plicamycin, poliglusam, polyestradiol phosphate, polysaccharide-K, porfimer sodium, pralatrexate, prednimustine, procarbazine, quinagolide, radium-223 chloride, raloxifene, raltitrexed, ranimustine, razoxane, regorafenib, risedronic acid, rituximab, 15 romidepsin, romiplostim, sargramostim, sipuleucel-T, sizofiran, sobuzoxane, sodium glycididazole, sorafenib, streptozocin, sunitinib, talaporfin, tamibarotene, tamoxifen, tasonermin, teceleukin, tegafur, tegafur + gimeracil + oteracil, temoporfin, temozolomide, temsirolimus, teniposide, testosterone, tetrofosmin, thalidomide, thiotepa, thymalfasin, tioguanine, tocilizumab, topotecan, toremifene, tositumomab, trabectedin, trastuzumab, treosulfan, tretinoin, trilostane, triptorelin, trofosfamide, 20 tryptophan, ubenimex, valrubicin, vandetanib, vapreotide, vemurafenib, vinblastine, vincristine, vindesine, vinflunine, vinorelbine, vorinostat, vorozole, yttrium-90 glass microspheres, zinostatin, zinostatin stimalamer, zoledronic acid, zorubicin.

Generally, the use of cytotoxic and/or cytostatic agents in combination with a compound or composition of the present invention will serve to:

- 25 (1) yield better efficacy in reducing the growth of a tumor or even eliminate the tumor as compared to administration of either agent alone,
- (2) provide for the administration of lesser amounts of the administered chemotherapeutic agents,
- (3) provide for a chemotherapeutic treatment that is well tolerated in the patient with fewer deleterious pharmacological complications than observed with single agent chemotherapies and 30 certain other combined therapies,
- (4) provide for treating a broader spectrum of different cancer types in mammals, especially humans,
- (5) provide for a higher response rate among treated patients,
- (6) provide for a longer survival time among treated patients compared to standard chemotherapy 35 treatments,
- (7) provide a longer time for tumor progression, and/or

- (8) yield efficacy and tolerability results at least as good as those of the agents used alone, compared to known instances where other cancer agent combinations produce antagonistic effects.

Furthermore, the compounds of formula (I) or (Ia) or (Ib) may be utilized, as such or in compositions, in research and diagnostics, or as analytical reference standards, and the like, which are well known in the art.

The compounds according to the invention can act systemically and/or locally. For this purpose, they can be administered in a suitable way, such as, for example, by the oral, parenteral, pulmonal, nasal, sublingual, lingual, buccal, rectal, dermal, transdermal, conjunctival or otic route, or as an implant or stent.

- For these administration routes, it is possible to administer the compounds according to the invention in suitable application forms.

Suitable for oral administration are administration forms which work as described in the prior art and deliver the compounds according to the invention rapidly and/or in modified form, which comprise the compounds according to the invention in crystalline and/or amorphous and/or dissolved form, such as, for example, tablets (coated or uncoated, for example tablets provided with enteric coatings or coatings whose dissolution is delayed or which are insoluble and which control the release of the compound according to the invention), tablets which rapidly decompose in the oral cavity, or films/wafers, films/lyophilizates, capsules (for example hard or soft gelatin capsules), sugar-coated tablets, granules, pellets, powders, emulsions, suspensions, aerosols or solutions.

- Parenteral administration can take place with avoidance of an absorption step (for example intravenously, intraarterially, intracardially, intraspinally or intralumbally) or with inclusion of absorption (for example intramuscularly, subcutaneously, intracutaneously, percutaneously or intraperitoneally). Administration forms suitable for parenteral administration are, inter alia, preparations for injection and infusion in the form of solutions, suspensions, emulsions, lyophilizates or sterile powders.

Examples suitable for the other administration routes are pharmaceutical forms for inhalation (inter alia powder inhalers, nebulizers), nasal drops/solutions/sprays; tablets to be administered lingually, sublingually or buccally, films/wafers or capsules, suppositories, preparations for the eyes or ears, vaginal capsules, aqueous suspensions (lotions, shaking mixtures), lipophilic suspensions, ointments, creams, transdermal therapeutic systems (such as plasters, for example), milk, pastes, foams, dusting powders, implants or stents.

The compounds according to the invention can be converted into the stated administration forms. This can take place in a manner known per se by mixing with inert, nontoxic, pharmaceutically suitable adjuvants. These adjuvants include, inter alia, carriers (for example microcrystalline cellulose, lactose,

mannitol), solvents (for example liquid polyethylene glycols), emulsifiers and dispersants or wetting agents (for example sodium dodecyl sulphate, polyoxysorbitan oleate), binders (for example polyvinylpyrrolidone), synthetic and natural polymers (for example albumin), stabilizers (for example antioxidants, such as, for example, ascorbic acid), colorants (for example inorganic pigments, such as, for example, iron oxides) and flavour- and/or odour-masking agents.

The present invention furthermore provides medicaments comprising at least one compound according to the invention, usually together with one or more inert, nontoxic, pharmaceutically suitable adjuvants, and their use for the purposes mentioned above.

When the compounds of the present invention are administered as pharmaceuticals, to humans or animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1% to 99,5% (more preferably 0.5% to 90%) of active ingredient in combination with one or more inert, nontoxic, pharmaceutically suitable adjuvants.

Regardless of the route of administration selected, the compounds of the invention of general formula (I) or (Ia) and/or the pharmaceutical composition of the present invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels and time course of administration of the active ingredients in the pharmaceutical compositions of the invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient without being toxic to the patient.

Materials and Methods:

The percentage data in the following tests and examples are percentages by weight unless otherwise indicated; parts are parts by weight. Solvent ratios, dilution ratios and concentration data of liquid/liquid solutions are in each case based on volume.

Examples were tested in selected biological assays one or more times. When tested more than once, data are reported as either average values or as median values, wherein

- the average value, also referred to as the arithmetic mean value, represents the sum of the values obtained divided by the number of times tested, and
- the median value represents the middle number of the group of values when ranked in ascending or descending order. If the number of values in the data set is odd, the median is the middle value. If the number of values in the data set is even, the median is the arithmetic mean of the two middle values.

•Examples were synthesized one or more times. When synthesized more than once, data from biological assays represent average values or median values calculated utilizing data sets obtained from testing of one or more synthetic batch.

5 The in vitro pharmacological and physico-chemical properties of the compounds can be determined according to the following assays and methods.

1. CDK9/CycT1 kinase assay:

CDK9/CycT1 -inhibitory activity of compounds of the present invention was quantified employing the CDK9/CycT1 TR-FRET assay as described in the following paragraphs:

10 Recombinant full-length His-tagged human CDK9 and CycT1, expressed in insect cells and purified by Ni-NTA affinity chromatography, were purchased from Invitrogen (Cat. No PV4131). As substrate for the kinase reaction biotinylated peptide biotin-Ttds-YISPLKSPYKISEG (C-terminus in amid form) was used which can be purchased e.g. form the company JERINI Peptide Technologies (Berlin, Germany).

For the assay 50 nl of a 100fold concentrated solution of the test compound in DMSO was pipetted into a
15 black low volume 384well microtiter plate (Greiner Bio-One, Frickenhausen, Germany), 2 µl of a solution of CDK9/CycT1 in aqueous assay buffer [50 mM Tris/HCl pH 8.0, 10 mM MgCl₂, 1.0 mM dithiothreitol, 0.1 mM sodium ortho-vanadate, 0.01% (v/v) Nonidet-P40 (Sigma)] were added and the mixture was incubated for 15 min at 22°C to allow pre-binding of the test compounds to the enzyme before the start of the kinase reaction. Then the kinase reaction was started by the addition of 3 µl of a
20 solution of adenosine-tri-phosphate (ATP, 16.7 µM => final conc. in the 5 µl assay volume is 10 µM) and substrate (1.67 µM => final conc. in the 5 µl assay volume is 1 µM) in assay buffer and the resulting mixture was incubated for a reaction time of 25 min at 22°C. The concentration of CDK9/CycT1 was adjusted depending of the activity of the enzyme lot and was chosen appropriate to have the assay in the linear range, typical concentrations were in the range of 1 µg/mL. The reaction was stopped by the
25 addition of 5 µl of a solution of TR-FRET detection reagents (0.2 µM streptavidine-XL665 [Cisbio Bioassays, Codolet, France] and 1 nM anti-RB(pSer807/pSer811)-antibody from BD Pharmingen [# 558389] and 1.2 nM LANCE EU-W1024 labeled anti-mouse IgG antibody [Perkin-Elmer, product no. AD0077]) in an aqueous EDTA-solution (100 mM EDTA, 0.2 % (w/v) bovine serum albumin in 100 mM HEPES/NaOH pH 7.0).

30 The resulting mixture was incubated 1 h at 22°C to allow the formation of complex between the phosphorylated biotinylated peptide and the detection reagents. Subsequently the amount of phosphorylated substrate was evaluated by measurement of the resonance energy transfer from the Eu-chelate to the streptavidine-XL. Therefore, the fluorescence emissions at 620 nm and 665 nm after excitation at 350 nm was measured in a HTRF reader, e.g. a Rubystar (BMG Labtechnologies,
35 Offenburg, Germany) or a Viewlux (Perkin-Elmer). The ratio of the emissions at 665 nm and at 622 nm was taken as the measure for the amount of phosphorylated substrate. The data were normalised (enzyme reaction without inhibitor = 0 % inhibition, all other assay components but no enzyme = 100 %

inhibition). Usually the test compounds were tested on the same microtiterplate in 11 different concentrations in the range of 20 μ M to 0.1 nM (20 μ M, 5.9 μ M, 1.7 μ M, 0.51 μ M, 0.15 μ M, 44 nM, 13 nM, 3.8 nM, 1.1 nM, 0.33 nM and 0.1 nM, the dilution series prepared separately before the assay on the level of the 100fold concentrated solutions in DMSO by serial 1:3.4 dilutions) in duplicate values for each concentration and IC₅₀ values were calculated by a 4 parameter fit.

2. CDK2/CycE kinase assay:

CDK2/CycE -inhibitory activity of compounds of the present invention was quantified employing the CDK2/CycE TR-FRET assay as described in the following paragraphs:

Recombinant fusion proteins of GST and human CDK2 and of GST and human CycE, expressed in insect cells (Sf9) and purified by Glutathion-Sepharose affinity chromatography, were purchased from ProQinase GmbH (Freiburg, Germany). As substrate for the kinase reaction biotinylated peptide biotin-Ttds-YISPLKSPYKISEG (C-terminus in amid form) was used which can be purchased e.g. from the company JERINI Peptide Technologies (Berlin, Germany).

For the assay 50 nl of a 100fold concentrated solution of the test compound in DMSO was pipetted into a black low volume 384well microtiter plate (Greiner Bio-One, Frickenhausen, Germany), 2 μ l of a solution of CDK2/CycE in aqueous assay buffer [50 mM Tris/HCl pH 8.0, 10 mM MgCl₂, 1.0 mM dithiothreitol, 0.1 mM sodium ortho-vanadate, 0.01% (v/v) Nonidet-P40 (Sigma)] were added and the mixture was incubated for 15 min at 22°C to allow pre-binding of the test compounds to the enzyme before the start of the kinase reaction. Then the kinase reaction was started by the addition of 3 μ l of a solution of adenosine-tri-phosphate (ATP, 16.7 μ M => final conc. in the 5 μ l assay volume is 10 μ M) and substrate (1.25 μ M => final conc. in the 5 μ l assay volume is 0.75 μ M) in assay buffer and the resulting mixture was incubated for a reaction time of 25 min at 22°C. The concentration of CDK2/CycE was adjusted depending of the activity of the enzyme lot and was chosen appropriate to have the assay in the linear range, typical concentrations were in the range of 130 ng/mL. The reaction was stopped by the addition of 5 μ l of a solution of TR-FRET detection reagents (0.2 μ M streptavidine-XL665 [Cisbio Bioassays, Codolet, France] and 1 nM anti-RB(pSer807/pSer811)-antibody from BD Pharmingen [# 558389] and 1.2 nM LANCE EU-W1024 labeled anti-mouse IgG antibody [Perkin-Elmer, product no. AD0077]) in an aqueous EDTA-solution (100 mM EDTA, 0.2 % (w/v) bovine serum albumin in 100 mM HEPES/NaOH pH 7.0).

The resulting mixture was incubated 1 h at 22°C to allow the formation of complex between the phosphorylated biotinylated peptide and the detection reagents. Subsequently the amount of phosphorylated substrate was evaluated by measurement of the resonance energy transfer from the Eu-chelate to the streptavidine-XL. Therefore, the fluorescence emissions at 620 nm and 665 nm after excitation at 350 nm was measured in a TR-FRET reader, e.g. a Rubystar (BMG Labtechnologies, Offenburg, Germany) or a Viewlux (Perkin-Elmer). The ratio of the emissions at 665 nm and at 622 nm was taken as the measure for the amount of phosphorylated substrate. The data were normalised (enzyme

reaction without inhibitor = 0 % inhibition, all other assay components but no enzyme = 100 % inhibition). Usually the test compounds were tested on the same microtiterplate in 11 different concentrations in the range of 20 μ M to 0.1 nM (20 μ M, 5.9 μ M, 1.7 μ M, 0.51 μ M, 0.15 μ M, 44 nM, 13 nM, 3.8 nM, 1.1 nM, 0.33 nM and 0.1 nM, the dilution series prepared separately before the assay on the level of the 100fold concentrated solutions in DMSO by serial 1:3.4 dilutions) in duplicate values for each concentration and IC50 values were calculated by a 4 parameter fit.

3. Proliferation Assay:

Cultivated tumour cells (NCI-H460, human non-small cell lung carcinoma cells, ATCC HTB-177; DU145, hormone-independent human prostate carcinoma cells, ATCC HTB-81; HeLa-MaTu, human cervical carcinoma cells, EPO-GmbH, Berlin; HeLa-MaTu-ADR, multidrug-resistant human cervical carcinoma cells, EPO-GmbH, Berlin; HeLa human cervical tumour cells, ATCC CCL-2; Caco-2 human colorectal carcinoma, ATCC HTB-37; B16F10 mouse melanoma cells, ATCC CRL-6475) were plated at a density of 5000 cells/well (DU145, HeLa-MaTu-ADR), 3000 cells/well (NCI-H460, HeLa-MaTu, HeLa), 1500 cells/well (Caco-2), or 1000 cells/well (B16F10) in a 96-well multititer plate in 200 μ L of their respective growth medium supplemented 10% fetal calf serum. After 24 hours, the cells of one plate (zero-point plate) were stained with crystal violet (see below), while the medium of the other plates was replaced by fresh culture medium (200 μ L), to which the test substances were added in various concentrations (0 μ M, as well as in the range of 0.001-10 μ M; the final concentration of the solvent dimethyl sulfoxide was 0.5%). The cells were incubated for 4 days in the presence of test substances. Cell proliferation was determined by staining the cells with crystal violet: the cells were fixed by adding 20 μ L/measuring point of an 11% glutaric aldehyde solution for 15 minutes at room temperature. After three washing cycles of the fixed cells with water, the plates were dried at room temperature. The cells were stained by adding 100 μ L/measuring point of a 0.1% crystal violet solution (pH 3.0). After three washing cycles of the stained cells with water, the plates were dried at room temperature. The dye was dissolved by adding 100 μ L/measuring point of a 10% acetic acid solution. The absorbance was determined by photometry at a wavelength of 595 nm. The change of cell number, in percent, was calculated by normalization of the measured values to the absorbance values of the zero-point plate (=0%) and the absorbance of the untreated (0 μ M) cells (=100%). The IC50 values were determined by means of a 4 parameter fit.

4. Equilibrium Shake Flask Solubility Assay:

The thermodynamic solubility of compounds in water was determined by an equilibrium shake flask method (see for example: E.H. Kerns, L. Di: Drug-like Properties: Concepts, Structure Design and Methods, 276-286, Burlington, MA, Academic Press, **2008**). A saturated solution of the drug was prepared and the solution was mixed for 24 h to ensure that equilibrium was reached. The solution was centrifuged to remove the insoluble fraction and the concentration of the compound in solution was determined using a standard calibration curve.

To prepare the sample, 2 mg solid compound was weighed in a 4 mL glass vial. 1 mL phosphate buffer pH 6.5 was added. The suspension was stirred for 24 hrs at room temperature. The solution was centrifuged afterwards. To prepare the sample for the standard calibration, 2 mg solid sample was dissolved in 30 mL acetonitrile. After sonification the solution was diluted with water to 50 mL. Sample and standards were quantified by HPLC with UV-detection. For each sample two injection volumes (5 and 50 µl) in triplicates were made. Three injection volumes (5 µl, 10 µl and 20 µl) were made for the standard.

Chromatographic conditions:

HPLC column: Xterra MS C18 2.5 µm 4.6 x 30 mm

Injection volume: Sample: 3x5µl and 3x50µl
Standard: 5µl, 10µl, 20µl

Flow: 1.5mL/min

Mobile phase: acidic gradient:
A: Water / 0.01% TFA
B: Acetonitrile / 0.01% TFA

0 min → 95%A 5%B

0-3 min → 35%A 65%B, linear gradient

3-5 min → 35%A 65%B, isocratic

5-6 min → 95%A 5%B, isocratic

UV detector: wavelength near the absorption maximum (between 200 and 400nm)

The areas of sample- and standard injections as well as the calculation of the solubility value (in mg/l) were determined by using HPLC software (Waters Empower 2 FR).

5. Carbonic anhydrase Assay

The principle of the assay is based on the hydrolysis of 4-nitrophenyl acetate by carbonic anhydrases (Pocker & Stone, Biochemistry, 1967, 6, 668), with subsequent photometric determination of the dye product 4-nitrophenolate at 400 nm by means of a 96-channel spectral photometer.


- 5 2 microL of the test compounds, dissolved in DMSO (100-fold final concentration), in a concentration range of 0.03-10 micromol/L (final), was pipetted as quadruplicates into the wells of a 96-hole microtiter plate. Wells that contained the solvent without test compounds were used as reference values (1. Wells without carbonic anhydrase for correction of the non-enzymatic hydrolysis of the substrate, and 2. Wells with carbonic anhydrase for determining the activity of the non-inhibited enzyme).
- 10 188 microL of assay buffer (10 millimol/L of Tris/HCl, pH 7.4, 80 millimol/L of NaCl), with or without 3 units/well of carbonic anhydrase-1 [= human carbonic anhydrase-1 (Sigma, #C4396)] in order to determine carbonic anhydrase-1 inhibition or 3 units/well of carbonic anhydrase-2 [= human carbonic anhydrase-2 (Sigma, #C6165)] for measuring carbonic anhydrase-2 inhibition, was pipetted into the wells of the microtiter plate. The enzymatic reaction was started by the addition of 10 microL of the
- 15 substrate solution (1 millimol/L of 4-nitrophenyl acetate (Fluka #4602), dissolved in anhydrous acetonitrile (final substrate concentration: 50 micromol/L). The plate was incubated at room temperature for 15 minutes. Absorption was measured by photometry at a wavelength of 400 nm. The enzyme inhibition was calculated after the measured values were normalized to the absorption of the reactions in the wells without enzyme (=100% inhibition) and to the absorption of reactions in the wells with non-
- 20 inhibited enzyme (=0% inhibition). IC50 values were determined by means of a 4 parameter fit.

Preparative Examples

Syntheses of compounds

The syntheses of the inventive disubstituted triazines according to the present invention is preferably carried out according to one of the general synthetic sequences, shown in schemes 1, 2, 3 or 4 below:

Scheme 1:



Scheme 1

In the first step 2,4-dichloro-1,3,5-triazine (**1**) is reacted with suitable anilines (**2**) to give the corresponding 4-chloro-N-phenyl-1,3,5-triazin-2-amines (**3**). The reaction is carried out with one equivalent of the aniline (**2**) in an inert solvent like DMF, THF, DME, dioxane or an alcohol like isopropanol, or mixtures of such solvents. Preferably, the reaction is carried out at a temperature below 0°C in such a way that the reaction mixture is kept homogenous. Preferred conditions use an additional base like triethylamine or N,N-diisopropylethylamine.

In the second step the intermediate 4-chloro-N-phenyl-1,3,5-triazin-2-amine (**3**) is reacted with a boronic acid derivative $R^2-B(OR)_2$ (**4**) to give compounds of formula (**5**). The boronic acid derivative (**4**) may be a boronic acid ($R = -H$) or an ester of the boronic acid, e.g. its isopropyl ester ($R = -CH(CH_3)_2$), preferably an ester derived from pinacol in which the boronic acid intermediate forms a 2-aryl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane ($R-R = -C(CH_3)_2-C(CH_3)_2-$).

The coupling reaction is catalyzed by Pd catalysts, e.g. by Pd(0) catalysts like tetrakis(triphenylphosphine)palladium(0) $[Pd(PPh_3)_4]$, tris(dibenzylideneacetone)di-palladium(0) $[Pd_2(dba)_3]$, or by Pd(II) catalysts like dichlorobis(triphenylphosphine)-palladium(II) $[Pd(PPh_3)_2Cl_2]$, palladium(II) acetate and triphenylphosphine or by [1,1'-bis(diphenylphosphino)ferrocene]palladium dichloride $[Pd(dppf)Cl_2]$.


The reaction is preferably carried out in a mixture of a solvent like 1,2-dimethoxyethane, dioxane, DMF, DME, THF or isopropanol with water and in the presence of a base like aqueous potassium carbonate, aqueous sodium bicarbonate or potassium phosphate.

Deprotection of compounds of formula (**5**) gives the corresponding N-unprotected sulfoximines of formula (**6**). The deprotection is preferably carried out with sodium ethanolate in ethanol at 60°C.

N-unprotected sulfoximines of formula (**6**) may be reacted to to give N-functionalized derivatives of formula (**I**).

Scheme 2

Another synthesis route to N-unprotected sulfoximines of formula (6) is shown in Scheme 2.



Scheme 2

In the first step 2,4-dichloro-1,3,5-triazine (**1**) is reacted with suitable anilines of formula (**13**) to give the corresponding 4-chloro-N-phenyl-1,3,5-triazin-2-amines of formula (**14**). The reaction is carried out with one equivalent of the aniline (**13**) in an inert solvent like DMF, THF, DME, dioxane or an alcohol like isopropanol, or mixtures of such solvents. Preferably, the reaction is carried out at a temperature below 0°C in such a way that the reaction mixture is kept homogenous. Preferred conditions use an additional base like triethylamine or N,N-diisopropylethylamine.

In the second step the intermediate 4-chloro-N-phenyl-1,3,5-triazin-2-amine of formula (**14**) is reacted with a boronic acid derivative $R^2-B(OR)_2$ (**4**) to give compounds of formula (**15**). The boronic acid derivative (**4**) may be a boronic acid ($R = -H$) or an ester of the boronic acid, e.g. its isopropyl ester ($R = -CH(CH_3)_2$), preferably an ester derived from pinacol in which the boronic acid intermediate forms a 2-aryl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane ($R-R = -C(CH_3)_2-C(CH_3)_2-$).


The coupling reaction is catalyzed by Pd catalysts, e.g. by Pd(0) catalysts like tetrakis(triphenylphosphine)palladium(0) [Pd(PPh₃)₄], tris(dibenzylideneacetone)di-palladium(0) [Pd₂(dba)₃], or by Pd(II) catalysts like dichlorobis(triphenylphosphine)-palladium(II) [Pd(PPh₃)₂Cl₂], palladium(II) acetate and triphenylphosphine or by [1,1'-bis(diphenylphosphino)ferrocene]palladium dichloride [Pd(dppf)Cl₂].

The reaction is preferably carried out in a mixture of a solvent like 1,2-dimethoxyethane, dioxane, DMF, DME, THF, or isopropanol with water and in the presence of a base like aqueous potassium carbonate, aqueous sodium bicarbonate or potassium phosphate.

Finally, the compound of formula (15) is reacted with sodium azide in trichloromethane and sulfuric acid at 45°C to give the N-unprotected sulfoximine of formula (6) (see for example: a) H. R. Bentley et al, J. Chem. Soc. **1952**, 1572; b) C. R. Johnson et al, J. Am. Chem. Soc. **1970**, 92, 6594; c) Satzinger et al, Angew. Chem. **1971**, 83, 83).

Scheme 3

Another synthesis route to N-cyanosulfoximines of formula (20), which can also be converted to unprotected sulfoximines of formula (6), is shown in Scheme 3.



Scheme 3

10 In the first step 2,4-dichloro-1,3,5-triazine (**1**) is reacted with suitable anilines of formula (**16**) to give the corresponding 4-chloro-N-phenyl-1,3,5-triazin-2-amines of formula (**17**). The reaction is carried out with

one equivalent of the aniline of formula (2) in an inert solvent like DMF, THF, DME, dioxane or an alcohol like isopropanol, or mixtures of such solvents. Preferably, the reaction is carried out at a temperature below 0°C in such a way that the reaction mixture is kept homogenous. Preferred conditions use an additional base like triethylamine or N,N-diisopropylethylamine.

- 5 In the second step the intermediate 4-chloro-N-phenyl-1,3,5-triazin-2-amine of formula (17) is reacted with a boronic acid derivative $R^2-B(OR)_2$ (4) to give compounds of formula (18). The boronic acid derivative (4) may be a boronic acid ($R = -H$) or an ester of the boronic acid, e.g. its isopropyl ester ($R = -CH(CH_3)_2$), preferably an ester derived from pinacol in which the boronic acid intermediate forms a 2-aryl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane ($R-R = -C(CH_3)_2-C(CH_3)_2-$).
- 10 The coupling reaction is catalyzed by Pd catalysts, e.g. by Pd(0) catalysts like tetrakis(triphenylphosphine)palladium(0) $[Pd(PPh_3)_4]$, tris(dibenzylideneacetone)di-palladium(0) $[Pd_2(dba)_3]$, or by Pd(II) catalysts like dichlorobis(triphenylphosphine)-palladium(II) $[Pd(PPh_3)_2Cl_2]$, palladium(II) acetate and triphenylphosphine or by [1,1'-bis(diphenylphosphino)ferrocene]palladium dichloride $[Pd(dppf)Cl_2]$.
- 15 The reaction is preferably carried out in a mixture of a solvent like 1,2-dimethoxyethane, dioxane, DMF, DME, THF, or isopropanol with water and in the presence of a base like aqueous potassium carbonate, aqueous sodium bicarbonate or potassium phosphate.


- In the next step, the sulfide of formula (18) is reacted with cyanogen amine as a nitrogen source to give the corresponding N-cyanosulfilimine of formula (19). Preferably, the reaction is carried out using NBS and potassium *tert*-butoxide in methanol at room temperature (see for example: a) C. Bolm et al, Org. Lett. **2007**, 9, 3809). Even more preferred is the use of iodobenzenediacetate in DCM at room temperature (see for example: a) J.M. Babcock, US 2009/0023782).
- 20

- Finally, N-cyanosulfilimine of formula (19) is oxidized to the corresponding N-cyanosulfoximine of formula (20). The reaction is preferably carried out using mCPBA and potassium carbonate in ethanol at room temperature (see for example: a) C. Bolm et al, Org. Lett. **2007**, 9, 3809). Even more preferred is the use of potassium permanganate in acetone at 50°C (see for example: a) C. Bolm et al, Adv. Synth. Catal. **2010**, 352, 309).
- 25

- The N-cyano group of compound (20) is cleaved upon treatment with TFAA affording the corresponding N-trifluoroacetylsulfoximine which is converted into the NH-free sulfoximine of formula (6) by methanolysis of the trifluoroacetyl moiety (see for example: C. Bolm et al, Org. Lett. **2007**, 9(19), 3809).
- 30

Scheme 4:

Another synthesis route to N-unprotected sulfoximines of formula (6) is shown in Scheme 4.



Scheme 4

In the first step the intermediate 4-chloro-N-phenyl-1,3,5-triazin-2-amine (**3**) is reacted with a suitable ortho-fluorine boronic acid derivative of formula (**21**) to give a compound of formula (**22**). The boronic acid derivative of formula (**21**) may be a boronic acid (R = -H) or an ester of the boronic acid, e.g. its isopropyl ester (R = -CH(CH₃)₂), or an ester derived from pinacol in which the boronic acid intermediate forms a 2-aryl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (R-R = -C(CH₃)₂-C(CH₃)₂-).

The coupling reaction is catalyzed by Pd catalysts, e.g. by Pd(0) catalysts like tetrakis(triphenylphosphine)palladium(0) [Pd(PPh₃)₄], tris(dibenzylideneacetone)di-palladium(0) [Pd₂(dba)₃], or by Pd(II) catalysts like dichlorobis(triphenylphosphine)-palladium(II) [Pd(PPh₃)₂Cl₂], palladium(II) acetate and triphenylphosphine or by [1,1'-bis(diphenylphosphino)ferrocene]palladium dichloride [Pd(dppf)Cl₂].

The reaction is preferably carried out in a mixture of a solvent like 1,2-dimethoxyethane, dioxane, DMF, DME, THF, or isopropanol with water and in the presence of a base like aqueous potassium carbonate, aqueous sodium bicarbonate or potassium phosphate.

In the second step the ortho-fluorine of the substituent in 4-position of the compound of formula (**22**) is replaced by a suitable alkoxy group -OR⁸. The reaction is preferably carried out by adding at least two equivalents of sodium hydride to a solution of compound (**22**) in alcohol (**23**) to give the desired N-unprotected sulfoximines of formula (**6**). The reactions are run at a temperature of 60°C or in the temperature range between 50°C and 70°C.

Preparation of compounds:


Abbreviations used in the description of the chemistry and in the Examples that follow are:

CDCl₃ (deuterated chloroform); cHex (cyclohexane); DCM (dichloromethane); DIPEA (di-isopropylethylamine); DME (1,2-dimethoxyethane), DMF (dimethylformamide); DMSO (dimethyl sulfoxide); eq (equivalent); ES (electrospray); EtOAc (ethyl acetate); EtOH (ethanol); iPrOH (isopropanol); mCPBA (meta-chloroperoxybenzoic acid), MeOH (methanol); MS (mass spectrometry); NBS (N-bromosuccinimide), NMR (nuclear magnetic resonance); Pd(dppf)Cl₂ ([1,1'-bis(diphenylphosphino)ferrocene]dichloro palladium(II) complex with dichloromethane); iPrOH (isopropanol); RT (room temperature); sat. aq. (saturated aqueous); SiO₂ (silica gel); TFA (trifluoroacetic acid); TFAA (trifluoroacetic anhydride), THF (tetrahydrofuran).

The IUPAC names of the examples were generated using the program 'ACD/Name batch version 12.01 from ACD LABS.


Example 1:

(rac)-Ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate




Preparation of Intermediate 1.1:

1-[(Methylsulfanyl)methyl]-3-nitrobenzene




Sodium methanethiolate (13.5 g; 192 mmol) was added in two portions to a stirred solution of 1-(chloromethyl)-3-nitrobenzene (30.0 g; 175 mmol; Aldrich) in ethanol (360 mL) at -15°C. The cold bath was removed and the batch was stirred at room temperature for 3 hours. The batch was diluted with brine and extracted with ethyl acetate (2x). The combined organic phases were washed with water, dried (sodium sulfate), filtered and concentrated to give the desired product (32.2 g) that was used without further purification.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.18 (m, 1H), 8.11 (m, 1H), 7.66 (m, 1H), 7.50 (m, 1H), 3.75 (s, 2H), 2.01 (s, 3H).

Preparation of Intermediate 1.2:**(rac)-1-[(Methylsulfinyl)methyl]-3-nitrobenzene**


Iron(III)chloride (0.55 g; 3.4 mmol) was added to a solution of 1-[(methylsulfonyl)methyl]-3-nitrobenzene (21.6 g; 117.9 mmol) in acetonitrile (280 mL) and the batch was stirred at room temperature for 10 minutes. Periodic acid (28.8 g; 126.1 mmol) was added under stirring in one portion and the temperature was kept below 30°C by cooling. The batch was stirred at room temperature for 90 minutes before it was added to a stirred solution of sodium thiosulfate pentahydrate (163 g; 660 mmol) in ice water (1500 mL). The batch was saturated with solid sodium chloride and extracted with THF (2x). The combined organic phases were washed with brine, dried (sodium sulfate), filtered and concentrated. The residue was purified by chromatography (DCM / ethanol 95:5) to give the desired product (16.6 g; 83.1 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.21 (m, 1H), 8.17 (m, 1H), 7.67 (m, 1H), 7.58 (m, 1H), 4.10 (d, 1H), 3.97 (d, 1H), 2.53 (s, 3H).

Preparation of Intermediate 1.3:**(rac)-2,2,2-Trifluoro-N-[methyl(3-nitrobenzyl)oxido-λ⁶-sulfanylidene]acetamide**

To a suspension of (rac)-1-[(methylsulfinyl)methyl]-3-nitrobenzene (16.6 g; 83.1 mmol), trifluoroacetamide (18.8 g; 166.1 mmol), magnesium oxide (13.4 g; 332.3 mmol) and rhodium(II)-acetate dimer (1.7 g; 8.3 mmol) in DCM (2290 mL) was added iodobenzene diacetate (40.1 g; 124.6 mmol) at room temperature. The batch was stirred for 16 hours at room temperature, filtered and concentrated. The residue was purified by chromatography (DCM / ethanol 97:3) to give the desired product (25.6 g; 82.4 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.36 (m, 1H), 8.31 (m, 1H), 7.80 (m, 1H), 7.69 (m, 1H), 4.91 (d, 1H), 4.79 (d, 1H), 3.28 (s, 3H).

Preparation of Intermediate 1.4:**(rac)-1-[(S-Methylsulfonimidoyl)methyl]-3-nitrobenzene**


Potassium carbonate (56.9 g; 411.8 mmol) was added to a solution of (rac)-2,2,2-trifluoro-N-[methyl(3-nitrobenzyl)oxido-λ⁶-sulfanylidene]acetamide (25.6 g; 82.4 mmol) in methanol (1768 mL) at room

temperature. The batch was stirred for 1 hour at room temperature before it was diluted with ethyl acetate and brine. After extraction with ethyl acetate (2x) the combined organic phases were dried (sodium sulfate), filtered and concentrated to give the desired product (13.9 g; 65.1 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.29 (m, 2H), 7.79 (m, 1H), 7.63 (m, 1H), 4.47 (d, 1H), 4.34 (d, 1H), 2.99 (s, 3H), 2.66 (br, 1H).

Preparation of Intermediate 1.5:

(rac)-Ethyl [methyl(3-nitrobenzyl)oxido- λ^6 -sulfanylidene]carbamate




10 Ethyl chlorocarbonate (8.1 mL; 84.6 mmol) was added dropwise to a stirred solution of (rac)-1-[(S-methylsulfonimidoyl)methyl]-3-nitrobenzene (13.9 g; 65.1 mmol) in pyridine (615 mL) at 0°C. The batch was slowly warmed to room temperature. After 24 hours the batch was concentrated and the residue was dissolved in ethyl acetate and washed with brine. The organic phase was filtered using a Whatman filter and concentrated to give the desired product (19.7 g) that was used without further purification.

¹H NMR (400MHz, d₆-CDCl₃, 300K) δ = 8.30 (m, 2H), 7.81 (m, 1H), 7.64 (m, 1H), 4.88 (d, 1H), 4.79 (d, 1H), 4.18 (q, 2H), 3.07 (s, 3H), 1.31 (tr, 3H).

Preparation of Intermediate 1.6:

20 **(rac)-Ethyl [(3-aminobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate**




25 Titanium(III)chloride solution (about 15% in about 10% hydrochloric acid, 118 mL; Merck Schuchardt OHG) was added to a stirred solution of (rac)-ethyl [methyl(3-nitrobenzyl)oxido- λ^6 -sulfanylidene]carbamate (5.0 g; 17.5 mmol) in THF (220 mL) at room temperature. The batch was stirred for 18 hours. By adding 2N sodium hydroxide solution the pH value of the reaction mixture, that was cooled with an ice bath, was raised to 8. The batch was saturated with solid sodium chloride and extracted with ethyl acetate (3x). The combined organic phases were washed with brine, dried (sodium sulfate), filtered and concentrated to give the desired product (4.2 g) that was used without further purification.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 7.00 (m, 1H), 6.53 (m, 3H), 5.18 (br, 2H), 4.62 (s, 2H), 3.95 (m, 2H), 3.08 (s, 3H), 1.13 (tr, 3H).

Preparation of Intermediate 1.7:

(rac)-Ethyl [{3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido- λ^6 -sulfanylidene]-carbamate :



5

DIPEA (3.1 mL; 17.8 mmol) was added to a stirred solution of 2,4-dichloro-1,3,5-triazine (1.34 g; 8.9 mmol) in THF / i-PrOH (1:1; 18 mL) at -40°C. Then a solution of (rac)-ethyl [(3-aminobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate (2.29 g; 8.9 mmol) in THF / i-PrOH (1:1; 9 mL) was added at this temperature. Under stirring the temperature of the reaction mixture was slowly raised over 3 hours to 0°C. The batch was concentrated to give the crude product (4.9 g) that was used without further purification.

10

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A1 = H ₂ O + 0.1% HCOOH
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A1 + B1 = C:\MassLynx\Mass_160_1000.flp
Retention:	0.88 min
MS(ES+):	m/z = 370 [M+H]

Preparation of end product:

15 A batch with crude (rac)-ethyl [{3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido- λ^6 -sulfanylidene]carbamate (400 mg), (4-fluoro-2-methoxyphenyl)boronic acid (276 mg; 1.62 mmol; Aldrich) and tetrakis(triphenylphosphin)palladium(0) (187 mg; 0.16 mmol) in 1,2-dimethoxyethane (5.0 mL) and 2M solution of potassium carbonate (1.1 mL) was degassed using argon. The batch was stirred under argon for 80 minutes at 100°C. After cooling the batch was diluted with ethyl acetate and washed with brine. The organic phase was filtered using a Whatman filter and concentrated. The residue was purified by chromatography (DCM / ethanol 95:5) to give the desired product (178 mg; 0.39 mmol).


20

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.94 (m, 1H), 7.84 (s, 1H), 7.74 (br, 1H), 7.53 (s, 1H), 7.42 (m, 1H), 7.16 (m, 1H), 6.77 (m, 2H), 4.74 (m, 2H), 4.17 (q, 2H), 3.93 (s, 3H), 3.00 (s, 3H), 1.30 (tr, 3H).

25

Example 2:

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 5 A freshly prepared 1.5M solution of sodium ethanolate in ethanol (2.9 mL; 4.35 mmol) was added under argon to a solution of (rac)-ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate (500 mg; 1.09 mmol) in ethanol (18.5 mL). The batch was stirred at 60°C for 2 hours. Further 1.5M solution of sodium ethanolate in ethanol (2.9 mL; 4.35 mmol) was added and the batch was stirred for additional 5 hours at 60°C. After cooling the batch was
- 10 diluted with brine and extracted with ethyl acetate (3x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by chromatography (DCM / ethanol 9:1) to give the desired product (378 mg; 0.98 mmol).


^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.95 (m, 1H), 7.77 (m, 2H), 7.55 (s, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 6.75 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H), 2.71 (s, 1H).

15

Alternative preparation of Example 2 ((rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine):

Preparation of Intermediate 2.1:

- 20 **(rac)-4-Chloro-N-{3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-**




amine

Intermediate 2.1 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using (rac)-3-[(methylsulfinyl)methyl]aniline (UkrOrgSynthesis Ltd.).

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A2 = H ₂ O + 0.2% NH ₃
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperature:	60°C
Injektion:	2.0 μL
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\MassLynx\NH3_Mass_100_1000.olp
System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Retention	1.13 min
MS(ES-):	m/z = 283 [M+H]

Preparation of Intermediate 2.2:

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine



Intermediate 2.2 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-4-chloro-N-{3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by column chromatography (DCM / EtOH 95:5) to give the desired product.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.95 (br, 1H), 7.74 (br, 1H), 7.65 (s, 1H), 7.40 (m, 2H), 7.05 (m, 1H), 6.77 (m, 2H), 4.07 (d, 1H), 3.95 (m, 4H), 2.49 (s, 3H).

Preparation of end product

Concentrated sulfuric acid (2.5 mL) was added dropwise to a stirred batch of sodium azide (0.61 g; 9.4 mmol) and ((rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine (1.75 g; 4.7 mmol) in trichloromethane (8.0 mL) at 0°C. The batch was stirred for 18 hours at 45°C. While cooling in an ice bath the batch was cautiously diluted with ice water. The batch was further diluted with saturated sodium chloride solution and THF before solid sodium bicarbonate was added under stirring to neutralize the acid. The batch was extracted with THF (3x). The combined organic phases were washed with saturated sodium chloride solution, dried (Na₂SO₄), filtered and concentrated to give the desired product (1.79 g; 4.6 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.95 (m, 1H), 7.77 (m, 2H), 7.55 (s, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 6.75 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H), 2.71 (s, 1H).

Example 3 and 4:

(-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (enantiomer 1) and (+)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (enantiomer 2)

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (see example 2) was separated into the enantiomers by preparative HPLC.

<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501		
<i>Column:</i>	Chiralpak IC 5µm 250x20 mm		
<i>Solvent:</i>	Hexane / Ethanol 60:40 + 0.1% Diethylamine		
<i>Flow:</i>	40 mL/min		
<i>Solution:</i>	2600 mg / 44 mL EtOH/DMSO 2:1		
<i>Injektion:</i>	55 x 0.8 mL		
<i>Temperature:</i>	RT		
<i>Detection:</i>	UV 254 nm		
	Retention time in min	purity in %	Optical rotation index

Example 3 Enantiomer 1	13.4 – 15.6	98.3	-5.2° +/- 0.31° (c = 1.0000 g/ 100 mL CHCl ₃) 20°C or 17.9° +/- 0.48° (c = 1.0000 g/ 100 mL DMSO) 20°C
Example 4 Enantiomer 2	15.6 – 17.8	95.5	2.3° +/- 0.06° (c = 1.0000 g/ 100 mL CHCl ₃) 20°C or -14.0° +/- 0.40° (c = 1.0000 g/ 100 mL DMSO) 20°C


Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.95 (m, 1H), 7.77 (m, 2H), 7.55 (s, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 6.75 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H), 2.71 (s, 1H).

5 Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.95 (m, 1H), 7.77 (m, 2H), 7.55 (s, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 6.75 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H), 2.71 (s, 1H).

10 Due to its negative optical rotation index in chloroform Enantiomer 1 is also referred to as (-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine.
Enantiomer 2 is also referred to as (+)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)-methyl]phenyl}-1,3,5-triazin-2-amine.

15 **Example 5:**

(rac)-Ethyl {[3-({4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl}amino)benzyl](methyl)oxido- λ^6 -sulfanylidene}carbamate



20 Example 5 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl[3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl](methyl)oxido- λ^6 -sulfanylidene]carbamate and [2-(benzyloxy)-4-fluorophenyl]boronic acid (ABCR GmbH & Co. KG).


The batch was purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H_2O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

25 ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.82 (s, 1H), 7.94 (m, 1H), 7.82 (s, 1H), 7.65 (m, 1H), 7.44 (m, 3H), 7.31 (m, 4H), 7.12 (m, 1H), 6.80 (m, 2H), 5.20 (s, 2H), 4.66 (s, 2H), 4.16 (q, 2H), 2.96 (s, 3H), 1.30 (tr, 3H).

Example 6:

(rac)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 5 Example 6 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl {3-({4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl} amino)benzyl}(methyl)oxido- λ^6 -sulfanylidene} carbamate. The batch was purified by column chromatography (DCM / EtOH 95:5).
 ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.82 (s, 1H), 7.95 (m, 1H), 7.70 (m, 2H), 7.50 (s, 1H), 7.44 (m, 2H), 7.32 (m, 4H), 7.12 (m, 1H), 6.79 (m, 2H), 5.20 (s, 2H), 4.33 (d, 1H), 4.21 (d, 1H), 2.92 (s, 3H), 2.69
 10 (s, 1H).

Example 7 and 8:

- 15 **(-)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (enantiomer 1) and (+)-4-[2-(benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)-methyl]phenyl}-1,3,5-triazin-2-amine (enantiomer 2)**

(rac)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

20


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501		
<i>Column:</i>	Chiralpak IC 5 μm 250x30 mm		
<i>Solvent:</i>	Hexane / Ethanol 50:50 + 0.1% Diethylamine		
<i>Flow:</i>	30 mL/min		
<i>Solution:</i>	52 mg / 1.5 mL EtOH/MeOH 1:1		
<i>Injektion:</i>	2 x 0.75 mL		
<i>Temperature:</i>	RT		
<i>Detection:</i>	UV 254 nm		
	Retention time in min	purity in %	Optical rotation index
Example 7	12.0 – 13.5	> 99.9	-7.1° +/- 0.11° (c = 1.0000 g / 100 mL CHCl_3) 20°C
Enantiomer 1			
Example 8	13.5 – 15.3	98.2	3.5° +/- 0.08° (c = 1.0000 g / 100 mL CHCl_3) 20°C
Enantiomer 2			

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.82 (s, 1H), 7.95 (m, 1H), 7.70 (m, 2H), 7.50 (s, 1H), 7.44 (m, 2H), 7.32 (m, 4H), 7.12 (m, 1H), 6.79 (m, 2H), 5.20 (s, 2H), 4.33 (d, 1H), 4.21 (d, 1H), 2.92 (s, 3H), 2.69 (s, 1H).

- Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.82 (s, 1H), 7.95 (m, 1H), 7.70 (m, 2H), 7.50 (s, 1H), 7.44 (m, 2H), 7.32 (m, 4H), 7.12 (m, 1H), 6.79 (m, 2H), 5.20 (s, 2H), 4.33 (d, 1H), 4.21 (d, 1H), 2.92 (s, 3H), 2.69 (s, 1H).
- 25

Example 9:

(rac)-Ethyl [(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate




5 Example 9 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-chloro-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (4,5-difluoro-2-methoxyphenyl)boronic acid (Aldrich). The batch was purified by column chromatography (DCM / EtOH 95:5).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.82 (m, 3H), 7.46 (m, 2H), 7.18 (m, 1H), 6.87 (m, 1H), 4.74 (m, 2H), 4.17 (q, 2H), 3.92 (s, 3H), 3.01 (s, 3H), 1.31 (tr, 3H).

Example 10:

15 (rac)-4-(4,5-Difluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 10 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. The batch was purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.2% NH ₃ B = Acetonitrile
Gradient:	0–1 min 15% B, 1-8 min 15-60% B
Flow:	50 mL/min
Solution:	48 mg / 2 mL DMSO
Injektion:	2 x 1 mL
Temperature:	RT
Detection:	DAD scan range 210–400 nm MS ESI+, scan range 160-1000 m/z ELSD
Retention	7.57 - 8.00 min

20 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.78 (m, 3H), 7.54 (s, 1H), 7.42 (m, 1H), 7.17 (m, 1H), 6.86 (m, 1H), 4.40 (d, 1H), 4.27 (d, 1H), 3.91 (s, 3H), 2.96 (s, 3H), 2.72 (s, 1H).

Example 11:

(rac)-Ethyl [(3-{[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate



5

Example 11 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-(4-chloro-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido- λ^6 -sulfanylidene]-carbamate and (4-chloro-2-methoxyphenyl)boronic acid (ABCR GmbH & Co. KG). The batch was purified by preparative HPLC.

10


System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1–8 min 1–99% B, 8–10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160–1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.85 (m, 2H), 7.75 (br, 1H), 7.57 (m, 1H), 7.42 (m, 1H), 7.17 (m, 1H), 7.04 (m, 2H), 4.73 (m, 2H), 4.16 (q, 2H), 3.93 (s, 3H), 2.99 (s, 3H), 1.30 (tr, 3H).

15

Example 12:

(rac)-4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-



amine

Example 12 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. The batch was purified by preparative HPLC.

20

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.80 (m, 4H), 7.39 (m, 1H), 7.15 (m, 1H), 7.03 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H). 2.39 (br, 1H).

5

Example 13 and 14:

Enantiomers of 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

(rac)-4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

10

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC	
<i>Column:</i>	Chiralpak IC 5µm 250x20 mm	
<i>Solvent:</i>	Ethanol / Methanol 65:35 +0,1% DEA	
<i>Flow:</i>	12 mL/min	
<i>Solution:</i>	34 mg / 1,5 mL MeOH/DMSO 2:1	
<i>Injektion:</i>	5 x 0,3 mL	
<i>Temperature:</i>	RT	
<i>Detection:</i>	MWD 254 nm	
	Retention time in min	purity in %
Example 13 Enantiomer 1	7,8 – 8,4	> 99.9
Example 14 Enantiomer 2	8,4 – 9,4	>95 %


Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.80 (m, 4H), 7.39 (m, 1H), 7.15 (m, 1H), 7.03 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H). 2.39 (br, 1H).

Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.80 (m, 4H), 7.39 (m, 1H), 7.15 (m, 1H), 7.03 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 3.92 (s, 3H), 2.96 (s, 3H). 2.39 (br, 1H).

15

Example 15:

(rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea



- 5 Isocyanatomethane (7.6 μ l; 0.13 mmol) was added to a solution of (rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (50 mg; 0.13 mmol) in DMF (2.0 ml) and triethylamine (18.0 μ l; 0.13 mmol) at room temperature. The batch was stirred for 5 hours before further isocyanatomethane (3.8 μ l; 0.07 mmol) was added. After 72 hours the batch was diluted with sodium bicarbonate and extracted with ethyl acetate (2x). The combined organic phases were filtered
- 10 using a Whatman filter and concentrated. The residue was purified by preparative HPLC to give the desired product (34 mg; 0.08 mmol)

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.80 (m, 4H), 7.41 (m, 1H), 7.17 (m, 1H), 6.78 (m, 2H), 5.04 (br, 1H), 4.83 (d, 1H), 4.64 (d, 1H), 3.94 (s, 3H), 3.00 (s, 3H), 2.78 (d, 3H).

15

Example 16 and 17:

- (-)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea (enantiomer 1) and (+)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea (enantiomer 2)
- 20

(rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea was separated into the enantiomers by preparative HPLC.


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501		
<i>Column:</i>	Chiralpak IA 5µm 250x30 mm		
<i>Solvent:</i>	Ethanol / Methanol 50:50		
<i>Flow:</i>	30 mL/min		
<i>Temperature:</i>	RT		
<i>Solution:</i>	26 mg / 1.5 mL EtOH/MeOH 1:1		
<i>Injektion:</i>	1 x 1.5 mL		
<i>Detection:</i>	UV 254 nm		
	Retention time in min	purity in %	Optical rotation index
Example 16 Enantiomer 1	27.5 – 35.5	98.9	-25.1° +/- 0.19° (c = 1.0000 g / 100 mL CHCl ₃) 20°C
Example 17 Enantiomer 2	37.0 – 50.3	99.2	18.7° +/- 0.10° (c = 1.0000 g / 100 mL CHCl ₃) 20°C

Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.80 (m, 4H), 7.41 (m, 1H), 7.17 (m, 1H), 6.78 (m, 2H), 5.04 (br, 1H), 4.83 (d, 1H), 4.64 (d, 1H), 3.94 (s, 3H), 3.00 (s, 3H), 2.78 (d, 3H).

Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.80 (m, 4H), 7.41 (m, 1H), 7.17 (m, 1H), 6.78 (m, 2H), 5.04 (br, 1H), 4.83 (d, 1H), 4.64 (d, 1H), 3.94 (s, 3H), 3.00 (s, 3H), 2.78 (d, 3H).

Example 18:

(rac)-Ethyl [(3-{[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-oxido-λ⁶-sulfanylidene]carbamate



Example 18 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-chloro-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (2,2-difluoro-1,3-benzodioxol-4-yl)boronic acid (Combi Blocks Inc.). The batch was purified by preparative HPLC.


System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.86 (s, 1H), 8.12 (m, 1H), 7.82 (m, 1H), 7.48 (m, 3H), 7.24 (m, 3H), 4.76 (m, 2H), 4.17 (q, 2H), 3.02 (s, 3H), 1.31 (tr, 3H).

5

Example 19:

(rac)-4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




10 Example 19 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl [(3-{[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate. The batch was purified by column chromatography (DCM / EtOH 95:5).
¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.13 (m, 1H), 7.87 (m, 2H), 7.52 (br, 1H), 7.45 (m, 1H), 7.22 (m, 3H), 4.44 (d, 1H), 4.29 (d, 1H), 2.97 (s, 3H), 2.70 (s, 1H).

15

Example 20:

(rac)-Ethyl [(3-{[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate



20


Example 20 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-chloro-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (5-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc.). The batch was
25 purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.84 (m, 1H), 7.73 (br, 1H), 7.60 (m, 1H), 7.51 (s, 1H), 7.43 (m, 1H), 7.19 (m, 2H), 7.00 (m, 1H), 4.74 (m, 2H), 4.18 (q, 2H), 3.91 (s, 3H), 3.00 (s, 3H), 1.30 (tr, 3H).

Example 21:

(rac)-4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 21 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl [(3-{[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate. The batch was purified by column chromatography (DCM / EtOH 95:5).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.73 (m, 2H), 7.63 (m, 1H), 7.43 (m, 2H), 7.19 (m, 2H), 7.02 (m, 1H), 4.38 (d, 1H), 4.28 (d, 1H), 3.91 (s, 3H), 2.95 (s, 3H), 2.70 (br, 1H).

Example 22 and 23:

Enantiomers of 4-(5-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

(rac)-4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:


<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC	
<i>Column:</i>	Chiralcel OJ-H 5µm 250x20 mm	
<i>Solvent:</i>	Ethanol / methanol 50:50 +0.1% DEA	
<i>Flow:</i>	16 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	57 mg / 0,6 mL EtOH/MeOH 1:1	
<i>Injektion:</i>	2 x 0,3 mL	
<i>Detection:</i>	MWD 254 nm	
	Retention time in min	purity in %
Example 22 Enantiomer 1	6,1 – 7,2	>99 %
Example 23 Enantiomer 2	8,6 – 10,2	>99 %

Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.73 (m, 2H), 7.63 (m, 1H), 7.43 (m, 2H), 7.19 (m, 2H), 7.02 (m, 1H), 4.38 (d, 1H), 4.28 (d, 1H), 3.91 (s, 3H), 2.95 (s, 3H), 2.70 (br, 1H).

Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.73 (m, 2H), 7.63 (m, 1H), 7.43 (m, 2H), 7.19 (m, 2H), 7.02 (m, 1H), 4.38 (d, 1H), 4.28 (d, 1H), 3.91 (s, 3H), 2.95 (s, 3H), 2.70 (br, 1H).

Example 24:

(rac)-N-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]acetamide



Acetyl chloride (10.1 µl; 0.14 mmol) was added to a solution of (rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (50 mg; 0.13 mmol) in DCM (1.5 ml) and triethylamine (45.0 µl; 0.32 mmol) at 0°C. The ice bath was removed and the batch was stirred for 23 hours before further acetyl chloride (4.0 µl; 0.06 mmol) was added. After 24 hours additional acetyl chloride (5.0 µl; 0.07 mmol) was added and the batch was stirred for 3 hours before it was diluted with water and extracted with ethyl acetate (2x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by preparative HPLC to give the desired product (35 mg; 0.08 mmol).


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.95 (m, 1H), 7.78 (m, 2H), 7.56 (m, 1H), 7.42 (m, 1H), 7.16 (m, 1H), 6.79 (m, 2H), 4.78 (d, 1H), 4.65 (d, 1H), 3.94 (s, 3H), 3.03 (s, 3H), 2.12 (s, 3H).

5

Example 25:


(rac)-Ethyl [(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate



- 10 Example 25 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc.). The batch was purified by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

- 15 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.85 (m, 2H), 7.71 (m, 1H), 7.56 (s, 1H), 7.48 (m, 1H), 7.42 (m, 1H), 7.17 (m, 1H), 7.07 (m, 2H), 4.73 (s, 2H), 4.17 (q, 2H), 3.93 (s, 3H), 2.98 (s, 3H), 1.30 (tr, 3H).

Example 26:**(rac)-4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine**

Example 26 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-carbamate. After aqueous work up no further purification was necessary.

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.86 (m, 1H), 7.75 (m, 2H), 7.54 (s, 1H), 7.49 (m, 1H), 7.41 (m, 1H), 7.16 (m, 1H), 7.05 (m, 2H), 4.38 (d, 1H), 4.26 (d, 1H), 3.93 (s, 3H), 2.94 (s, 3H). 2.70 (s, 1H).

Example 27 and 28:

Enantiomers of 4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine(rac)-4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC	
<i>Column:</i>	Chiralcel OJ-H 5 μm 250x20 mm	
<i>Solvent:</i>	Ethanol / methanol 50:50 +0.1% DEA	
<i>Flow:</i>	16 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	59 mg / 0,6 mL EtOH	
<i>Injektion:</i>	2 x 0,3 mL	
<i>Detection:</i>	MWD 254 nm	
	Retention time in min	purity in %
Example 27 Enantiomer 1	11,7 – 12,2	>99 %
Example 28 Enantiomer 2	7,5 – 7,9	>99 %


Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.86 (m, 1H), 7.75 (m, 2H), 7.54 (s, 1H), 7.49 (m, 1H), 7.41 (m, 1H), 7.16 (m, 1H), 7.05 (m, 2H), 4.38 (d, 1H), 4.26 (d, 1H), 3.93 (s, 3H), 2.94 (s, 3H). 2.70 (s, 1H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.86 (m, 1H), 7.75 (m, 2H), 7.54 (s, 1H), 7.49 (m, 1H), 7.41 (m, 1H), 7.16 (m, 1H), 7.05 (m, 2H), 4.38 (d, 1H), 4.26 (d, 1H), 3.93 (s, 3H), 2.94 (s, 3H), 2.70 (s, 1H).

5

Example 29:

(rac)-Ethyl [(3-{[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate



10

Example 29 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-chloro-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 3,4-dihydro-2H-chromen-8-ylboronic acid (Parkway Scientific LLC). The batch was purified by preparative HPLC.

15


System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H_2O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.84 (s, 1H), 7.85 (m, 1H), 7.74 (br, 1H), 7.65 (m, 1H), 7.42 (m, 2H), 7.19 (m, 2H), 6.94 (m, 1H), 4.73 (s, 2H), 4.30 (tr, 2H), 4.17 (q, 2H), 2.98 (s, 3H), 2.89 (tr, 2H), 2.10 (m, 2H), 1.31 (tr, 3H).

20

Example 30:

(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



25

Example 30 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. After aqueous work up no further purification was necessary.

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.74 (m, 3H), 7.49 (br, 1H), 7.40 (m, 1H), 7.17 (m, 2H), 6.94 (m, 1H), 4.30 (m, 4H), 2.94 (s, 3H), 2.88 (tr, 2H), 2.69 (s, 1H), 2.10 (m, 2H).

5 **Example 31 and 32:**


Enantiomers of 4-(3,4-dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC		
<i>Column:</i>	Chiralcel OJ-H 5 μm 250x20 mm		
<i>Solvent:</i>	Ethanol / methanol 50:50 +0.1% DEA		
<i>Flow:</i>	16 mL/min		
<i>Temperature:</i>	RT		
<i>Solution:</i>	46 mg / 0,6 mL MeOH		
<i>Injektion:</i>	2 x 0,3 mL		
<i>Detection:</i>	MWD 254 nm		
	Retention time in min	purity in %	
Example 31 Enantiomer 1	7,0 – 8,1	>99 %	
Example 32 Enantiomer 2	10,0 – 11,3	>99 %	

- 10 Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.74 (m, 3H), 7.49 (br, 1H), 7.40 (m, 1H), 7.17 (m, 2H), 6.94 (m, 1H), 4.30 (m, 4H), 2.94 (s, 3H), 2.88 (tr, 2H), 2.69 (s, 1H), 2.10 (m, 2H).
 Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.83 (s, 1H), 7.74 (m, 3H), 7.49 (br, 1H), 7.40 (m, 1H), 7.17 (m, 2H), 6.94 (m, 1H), 4.30 (m, 4H), 2.94 (s, 3H), 2.88 (tr, 2H), 2.69 (s, 1H), 2.10 (m, 2H).

Example 33:

(rac)-Ethyl [(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate



5

Example 33 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (2,3-dihydro-1-benzofuran-7-yl)boronic acid (ChemBridge Corporation). The batch was purified by preparative HPLC.

10


System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.10 (m, 1H), 7.96 (s, 1H), 7.75 (m, 1H), 7.42 (m, 3H), 7.18 (m, 1H), 6.98 (m, 1H), 4.80 (tr, 2H), 4.76 (s, 2H), 4.17 (q, 2H), 3.29 (tr, 2H), 3.01 (s, 3H), 1.31 (tr, 3H).

15

Example 34:

(rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



20

Example 34 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. After aqueous work up no further purification was necessary.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.11 (m, 1H), 7.87 (s, 1H), 7.76 (br, 1H), 7.44 (m, 2H), 7.39 (m, 1H), 7.17 (m, 1H), 6.98 (m, 1H), 4.81 (tr, 2H), 4.40 (d, 1H), 4.30 (d, 1H), 3.29 (tr, 2H), 2.95 (s, 3H), 2.72 (s, 1H).

25

Example 35 and 36:**Enantiomers of 4-(2,3-dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine**


5 (rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC	
<i>Column:</i>	Chiralpak IA 5µm 250x20 mm	
<i>Solvent:</i>	Ethanol / Methanol 50:50 +0,1% DEA	
<i>Flow:</i>	15 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	74 mg / 0,9 mL DMSO	
<i>Injektion:</i>	3 x 0,3 mL	
<i>Detection:</i>	MWD 254 nm	
	Retention time in min	purity in %
Example 35 Enantiomer 1	9,1 – 10,5	>97 %
Example 36 Enantiomer 2	10,8 – 15,8	>95 %

Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.11 (m, 1H), 7.87 (s, 1H), 7.76 (br, 1H), 7.44 (m, 2H), 7.39 (m, 1H), 7.17 (m, 1H), 6.98 (m, 1H), 4.81 (tr, 2H), 4.40 (d, 1H), 4.30 (d, 1H), 3.29 (tr, 2H), 2.95 (s, 3H), 2.72 (s, 1H).

10 Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.11 (m, 1H), 7.87 (s, 1H), 7.76 (br, 1H), 7.44 (m, 2H), 7.39 (m, 1H), 7.17 (m, 1H), 6.98 (m, 1H), 4.81 (tr, 2H), 4.40 (d, 1H), 4.30 (d, 1H), 3.29 (tr, 2H), 2.95 (s, 3H), 2.72 (s, 1H).

15 **Example 37:**


(rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-oxido-λ⁶-sulfanylidene]carbamate

Example 37 was prepared under similar conditions as described in the preparation of Example 1 using crude (rac)-ethyl [3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl] (methyl)oxido- λ^6 -sulfanylidene]carbamate and 2,3-dihydro-1,4-benzodioxin-5-ylboronic acid (Combi Blocks Inc.). The batch was purified by column chromatography (DCM / EtOH 95:5).

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.85 (s, 1H), 7.87 (s, 1H), 7.71 (m, 1H), 7.48 (m, 2H), 7.42 (m, 1H), 7.17 (m, 1H), 7.04 (m, 1H), 6.95 (m, 1H), 4.74 (m, 2H), 4.37 (m, 4H), 4.17 (q, 2H), 3.00 (s, 3H), 1.31 (tr, 3H).

Example 38:

(rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 38 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. After aqueous work up no further purification was necessary.

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.84 (s, 1H), 7.75 (m, 2H), 7.56 (s, 1H), 7.49 (m, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 7.04 (m, 1H), 6.94 (m, 1H), 4.38 (m, 5H), 4.26 (d, 1H), 2.95 (s, 3H), 2.72 (s, 1H).

Example 39 and 40:

Enantiomers of 4-(2,3-dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, Prep FC	
<i>Column:</i>	Chiralcel OJ-H 5µm 250x20 mm	
<i>Solvent:</i>	Ethanol / methanol 50:50 +0.1% DEA	
<i>Flow:</i>	16 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	140 mg / 1,5 mL MeOH	
<i>Injektion:</i>	5 x 0,3 mL	
<i>Detection:</i>	MWD 254 nm	
	Retention time in min	purity in %
Example 39 Enantiomer 1	7,5 – 9,0	>99 %
Example 40 Enantiomer 2	9,4 – 11,3	>98 %

Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.75 (m, 2H), 7.56 (s, 1H), 7.49 (m, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 7.04 (m, 1H), 6.94 (m, 1H), 4.38 (m, 5H), 4.26 (d, 1H), 2.95 (s, 3H), 2.72 (s, 1H).


5

Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.75 (m, 2H), 7.56 (s, 1H), 7.49 (m, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 7.04 (m, 1H), 6.94 (m, 1H), 4.38 (m, 5H), 4.26 (d, 1H), 2.95 (s, 3H), 2.72 (s, 1H).

10

Example 41:

(rac)-N-{3-[(N,S-Dimethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine




15 Formaldehyde (17.9 µl; 0.65 mmol) was added to a solution of (rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (50 mg; 0.13 mmol) in formic acid (1.0 ml) at room temperature. The batch was stirred at 80°C for 24 hours. After cooling the batch was diluted with aqueous sodium bicarbonate solution and extracted with ethyl acetate (2x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by
20 preparative HPLC to give the desired product (5 mg; 0.01 mmol).

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBridge C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.2% NH ₃
	B = MeOH
Gradient:	0–1 min 15% B, 1-8 min 15-60% B, 8-8.1 min 60-100% B, 8.1–10 min 100% B
Flow:	50 mL/min
Temperature:	RT
Solution:	59mg / 4.5mL DMSO
Injektion:	9x 0.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	5.6– 5.8 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.94 (m, 1H), 7.71 (m, 2H), 7.58 (m, 1H), 7.40 (m, 1H), 7.13 (m, 1H), 6.75 (m, 2H), 4.33 (s, 2H), 3.93 (s, 3H), 2.86 (s, 3H), 2.74 (s, 3H).

Example 42:

(rac)-Ethyl [{3-[(4-{2-[(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido-λ⁶-sulfanylidene]carbamate




- 10 A batch with crude (rac)-ethyl [{3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido-λ⁶-sulfanylidene]carbamate (400 mg), {2-[(4-fluorobenzyl)oxy]phenyl}boronic acid (266 mg; 1.08 mmol; Aldrich Chemical Company Inc.) and [1,1'-bis(diphenylphosphino)ferrocene]palladium dichloride (132 mg; 0.16 mmol) in 1,2-dimethoxyethane (2.5 mL) and 2M solution of potassium carbonate (1.1 mL) was degassed using argon. The batch was stirred under argon for 90 minutes at 100°C. After cooling the
- 15 batch was diluted with DCM. The organic phase was filtered using a Whatman filter and concentrated. The residue was purified by chromatography (DCM / ethanol 95:5) to give the desired product (134 mg; 0.25 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.86 (m, 2H), 7.62 (m, 1H), 7.42 (m, 5H), 7.12 (m, 3H), 7.09 (m, 2H), 5.16 (s, 2H), 4.67 (s, 2H), 4.16 (q, 2H), 2.95 (s, 3H), 1.30 (tr, 3H).

Example 43:

(rac)-4-{2-[(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




5 Example 43 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[3-[(4-{2-[(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl](methyl)oxido- λ^6 -sulfanylidene]carbamate. After aqueous work up no further purification was necessary.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.87 (m, 1H), 7.76 (s, 1H), 7.68 (m, 1H), 7.42 (m, 5H), 7.04 (m, 5H), 5.16 (s, 2H), 4.34 (d, 1H), 4.22 (d, 1H), 2.92 (s, 3H), 2.70 (s, 1H).

Example 44:

(rac)-N-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]methanesulfonamide




15 Methanesulfonyl chloride (12.0 μ l; 0.16 mmol) was added to a solution of (rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine (50.0 mg; 0.13 mmol), triethylamine (21.6 μ l; 0.16 mmol) and 4-dimethylaminopyridine (1.6 mg; 0.01 mmol) in DCM (2.0 ml) at room temperature. The batch was stirred for 23 hours before additional methanesulfonyl chloride (8.0 μ l; 0.10 mmol) was added. After 23 hours additional methanesulfonyl chloride (8.0 μ l; 0.10 mmol) was added. After 24 hours additional methanesulfonyl chloride (12.0 μ l; 0.16 mmol) was added. Finally, after 48 hours additional methanesulfonyl chloride (20.0 μ l; 0.26 mmol) was added and the batch was stirred for 2 additional hours. The batch was diluted with aqueous water and extracted with DCM (2x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by preparative HPLC to give the desired product (21 mg; 0.05 mmol).

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.94 (m, 1H), 7.82 (m, 3H), 7.43 (m, 1H), 7.21 (m, 1H), 6.76 (m, 2H), 4.74 (s, 2H), 3.92 (s, 3H), 3.14 (s, 3H), 3.06 (s, 3H).

5 **Example 45:**

(rac)-Ethyl [(3-{[4-(3-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate




- 10 Example 45 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [(3-{[4-chloro-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (3-chloro-2-methoxyphenyl)boronic acid (Aalen Chemical Co., Ltd.). The batch was purified by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

- 15 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 7.97 (s, 1H), 7.81 (m, 1H), 7.58 (m, 3H), 7.43 (m, 1H), 7.18 (m, 2H), 4.75 (m, 2H), 4.17 (q, 2H), 3.87 (s, 3H), 3.02 (s, 3H), 1.30 (tr, 3H).

Example 46:

(rac)-Ethyl {[3-({4-[5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]-1,3,5-triazin-2-yl}-amino)benzyl](methyl)oxido- λ^6 -sulfanylidene}carbamate




- 5 Example 46 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl](methyl)oxido- λ^6 -sulfanylidene]-carbamate and [5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]boronic acid (FCH Group Company). The batch was purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

- 10 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.89 (br, 1H), 7.70 (m, 3H), 7.43 (m, 1H), 7.15 (m, 2H), 6.93 (m, 1H), 4.74 (m, 2H), 4.18 (q, 2H), 3.96 (m, 4H), 3.41 (m, 2H), 3.01 (s, 3H), 2.02 (br, 1H), 1.64 (m, 4H). 1.31 (tr, 3H).

Example 47:

- 15 (rac)-Ethyl [methyl(oxido)(3-{[4-(2-phenoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)- λ^6 -sulfanylidene]carbamate



- 20 Example 47 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl](methyl)oxido- λ^6 -sulfanylidene]-carbamate and (2-phenoxyphenyl)boronic acid (ABCR GmbH & CO. KG). The batch was purified by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.35 (s, 1H), 8.72 (s, 1H), 7.89 (m, 2H), 7.65 (br, 1H), 7.52 (m, 1H), 7.26 (m, 4H), 7.03 (m, 3H), 6.89 (m, 2H), 4.78 (s, 2H), 3.93 (q, 2H), 3.12 (s, 3H), 1.09 (tr, 3H).

5

Example 48:


(rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]cyanamide



10

Preparation of Intermediate 48.1:

4-Chloro-N-{3-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine




15 Intermediate 48.1 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3-[(methylsulfanyl)methyl]aniline (UkrOrgSynthesis Ltd.).

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A2 = H ₂ O + 0.2% NH ₃
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\MassLynx\Mass_160_1000_BasicReport.flp
System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Retention	1.13 min
MS(ES-):	m/z = 268 [M+H]

Preparation of Intermediate 48.2:

4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine



5


Intermediate 48.2 was prepared under similar conditions as described in the preparation of Example 1 using crude 4-chloro-N-{3-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by column chromatography (DCM / EtOH 95:5) to give the desired product.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.97 (br, 1H), 7.83 (m, 2H), 7.50 (s, 1H), 7.31 (m, 1H), 7.07 (m, 1H), 6.77 (m, 2H), 3.93 (s, 3H), 3.69 (s, 2H), 2.02 (s, 3H).

Preparation of Intermediate 48.3:

(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-λ⁶-sulfanylidene]cyanamide

15



20

2-Bromo-1H-isoindole-1,3(2H)-dione (150 mg; 0.84 mmol) was added to a solution of 4-(4-fluoro-2-methoxyphenyl)-N-{3-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine (200 mg; 0.56 mmol), cyanamide (31 mg; 0.73 mmol) and potassium 2-methylpropan-2-olate (76 mg; 0.67 mmol) in methanol (3.0 ml) at room temperature. The batch was stirred for 2 hours before it was diluted with aqueous sodium thiosulfate solution and extracted with DCM (2x). The combined organic phases were filtered

using a Whatman filter and concentrated. The residue was purified by preparative HPLC to give the desired product.

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.94 (m, 2H), 7.76 (m, 2H), 7.40 (m, 1H), 7.08 (m, 1H), 6.77 (m, 2H), 4.41 (d, 1H), 4.20 (d, 1H), 3.92 (s, 3H), 2.74 (s, 3H).

Preparation of end product:

Potassium carbonate (84 mg; 0.61 mmol) and 3-chlorobenzenecarboperoxoic acid (75 mg; 0.30 mmol) were added to a stirred solution of (rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-λ⁶-sulfanylidene]cyanamide (80 mg; 0.20 mmol) in ethanol (2.0 ml) at 0°C. The ice bath was removed and the batch was slowly warmed to room temperature. The batch was stirred for 22 hours at room temperature. The batch was diluted with aqueous sodium chloride solution and extracted with ethyl acetate (2x) and DCM (1x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by preparative HPLC to give the desired product (10 mg; 0.03 mmol).

<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501
<i>Column:</i>	Chiralpak IA 5µm 250x30 mm
<i>Solvent:</i>	Methanol + 0.1% diethylamine
<i>Flow:</i>	40 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	55 mg / 2 mL DMSO/MeOH 1:1
<i>Injektion:</i>	4 x 0.5 mL
<i>Detection:</i>	UV 280 nm
<i>Retention:</i>	8.0 – 10.3 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 7.93 (m, 2H), 7.76 (m, 1H), 7.48 (m, 2H), 7.19 (m, 1H), 6.79 (m, 2H), 4.63 (m, 2H), 3.94 (s, 3H), 3.04 (s, 3H).

Example 49 and 50:

Enantiomers of [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide

(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -

5 sulfanylidene]cyanamide was separated into the enantiomers by preparative HPLC:


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IA 5 μ m 250x20 mm	
<i>Solvent:</i>	Ethanol 100 (v/v)	
<i>Flow:</i>	20 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	48 mg / 1.2 mL DCM/MeOH 1:1	
<i>Injektion:</i>	2 x 0.6 mL	
<i>Detection:</i>	UV 254 nm	
	Retention time in min	purity in %
Example 49 Enantiomer 1	5.9	>99
Example 50 Enantiomer 2	9.5	>99


Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.84 (s, 1H), 7.93 (m, 2H), 7.76 (m, 1H), 7.48 (m, 2H), 7.19 (m, 1H), 6.79 (m, 2H), 4.63 (m, 2H), 3.94 (s, 3H), 3.04 (s, 3H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.84 (s, 1H), 7.93 (m, 2H), 7.76 (m, 1H), 7.48 (m, 2H), 7.19 (m, 1H), 6.79 (m, 2H), 4.63 (m, 2H), 3.94 (s, 3H), 3.04 (s, 3H).

Example 51:


(rac)-Ethyl [(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate



Preparation of Intermediate 51.1:**1-Fluoro-3-[(methylsulfanyl)methyl]-5-nitrobenzene**


Intermediate 51.1 was prepared under similar conditions as described in the preparation of Intermediate 1.1 using 1-(chloromethyl)-3-fluoro-5-nitrobenzene (Hansa Fine Chemicals GmbH).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.00 (m, 1H), 7.82 (m, 1H), 7.44 (m, 1H), 3.74 (s, 2H), 2.03 (s, 3H).

Preparation of Intermediate 51.2:**(rac)-1-Fluoro-3-[(methylsulfinyl)methyl]-5-nitrobenzene**


Intermediate 51.2 was prepared under similar conditions as described in the preparation of Intermediate 1.2 using 1-fluoro-3-[(methylsulfanyl)methyl]-5-nitrobenzene.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 8.06 (m, 2H), 7.63 (m, 1H), 4.32 (d, 1H), 4.08 (d, 1H), 2.45 (s, 3H).


Preparation of Intermediate 51.3:**(rac)-2,2,2-trifluoro-N-[(3-fluoro-5-nitrobenzyl)(methyl)oxido-λ⁶-sulfanylidene]acetamide**

Intermediate 51.3 was prepared under similar conditions as described in the preparation of Intermediate 1.3 using (rac)-1-fluoro-3-[(methylsulfinyl)methyl]-5-nitrobenzene.


¹H NMR (400MHz, CDCl₃, 300K) δ = 8.13 (m, 1H) 8.07 (m, 1H), 7.56 (m, 1H), 4.92 (d, 1H), 4.76 (d, 1H), 3.33 (s, 3H).

Preparation of Intermediate 51.4:**(rac)-1-Fluoro-3-[(S-methylsulfonimidoyl)methyl]-5-nitrobenzene**

- 5 Intermediate 51.4 was prepared under similar conditions as described in the preparation of Intermediate 1.4 using (rac)-2,2,2-trifluoro-N-[(3-fluoro-5-nitrobenzyl)(methyl)oxido- λ^6 -sulfanylidene]acetamide.
 ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.19 (m, 1H), 8.08 (m, 1H), 7.76 (m, 1H), 4.60 (d, 1H), 4.49 (d, 1H), 3.85 (s, 1H), 2.79 (s, 3H).

10 Preparation of Intermediate 51.5:**(rac)-Ethyl [(3-fluoro-5-nitrobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate**


- Intermediate 51.5 was prepared under similar conditions as described in the preparation of Intermediate 1.5 using (rac)-1-fluoro-3-[(S-methylsulfonimidoyl)methyl]-5-nitrobenzene.
 15 ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.11 (m, 1H), 8.02 (m, 1H), 7.57 (m, 1H), 4.90 (d, 1H), 4.759 (d, 1H), 4.18 (q, 2H), 3.12 (s, 3H), 1.31 (tr, 3H).

Preparation of Intermediate 51.6:**(rac)-Ethyl [(3-amino-5-fluorobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate**

- 20 Intermediate 51.6 was prepared under similar conditions as described in the preparation of Intermediate 1.6 using (rac)-ethyl [(3-fluoro-5-nitrobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate.
 ^1H NMR (400MHz, CDCl_3 , 300K) δ = 6.49 (m, 3H), 4.58 (m, 2H), 4.17 (q, 2H), 3.91 (s, 2H), 3.00 (s, 3H), 1.31 (tr, 3H).

Preparation of Intermediate 51.7:

(rac)-Ethyl [{3-[(4-chloro-1,3,5-triazin-2-yl)amino]-5-fluorobenzyl}(methyl)oxido- λ^6 -sulfanylidene]carbamate



- 5 Intermediate 51.7 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using (rac)-ethyl [(3-amino-5-fluorobenzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate.

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A1 = H ₂ O + 0.1% HCOOH
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A1 + B1 = C:\ MassLynx\NH3_Mass_100_1000.olp
Retention:	0.94 min
MS(ES+):	m/z = 388 [M+H]

Preparation of end product:

- 10 Example 51 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [{3-[(4-chloro-1,3,5-triazin-2-yl)amino]-5-fluorobenzyl}(methyl)oxido- λ^6 -sulfanylidene]carbamate and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitril
<i>Gradient:</i>	0–1 min 30% B, 1-8 min 30-70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	664 mg / 7 mL acetone
<i>Injektion:</i>	7 x 1 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	1.12 min

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.65 (s, 1H), 8.86 (s, 1H), 8.32 (br, 1H), 8.00 (br, 1H), 7.48 (br, 1H), 7.12 (m, 1H), 6.98 (m, 1H), 6.92 (m, 1H), 4.87 (m, 2H), 3.98 (m, 2H), 3.92 (s, 3H), 3.22 (s, 3H), 1.15 (tr, 3H).

5

Example 52:

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



10 Example 52 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl[(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.79 (s, 1H), 8.10 (br, 2H), 7.92 (br, 1H), 7.20 (br, 1H), 6.85 (m, 1H), 6.76 (m, 2H), 4.36 (d, 1H), 4.22 (s, 1H), 3.95 (s, 3H), 3.01 (s, 3H).

15

Example 53 and 54:

Enantiomers of 4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

20 (rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IC 5 μ m 250x30 mm	
<i>Solvent:</i>	Hexane / ethanol 70:30 (v/v)	
<i>Flow:</i>	40 mL/min	
<i>Temperature:</i>	RT	
<i>Solution</i>	1860 mg / 10.8 mL THF/DMSO 9:1	
<i>Injection</i>	36 x 0.3 mL	
<i>Detection:</i>	UV 280 nm	
	Retention time in min	purity in %
Example 53 Enantiomer 1	13.7 – 15.4	99.8
Example 54 Enantiomer 2	15.4 – 17.2	95.4

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 8.10 (br, 2H), 7.92 (br, 1H), 7.20 (br, 1H), 6.85 (m, 1H), 6.76 (m, 2H), 4.36 (d, 1H), 4.22 (s, 1H), 3.95 (s, 3H), 3.01 (s, 3H).

5 Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 8.10 (br, 2H), 7.92 (br, 1H), 7.20 (br, 1H), 6.85 (m, 1H), 6.76 (m, 2H), 4.36 (d, 1H), 4.22 (s, 1H), 3.95 (s, 3H), 3.01 (s, 3H).


Example 55:

10 **(rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine**



Preparation of Intermediate 55.1:

(rac)-Ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate



5 Intermediate 55.1 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (2,4-difluorophenyl)boronic acid (Aldrich Chemical Company Inc.). The batch was purified by column chromatography (DCM / EtOH 95:5).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 8.31 (m, 1H), 8.05 (br, 1H), 7.61 (m, 1H), 7.45 (m, 10 2H), 7.21 (m, 1H), 7.00 (m, 2H), 4.76 (dd, 2H), 4.17 (q, 2H), 3.02 (s, 3H), 1.31 (tr, 3H).

Preparation of end product:


Sodium hydride (60%; 3.2 mg; 0.08 mmol) was added under stirring to a solution of (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate (30.0 15 mg; 0.07 mmol) in cyclohexylmethanol (0.5 ml) at room temperature. The batch was stirred under argon at 50°C for 2 hours before additional sodium hydride (60%; 2.7 mg; 0.07 mmol) was added. After 20 hours additional sodium hydride (60%; 2.7 mg; 0.07 mmol) was added and the batch was stirred for further 5 hours. After cooling, the batch was diluted with ethyl acetate and diluted sodium chloride solution. The organic phase was filtered using a Whatman filter and concentrated. The desired product 20 (10 mg; 0.02 mmol) was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1–8 min 1–99% B, 8–10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160–1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.88 (m, 1H), 7.84 (br, 1H), 7.72 (br, 1H), 7.44 (m, 1H), 7.36 (s, 1H), 7.19 (m, 1H), 6.77 (m, 2H), 4.43 (d, 1H), 4.28 (d, 1H), 3.85 (d, 2H), 2.98 (s, 3H), 2.73 (br, 1H), 1.78 (m, 6H), 1.15 (m, 5H).

Example 56:

(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 5 Example 56. was prepared under similar conditions as described in the preparation of Example 55 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (4-fluorophenyl)methanol.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Methanol
<i>Gradient:</i>	0–1 min 50% B, 1-8 min 50-90% B, 8-8.1 min 90-100% B, 8.1–10 min 100% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	620mg /4.5mL DMSO
<i>Injektion:</i>	9x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	5.0 – 5.6 min

- 10 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.95 (m, 1H), 7.75 (br, 1H), 7.68 (m, 1H), 7.38 (m, 4H), 7.13 (m, 1H), 7.01 (m, 2H), 6.80 (m, 2H), 5.15 (s, 2H), 4.36 (d, 1H), 4.22 (d, 1H), 2.93 (s, 3H), 2.68 (br, 1H).

15 **Example 56.a and 56.b:**

Enantiomers of 4-{4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IA 5µm 250x30 mm	
<i>Solvent:</i>	Ethanol / methanol 50:50 (v/v)	
<i>Flow:</i>	30 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	190mg / 3 mL EtOH/MeOH	
<i>Injektion:</i>	2 x 1.5 mL	
<i>Detection:</i>	UV 280 nm	
	Retention time in min	purity in %
Example 56.a Enantiomer 1	17.2 – 23.5	> 99.9
Example 56.b Enantiomer 2	35.8 – 52.5	99.3

Enantiomer 1: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.95 (m, 1H), 7.75 (br, 1H), 7.68 (m, 1H), 7.38 (m, 4H), 7.13 (m, 1H), 7.01 (m, 2H), 6.80 (m, 2H), 5.15 (s, 2H), 4.36 (d, 1H), 4.22 (d, 1H), 2.93 (s, 3H), 2.68 (br, 1H).

- 5 Enantiomer 2: ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.95 (m, 1H), 7.75 (br, 1H), 7.68 (m, 1H), 7.38 (m, 4H), 7.13 (m, 1H), 7.01 (m, 2H), 6.80 (m, 2H), 5.15 (s, 2H), 4.36 (d, 1H), 4.22 (d, 1H), 2.93 (s, 3H), 2.68 (br, 1H).

10 **Example 57:**

(rac)-4-{4-Fluoro-2-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)-methyl]phenyl}-1,3,5-triazin-2-amine



- 15 Example 57. was prepared under similar conditions as described in the preparation of Example 55 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanyl-idene]carbamate and 2-(tetrahydro-2H-pyran-4-yl)ethanol.


<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Methanol
<i>Gradient:</i>	0–1 min 50% B, 1-8 min 50-90% B, 8-8.1 min 90-100% B, 8.1–10 min 100% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	603mg / 4.5mL DMSO
<i>Injektion:</i>	9x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	3.8 – 4.2 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.78 (m, 3H), 7.49 (s, 1H), 7.42 (m, 1H), 7.16 (m, 1H), 6.76 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 4.10 (tr, 2H), 3.91 (m, 2H), 3.32 (m, 2H), 2.95 (s, 3H), 2.71 (br, 1H), 1.75 (m, 3H), 1.33 (m, 4H).

5

Example 58:


(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-(3-{[S-(tetrahydro-2H-pyran-4-yl)sulfonimidoyl]methyl}-phenyl)-1,3,5-triazin-2-amine



10

Preparation of Intermediate 58.1:

(rac)-4-Chloro-N-{3-[(tetrahydro-2H-pyran-4-ylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine




15

Intermediate 58.1 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using (rac)-3-[(tetrahydro-2H-pyran-4-ylsulfinyl)methyl]aniline (UkrOrgSynthesis Ltd.).

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A2 = H ₂ O + 0.2% NH ₃
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\ MassLynx\NH3_Mass_100_1000.olp
Retention	0.81 min
MS(ES-):	m/z = 351 [M-H]

Preparation of Intermediate 58.2:

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(tetrahydro-2H-pyran-4-ylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine



Intermediate 58.2 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-4-chloro-N-{3-[(tetrahydro-2H-pyran-4-ylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by column chromatography (DCM / EtOH 95:5) to give the desired product.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.96 (m, 1H), 7.70 (m, 2H), 7.40 (m, 2H), 7.06 (m, 1H), 6.78 (m, 2H), 4.03 (m, 7H), 3.39 (m, 2H), 2.73 (m, 1H), 1.83 (m, 4H).

Preparation and end product


Example 58 was prepared under similar conditions as described in the alternative preparation of Example 2 using (rac)-4-(4-fluoro-2-methoxyphenyl)-N-{3-[(tetrahydro-2H-pyran-4-ylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine, sodium azide, sulfuric acid and trichloromethane.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% Vol. HCOOH (99%)
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 10% B, 1-8 min 10-45% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	200 mg / 2 mL DMSO
<i>Injektion:</i>	2 x 1 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.5 – 6.9 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.97 (br, 1H), 7.76 (m, 2H), 7.53 (s, 1H), 7.41 (m, 1H), 7.16 (m, 1H), 6.77 (m, 2H), 4.31 (d, 1H), 4.09 (m, 3H), 3.93 (s, 3H), 3.36 (m, 2H), 3.15 (m, 1H), 2.61 (s, 1H), 2.01 (m, 4H).


Example 59:

(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



Preparation of Intermediate 59.1:

(rac)-4-Chloro-N-{4-chloro-3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine




Intermediate 59.1 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using (rac)-4-chloro-3-[(methylsulfinyl)methyl]aniline (UkrOrgSynthesis Ltd.).

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A2 = H ₂ O + 0.2% NH ₃
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\ MassLynx\NH3_Mass_100_1000.olp
System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Retention	0.85 min
MS(ES-):	m/z = 317 [M+H]

Preparation of Intermediate 59.2:

(rac)-N-{4-Chloro-3-[(methylsulfinyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



Intermediate 59.2 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-4-chloro-N-{4-chloro-3-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by column chromatography (DCM / EtOH 95:5) to give the desired product.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.85 (m, 3H), 7.64 (m, 1H), 7.39 (m, 1H), 6.76 (m, 2H), 4.19 (d, 1H), 4.13 (d, 1H), 3.93 (s, 3H), 2.57 (s, 3H),

- 15 **Preparation of end product** Example 59 was prepared under similar conditions as described in the alternative preparation of Example 2 using ((rac)-N-{4-chloro-3-[(methylsulfinyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine, sodium azide, sulfuric acid and trichloromethane.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 15% B, 1-8 min 15-60% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	55 mg / 1 mL DMSO/MeOH 1:1
<i>Injektion:</i>	1 x 1 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.2 – 6.5 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.87 (m, 3H), 7.57 (s, 1H), 7.42 (m, 1H), 6.78 (m, 2H), 4.60 (d, 1H), 4.51 (d, 1H), 3.93 (s, 3H), 2.97 (s, 3H), 2.84 (s, 1H).

5

Example 59.a and 59.b:

Enantiomers of N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:

10

<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501
<i>Column:</i>	Chiralpak IA 5µm 250x20 mm
<i>Solvent:</i>	Methanol 100 (v/v)
<i>Flow:</i>	20 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	14 mg / 1 mL DCM/MeOH 1:1
<i>Injektion:</i>	2 x 0.5 mL
<i>Detection:</i>	UV 280 nm


	Retention time in min	purity in %
Example 59.a Enantiomer 1	5.3	98.7
Example 59.b Enantiomer 2	7.3	96.8

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.87 (m, 3H), 7.57 (s, 1H), 7.42 (m, 1H), 6.78 (m, 2H), 4.60 (d, 1H), 4.51 (d, 1H), 3.93 (s, 3H), 2.97 (s, 3H), 2.84 (s, 1H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.87 (m, 3H), 7.57 (s, 1H), 7.42 (m, 1H), 6.78 (m, 2H), 4.60 (d, 1H), 4.51 (d, 1H), 3.93 (s, 3H), 2.97 (s, 3H), 2.84 (s, 1H).

Example 60:

(rac)-Ethyl [(3-[(4-{2-[(3,4-dichlorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl)(methyl)-oxido- λ^6 -sulfanylidene]carbamate




Example 60 was prepared under similar conditions as described in the preparation of Example 42 using crude (rac)-ethyl [(3-[(4-chloro-1,3,5-triazin-2-yl)amino]benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and {2-[(3,4-dichlorobenzyl)oxy]phenyl}boronic acid (Combi Blocks Inc.). The batch was purified by preparative HPLC.

<i>System:</i>	Agilent: Prep 1200, 2 x Prep Pump, DLA, MWD, ELSD, Prep FC
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H_2O +0,2% NH_3 B = Acetonitrile
<i>Gradient:</i>	0-17,5 min 40-80% B; 17,5-20 min 80-100% B
<i>Flow:</i>	38 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	120 mg / 1,6 mL DMSO/ACN 1:1
<i>Injektion:</i>	2 x 0,8 mL
<i>Detection:</i>	MWD 210 nm
<i>Retention:</i>	12.2 – 13.3 min

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.87 (s, 1H), 7.91 (m, 1H), 7.85 (s, 1H), 7.66 (m, 2H), 7.46 (m, 2H), 7.38 (m, 2H), 7.22 (m, 1H), 7.15 (m, 2H), 7.06 (m, 1H), 5.15 (s, 2H), 4.70 (s, 2H), 4.16 (q, 2H), 2.97 (s, 3H), 1.30 (tr, 3H).

Example 61:


(rac)-4-{2-[(3,4-Dichlorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 5 Example 61 was prepared under similar conditions as described in the preparation of Example 2 using (rac)-ethyl [(3-[(4-{2-[(3,4-dichlorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate. After aqueous work up no further purification was necessary.
- ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.87 (s, 1H), 7.93 (m, 1H), 7.78 (s, 1H), 7.70 (m, 2H), 7.48 (m, 1H), 7.36 (m, 3H), 7.24 (m, 1H), 7.14 (m, 2H), 7.06 (m, 1H), 5.15 (s, 2H), 4.37 (d, 1H), 4.25 (d, 1H),
 10 2.93 (s, 3H), 2.68 (s, 1H),

Example 62:

- 15 (rac)-4-(4-Fluoro-2-{[($^2\text{H}_5$)phenyl($^2\text{H}_2$)methyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine




- Example 62 was prepared under similar conditions as described in the preparation of Example 55 using (rac)-ethyl [(3-[(4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl)amino]benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and ($^2\text{H}_5$)phenyl($^2\text{H}_2$)methanol (Aldrich Chemical Company Inc). The batch was
 20 purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H_2O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.81 (s, 1H), 7.94 (m, 1H), 7.68 (m, 3H), 7.32 (br, 1H), 7.11 (m, 1H), 6.79 (m, 2H), 4.33 (d, 1H), 4.21 (d, 1H), 2.92 (s, 3H).

Example 63:

4-[2-(1-Cyclopentylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of all 4 stereoisomers




Example 63. was prepared under similar conditions as described in the preparation of Example 55 using (rac)-ethyl [(3- {[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (rac)-1-cyclopentylethanol (ChemSampCo, Inc.).


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1–8 min 1–99% B, 8–10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm MS ESI+, ESI-, scan range 160–1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.78 (s, 1H), 7.76 (m, 3H), 7.54 (s, 1H), 7.40 (m, 1H), 7.15 (m, 1H), 6.72 (m, 2H), 4.39 (d, 1H), 4.24 (m, 2H), 2.95 (s, 3H), 2.70 (br, 1H), 2.09 (m, 1H), 1.47 (m, 11H).

Example 64:


(rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



Preparation of Intermediate 64.1:**1-Chloro-3-[(methylsulfanyl)methyl]-5-nitrobenzene**


- 5 A suspension of 3-bromomethyl-1-chloro-5-nitrobenzene (10.0 g) in ethanol (200 mL) at -20°C was treated with sodiummethanethiolate (3.3 g) in 3 portions, during 3 hours the temperature was increased from -20°C to room temperature. Then brine was added, extracted with ethyl acetate (3x), the combined organic phases were washed with water to neutrality, dried with sodium sulfate, filtered and concentrated. The title compound (8.6 g) was thus obtained and used without further purification.

10

Preparation of Intermediate 64.2:**1-Chloro-3-[(methylsulfinyl)methyl]-5-nitrobenzene**

- 15 To a solution of 1-chloro-3-[(methylsulfanyl)methyl]-5-nitrobenzene (4.3 g) in methanol (340 mL) water (18.7 mL) and sodium periodat (4.4 g) was added and stirred for 24 hours at room temperature. Then the reaction mixture was concentrated under reduced pressure to 30% of the volume, diluted with water, extracted with ethyl acetate (3x), washed with disodium sulfurothioate and brine, dried with sodium sulfate, filtered and concentrated. The title compound (4.5 g) was thus obtained and used without further purification.

- 20 ¹H-NMR (600MHz, CDCl₃): δ □ = 8.23 (t, 1H), 8.08 (t, 1H), 7.67 (t, 1H), 4.07 (d, 1H), 3.92 (d, 1H), 2.57 (s, 3H).

Preparation of Intermediate 64.3:**3-Chloro-5-[(methylsulfinyl)methyl]aniline**

25


- A solution of 1-chloro-3-[(methylsulfinyl)methyl]-5-nitrobenzene (4.4 g) in methanol (40 mL) and water (11.6 mL) was treated with ammoniumchloride (5.1 g), cooled to 0°C, treated with portions of zinc powder (6.2 g) and stirred for 5 hours at room temperature. The reaction mixture was filtered over cellite, washed with THF / ethyl acetate, washed with brine, dried with sodium sulfate and condensed to dryness.

30

Crystallization of the crude product (4.9 g) from diethyl ether furnished the pure title compound (3.4 g).
¹H-NMR (600MHz, CDCl₃) δ : 6.63 (m, 2H), 6.49 (m, 1H), 3.90 (d, 1H), 3.80 (m, 3H), 2.48 (s, 3H).

Preparation of Intermediate 64.4:


5 4-Chloro-N-{3-chloro-5-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine



10 A solution of 2,4-dichloro-triazine (250 mg) in THF (2.2 mL) and 2-propanol (2.2 mL) at -40°C was treated with N,N-diisopropylethylamine (0.55 mL) and 3-chloro-5-[(methylsulfinyl)methyl]aniline (322.6 mg) and then stirred and gradually warmed from -40°C to 0°C for 3 hours. The reaction mixture was then concentrated in vacuo to give the crude title compound (934 mg) which was used without further purification.

15 Preparation of Intermediate 64.5:

N-{3-Chloro-5-[(methylsulfinyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



20 A solution of 4-chloro-N-{3-chloro-5-[(methylsulfinyl)methyl]phenyl}-1,3,5-triazin-2-amine (934 mg) in 1,2-dimethoxyethane (5.2 mL) and a 2 M solution of sodium carbonate (1.6 mL) is treated with 4-fluoro-2-methoxyphenylboronic acid (269 mg) and Pd(dppf)Cl₂ (129 mg) and then heated for 3 hours at 100°C. The reaction mixture was then allowed to cool to room temperature, taken up in ethyl acetate (100 mL) and water (50 mL), washed with saturated brine, dried over sodium sulfate, and condensed in vacuo to give the crude product that was purified by flash column chromatography on SiO₂ with DCM / acetone (5% – 60%) to give analytically pure product (340 mg).

25 ¹H-NMR (500MHz, CDCl₃): δ = 8.83 (s, 1H), 8.18 (br. s., 1H), 7.99 (br. s., 1H), 7.33 (br. s., 1H), 7.64 (s, 1H), 7.02 (s, 1H), 6.78 (m, 2H), 3.96 (m, 5H), 2.53 (s, 3H).

Preparation of end product: To a solution of N-{3-chloro-5-[(methylsulfinyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine (75 mg) in chloroform (0.8 mL) sodiumazide (48.4 mg) was added and treated at 0°C dropwise with conc. sulfuric acid (0.35 mL) and stirred for 48 hours in a sealed tube. Then the reaction mixture was poured into ice water, alkalized with sodium bicarbonate, extracted with ethyl acetate / THF (9:1), washed, dried over sodium sulfate and evaporated to dryness.

30


The crude product was purified by flash column chromatography on SiO₂ with DCM / ethanol (0% - 5%) to give the desired product (42 mg).

¹H-NMR (600MHz, CDCl₃): δ = 8.81 (s, 1H), 7.99 (m, 3H), 7.41 (br, 1H), 7.10 (m, 1H), 6.76 (m, 2H), 4.35 (d, 1H), 4.20 (d, 1H), 3.94 (s, 3H), 3.01 (s, 3H).

5

Example 65:

(rac)-4-[4-Fluoro-2-(3,3,3-trifluoropropoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



10

Sodium hydride (60%; 26.8 mg; 0.67 mmol) was added under stirring to a solution of (rac)-ethyl[(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate (75.0 mg; 0.17 mmol) in 3,3,3-trifluoropropan-1-ol (0.5 ml; ABCR GmbH & CO. KG) at room temperature. The batch was stirred under argon at 70°C for 19 hours. After UPLC MS check, additional sodium hydride (60%; 13.4 mg; 0.34 mmol) was added and the batch was stirred for additional 22 hours at 70°C. After cooling, the batch was diluted with ethyl acetate and saturated aqueous sodium chloride solution. The organic phase was filtered using a Whatman filter and concentrated. The desired product (12 mg; 0.03 mmol) was isolated by preparative HPLC.

15


System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% Vol. HCOOH (99%) B = methanol
Gradient:	0–1 min 20% B, 1–8 min 20–90% B
Flow:	50 mL/min
Temperature:	RT
Solution:	59 mg / 2 mL DMSO/MeOH 1:1
Injektion:	1 x 2 mL
Detection:	DAD scan range 210–400 nm MS ESI+, ESI-, scan range 160–1000 m/z
Retention:	6.5 - 6.9 min

20

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.78 (s, 1H), 7.88 (m, 1H), 7.71 (m, 3H), 7.39 (m, 1H), 7.15 (m, 1H), 6.82 (m, 1H), 6.73 (m, 1H), 4.39 (d, 1H), 4.25 (m, 3H), 2.96 (s, 3H), 2.60 (m, 3H),

Example 66:

(rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 66 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and pyridin-3-ylmethanol (Aldrich Chemical Company Inc.). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B = Methanol
<i>Gradient:</i>	0–1 min 30% B, 1–8 min 30–70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	233 mg / 2.3 mL DMSO/MeOH 1:1
<i>Injektion:</i>	3 x 0.75 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160–1000 m/z
	ELSD
<i>Retention:</i>	6.3 - 6.7 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 8.76 (br, 1H), 8.55 (m, 1H), 7.98 (m, 1H), 7.77 (m, 2H), 7.69 (m, 1H), 7.39 (m, 2H), 7.28 (m, 1H), 7.14 (m, 1H), 6.82 (m, 2H), 5.20 (s, 2H), 4.37 (d, 1H), 4.23 (d, 1H), 2.94 (s, 3H), 2.70 (br, 1H).

Example 67:

(rac)-4-[4-Fluoro-2-(pyridin-2-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 67. was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and pyridin-2-ylmethanol (Aldrich Chemical Company Inc.). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B = Methanol
<i>Gradient:</i>	0–1 min 30% B, 1-8 min 30-70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	139 mg / 1.5 mL DMSO/MeOH 1:1
<i>Injektion:</i>	2 x 0.75 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.7 – 7.0 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.59 (m, 1H), 8.00 (m, 1H), 7.75 (m, 2H), 7.63 (m, 2H), 7.51 (br, 1H), 7.36 (m, 1H), 7.21 (m, 1H), 7.13 (m, 1H), 6.81 (m, 2H), 5.32 (s, 2H), 4.37 (d, 1H), 4.23 (d, 1H), 2.95 (s, 3H), 2.68 (s, 1H).

Example 68:

(rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 68. was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and pyridin-4-ylmethanol (Aldrich Chemical Company Inc.). The desired product was isolated by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B = Methanol
<i>Gradient:</i>	0–1 min 30% B, 1-8 min 30-70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	123 mg / 1.5 mL DMSO/MeOH 1:1
<i>Injektion:</i>	2 x 0.75 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.5 - 6.8 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.57 (br, 2H), 8.01 (m, 1H), 7.77 (br, 1H), 7.70 (br, 1H), 7.36 (m, 4H), 7.13 (m, 1H), 6.85 (m, 1H), 6.75 (m, 1H), 5.20 (s, 2H), 4.37 (d, 1H), 4.22 (d, 1H), 2.94 (s, 3H), 2.70 (br, 1H).

5

Example 69:

4-{4-Fluoro-2-[1-(4-fluorophenyl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of 4 stereoisomers



10

Example 69 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (rac)-1-(4-fluorophenyl)ethanol (Aldrich Chemical Company Inc.). The desired product was isolated by preparative HPLC.

15



System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
Gradient:	0–1 min 30% B, 1-8 min 30-70% B, 8-8.1 min 70-100% B, 8.1–10 min 100% B
Flow:	50 mL/min
Temperature:	RT
Solution:	187mg /4mL DMSO
Injektion:	4x 1 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	5.8 – 6.2 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.78 (m, 3H), 7.48 (br, 1H), 7.34 (m, 3H), 7.15 (m, 1H), 6.99 (m, 2H), 6.73 (m, 1H), 6.58 (m, 1H), 5.32 (q, 1H), 4.38 (d, 1H), 4.25 (d, 1H), 2.95 (s, 3H), 1.59 (d, 3H).

20


Example 70:

(rac)-[3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide

**Preparation of Intermediate 70.1:****3-Fluoro-5-[(methylsulfonyl)methyl]aniline**

Intermediate 70.1 was prepared under similar conditions as described in the preparation of Intermediate 1.6 using 1-fluoro-3-[(methylsulfonyl)methyl]-5-nitrobenzene.

¹H NMR (400MHz, CDCl₃, 300K) δ = 6.42 (m, 2H), 6.26 (m, 1H), 3.74 (br, 2H), 3.55 (s, 2H), 2.01 (s, 3H).


Preparation of Intermediate 70.2:**4-Chloro-N-{3-fluoro-5-[(methylsulfonyl)methyl]phenyl}-1,3,5-triazin-2-amine**

Intermediate 70.2 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3-fluoro-5-[(methylsulfonyl)methyl]aniline.

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD 3001
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A2 = H ₂ O + 0.2% NH ₃
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 μ l
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\MassLynx\Mass_100_1000_BasicReport.flp
Retention	1.20 min
MS(ES-):	m/z = 285 [M+H]

Preparation of Intermediate 70.3:


4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 5 Intermediate 70.3 was prepared under similar conditions as described in the preparation of Example 42 using crude 4-chloro-N-{3-fluoro-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by chromatography (hexane / ethyl acetate 6:4) and finally recrystallized from ethyl acetate to give the desired product.
- 10 ¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.47 (s, 1H), 8.81 (s, 1H), 7.95 (br, 2H), 7.38 (br, 1H), 7.09 (m, 1H), 6.89 (m, 1H), 6.81 (m, 1H), 3.87 (s, 3H), 3.64 (s, 2H), 1.94 (s, 3H).

Preparation of Intermediate 70.4:

(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ⁴-sulfanylidene]cyanamide



- A mixture of 4-(4-fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine (500 mg; 1.33 mmol), cyanamide (112 mg; 2.67 mmol) and bis(acetyloxy)(phenyl)-λ³-iodane (473 mg; 1.47 mmol) in DCM (7.5 ml) was stirred at a temperature between 0-5°C for 3 hours. The batch was concentrated and the residue was purified by chromatography (DCM / EtOH 92:8) to give the desired product (494 mg; 1.19 mmol).
- 20 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.06 (br, 2H), 7.79 (s, 1H), 7.24 (m, 1H), 6.80 (m, 3H), 4.40 (d, 1H), 4.15 (d, 1H), 3.98 (s, 3H), 2.80 (s, 3H).

Preparation of end product:

- At room temperature sodium metaperiodate (380 mg; 1.774 mmol) was dissolved in water (4.5 ml). DCM (6.0 ml) and ruthenium(III)chloride (2 mg; 0.009 mmol) were added under stirring. A suspension of (rac)-[(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ⁴-sulfanylidene]cyanamide (490 mg; 1.183 mmol) in DCM (8.0 ml) was added dropwise and the batch was stirred at room temperature. After 18 hours additional ruthenium(III)chloride (2 mg; 0.009 mmol) was added and the batch was stirred for 5 hours. Finally, further ruthenium(III)chloride (2 mg; 0.009 mmol) was added and the batch was stirred over night. The batch was filtered and the filtrate was extracted with
- 30

DCM (3x). The combined organic phases were filtered using a Whatman filter and concentrated. The residue was purified by chromatography (DCM / EtOH 9:1) to give the desired product (146 mg; 0.340 mmol).

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.66 (s, 1H), 8.83 (s, 1H), 8.27 (br, 1H), 7.93 (br, 1H), 7.47 (br, 1H), 7.09 (m, 1H), 6.97 (m, 1H), 6.89 (m, 1H), 4.99 (m, 2H), 3.87 (s, 3H), 3.38 (s, 3H).

Example 71 and 72:

Enantiomers of (rac)-[(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)-(methyl)oxido- λ^6 -sulfanylidene]cyanamide

(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide was separated into the enantiomers by preparative HPLC.


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IA 5 μ m 250x30 mm	
<i>Solvent:</i>	Ethanol / methanol 50:50 (v/v)	
<i>Flow:</i>	30 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	143 mg / 3.3 mL EtOH/MeOH/DMSO 1.5:1.5:0.3	
<i>Injektion:</i>	3 x 1.1 mL	
<i>Detection:</i>	UV 254 nm	
	Retention time in min	purity in %
Example 71	11.8 – 15.2	> 99.9
Enantiomer 1		
Example 72	15.2 – 25.1	96.9
Enantiomer 2		

Enantiomer 1: ¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.66 (s, 1H), 8.83 (s, 1H), 8.27 (br, 1H), 7.93 (br, 1H), 7.47 (br, 1H), 7.09 (m, 1H), 6.97 (m, 1H), 6.89 (m, 1H), 4.99 (m, 2H), 3.87 (s, 3H), 3.38 (s, 3H).

Enantiomer 2: ¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.66 (s, 1H), 8.83 (s, 1H), 8.27 (br, 1H), 7.93 (br, 1H), 7.47 (br, 1H), 7.09 (m, 1H), 6.97 (m, 1H), 6.89 (m, 1H), 4.99 (m, 2H), 3.87 (s, 3H), 3.38 (s, 3H).

Example 73:

(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 73 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and but-2-yn-1-ol (Aldrich Chemical Company Inc.). The compound was purified by chromatography (DCM / EtOH 97:3).

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 7.96 (m, 1H), 7.80 (m, 2H), 7.60 (m, 1H), 7.41 (m, 1H), 7.15 (m, 1H), 6.96 (m, 1H), 6.80 (m, 1H), 4.79 (d, 2H), 4.42 (d, 1H), 4.29 (d, 1H), 2.95 (s, 3H), 2.72 (br, 1H), 1.85 (tr, 3H).

Example 74 and 75:

Enantiomers of 4-[2-(but-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IC 5 μm 250x30 mm	
<i>Solvent:</i>	Hexan / ethanol 70:30 (v/v)	
<i>Flow:</i>	40 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	360 mg / 6 mL EtOH/MeOH/DCM	
<i>Injektion:</i>	5 x 1.2 mL	
<i>Detection:</i>	UV 280 nm	
	Retention time in min	purity in %
Example 74 Enantiomer 1	20.4 – 23.8	99.9
Example 75 Enantiomer 2	24.1 – 28.1	98.9

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 7.96 (m, 1H), 7.80 (m, 2H), 7.60 (m, 1H), 7.41 (m, 1H), 7.15 (m, 1H), 6.96 (m, 1H), 6.80 (m, 1H), 4.79 (d, 2H), 4.42 (d, 1H), 4.29 (d, 1H), 2.95 (s, 3H), 2.72 (br, 1H), 1.85 (tr, 3H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 7.96 (m, 1H), 7.80 (m, 2H), 7.60 (m, 1H), 7.41 (m, 1H), 7.15 (m, 1H), 6.96 (m, 1H), 6.80 (m, 1H), 4.79 (d, 2H), 4.42 (d, 1H), 4.29 (d, 1H), 2.95 (s, 3H), 2.72 (br, 1H), 1.85 (tr, 3H).

Example 76:

(rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 76. was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 2-cyclopropylethanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5 μm 100x30 mm
Solvent:	A = H_2O + 0.2% NH_3 B = Acetonitrile
Gradient:	0-8 min 30 -70% B
Flow:	50 mL/min
Temperature:	RT
Solution:	87 mg /2 mL DMSO
Injektion:	4 x 0.5 mL
Detection:	DAD scan range 210–400 nm MS ESI+, ESI-, scan range 160-1000 m/z ELSD
Retention:	5.1 – 5.4 min

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.79 (s, 1H), 7.85 (m, 1H), 7.73 (m, 2H), 7.50 (br, 1H), 7.40 (m, 1H), 7.16 (m, 1H), 6.75 (m, 2H), 4.39 (d, 1H), 4.25 (d, 1H), 4.11 (tr, 2H), 2.95 (s, 3H), 2.71 (s, 1H), 1.67 (m, 2H), 0.79 (m, 1H), 0.43 (m, 2H), 0.07 (m, 2H).

Example 77:

(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 77. was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and prop-2-yn-1-ol (Aldrich Chemical Company Inc.). The compound was purified by chromatography (DCM / EtOH 97:3).

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.98 (m, 1H), 7.76 (m, 2H), 7.62 (br, 1H), 7.42 (m, 1H), 7.15 (m, 1H), 6.93 (m, 1H), 6.64 (m, 1H), 4.82 (d, 2H), 4.42 (d, 1H), 4.28 (d, 1H), 2.96 (s, 3H), 2.72 (s, 1H), 2.60 (br, 1H).

Example 78 and 79:

Enantiomers of 4-[4-fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.

<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501
<i>Column:</i>	Chiralpak IC 5 μm 250x30 mm
<i>Solvent:</i>	Hexane / ethanol 70:30 (v/v)
<i>Flow:</i>	40 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	228 mg / 3 mL EtOH/MeOH
<i>Injektion:</i>	3 x 1.0 mL
<i>Detection:</i>	UV 280 nm


	Retention time in min	purity in %
Example 78	19.1 – 22.9	98.7
Enantiomer 1		
Example 79	22.9 – 27.8	98.7
Enantiomer 2		

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.98 (m, 1H), 7.76 (m, 2H), 7.62 (br, 1H), 7.42 (m, 1H), 7.15 (m, 1H), 6.93 (m, 1H), 6.64 (m, 1H), 4.82 (d, 2H), 4.42 (d, 1H), 4.28 (d, 1H), 2.96 (s, 3H), 2.72 (s, 1H), 2.60 (br, 1H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.98 (m, 1H), 7.76 (m, 2H), 7.62 (br, 1H), 7.42 (m, 1H), 7.15 (m, 1H), 6.93 (m, 1H), 6.64 (m, 1H), 4.82 (d, 2H), 4.42 (d, 1H), 4.28 (d, 1H), 2.96 (s, 3H), 2.72 (s, 1H), 2.60 (br, 1H).

Example 80:

(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine




Example 80 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (3,4-difluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H_2O + 0.1% Vol. HCOOH (99%)
	B = Acetonitrile
Gradient:	0-8 min 15-60% B
Flow:	50 mL/min
Temperature:	RT
Solution:	238 mg / 3.9 mL DMSO/MeOH 1:1
Injektion:	3 x 1.3 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	8.0 – 8.5 min

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.80 (s, 1H), 7.96 (m, 1H), 7.69 (m, 3H), 7.33 (m, 2H), 7.09 (m, 3H), 6.83 (m, 2H), 5.25 (s, 2H), 4.37 (d, 1H), 4.23 (d, 1H), 2.95 (s, 3H), 2.70 (br, 1H),

Example 81:

(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine




Example 81 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 1,3-thiazol-5-ylmethanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 15 -60% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	579 mg /3.5 mL DMSO
<i>Injektion:</i>	7 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	5.0 – 5.2 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.79 (m, 2H), 7.96 (m, 1H), 7.88 (m, 1H), 7.74 (br, 1H), 7.68 (m, 1H), 7.51 (br, 1H), 7.37 (m, 1H), 7.14 (m, 1H), 6.86 (m, 2H), 5.37 (s, 2H), 4.38 (d, 1H), 4.23 (d, 1H), 2.95 (s, 3H).

Example 82:

(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)-methyl]phenyl}-1,3,5-triazin-2-amine



Example 82 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (2-fluoropyridin-4-yl)methanol (Activate Scientific GmbH). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 15 -60% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	312 mg /3 mL DMSO
<i>Injektion:</i>	6 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	5.7 – 6.0 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.17 (m, 1H), 8.04 (m, 1H), 7.76 (br, 1H), 7.70 (m, 1H), 7.61 (br, 1H), 7.36 (br, 1H), 7.22 (m, 2H), 7.13 (m, 1H), 6.86 (m, 1H), 6.74 (m, 1H), 5.20 (s, 2H), 4.37 (d, 1H), 4.24 (d, 1H), 2.95 (s, 3H), 2.73 (br, 1H).

5

Example 83 and 84:

Enantiomers of 4-{4-fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

10

(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC.


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IA 5 μ m 250x30 mm	
<i>Solvent:</i>	Ethanol 100%	
<i>Flow:</i>	25 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	67 mg / 4 mL EtOH/MeOH	
<i>Injektion:</i>	1 x 4.0 mL	
<i>Detection:</i>	UV 254 nm	
	Retention time in min	purity in %
Example 83 Enantiomer 1	19.0 – 31.0	99.7
Example 84 Enantiomer 2	42.0 – 65.0	99.1

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.85 (s, 1H), 8.17 (m, 1H), 8.04 (m, 1H), 7.76 (br, 1H), 7.70 (m, 1H), 7.61 (br, 1H), 7.36 (br, 1H), 7.22 (m, 2H), 7.13 (m, 1H), 6.86 (m, 1H), 6.74 (m, 1H), 5.20 (s, 2H), 4.37 (d, 1H), 4.24 (d, 1H), 2.95 (s, 3H), 2.73 (br, 1H).

- 5 Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.85 (s, 1H), 8.17 (m, 1H), 8.04 (m, 1H), 7.76 (br, 1H), 7.70 (m, 1H), 7.61 (br, 1H), 7.36 (br, 1H), 7.22 (m, 2H), 7.13 (m, 1H), 6.86 (m, 1H), 6.74 (m, 1H), 5.20 (s, 2H), 4.37 (d, 1H), 4.24 (d, 1H), 2.95 (s, 3H), 2.73 (br, 1H).

10 **Example 85:**

(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 15 Example 85 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and prop-2-en-1-ol (Aldrich Chemical Company Inc.). The desired product was isolated by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 30 -70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	50 mg /2.5 mL DMSO
<i>Injektion:</i>	5 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	3.2 - 3.4 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.92 (m, 1H), 7.75 (m, 2H), 7.49 (s, 1H), 7.40 (m, 1H), 7.16 (m, 1H), 6.76 (m, 2H), 6.03 (m, 1H), 5.46 (m, 1H), 5.28 (m, 1H), 4.85 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 2.94 (s, 3H), 2.70 (br, 1H).

5

Example 86:

(rac)-4-(4-Fluoro-2-{[4-(trifluoromethyl)benzyl]oxy} phenyl)-N-{3-[(S-methylsulfonimidoyl)-methyl]phenyl}-1,3,5-triazin-2-amine



10

Example 86 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and [4-(trifluoromethyl)phenyl]methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 30 -70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	389 mg /4.5 mL DMSO
<i>Injektion:</i>	9 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.2 – 6.6 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.99 (m, 1H), 7.90 (br, 1H), 7.60 (m, 5H), 7.45 (m, 1H), 7.34 (m, 1H), 7.13 (m, 1H), 6.81 (m, 2H), 5.22 (s, 2H), 4.36 (d, 1H), 4.23 (d, 1H), 2.94 (s, 3H), 2.73 (br, 1H).

Example 87:

(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 87 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (4-chlorophenyl)methanol (Aldrich Chemical Company Inc). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1–8 min 1–99% B, 8–10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160–1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.94 (s, 1H), 7.99 (m, 1H), 7.75 (m, 2H), 7.34 (m, 6H), 7.16 (m, 1H), 6.82 (m, 2H), 5.18 (s, 2H), 4.39 (d, 1H), 4.25 (d, 1H), 2.96 (s, 3H), 2.70 (br, 1H).

Example 88:

(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 88 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and ethanol. The desired product was isolated by preparative HPLC.


System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
Gradient:	0-8 min 15 -60% B
Flow:	50 mL/min
Temperature:	RT
Solution:	129 mg /3 mL DMSO
Injektion:	6 x 0.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	5.8 – 6.2 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.81 (m, 3H), 7.53 (s, 1H), 7.43 (m, 1H), 7.18 (m, 1H), 6.78 (m, 2H), 4.41 (d, 1H), 4.28 (d, 1H), 4.16 (q, 2H), 2.97 (s, 3H), 2.73 (s, 1H), 1.44 (tr, 3H).

5

Example 89:

(rac)-4-(4-Fluoro-2-([3-fluoro-5-(trifluoromethyl)benzyl]oxy)phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




- 10 Example 89 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and [3-fluoro-5-(trifluoromethyl)phenyl]methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

- 15 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.04 (s, 1H), 7.77 (br, 1H), 7.67 (m, 2H), 7.42 (m, 3H), 7.26 (m, 1H), 7.15 (m, 1H), 6.83 (m, 2H), 5.21 (s, 2H), 4.38 (d, 1H), 4.25 (d, 1H), 2.95 (s, 3H).

Example 90:

(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



5

Example 90 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (3-fluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


10

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1–8 min 1–99% B, 8–10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160–1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.86 (s, 1H), 8.02 (m, 1H), 7.75 (m, 2H), 7.54 (s, 1H), 7.35 (m, 3H), 7.18 (m, 2H), 7.01 (m, 1H), 6.82 (m, 2H), 5.21 (s, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 2.96 (s, 3H).

Example 91:

(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 91 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and propan-1-ol. The desired product was isolated by preparative HPLC.

20


<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 30 -70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	126 mg / 4 mL DMSO
<i>Injektion:</i>	8 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.6 – 6.9 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.85 (m, 1H), 7.75 (m, 2H), 7.42 (m, 2H), 7.16 (m, 1H), 8.74 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 4.01 (tr, 2H), 2.95 (s, 3H), 2.70 (s, 1H), 1.81 (m, 2H), 1.00 (tr, 3H).

5

Example 92:

(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



10

Example 92 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (3-chlorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 30 -70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	233 mg / 3 mL DMSO
<i>Injektion:</i>	6 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	5.9 – 6.2 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.00 (m, 1H), 7.68 (m, 3H), 7.53 (s, 1H), 7.35 (m, 1H), 7.26 (m, 3H), 7.14 (m, 1H), 6.81 (m, 2H), 5.16 (s, 2H), 4.36 (d, 1H), 4.23 (d, 1H), 2.94 (s, 3H), 2.70 (br, 1H).

Example 93:

(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine




- 5 Example 93 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 1,2-oxazol-3-ylmethanol (UkrOrgSynthesis Ltd.). The desired product was isolated by preparative HPLC.

System:	Agilent: Prep 1200, 2xPrep Pump, DLA, MWD, Prep FC,
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O +0,1% TFA
	B = Methanol
Gradient:	0-17,5 min 25-55% B, 17,5-20 min 55-100% B
Flow:	38 mL/min
Temperature:	RT
Solution:	29 mg / 0,8 mL DMSO
Injektion:	1 x 0,8 mL
Detection:	UV 210 nm
Retention:	8.3 – 10.2 min

- 10 ¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.41 (s, 1H), 8.80 (m, 2H), 7.80 (m, 3H), 7.37 (m, 1H), 7.18 (m, 2H), 6.94 (m, 1H), 6.55 (br, 1H), 5.33 (s, 2H), 4.99 (m, 2H), 3.37 (s, 3H).

Example 94:

- 15 (rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine



- 20 Example 94 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (3-chloro-5-fluorophenyl)methanol (Apollo Scientific Ltd.). The desired product was isolated by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.85 (s, 1H), 8.03 (m, 1H), 7.77 (s, 1H), 7.69 (m, 1H), 7.46 (m, 1H), 7.37 (m, 2H), 7.16 (m, 2H), 7.01 (m, 1H), 6.85 (m, 1H), 6.76 (m, 1H), 5.14 (s, 2H), 4.38 (d, 1H), 4.24 (d, 1H), 2.95 (s, 3H).

5

Example 95:

(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



10

Example 95 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and 2,2-difluoroethanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
<i>Gradient:</i>	0–8 min 15-50% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	68 mg / 3 mL DMSO/MeOH 1:1
<i>Injektion:</i>	3 x 1 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	6.18 - 6.54 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.95 (m, 1H), 7.77 (br, 1H), 7.69 (m, 1H), 7.43 (m, 1H), 7.34 (br, 1H), 7.18 (m, 1H), 6.88 (m, 1H), 6.76 (m, 1H), 6.05 (m, 1H), 4.40 (d, 1H), 4.27 (m, 3H), 2.95 (s, 3H), 2.68 (s, 1H).

Example 96:

(rac)-4-{4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine




- 5 Example 96 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (4-fluoro-3-methylphenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm MS ESI+, ESI-, scan range 160-1000 m/z

- 10 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 7.94 (m, 1H), 7.70 (m, 2H), 7.51 (br, 1H), 7.32 (m, 2H), 7.20 (m, 1H), 7.13 (m, 1H), 6.94 (m, 1H), 6.79 (m, 2H), 5.11 (s, 2H), 4.35 (d, 1H), 4.33 (d, 1H), 2.93 (s, 3H), 2.61 (s, 1H), 2.23 (s, 3H).

15 **Example 97:**

(rac)-4-{2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine



- 20 Example 97 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (3-chloro-4-fluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.00 (m, 1H), 7.76 (br, 1H), 7.69 (m, 2H), 7.48 (br, 1H), 7.36 (m, 1H), 7.27 (m, 1H), 7.11 (m, 2H), 6.79 (m, 2H), 5.12 (s, 2H), 4.37 (d, 1H), 4.24 (d, 1H), 2.95 (s, 3H), 2.71 (br, 1H).

5

Example 98:

(rac)-3-((5-Fluoro-2-[4-((3-((S-methylsulfonimidoyl)methyl)phenyl)amino)-1,3,5-triazin-2-yl]-phenoxy)methyl)benzonitrile



10

Example 98 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-([4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino)benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and 3-(hydroxymethyl)benzonitrile (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.2% NH ₃
	B = Acetonitrile
Gradient:	0-8 min 30 -70% B
Flow:	50 mL/min
Temperature:	RT
Solution:	473 mg / 4.5 mL DMSO
Injektion:	9 x 0.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	4.7 – 4.9 min

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.35 (s, 1H), 8.83 (s, 1H), 7.85 (m, 6H), 7.53 (br, 1H), 7.28 (br, 1H), 7.16 (m, 1H), 7.11 (br, 1H), 6.95 (m, 1H), 5.32 (s, 2H), 4.32 (br, 2H), 3.55 (s, 1H), 2.79 (s, 3H).

Example 99:

(rac)-4-{4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




Example 99 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 2-methylprop-2-en-1-ol (Aldrich Chemical Company Inc). The desired product was isolated by preparative HPLC.

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5 μ m 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.83 (s, 1H), 7.94 (m, 1H), 7.76 (m, 2H), 7.55 (br, 1H), 7.42 (m, 1H), 7.18 (m, 1H), 6.78 (m, 2H), 5.18 (br, 1H), 5.01 (br, 1H), 4.55 (s, 2H), 4.41 (d, 1H), 4.28 (d, 1H), 2.97 (s, 3H), 2.70 (br, 1H).1.82 (s, 3H).

Example 100:

(rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



Example 100 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and 4,4,4-trifluorobutan-1-ol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.82 (s, 1H), 7.95 (m, 1H), 7.80 (br, 1H), 7.70 (m, 1H), 7.46 (m, 2H), 7.19 (m, 1H), 6.82 (m, 1H), 6.73 (m, 1H), 4.42 (d, 1H), 4.29 (d, 1H), 4.12 (tr, 2H), 2.98 (s, 3H), 2.74 (br, 1H), 2.38 (m, 2H), 2.07 (m, 2H).

5

Example 101:

(rac)-4-{4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine



10

Example 101 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-sulfanyl-idene]carbamate and (2,3,5-trifluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.87 (s, 1H), 8.07 (m, 1H), 7.80 (s, 1H), 7.72 (m, 1H), 7.40 (m, 3H), 7.18 (m, 1H), 6.89 (m, 3H), 5.26 (s, 2H), 4.42 (d, 1H), 4.28 (d, 1H), 2.98 (s, 3H), 2.71 (br, 1H).

Example 102:

(rac)-4-{2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine




- 5 Example 102 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (2Z)-but-2-en-1-ol (ChemSampCo, Inc.). The desired product was isolated by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5 μ m 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

- 10 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.89 (m, 1H), 7.76 (m, 2H), 7.57 (br, 1H), 7.39 (m, 1H), 7.15 (m, 1H), 6.75 (m, 2H), 5.70 (m, 2H), 4.72 (m, 2H), 4.39 (d, 1H), 4.26 (d, 1H), 2.95 (s, 3H), 2.71 (br, 1H). 1.73 (d, 3H).

15 **Example 103:**

(rac)-4-{4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine



- 20 Example 103 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate and (2,4,5-trifluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


<i>System:</i>	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	Max. 250 mg / max. 2.5 mL DMSO o. DMF
<i>Injektion:</i>	1 x 2.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.03 (m, 1H), 7.77 (br, 1H), 7.65 (m, 2H), 7.48 (br, 1H), 7.39 (m, 1H), 7.15 (m, 1H), 6.94 (m, 1H), 6.83 (m, 2H), 5.17 (s, 2H), 4.39 (d, 1H), 4.25 (d, 1H), 2.96 (s, 3H), 2.70 (br, 1H).

5

Example 104:

(rac)-4-{4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]-phenyl}-1,3,5-triazin-2-amine



10

Example 104 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-{[4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and (3,4,5-trifluorophenyl)methanol (ABCR GmbH & CO. KG). The desired product was isolated by preparative HPLC.


15

<i>System:</i>	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
<i>Column:</i>	XBrigde C18 5µm 100x30 mm
<i>Solvent:</i>	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
<i>Gradient:</i>	0-8 min 30 -70% B
<i>Flow:</i>	50 mL/min
<i>Temperature:</i>	RT
<i>Solution:</i>	431 mg /4mL DMSO
<i>Injektion:</i>	8 x 0.5 mL
<i>Detection:</i>	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
<i>Retention:</i>	5.6 – 6.0 min

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.84 (s, 1H), 8.02 (m, 1H), 7.77 (br, 1H), 7.69 (m, 1H), 7.39 (m, 2H), 7.17 (m, 3H), 6.85 (m, 1H), 6.74 (m, 1H), 5.10 (s, 2H), 4.39 (d, 1H), 4.25 (d, 1H), 2.95 (s, 3H). 2.69 (br, 1H).

Example 105:


(rac)-[(2,3-Difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-oxido- λ^6 -sulfanylidene]cyanamide



5

Preparation of Intermediate 105.1:

(2,3-Difluoro-5-nitrophenyl)methanol



10


Borane-tetrahydrofuran complex (1.0M solution in THF; 177 mL) was added under stirring to an ice-cold solution of 2,3-difluoro-5-nitrobenzoic acid (9.0 g; 44.3 mmol; Butt Park Ltd.) in THF (85 mL). The ice bath was removed and the batch was stirred for 18 hours at room temperature. The batch was cautiously diluted with methanol under stirring at 0°C. Ethyl acetate was added and the batch was washed with aqueous 1 M solution of sodium hydroxide and half-concentrated aqueous sodium chloride solution. The organic phase was dried (Na₂SO₄), filtered and concentrated. The residue was purified by chromatography (hexane / ethyl acetate 1:1) to give the desired product (8.2 g; 43.3 mmol).

15

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.26 (m, 1H), 8.03 (m, 1H), 4.89 (s, 2H), 2.13 (br, 1H).

Preparation of Intermediate 105.2:

(5-Amino-2,3-difluorophenyl)methanol




20


Intermediate 105.2 was prepared under similar conditions as described in the preparation of Intermediate 1.6 using (2,3-difluoro-5-nitrophenyl)methanol.

¹H NMR (400MHz, CDCl₃, 300K) δ = 6.46 (m, 1H), 6.39 (m, 1H), 4.67 (s, 2H), 3.02 (br, 3H).

25


Preparation of Intermediate 105.3:**3-(Chloromethyl)-4,5-difluoroaniline**

Thionylchloride (7.7 g; 64.9 mmol) was added dropwise under stirring to a water-cooled solution of (5-amino-2,3-difluorophenyl)methanol (4.1 g; 26.0 mmol) in DCM (78 ml) and 1-methylpyrrolidin-2-one (11 ml). The batch was stirred at room temperature for 18 hours before it was diluted with ice-water, aqueous sodium bicarbonate solution and brine. The batch was stirred for 2 hours at room temperature and finally it was extracted with ethyl acetate (2x). The combined organic phases were filtered using a Whatman filter and concentrated to give the crude product (6.5 g) that was used without further purification.

Preparation of Intermediate 105.4:**3,4-Difluoro-5-[(methylsulfanyl)methyl]aniline**

Intermediate 105.4 was prepared under similar conditions as described in the preparation of Intermediate 1.1 using crude 3-(chloromethyl)-4,5-difluoroaniline. The batch was purified by chromatography (hexane / ethyl acetate 1:1).

^1H NMR (400MHz, CDCl_3 , 300K) δ = 6.35 (m, 2H), 3.62 (br, 4H), 2.99 (s, 3H).


Preparation of Intermediate 105.5:**4-Chloro-N-{3,4-difluoro-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine**

Intermediate 105.5 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3,4-difluoro-5-[(methylsulfanyl)methyl]aniline.

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A1 = H ₂ O + 0.1% Vol. HCOOH (99%) A2 = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B1 = Acetonitril
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\MassLynx\NH3_Mass_100_1000.olp
Retention	1.19 min
MS(ES-):	m/z = 303 [M+H]

Preparation of Intermediate 105.6:

N-{3,4-Difluoro-5-[(methylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine




Intermediate 105.6 was prepared under similar conditions as described in the preparation of Example 42 using crude 4-chloro-N-{3,4-difluoro-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by chromatography (hexane / ethyl acetate 1:1) to give the desired product.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 8.15 (br, 2H), 7.35 (m, 1H), 7.08 (br, 1H), 6.78 (m, 2H), 3.98 (s, 3H), 3.73 (s, 2H), 2.08 (s, 3H).

Preparation of Intermediate 105.7:

(rac)-[(2,3-Difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-λ⁴-sulfanylidene]cyanamide



Bis(acetyloxy)(phenyl)-λ³-iodane (469 mg; 1.46 mmol) was added to a stirred solution of N-{3,4-difluoro-5-[(methylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine (520 mg; 1.33 mmol) and cyanamide (111 mg; 2.65 mmol) in DCM (7.5 ml) at 0°C. The batch was stirred at

this temperature for 4 hours. The batch was concentrated and the residue was purified by chromatography (DCM / EtOH 8:2) to give the desired product (300 mg; 0.69 mmol).

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.81 (s, 1H), 8.49 (br, 1H), 8.06 (br, 1H), 7.87 (br, 1H), 7.22 (br, 1H), 6.79 (m, 2H), 4.34 (m, 2H), 3.96 (s, 3H), 2.86 (s, 3H).

5

Preparation of end product:

Potassium permanganate (217 mg; 1.37 mmol) was added under stirring to a solution of (rac)-[(2,3-difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)-λ⁴-sulfanylidene]cyanamide (297 mg; 0.69 mmol) in acetone (6.9 mL) at room temperature. The batch was stirred at 50°C for 1 hour. The batch was concentrated and the residue was purified by chromatography (DCM / EtOH 9:1) to give the desired product (153 mg; 0.34 mmol).


10

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.62 (s, 1H), 8.81 (s, 1H), 8.55 (br, 1H), 7.96 (br, 1H), 7.43 (br, 1H), 7.08 (m, 1H), 6.89 (m, 1H), 5.06 (m, 2H), 3.87 (s, 3H), 3.50 (s, 3H).

15

Example 106:

(rac)-N-{3,4-Difluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



20

Trifluoroacetic anhydride (0.32 mL; 0.90 mmol) was added under stirring to an ice-cooled suspension of (rac)-[(2,3-difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ⁶-sulfanylidene]cyanamide (134 mg; 0.30 mmol) in DCM (13 mL). The batch was stirred for 2 hours at room temperature before it was concentrated in vacuo. The residue was taken up in methanol (2.1 mL) and potassium carbonate (206 mg; 1.49 mmol) was added under stirring at room temperature. After 2 hours the batch was diluted with ethyl acetate and THF. The batch was washed with aqueous sodium chloride solution. The organic phase was filtered using a Whatman filter and concentrated in vacuo. The residue was purified by chromatography (DCM / EtOH 9:1) to give the desired product (68 mg; 0.16 mmol).


25

30

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.78 (s, 1H), 8.49 (br, 1H), 8.08 (br, 1H), 7.69 (br, 1H), 7.18 (br, 1H), 6.78 (m, 2H), 4.46 (d, 1H), 4.38 (d, 1H), 3.94 (s, 3H), 3.03 (s, 3H).

Example 107:


(rac)-[Ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido- λ^6 -sulfanylidene]cyanamide



5

Preparation of Intermediate 107.1:

1-[(Ethylsulfanyl)methyl]-3-nitrobenzene




10 Intermediate 107.1 was prepared under similar conditions as described in the preparation of Intermediate 1.1 using 1-(chloromethyl)-3-nitrobenzene and sodium ethanethiolate.

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.19 (m, 1H), 8.11 (m, 1H), 7.67 (m, 1H), 7.49 (m, 1H), 3.80 (s, 2H), 2.45 (q, 2H), 1.25 (tr, 3H).

15 Preparation of Intermediate 107.2:

3-[(Ethylsulfanyl)methyl]aniline




20 Intermediate 107.2 was prepared under similar conditions as described in the preparation of Intermediate 1.6 using 1-[(ethylsulfanyl)methyl]-3-nitrobenzene.

¹H NMR (400MHz, CDCl₃, 300K) δ = 7.09 (m, 1H), 6.68 (m, 2H), 6.57 (m, 1H), 3.63 (s, 2H), 3.42 (br, 2H), 2.45 (q, 2H), 1.23 (tr, 3H).

Preparation of Intermediate 107.3:

25 **4-Chloro-N-{3-[(ethylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine**




Intermediate 107.3 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3-[(ethylsulfanyl)methyl]aniline.

30

System:	Waters Acquity UPLC-MS: Binary Solvent Manager, Sample Manager/Organizer, Column Manager, PDA, ELSD, SQD
Column:	Acquity UPLC BEH C18 1.7 50x2.1mm
Solvent:	A1 = H ₂ O + 0.1% Vol. HCOOH (99%) A2 = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B1 = Acetonitrile
Gradient:	0-1.6 min 1-99% B, 1.6-2.0 min 99% B
Flow:	0.8 mL/min
Temperatuer:	60°C
Injektion:	2.0 µl
Detection:	DAD scan range 210-400 nm -> Peaktable
	ELSD
Method:	MS ESI+, ESI- Switch A2 + B1 = C:\MassLynx\NH3_Mass_100_1000.olp
Retention	1.22 min
MS(ES-):	m/z = 281 [M+H]

Preparation of Intermediate 107.4:

N-{3-[(Ethylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



5

Intermediate 107.4 was prepared under similar conditions as described in the preparation of Example 42 using crude 4-chloro-N-{3-[(ethylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc.). The batch was purified by column chromatography (hexane / ethyl acetate 1:1) to give the desired product.


10

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.80 (s, 1H), 7.98 (br, 1H), 7.62 (m, 2H), 7.46 (br, 1H), 7.31 (m, 1H), 7.08 (m, 1H), 6.76 (m, 2H), 3.93 (s, 3H), 3.73 (s, 2H), 2.46 (q, 2H), 1.23 (tr, 3H).

Preparation of Intermediate 107.5:

(rac)-[Ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl-λ⁴-sulfanylidene]-cyanamide

15



Intermediate 107.5 was prepared under similar conditions as described in the preparation of Intermediate 105.7 using N-{3-[(ethylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine.

20

¹H NMR (400MHz, CDCl₃, 300K) δ = 8.63 (s, 1H), 7.95 (m, 1H), 7.75 (m, 3H), 7.42 (m, 1H), 7.10 (m, 1H), 6.79 (m, 2H), 4.37 (d, 1H), 4.18 (d, 1H), 3.94 (s, 3H), 3.11 (m, 1H), 2.89 (m, 1H), 1.43 (tr, 3H).


Preparation of end product:

Example 107 was prepared under similar conditions as described in the preparation of Example 105 using (rac)-[ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl- λ^4 -sulfanylidene]-cyanamide.

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.85 (s, 1H), 7.95 (m, 2H), 7.75 (br, 1H), 7.66 (m, 1H), 7.45 (m, 1H), 7.18 (m, 1H), 6.79 (m, 2H), 4.60 (m, 2H), 3.94 (s, 3H), 3.16 (q, 2H), 1.44 (tr, 3H).

Example 108:

(rac)-N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine



Example 108 was prepared under similar conditions as described in the preparation of Example 106 using (rac)-[ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)oxido- λ^6 -sulfanylidene]cyanamide.

^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.81 (s, 1H), 7.96 (m, 1H), 7.78 (m, 2H), 7.41 (m, 2H), 7.16 (m, 1H), 6.78 (m, 2H), 4.33 (d, 1H), 4.18 (d, 1H), 3.94 (s, 3H), 3.04 (q, 2H), 1.43 (tr, 3H).

Example 109 and 110:

Enantiomers of N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine

(rac)-N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine was separated into the enantiomers by preparative HPLC:


<i>System:</i>	Dionex: Pump P 580, Gilson: Liquid Handler 215, Knauer: UV-Detektor K-2501	
<i>Column:</i>	Chiralpak IC 5µm 250x30 mm	
<i>Solvent:</i>	Hexane / ethanol 70:30 (v/v)	
<i>Flow:</i>	50 mL/min	
<i>Temperature:</i>	RT	
<i>Solution:</i>	205 mg / 4.9 mL DCM/MeOH	
<i>Injektion:</i>	7 x 0.7 mL	
<i>Detection:</i>	UV 280 nm	
	Retention time in min	purity in %
Example 109 Enantiomer 1	17.8 – 19.2	> 99.9
Example 110 Enantiomer 2	19.2 – 21.6	96.5

Enantiomer 1: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.81 (s, 1H), 7.96 (m, 1H), 7.78 (m, 2H), 7.41 (m, 2H), 7.16 (m, 1H), 6.78 (m, 2H), 4.33 (d, 1H), 4.18 (d, 1H), 3.94 (s, 3H), 3.04 (q, 2H), 1.43 (tr, 3H).

Enantiomer 2: ^1H NMR (400MHz, CDCl_3 , 300K) δ = 8.81 (s, 1H), 7.96 (m, 1H), 7.78 (m, 2H), 7.41 (m, 2H), 7.16 (m, 1H), 6.78 (m, 2H), 4.33 (d, 1H), 4.18 (d, 1H), 3.94 (s, 3H), 3.04 (q, 2H), 1.43 (tr, 3H).


Example 111:

(rac)-[3-([4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino)-5-methylbenzyl](methyl)oxido- λ^6 -sulfanylidene]cyanamide



Preparation of Intermediate 111.1:


3-(Chloromethyl)-5-methylaniline



Thionylchloride (15.3 g; 128.5 mmol) was added dropwise under stirring to an ice-cooled solution of (3-amino-5-methylphenyl)methanol (6.3 g; 42.8 mmol; GLSyntech, LLC) in DCM (140 ml). The batch was stirred at room temperature for 18 hours before it was concentrated in vacuo. The residue was taken up in DCM and concentrated once more in vacuo to give the crude product that was used without further purification.

Preparation of Intermediate 111.2:

3-Methyl-5-[(methylsulfanyl)methyl]aniline




Intermediate 111.2 was prepared under similar conditions as described in the preparation of Intermediate 1.1 using crude 3-(chloromethyl)-5-methylaniline. The batch was purified by chromatography (hexane / ethyl acetate 6:4).

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 6.26 (m, 1H), 6.21 (m, 2H), 3.43 (s, 2H), 2.07 (s, 3H), 1.89 (s, 3H).

Preparation of Intermediate 111.3:

4-Chloro-N-{3-methyl-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine




Intermediate 111.3 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3-methyl-5-[(methylsulfanyl)methyl]aniline. The crude product was purified by chromatography (hexane / ethylacetate 2:1) to give the desired product.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.63 (s, 1H), 8.59 (br, 1H), 7.37 (br, 1H), 7.30 (m, 1H), 6.87 (br, 1H), 3.60 (s, 2H), 2.25 (s, 3H), 1.95 (s, 3H).

Preparation of Intermediate 111.4:

4-(4-Fluoro-2-methoxyphenyl)-N-{3-methyl-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine




Intermediate 111.4 was prepared under similar conditions as described in the preparation of Example 42 using 4-chloro-N-{3-methyl-5-[(methylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by chromatography (hexane / ethyl acetate 1:1) to give the desired product.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.17 (s, 1H), 8.75 (s, 1H), 7.78 (br, 1H), 7.52 (m, 2H), 7.05 (m, 1H), 6.87 (m, 1H), 6.79 (m, 1H), 3.84 (s, 3H), 3.59 (s, 2H), 2.25 (s, 3H), 1.93 (s, 3H).

Preparation of Intermediate 111.5:

- 5 (rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)-λ⁴-sulfanylidene]cyanamide



- 10 Intermediate 111.5 was prepared under similar conditions as described in the preparation of Example 105.7 using 4-(4-fluoro-2-methoxyphenyl)-N-{3-methyl-5-[(methylsulfonyl)methyl]phenyl}-1,3,5-triazin-2-amine. The batch was purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.32 (s, 1H), 8.77 (s, 1H), 7.68 (m, 3H), 7.07 (m, 1H), 6.87 (m, 2H), 4.39 (d, 1H), 4.20 (d, 1H), 3.84 (s, 3H), 2.81 (s, 3H), 2.29 (s, 3H).

Preparation of end product:


Example 111 was prepared under similar conditions as described in the preparation of Example 105 using (rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)-λ⁴-sulfanylidene]cyanamide The batch was purified by preparative HPLC.

System:	Waters Autopurificationsystem: Pump 2545, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3001
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.1% HCOOH
	B = Acetonitrile
Gradient:	0–1 min 1% B, 1-8 min 1-99% B, 8-10 min 99% B
Flow:	50 mL/min
Temperature:	RT
Solution:	Max. 250 mg / max. 2.5 mL DMSO o. DMF
Injektion:	1 x 2.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.34 (s, 1H), 8.77 (s, 1H), 7.76 (br, 2H), 7.62 (m, 1H), 7.07 (m, 1H), 6.96 (m, 1H), 6.87 (m, 1H), 4.89 (m, 2H), 3.83 (s, 3H), 3.34 (s, 3H), 2.30 (s, 3H).

5 **Example 112:**

(rac)-2-((5-Fluoro-2-[4-((3-((S-methylsulfonimidoyl)methyl)phenyl)amino)-1,3,5-triazin-2-yl]phenoxy)methyl)prop-2-en-1-ol




- 10 Example 112 was prepared under similar conditions as described in the preparation of Example 65 using (rac)-ethyl [(3-([4-(2,4-difluorophenyl)-1,3,5-triazin-2-yl]amino)benzyl)(methyl)oxido-λ⁶-sulfanylidene]carbamate and 2-methylidenepropane-1,3-diol (Aldrich Chemical Company Inc.).

System:	Waters Autopurificationsystem: Pump 254, Sample Manager 2767, CFO, DAD 2996, ELSD 2424, SQD 3100
Column:	XBrigde C18 5µm 100x30 mm
Solvent:	A = H ₂ O + 0.2% Vol. NH ₃ (32%)
	B = Acetonitrile
Gradient:	0-8 min 20-30% B
Flow:	50 mL/min
Temperature:	RT
Solution:	40 mg / 1.5 mL DMSO/MeOH 1:1
Injektion:	3 x 0.5 mL
Detection:	DAD scan range 210–400 nm
	MS ESI+, ESI-, scan range 160-1000 m/z
	ELSD
Retention:	7.2 – 7.8 min


- 15 ¹H NMR (400MHz, CDCl₃, 300K) δ = 8.77 (s, 1H), 8.53 (br, 1H), 8.36 (br, 1H), 7.81 (s, 1H), 7.64 (m, 1H), 7.42 (m, 1H), 7.18 (m, 1H), 6.83 (m, 1H), 6.76 (m, 1H), 5.29 (m, 2H), 4.78 (s, 2H), 4.40 (m, 3H), 4.28 (m, 1H), 2.97 (s, 3H).

Example 113:

(rac)-[Cyclopropyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido- λ^6 -sulfanylidene]cyanamide


**Preparation of Intermediate 113.1:**

1-[(Cyclopropylsulfanyl)methyl]-3-nitrobenzene




Sulfur (3.63 g; 25.0 mmol) was added portionswise to a stirred 0.5 M solution of bromo(cyclopropyl)magnesium in THF (50.0 ml; 25.0 mmol). The batch was stirred at 50°C for 1 hour and then cooled to 0°C. Lithium tetrahydridoaluminate(1-) (522 mg; 13.8 mmol) was cautiously added under stirring. The batch was stirred for 30 minutes at 50°C and cooled to 0°C again. Water (2 ml) was cautiously added under stirring. Finally, sulfuric acid (5%; 100 ml) was cautiously added and the batch was stirred for 10 minutes. The organic phase was separated and the aqueous phase was extracted with diethyl ether (2x). The combined organic phases were washed with saturated aqueous ammonium chloride solution (2x), aqueous sodium bicarbonate solution (5%, 2x), water (2x) and saturated aqueous sodium chloride solution (2x). The organic phase was dried (Na₂SO₄) and filtered before it was slowly added to a stirred batch of 1-(chloromethyl)-3-nitrobenzene (2.15 g; 12.5 mmol) and potassium carbonate (2.59 g; 18.8 mmol) in DMF (40 ml). The batch was stirred at 85°C over night. After cooling, the batch was filtered over celite and concentrated in vacuo. The residue was taken up in ethyl acetate and washed with water (2x) and saturated aqueous sodium chloride solution (2x). The organic phase was dried (Na₂SO₄), filtered and concentrated. The residue was purified by chromatography (hexane / ethyl acetate 8:2) to give the desired product (2.38 g; 11.4 mmol).

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 8.16 (m, 1H), 8.06 (m, 1H), 7.75 (m, 1H), 7.56 (m, 1H), 3.90 (s, 2H), 1.72 (m, 1H), 0.77 (m, 2H), 0.39 (m, 2H).


Preparation of Intermediate 113.2:**3-[(Cyclopropylsulfanyl)methyl]aniline**

Intermediate 113.2 was prepared under similar conditions as described in the preparation of Intermediate 1.6 using 1-[(cyclopropylsulfanyl)methyl]-3-nitrobenzene. The batch was purified by chromatography.
¹H NMR (400MHz, d₆-DMSO, 300K) δ = 6.89 (m, 1H), 6.49 (m, 1H), 6.38 (m, 2H), 4.96 (s, 2H), 3.56 (s, 2H), 1.75 (m, 1H), 0.78 (m, 2H), 0.42 (m, 2H).

Preparation of Intermediate 113.3:**4-Chloro-N-{3-[(cyclopropylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine**

Intermediate 113.3 was prepared under similar conditions as described in the preparation of Intermediate 1.7 using 3-[(cyclopropylsulfanyl)methyl]aniline. The crude product was purified by chromatography (hexane / ethylacetate 4:1) to give the desired product.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.69 (s, 1H), 8.59 (br, 1H), 7.59 (br, 1H), 7.48 (m, 1H), 7.27 (m, 1H), 7.05 (m, 1H), 3.72 (s, 2H), 1.76 (m, 1H), 0.80 (m, 2H), 0.44 (m, 2H).


Preparation of Intermediate 113.4:**N-{3-[(Cyclopropylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine**

Intermediate 113.4 was prepared under similar conditions as described in the preparation of Example 42 using 4-chloro-N-{3-[(cyclopropylsulfanyl)methyl]phenyl}-1,3,5-triazin-2-amine and (4-fluoro-2-methoxyphenyl)boronic acid (Aldrich Chemical Company Inc). The batch was purified by chromatography (hexane / ethyl acetate 2:1) to give the desired product.

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.24 (s, 1H), 8.75 (s, 1H), 7.75 (m, 3H), 7.25 (m, 1H), 7.07 (m, 1H), 6.99 (m, 1H), 6.86 (m, 1H), 3.84 (s, 3H), 3.72 (s, 2H), 1.74 (m, 1H), 0.75 (m, 2H), 0.42 (m, 2H).

Preparation of Intermediate 113.5:

(rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)- λ^4 -sulfanylidene]cyanamide



5

Intermediate 113.5 was prepared under similar conditions as described in the preparation of Example 105.7 using N-{3-[(cyclopropylsulfanyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine. The batch was purified by chromatography (ethyl acetate / methanol 9:1).

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.38 (s, 1H), 8.78 (s, 1H), 7.63 (m, 3H), 7.37 (m, 1H), 7.14 (m, 1H), 7.06 (m, 1H), 6.88 (m, 1H), 4.53 (d, 1H), 4.38 (d, 1H), 3.85 (s, 3H), 2.70 (m, 1H), 1.07 (m, 3H), 0.84 (m, 1H).

10

Preparation of end product:

Example 113 was prepared under similar conditions as described in the preparation of Example 105 using (rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)- λ^4 -sulfanylidene]cyanamide. The batch was purified by chromatography (gradient: ethyl acetate / hexanes 4 : 1 $\rightarrow$ ethyl acetate).

15

¹H NMR (400MHz, d₆-DMSO, 300K) δ = 10.44 (s, 1H), 8.81 (s, 1H), 7.93 (m, 3H), 7.43 (m, 1H), 7.20 (m, 1H), 7.10 (m, 1H), 6.91 (m, 1H), 5.01 (m, 2H), 3.88 (br, 3H), 2.97 (m, 1H), 1.18 (m, 3H), 0.90 (m, 1H).

20

The following Table 1 provides an overview on the compounds of the invention:

Table 1

Compound No.	Structure	Nomenclature
1		(rac)-Ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
2		(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
3		(-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
4		(+)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
5		(rac)-Ethyl {[3-(4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl]amino)benzyl](methyl)oxido-λ ⁶ -sulfanylidene}carbamate
6		(rac)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
7		(-)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
8		(+)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
9		(rac)-Ethyl [(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate

Compound No.	Structure	Nomenclature
10		(rac)- 4-(4,5-Difluoro-2-methoxyphenyl)-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} - 1,3,5-triazin-2-amine
11		(rac)-Ethyl [(3-{[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
12		(rac) 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl} - 1,3,5-triazin-2-amine
13		4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl} - 1,3,5-triazin-2-amine; enantiomer 1
14		4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl} - 1,3,5-triazin-2-amine; enantiomer 2
15		(rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]-3-methylurea
16		(-)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]-3-methylurea; enantiomer 1
17		(+)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]-3-methylurea; enantiomer 2
18		(rac)-Ethyl [(3-{[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
19		(rac)-4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl} - 1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
20		(rac)-Ethyl [(3- {[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
21		(rac)-4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
22		4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
23		4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
24		(rac)-N-[(3- {[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]acetamide
25		(rac)-Ethyl [(3- {[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
26		(rac)-4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
27		4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
28		4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
29		(rac)-Ethyl [(3- {[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate

Compound No.	Structure	Nomenclature
30		(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
31		4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
32		4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
33		(rac)-Ethyl [(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
34		(rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
35		4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
36		4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
37		(rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
38		(rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
39		4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
40		4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
41		(rac)-N-{3-[(N,S-Dimethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
42		(rac)-Ethyl [{3-[(4-{2-[(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
43		(rac)-4-{2-[(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
44		(rac)-N-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]methanesulfonamide
45		(rac)-Ethyl [(3-{[4-(3-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
46		(rac)-Ethyl {[3-({4-[5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]-1,3,5-triazin-2-yl}amino)benzyl}(methyl)oxido-λ ⁶ -sulfanylidene]carbamate
47		(rac)-Ethyl [methyl(oxido)(3-{[4-(2-phenoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)-λ ⁶ -sulfanylidene]carbamate
48		(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]cyanamide
49		[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]cyanamide; enantiomer 1
50		[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido-λ ⁶ -sulfanylidene]cyanamide; enantiomer 2

Compound No.	Structure	Nomenclature
51		(rac)-Ethyl [(3-fluoro-5-[[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate
52		(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
53		4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
54		4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
55		(rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
56		(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
56.a		4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
56.b		4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
57		(rac)-4-{4-Fluoro-2-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
58		(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-(3-([S-(tetrahydro-2H-pyran-4-yl)sulfonimidoyl)methyl]phenyl)-1,3,5-triazin-2-amine
59		(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine
59.a		N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 1
59.b		N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 2
60		(rac)-Ethyl [{3-[(4-{2-[(3,4-dichlorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl}(methyl)oxido-λ⁶-sulfanylidene]carbamate
61		(rac)-4-{2-[(3,4-Dichlorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
62		(rac)-4-(4-Fluoro-2-[(²H₅)phenyl(²H₂)methyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
63		4-[2-(1-cyclopentylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of all 4 stereoisomers
64		(rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
65		(rac)-4-[4-Fluoro-2-(3,3,3-trifluoropropoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
66		(rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
67		(rac)-4-[4-Fluoro-2-(pyridin-2-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
68		(rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
69		4-{4-Fluoro-2-[1-(4-fluorophenyl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of 4 stereoisomers
70		(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ ⁶ -sulfanylidene)cyanamide
71		[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ ⁶ -sulfanylidene)cyanamide; enantiomer 1
72		[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ ⁶ -sulfanylidene)cyanamide; enantiomer 2
73		(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
74		4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
75		4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
76		(rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
77		(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
78		4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
79		4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2
80		(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
81		(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
82		(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
83		4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1
84		4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2

Compound No.	Structure	Nomenclature
85		(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
86		(rac)-4-(4-Fluoro-2-{[4-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
87		(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
88		(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
89		(rac)-4-(4-Fluoro-2-{[3-fluoro-5-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
90		(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
91		(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
92		(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
93		(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
94		(rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
95		(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine

Compound No.	Structure	Nomenclature
96		(rac)-4-{4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
97		(rac)-4-{2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
98		(rac)-3-({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl} amino)-1,3,5-triazin-2-yl]phenoxy} methyl)benzonitrile
99		(rac)-4-{4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
100		(rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
101		(rac)-4-{4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
102		(rac)-4-{2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
103		(rac)-4-{4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
104		(rac)-4-{4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine
105		(rac)-[(2,3-Difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide

Compound No.	Structure	Nomenclature
106		(rac)-N-{3,4-Difluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine
107		(rac)-[Ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido-λ ⁶ -sulfanylidene]cyanamide
108		(rac)-N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine
109		N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 1
110		N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 2
111		(rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)oxido-λ ⁶ -sulfanylidene]cyanamide
112		(rac)-2-({5-Fluoro-2-[4-(3-[(S-methylsulfonimidoyl)methyl]phenyl)amino]-1,3,5-triazin-2-yl]phenoxy}methyl)prop-2-en-1-ol
113		(rac)-[Cyclopropyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido-λ ⁶ -sulfanylidene]cyanamide

Results:**Table 2:** Inhibition of CDK9 and CDK2 of compounds according to the present invention

The IC₅₀ (inhibitory concentration at 50% of maximal effect) values are indicated in nM or μ M, “n.t.” means that the compounds have not been tested in this assay.

①: Compound Number

②: CDK9 CDK9/CycT1 kinase assay as described under Method 1. of Materials and Methods

③: CDK2 CDK2/CycE kinase assay as described under Method 2. of Materials and Methods

①	Nomenclature	②	③
1	(rac)-Ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	16 nM	2200 nM
2	(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	25 nM	2100 nM
3	(-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	16 nM	1700 nM
4	(+)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	13 nM	1300 nM
5	Ethyl {3-([4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl]amino)benzyl}(methyl)oxido- λ^6 -sulfanylidene} carbamate	4 nM	250 nM
6	4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	3 nM	220 nM
9	(rac)-Ethyl [(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	8 nM	690 nM
10	(rac)- 4-(4,5-Difluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	28 nM	1300 nM
11	(rac)-Ethyl [(3-{[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	23 nM	5500 nM
12	(rac) 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	74 nM	6800 nM
13	4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	93 nM	9500 nM
14	4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	160 nM	13000 nM
15	(rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea	7 nM	930 nM

①	Nomenclature	②	③
16	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 1	19 nM	1100 nM
17	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 2	14 nM	1200 nM
18	(rac)-Ethyl [(3-{[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	33 nM	3900 nM
19	(rac)-4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	59 nM	2100 nM
20	(rac)-Ethyl [(3-{[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	22 nM	2000 nM
21	(rac)-4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	56 nM	3500 nM
22	4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	91 nM	2800 nM
23	4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	110 nM	5600 nM
24	(rac)-N-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]acetamide	11 nM	1800 nM
25	(rac)-Ethyl [(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	28 nM	6300 nM
26	(rac)-4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	35 nM	4600 nM
27	4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	100 nM	7600 nM
28	4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	89 nM	5900 nM
29	(rac)-Ethyl [(3-{[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	13 nM	1200 nM
30	(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	21 nM	940 nM
31	4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	22 nM	670 nM
32	4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	40 nM	1600 nM
33	(rac)-Ethyl [(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	21 nM	3100 nM

①	Nomenclature	②	③
34	(rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	68 nM	2200 nM
35	4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	64 nM	3100 nM
36	4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	50 nM	2900 nM
37	(rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	9 nM	2300 nM
38	(rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	14 nM	1600 nM
39	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	29 nM	1500 nM
40	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	25 nM	1800 nM
41	(rac)-N-{3-[(N,S-Dimethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	36 nM	2600 nM
42	(rac)-Ethyl [(3-{[4-(2-{[4-(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	5 nM	1600 nM
43	(rac)-4-{2-[(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	9 nM	1900 nM
44	(rac)-N-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]methanesulfonamide	20 nM	1800 nM
45	(rac)-Ethyl [(3-{[4-(3-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	820 nM	9000 nM
46	(rac)-Ethyl [(3-{[4-[5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	120 nM	11000 nM
47	(rac)-Ethyl [methyl(oxido)(3-{[4-(2-phenoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)- λ^6 -sulfanylidene]carbamate	35 nM	6500 nM
48	(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	13 nM	630 nM
49	[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 1	7 nM	650 nM
50	[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2	7 nM	430 nM
51	(rac)-Ethyl [(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	10 nM	430 nM

①	Nomenclature	②	③
52	(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	17 nM	1000 nM
53	4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	16 nM	540 nM
54	4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	16 nM	850 nM
55	(rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	10 nM	2200 nM
56	(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	6 nM	410 nM
56.a	4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	4 nM	560 nM
56.b	4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	5 nM	560 nM
57	(rac)-4-{4-Fluoro-2-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	370 nM	8300 nM
58	(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-(3-{[S-(tetrahydro-2H-pyran-4-yl)sulfonimidoyl]methyl}phenyl)-1,3,5-triazin-2-amine	89 nM	5200 nM
59	(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	26 nM	1400 nM
59.a	N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 1	25 nM	2900 nM
59.b	N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 2	33 nM	1800 nM
60	(rac)-Ethyl [3-[(4-{2-[(3,4-dichlorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl] (methyl)oxido- λ^6 -sulfanylidene]carbamate	13 nM	3300 nM
61	(rac)-4-{2-[(3,4-Dichlorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	15 nM	1200 nM
62	(rac)-4-(4-Fluoro-2-{[($^2\text{H}_5$)phenyl($^2\text{H}_2$)methyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	7 nM	220 nM
63	4-[2-(1-cyclopentylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of all 4 stereoisomers	99 nM	n.t.
64	(rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	15 nM	250 nM
65	(rac)-4-[4-Fluoro-2-(3,3,3-trifluoropropoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	79 nM	6000 nM
66	(rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	24 nM	1600 nM

①	Nomenclature	②	③
67	(rac)-4-[4-Fluoro-2-(pyridin-2-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	97 nM	6900 nM
68	(rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	14 nM	1500 nM
69	4-{4-Fluoro-2-[1-(4-fluorophenyl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of 4 stereoisomers	22 nM	3500 nM
70	(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	6 nM	180 nM
71	[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 1	6 nM	230 nM
72	[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2	5 nM	240 nM
73	(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	8 nM	820 nM
74	4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	3 nM	590 nM
75	4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	4 nM	450 nM
76	(rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	30 nM	3200 nM
77	(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	15 nM	1600 nM
78	4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	9 nM	580 nM
79	4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	5 nM	440 nM
80	(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	13 nM	1000 nM
81	(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	23 nM	1500 nM
82	(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	7 nM	360 nM
83	4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	2 nM	270 nM
84	4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	3 nM	220 nM

①	Nomenclature	②	③
85	(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	9 nM	380 nM
86	(rac)-4-(4-Fluoro-2-{4-(trifluoromethyl)benzyl}oxy)phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	8 nM	1400 nM
87	(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	3 nM	680 nM
88	(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	11 nM	1300 nM
89	(rac)-4-(4-Fluoro-2-{3-fluoro-5-(trifluoromethyl)benzyl}oxy)phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	4 nM	280 nM
90	(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	2 nM	260 nM
91	(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	29 nM	2300 nM
92	(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	4 nM	250 nM
93	(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	7 nM	2000 nM
94	(rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	3 nM	230 nM
95	(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	21 nM	1800 nM
96	(rac)-4-{4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	3 nM	350 nM
97	(rac)-4-{2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	3 nM	200 nM
98	(rac)-3-(5-Fluoro-2-[4-(3-[(S-methylsulfonimidoyl)methyl]phenyl)amino]-1,3,5-triazin-2-yl)phenoxy)methyl)benzonitrile	3 nM	180 nM
99	(rac)-4-{4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	2 nM	250 nM
100	(rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	6 nM	1300 nM
101	(rac)-4-{4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	4 nM	500 nM
102	(rac)-4-{2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	7 nM	800 nM

①	Nomenclature	②	③
103	(rac)-4-{4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	7 nM	1200 nM
104	(rac)-4-{4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	2 nM	54 nM
105	(rac)-[(2,3-Difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	3 nM	140 nM
106	(rac)-N-{3,4-Difluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	7 nM	400 nM
107	(rac)-[Ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido- λ^6 -sulfanylidene]cyanamide	4 nM	360 nM
108	(rac)-N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	8 nM	1000 nM
109	N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 1	24	780
110	N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 2	9	1100
111	(rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	3 nM	250 nM
112	(rac)-2-(5-Fluoro-2-[4-(3-[(S-methylsulfonimidoyl)methyl]phenyl)amino]-1,3,5-triazin-2-yl]phenoxy)methylprop-2-en-1-ol	4 nM	730 nM
113	(rac)-[Cyclopropyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido- λ^6 -sulfanylidene]cyanamide	2nM	300nM

Table 3: Inhibition of proliferation of HeLa, HeLa/MaTu/ADR, NCI-H460, DU145, Caco-2 and B16F10 cells by compounds according to the present invention. Determined as described above (Method 3. of Materials and Methods section). All IC₅₀ (inhibitory concentration at 50% of maximal effect) values are indicated in nM, “n.t.” means that the compounds have not been tested in this assay.

- ①: Compound Number
- ②: Inhibition of HeLa cell proliferation
- ③: Inhibition of HeLa/MaTu/ADR cell proliferation
- ④: Inhibition of NCI-H460 cell proliferation
- ⑤: Inhibition of DU145 cell proliferation
- ⑥: Inhibition of Caco-2 cell proliferation
- ⑦: Inhibition of B16F10 cell proliferation

①	Nomenclature	②	③	④	⑤	⑥	⑦
1	(rac)-Ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	1000	380	360	390	390	360
2	(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
3	(-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	1100	400	1300	740	1300	1000
4	(+)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	970	440	1200	670	1300	1000
5	Ethyl {[3-(4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl]amino)benzyl](methyl)oxido- λ^6 -sulfanylidene} carbamate	100	270	210	260	250	270
6	4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	<100	200	160	180	270	250
9	(rac)-Ethyl [(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	400	960	980	900	800	1000
10	(rac)-4-(4,5-Difluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
15	(rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea	390	940	1000	370	1000	620
16	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 1	950	n.t.	n.t.	n.t.	n.t.	n.t.
17	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 2	580	600	540	580	800	410
24	(rac)-N-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]acetamide	490	520	760	680	710	940
25	(rac)-Ethyl [(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	1100	n.t.	n.t.	n.t.	n.t.	n.t.
29	(rac)-Ethyl [(3-{[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	510	760	720	540	760	960
30	(rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1200	n.t.	n.t.	n.t.	n.t.	n.t.
37	(rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	630	750	920	500	810	720

①	Nomenclature	②	③	④	⑤	⑥	⑦
39	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	1100	n.t.	n.t.	n.t.	n.t.	n.t.
40	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	1100	n.t.	n.t.	n.t.	n.t.	n.t.
42	(rac)-Ethyl [(3-[(4-{2-[(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl)amino]benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	610	520	370	590	680	680
43	(rac)-4-{2-[(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	590	1100	530	800	1100	800
44	(rac)-N-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]methanesulfonamide	1000	n.t.	n.t.	n.t.	n.t.	n.t.
48	(rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	330	400	410	280	420	320
50	[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2	260	190	310	140	350	250
51	(rac)-Ethyl [(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate	300	240	230	240	290	340
52	(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	410	140	650	340	690	550
55	(rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1000	850	980	990	890	n.t.
56	(rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	370	780	350	370	860	470
56.a	4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	350	270	340	340	430	340
56.b	4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	540	540	470	400	540	390
59	(rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	860	n.t.	n.t.	n.t.	n.t.	n.t.
59.a	N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 1	1100	n.t.	n.t.	n.t.	n.t.	n.t.

59.b	N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine; enantiomer 2	1100	n.t.	n.t.	n.t.	n.t.	n.t.
64	(rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	400	280	360	290	350	320
66	(rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1700	n.t.	n.t.	n.t.	n.t.	n.t.
68	(rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	410	3000	1100	460	1300	740
70	(rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ^6 -sulfanylidene)cyanamide	150	120	230	79	170	130
71	[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ^6 -sulfanylidene)cyanamide; enantiomer 1	100	120	160	120	130	110
72	[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl- λ^6 -sulfanylidene)cyanamide; enantiomer 2	71	120	160	110	190	160
73	(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	240	280	260	200	300	150
74	4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	310	340	330	360	390	340
75	4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	260	210	220	170	310	240
76	(rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
77	(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	470	630	560	480	300	520
78	4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	650	260	390	370	760	500
79	4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	440	350	420	380	390	370
80	(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	730	n.t.	n.t.	n.t.	n.t.	n.t.

81	(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	860	n.t.	n.t.	n.t.	n.t.	n.t.
82	(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	180	190	180	130	180	150
83	4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	110	640	180	180	210	220
84	4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	120	340	180	180	180	210
85	(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	330	560	230	190	51	280
86	(rac)-4-(4-Fluoro-2-{[4-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
87	(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	560	450	570	610	800	590
88	(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	380	700	890	730	780	810
89	(rac)-4-(4-Fluoro-2-{[3-fluoro-5-(trifluoromethyl)benzyl]oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	470	450	370	420	520	470
90	(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	170	100	110	110	80	100
91	(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
92	(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	470	360	170	380	430	390
93	(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	680	n.t.	n.t.	n.t.	n.t.	n.t.
94	(rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	110	190	200	140	220	180

95	(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
96	(rac)-4-{4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1000	n.t.	n.t.	n.t.	n.t.	n.t.
97	(rac)-4-{2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	940	n.t.	n.t.	n.t.	n.t.	n.t.
98	(rac)-3-({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl} amino)-1,3,5-triazin-2-yl]phenoxy} methyl)benzonitrile	140	260	120	120	150	130
99	(rac)-4-{4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	120	180	150	120	230	200
100	(rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	920	n.t.	n.t.	n.t.	n.t.	n.t.
101	(rac)-4-{4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	330	350	360	340	240	340
102	(rac)-4-{2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	340	140	150	95	160	140
103	(rac)-4-{4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1100	n.t.	n.t.	n.t.	n.t.	n.t.
104	(rac)-4-{4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	110	240	110	120	150	120
105	(rac)-[(2,3-Difluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	110	71	130	120	81	140
107	(rac)-[Ethyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)oxido- λ^6 -sulfanylidene]cyanamide	230	340	450	280	500	300
108	(rac)-N-{3-[(S-ethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine	1000	n.t.	n.t.	n.t.	n.t.	n.t.
111	(rac)-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}-5-methylbenzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide	100	170	270	110	220	130
112	(rac)-2-({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl} amino)-1,3,5-triazin-2-yl]phenoxy} methyl)prop-2-en-1-ol	330	390	320	300	350	320

Table 4: Thermodynamic solubility of compounds according to the present invention in water at pH 6.5 as determined by the equilibrium shake flask method described under Method 4. of Materials and Methods.

5 ①: Compound Number

②: Solubility in mg/l.

①	Nomenclature	②
3	(-)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	601
4	(+)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	479
10	(rac)- 4-(4,5-Difluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	130
16	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 1	323
17	1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 2	429
22	4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	787
23	4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	888
31	4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	1000
32	4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	1000
39	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	819
40	4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	865
49	[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 1	88
50	[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2	126
54	4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	120
73	(rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	1220

①	Nomenclature	②
77	(rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	256
88	(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	884
91	(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine	104

Table 5: Inhibition of Carbonic anhydrase-1 and Carbonic anhydrase-2 as determined by the Carbonic anhydrase Assay described above

5


- ①: Compound Number
- ②: Inhibition of Carbonic anhydrase-1: the IC₅₀ (inhibitory concentration at 50% of maximal effect) values are indicated in (mol/l)
- ③: Inhibition of Carbonic anhydrase-2: the IC₅₀ (inhibitory concentration at 50% of maximal effect) values are indicated in (mol/l)

10

①	Nomenclature	②	③
3	4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1	>1.0E-05	>1.0E-05
4	4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2	>1.0E-05	>1.0E-05

Patent claims

1. A compound of general formula (I)




(I), wherein

R^1 represents a group selected from C_1 - C_6 -alkyl-, C_3 - C_7 -cycloalkyl-, heterocyclyl-, phenyl, heteroraryl, phenyl- C_1 - C_3 -alkyl- or heteroaryl- C_1 - C_3 -alkyl-,

wherein said group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of hydroxy, cyano, halogen, halo- C_1 - C_3 -alkyl-, C_1 - C_6 -alkoxy-, C_1 - C_3 -fluoroalkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines;

R^2 represents a group selected from



- R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;
- R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, C_1 - C_6 -alkyl-, C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl, heteroaryl
 5 wherein said C_1 - C_6 -alkyl- C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, cyano, hydroxy, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-,
 10 cyclic amines, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;
- R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;
- R^8 represents a group selected from
 15 a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-, C_1 - C_3 -alkoxy-,
 20 C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-, C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl, heteroaryl, wherein said C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, cyano, C_1 - C_3 -alkyl-, C_1 - C_3 -alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-;
- b) a C_3 - C_7 -cycloalkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-, C_1 - C_3 -alkoxy-,
 25 C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-;
- c) a heterocycyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-, C_1 - C_3 -fluoroalkoxy-, C_1 - C_3 -alkoxy-,
 30 C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-;

d) a phenyl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

5 e) a heteroaryl group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

10 f) a phenyl-C₁-C₃-alkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

15 g) a heteroaryl-C₁-C₃-alkyl- group, which is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, hydroxy, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, cyano, C₁-C₃-alkyl-, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-, C₁-C₃-alkoxy-;

20 R⁹ represents a group selected from C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl, benzyl or heteroaryl wherein said group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-;

25 R¹⁰, R¹¹ represent, independently from each other, a group selected from hydrogen, C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl or heteroaryl wherein said C₁-C₆-alkyl-, C₃-C₇-cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, hydroxy, C₁-C₃-alkyl-, C₁-C₃-alkoxy-, amino, alkylamino-, dialkylamino-, acetylamino-, N-methyl-N-acetylamino-, cyclic amines, halo-C₁-C₃-alkyl-, C₁-C₃-fluoroalkoxy-;


or its salts, solvates or salts of solvates.

30

2. The compound according to claim 1, wherein

R¹ represents a group selected from C₁-C₆-alkyl-, C₃-C₇-cycloalkyl- or heterocycyl-;

R^2 represents a group selected from



R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom or a C_1 - C_3 -alkyl-;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$, C_1 - C_6 -alkyl-;

R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two or three substituents, selected from the group consisting of halogen, hydroxy, C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-, C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl, heteroaryl, wherein said C_3 - C_7 -cycloalkyl-, heterocycyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-;

b) a phenyl group;

c) a phenyl- C_1 - C_3 -alkyl- group, which phenyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, cyano, C_1 - C_3 -alkyl-, halo- C_1 - C_3 -alkyl-;


d) a heteroaryl- C_1 - C_3 -alkyl- group, which heteroaryl group is optionally substituted with one substituent selected from the group consisting of halogen;

R^9 represents a C_1 - C_6 -alkyl group;

R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen or C_1 - C_6 -alkyl-;

or its salts, solvates or salts of solvates.

3. The compound according to claim 1 or 2 of general formula (Ia)




(Ia)

, wherein

R^1 represents a C_1 - C_6 -alkyl- or C_3 - C_7 -cycloalkyl- group;

R^2 represents a group selected from



R^3 , R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom, C_1 - C_3 -alkyl-;

R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^{10}R^{11}$;

R^6 , R^7 represent, independently from each other, a hydrogen atom or fluoro atom;

R^8 represents a group selected from

a) a C_1 - C_{10} -alkyl group, which is optionally substituted with one or two substituents, selected from the group consisting of hydroxy, C_2 - C_3 -alkenyl-, C_2 - C_3 -alkynyl-, C_3 - C_7 -cycloalkyl-, phenyl, heteroaryl, wherein said C_3 - C_7 -cycloalkyl-, phenyl or heteroaryl group is optionally substituted with one, two or three substituents, identically or differently, selected from halogen, cyano, halo- C_1 - C_3 -alkyl-;

b) a phenyl- C_1 - C_3 -alkyl- group, which phenyl group is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of halogen, cyano, halo- C_1 - C_3 -alkyl-;


c) a heteroaryl-C₁-C₃-alkyl- group, which heteroaryl group is optionally substituted with one substituent selected from the group consisting of halogen;

R⁹ represents a C₁-C₆-alkyl group;

R¹⁰, R¹¹ represent, independently from each other, a group selected from hydrogen, C₁-C₆-alkyl-;

5 or its salts, solvates or salts of solvates.

4. The compound according to claim 1, 2 or 3 of general formula (Ia)



, wherein

R¹ represents a group selected from methyl;

10 R² represents a group selected from 4-fluoro-2-methoxyphenyl-, 2-(benzyloxy)-4-fluorophenyl-, 4,5-difluoro-2-methoxyphenyl-, 2-[(4-fluorobenzyl)oxy]phenyl-, 4-fluoro-2-[(4-fluorobenzyl)oxy]phenyl ;

R³ represents a group selected from a hydrogen atom, fluoro atom, chloro atom;


R⁴ represents a group selected from a hydrogen atom, chloro atom;

15 R⁵ represents a group selected from a hydrogen atom, -C(O)R⁹, -C(O)OR⁹;

R⁹ represents a group selected from methyl or ethyl;

or its salts, solvates or salts of solvates.

5. The compound according to claim 1, 2, 3 or 4 of general formula (Ia)




, wherein

- R^1 represents a group selected from methyl;
- R^2 represents a group selected from 4-fluoro-2-methoxyphenyl- or 4,5-difluoro-2-methoxyphenyl-;
- R^3, R^4 represent, independently from each other, a hydrogen atom;
- 5 R^5 represents a hydrogen atom;
- or its salts, solvates or salts of solvates.

6. The compound according to claim 1 or 2, wherein

- R^1 represents a group selected from methyl, ethyl, cyclopropyl, tetrahydro-2H-pyran-yl-;
- 10 R^2 represents a group selected from



- R^3, R^4 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom or methyl;

- R^5 represents a group selected from a hydrogen atom, cyano, $-C(O)R^9$, $-C(O)OR^9$, $-S(O)_2R^9$, $-C(O)NR^{10}R^{11}$ or C_1 - C_6 -alkyl-;

- 15 R^6, R^7 represent, independently from each other, a group selected from a hydrogen atom, fluoro atom, chloro atom;

- R^8 represents a group selected from

a) a methyl, ethyl propyl or butyl group, which group is optionally substituted with one or two or three substituents, selected from the group consisting of halogen, hydroxy, ethenyl, propenyl, ethynyl-, propynyl-, cyclopentyl-, cyclohexyl, tetrahydro-2H-pyran-yl-, phenyl,

pyridinyl-, thiazolyl-, oxazolyl, wherein said phenyl or pyridinyl- group is optionally substituted with one, two or three substituents, identically or differently, selected from a fluoro or chloro atom, cyano, methyl, or trifluoromethyl;

b) a ($^2\text{H}_2$)methyl group substituted with a ($^2\text{H}_5$)phenyl group;

5 c) a phenyl group;

d) a benzyl group, which phenyl ring is optionally substituted with one or two or three substituents, identically or differently, selected from the group consisting of a fluoro atom, chloro atom, cyano, methyl or trifluoromethyl-;

10 e) a pyridin-2-ylmethyl- group, which pyridin is optionally substituted with one fluoro atom;

f) a pyridin-3-ylmethyl- group, which pyridin is optionally substituted with one fluoro atom;

g) a pyridin-4-ylmethyl- group, which pyridin is optionally substituted with one fluoro atom;

15 h) a thiazolylmethyl- group;

i) an oxazolylmethyl- group

R^9 represents a methyl or ethyl group;

R^{10} , R^{11} represent, independently from each other, a group selected from hydrogen, methyl;


or its salts, solvates or salts of solvates.

20

7. The compound according to any one of claims 1, 2 or 3, wherein

R^1 represents a methyl or ethyl or cyclopropyl group;

R^2 represents a group selected from



R^3 represents a group selected from a hydrogen atom, fluoro atom, chloro atom or methyl;

R⁴ represents a group selected from a hydrogen atom or fluoro atom

R⁵ represents a group selected from a hydrogen atom, cyano, -C(O)R⁹, -C(O)OR⁹,
-C(O)NR¹⁰R¹¹;

R⁶, R⁷ represent, independently from each other, a group selected from a hydrogen atom or fluoro
atom;

R⁸ represents a group selected from

methyl, ethyl, prop-2-en-1-yl-, 2-methylprop-2-en-1-yl-, 2-(hydroxymethyl)prop-2-en-1-yl-,
(2Z)-but-2-en-1-yl-, prop-2-yn-1-yl-, but-2-yn-1-yl-, cyclohexylmethyl-, benzyl-,
3-cyanobenzyl-, 3-fluorobenzyl-, 3-chlorobenzyl-, 4-fluorobenzyl-, 4-chlorobenzyl-,
3-fluoro-5-(trifluoromethyl)benzyl-, 3-chloro-5-fluorobenzyl-, pyridin-4-yl-,
2-fluoropyridin-4-yl-, 2,3,5-trifluorobenzyl-, 3,4,5-trifluorobenzyl-.

R⁹ represents a methyl or ethyl group;

R¹⁰, R¹¹ represent, independently from each other, a group selected from hydrogen, C₁-C₆-alkyl-;

or its salts, solvates or salts of solvates.

8. The compound according to claim 1, which is selected from

(rac)-Ethyl [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido-λ⁶-
sulfanylidene]carbamate,

(rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-
amine,

4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine;
enantiomer 1,

4-(4-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine;
enantiomer 2,

(rac)-Ethyl {[3-({4-[2-(benzyloxy)-4-fluorophenyl]-1,3,5-triazin-2-yl} amino)benzyl](methyl)oxido-λ⁶-
sulfanylidene} carbamate,

(rac)-4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-
amine,

4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-
amine; enantiomer 1,

- 4-[2-(Benzyloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-Ethyl [(3-{[4-(4,5-difluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)- 4-(4,5-Difluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-Ethyl [(3-{[4-(4-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac) 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(4-Chloro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea,
- 1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 1,
- 1-[(3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]-3-methylurea; enantiomer 2,
- (rac)-Ethyl [(3-{[4-(2,2-difluoro-1,3-benzodioxol-4-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(2,2-Difluoro-1,3-benzodioxol-4-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-Ethyl [(3-{[4-(5-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(5-Fluoro-2-methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-N-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]acetamide,
- (rac)-Ethyl [(3-{[4-(2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

- 4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(2-Methoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-Ethyl [(3-{[4-(3,4-dihydro-2H-chromen-8-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(3,4-Dihydro-2H-chromen-8-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-Ethyl [(3-{[4-(2,3-dihydro-1-benzofuran-7-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(2,3-Dihydro-1-benzofuran-7-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-Ethyl [(3-{[4-(2,3-dihydro-1,4-benzodioxin-5-yl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
- 4-(2,3-Dihydro-1,4-benzodioxin-5-yl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
- (rac)-N-{3-[(N,S-Dimethylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
- (rac)-Ethyl [(3-{[4-{2-[(4-fluorobenzyl)oxy]phenyl}-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-{2-[(4-Fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-N-{[3-{[4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]methanesulfonamide,

- (rac)-Ethyl [(3-{[4-(3-chloro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-Ethyl {[3-({4-[5-fluoro-2-(tetrahydro-2H-pyran-4-ylmethoxy)phenyl]-1,3,5-triazin-2-yl} amino)benzyl](methyl)oxido- λ^6 -sulfanylidene} carbamate,
- (rac)-Ethyl [methyl(oxido)(3-{[4-(2-phenoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)- λ^6 -sulfanylidene]carbamate,
- (rac)-[(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,
- [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 1,
- [(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide; enantiomer 2,
- (rac)-Ethyl [(3-fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)-oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- 4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, enantiomer 1,
- 4-(4-Fluoro-2-methoxyphenyl)-N-{3-fluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, enantiomer 2,
- (rac)-4-[2-(Cyclohexylmethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-4-{4-Fluoro-2-[(4-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-4-{4-Fluoro-2-[2-(tetrahydro-2H-pyran-4-yl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
- (rac)-4-(4-Fluoro-2-methoxyphenyl)-N-(3-{[S-(tetrahydro-2H-pyran-4-yl)sulfonimidoyl]methyl} phenyl)-1,3,5-triazin-2-amine,
- (rac)-N-{4-Chloro-3-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
- (rac)-Ethyl [(3-{[4-(2-(3,4-dichlorobenzyl)oxy]phenyl)-1,3,5-triazin-2-yl]amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]carbamate,
- (rac)-4-{2-[(3,4-Dichlorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-{{(2H₅)phenyl(2H₂)methyl}oxy}phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 4-[2-(1-cyclopentylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 (rac)-N-{3-Chloro-5-[(S-methylsulfonimidoyl)methyl]phenyl}-4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
 (rac)-4-[4-Fluoro-2-(3,3,3-trifluoropropoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 (rac)-4-[4-Fluoro-2-(pyridin-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 (rac)-4-[4-Fluoro-2-(pyridin-2-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 (rac)-4-[4-Fluoro-2-(pyridin-4-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 4-{4-Fluoro-2-[1-(4-fluorophenyl)ethoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine, mixture of 4 stereoisomers,
 (rac)-[(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ6-sulfanylidene)cyanamide,
 [(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ6-sulfanylidene)cyanamide; enantiomer 1,
 [(3-Fluoro-5-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)(methyl-λ6-sulfanylidene)cyanamide; enantiomer 2,
 (rac)-4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,
 4-[2-(But-2-yn-1-yloxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,
 (rac)-4-[2-(2-Cyclopropylethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 (rac)-4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,
 4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,

4-[4-Fluoro-2-(prop-2-yn-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,

(rac)-4-{2-[(3,4-Difluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[4-Fluoro-2-(1,3-thiazol-5-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 1,

4-{4-Fluoro-2-[(2-fluoropyridin-4-yl)methoxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine; enantiomer 2,

(rac)-4-[4-Fluoro-2-(prop-2-en-1-yloxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-{4-(trifluoromethyl)benzyl}oxy)phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(4-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(2-Ethoxy-4-fluorophenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-{3-fluoro-5-(trifluoromethyl)benzyl}oxy)phenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{4-Fluoro-2-[(3-fluorobenzyl)oxy]phenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-(4-Fluoro-2-propoxyphenyl)-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(3-Chlorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[4-Fluoro-2-(1,2-oxazol-3-ylmethoxy)phenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-{2-[(3-Chloro-5-fluorobenzyl)oxy]-4-fluorophenyl}-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

(rac)-4-[2-(2,2-Difluoroethoxy)-4-fluorophenyl]-N-{3-[(S-methylsulfonimidoyl)methyl]phenyl}-1,3,5-triazin-2-amine,

- (rac)-4- {4-Fluoro-2-[(4-fluoro-3-methylbenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4- {2-[(3-Chloro-4-fluorobenzyl)oxy]-4-fluorophenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-3- ({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl} amino)-1,3,5-triazin-2-yl]phenoxy} methyl)benzonitrile,
- (rac)-4- {4-Fluoro-2-[(2-methylprop-2-en-1-yl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4-[4-Fluoro-2-(4,4,4-trifluorobutoxy)phenyl]-N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4- {4-Fluoro-2-[(2,3,5-trifluorobenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4- {2-[(2Z)-But-2-en-1-yloxy]-4-fluorophenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4- {4-Fluoro-2-[(2,4,5-trifluorobenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-4- {4-Fluoro-2-[(3,4,5-trifluorobenzyl)oxy]phenyl} -N- {3-[(S-methylsulfonimidoyl)methyl]phenyl} -1,3,5-triazin-2-amine,
- (rac)-[(2,3-Difluoro-5- {4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl} amino} benzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,
- (rac)-N- {3,4-Difluoro-5-[(S-methylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine,
- (rac)-[Ethyl(3- {4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl} amino} benzyl)oxido- λ^6 -sulfanylidene]cyanamide,
- (rac)-N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine
N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine;
enantiomer 1,
- N- {3-[(S-ethylsulfonimidoyl)methyl]phenyl} -4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-amine;
enantiomer 2,
- (rac)-[(3- {4-(4-Fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl} amino} -5-methylbenzyl)(methyl)oxido- λ^6 -sulfanylidene]cyanamide,

(rac)-2-({5-Fluoro-2-[4-({3-[(S-methylsulfonimidoyl)methyl]phenyl}amino)-1,3,5-triazin-2-yl]phenoxy}methyl)prop-2-en-1-ol,
 (rac)-[Cyclopropyl(3-{[4-(4-fluoro-2-methoxyphenyl)-1,3,5-triazin-2-yl]amino}benzyl)oxido- λ^6 -sulfanylidene]cyanamide,
 or its salts, solvates or salts of solvates.

9. A pharmaceutical combination comprising a compound according to any one of claims 1 to 8 in combination with at least one or more further active ingredients.

5

10. A pharmaceutical composition comprising a compound according to any one of claims 1 to 8 in combination with an inert, nontoxic, pharmaceutically suitable adjuvant.


10

11. The pharmaceutical combination according to claim 9 for use in the treatment and/or prophylaxis of hyper-proliferative disorders, virally induced infectious diseases and/or of cardiovascular diseases.

12. The pharmaceutical composition according to claim 10 for use in the treatment and/or prophylaxis of hyper-proliferative disorders, virally induced infectious diseases and/or of cardiovascular diseases.

15

13. A compound of general formula (3)




, wherein

R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) according to any one of claims 1 to 7.

20


14. A compound of general formula (2)



R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) according to any one of claims 1 to 7.

5


15. A compound of general formula (11)



R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) according to any one of claims 1 to 7.

10

16. A compound of general formula (12)



R^1 , R^3 and R^4 are as defined for the compound of general formula (I) or (Ia) according to any one of claims 1 to 7.

15