

METHODS OF MAKING CHIMERIC ANTIGEN RECEPTOR –EXPRESSING CELLS

This application claims priority to U.S. Serial No. 62/097,375 filed December 29, 2014 and U.S. Serial No. 62/133,137 filed March 13, 2015, the contents of which are incorporated
5 herein by reference in their entireties.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said
10 ASCII copy, created on December 18, 2015, is named N2067-7067WO_SL.txt and is 246,147 bytes in size.

FIELD OF THE INVENTION

The present invention relates generally to methods of making immune effector cells (e.g., T cells, NK cells) engineered to express a Chimeric Antigen Receptor (CAR), and
15 compositions comprising the same.

BACKGROUND OF THE INVENTION

Adoptive cell transfer (ACT) therapy with autologous T-cells, especially with T-cells transduced with Chimeric Antigen Receptors (CARs), has shown promise in several
20 hematologic cancer trials.

SUMMARY OF THE INVENTION

The present disclosure pertains to methods of making immune effector cells (e.g., T cells, NK cells) that can be engineered to express a CAR, and compositions comprising the
25 same.

Accordingly, in one aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells) that can be engineered to express a CAR, the method comprising providing a population of immune effector cells (e.g., T cells), removing T

regulatory cells, e.g., CD25+ T cells, from the population, to thereby provide a population of T regulatory-depleted cells, e.g., CD25+ depleted cells, that are suitable for expression of a CAR.

In one embodiment, the population of T regulatory-depleted cells contains less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells.

5 In one embodiment, the population of immune effector cells are cells of a subject having cancer, e.g., a subject having a CD25 expressing cancer such as, e.g., chronic lymphocytic leukemia (CLL). In one embodiment, the population of T regulatory-depleted cells contains less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of
10 tumor cells.

In one embodiment, the population of immune effector cells are autologous to the subject who the cells will be administered to for treatment. In one embodiment, the population of immune effector cells are allogeneic to the subject who the cells will be administered to for treatment.

15 In one embodiment, the T regulatory cells, e.g., CD25+ T cells, are removed from the population using an anti-CD25 antibody, or fragment thereof, or a CD25-binding ligand, e.g. IL-2. In one embodiment, the anti-CD25 antibody, or fragment thereof, or CD25-binding ligand is conjugated to a substrate, e.g., a bead, or is otherwise coated on a substrate, e.g., a bead. In one embodiment, the anti-CD25 antibody, or fragment thereof, is conjugated to a
20 substrate as described herein. In one embodiment, the T regulatory cells, e.g., CD25+ T cells, are removed from the population using an anti-CD25 antibody molecule, or fragment thereof.

In one embodiment, the T regulatory cells, e.g., CD25+ T cells, are removed from the population using CD25 depletion reagent from MiltenyiTM. In one embodiment, the ratio of cells to CD25 depletion reagent is 1e7 cells to 20 uL, or 1e7 cells to 15 uL, or 1e7 cells to 10
25 uL, or 1e7 cells to 5 uL, or 1e7 cells to 2.5 uL, or 1e7 cells to 1.25 uL.

In one embodiment, the population of T regulatory-depleted cells, e.g., CD25+ depleted cells, are suitable for expression of a CAR described herein, e.g., a CD19 CAR described herein. In one embodiment, the population of immune effector cells are obtained from a subject having a haematological cancer, e.g., a leukemia, e.g., chronic lymphocytic leukemia
30 (CLL), acute lymphocytic leukemia (ALL), or a lymphoma, e.g., mantle cell lymphoma (MCL)

or Hodgkin lymphoma (HL). In one embodiment, the population of T regulatory-depleted cells contains less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of the leukemia cells, e.g., CLL cells, ALL cells, or lymphoma cells, e.g., MCL cells or HL cells. In one embodiment, the population of immune effector cells are obtained from a subject having CLL, and the population of T regulatory-depleted cells, e.g., CD25+ depleted cells, contains less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of the leukemia cells, e.g., CLL cells and are suitable for expression of a CD19 CAR described herein. In one embodiment, the population of T regulatory-depleted cells contains less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the population of T regulatory-depleted cells contains less than 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells.

In one embodiment, the population of immune effector cells are T cells isolated from peripheral blood lymphocytes. In an embodiment, the population of T cells are obtained by lysing the red blood cells and/or by depleting the monocytes. In an embodiment, the population of T cells is isolated from peripheral lymphocytes using, e.g., a method described herein.

In one embodiment, the population of immune effector cells can be obtained from a blood sample from a subject, e.g., obtained by apheresis. In one embodiment, the cells collected by apheresis are washed to remove the plasma fraction and, optionally, the cells are provided in an appropriate buffer or media for subsequent processing steps. In one embodiment, the cells are washed with a buffer such as, e.g., phosphate buffered saline (PBS). In an embodiment, the cells are washed in a wash solution that lacks one or more divalent cation such as calcium and magnesium, e.g., lacks both calcium and magnesium. In one embodiment, the cells are washed in a buffer that has substantially no divalent cations.

In one embodiment, the method further comprises removing cells from the population which express a tumor antigen, e.g., a tumor antigen that does not comprise CD25, e.g., CD19, CD30, CD38, CD123, CD20, CD14 or CD11b, to thereby provide a population of T regulatory depleted, e.g., CD25+ depleted, and tumor antigen depleted cells that are suitable for expression of a CAR, e.g., a CAR described herein. In one embodiment, tumor antigen expressing cells are removed simultaneously with the T regulatory, e.g., CD25+ cells. For

example, an anti-CD25 antibody, or fragment thereof, and an anti-tumor antigen antibody, or fragment thereof, can be attached to the same substrate, e.g., bead, which can be used to remove the cells or an anti-CD25 antibody, or fragment thereof, or the anti-tumor antigen antibody, or fragment thereof, can be attached to separate beads, a mixture of which can be used to remove the cells. In other embodiments, the removal of T regulatory cells, e.g., CD25+ cells, and the removal of the tumor antigen expressing cells is sequential, and can occur, e.g., in either order.

In one embodiment, the method further comprises removing cells from the population which express a check point inhibitor, e.g., a check point inhibitor described herein, e.g., one or more of (e.g., 2 or 3 of) PD1+ cells, LAG3+ cells, and TIM3+ cells, to thereby provide a population of T regulatory depleted, e.g., CD25+ depleted cells, and check point inhibitor depleted cells, e.g., PD1+, LAG3+ and/or TIM3+ depleted cells. In embodiments, PD1+ cells and LAG3+ cells are removed; PD1+ cells and TIM3+ cells are removed; or LAG3+ and TIM3+ cells are removed. In one embodiment, check point inhibitor expressing cells are removed simultaneously with the T regulatory, e.g., CD25+ cells. For example, an anti-CD25 antibody, or fragment thereof, and an anti-check point inhibitor antibody, or fragment thereof, can be attached to the same bead which can be used to remove the cells, or an anti-CD25 antibody, or fragment thereof, and the anti-check point inhibitor antibody, or fragment there, can be attached to separate beads, a mixture of which can be used to remove the cells. In other embodiments, the removal of T regulatory cells, e.g., CD25+ cells, and the removal of the check point inhibitor expressing cells is sequential, and can occur, e.g., in either order.

In one embodiment, the population of cells to be removed are neither the regulatory T cells or tumor cells, but cells that otherwise negatively affect the expansion and/or function of CART cells, e.g. cells expressing CD14, CD11b, CD33, CD15, or other markers expressed by potentially immune suppressive cells. In one embodiment, such cells are envisioned to be removed concurrently with regulatory T cells and/or tumor cells, or following said depletion, or in another order.

In one embodiment, the method further comprises removing cells from the population which express CD14, to thereby provide a population of T regulatory-depleted, e.g., CD25+ depleted cells, and CD14+ depleted cells. In one embodiment, CD14+ cells are removed simultaneously with the T regulatory, e.g., CD25+ cells. For example, an anti-CD25 antibody,

or fragment thereof, and an anti-CD14 antibody, or fragment thereof, can be attached to the same bead which can be used to remove the cells; or an anti-CD25 antibody, or fragment thereof, and the anti-CD14 antibody, or fragment thereof, can be attached to separate beads, a mixture of which can be used to remove the cells. In other embodiments, the removal of T regulatory cells, e.g., CD25+ cells, and the removal of the CD14+ cells is sequential, and can occur, e.g., in either order. In one embodiment, the CD14+ cells are removed using a CD14 antibody molecule or fragment thereof.

In one embodiment, the population of immune effector cells provided have been selected based upon the expression of one or more markers (e.g., 2, 3, 4, 5, 6, 7, or more markers), e.g., CD3, CD28, CD4, CD8, CD27, CD127, CD45RA, and CD45RO, e.g., the provided population of immune effector cells (e.g., T cells) are CD3+ and/or CD28+.

In one embodiment, the method further comprises obtaining a population of immune effector cells, e.g., T cells, enriched for the expression of one or more markers (e.g., 2, 3, 4, 5, 6, 7, or more markers), e.g., CD3, CD28, CD4, CD8, CD27, CD127, CD45RA, and CD45RO. In an embodiment, population of immune effector cells are enriched for CD3+ and/or CD28+ cells. For example, T cells isolated by incubation with anti-CD3/anti-CD28 conjugated beads are obtained. In one embodiment, the method further comprises selecting cells from the population of T regulatory- depleted cells, e.g., CD25+ depleted cells, which express one or more markers (e.g., 2, 3, 4, 5, or more markers), e.g., CD3, CD28, CD4, CD8, CD45RA, and CD45RO.

In one embodiment, the method further comprises activating the population of T regulatory depleted cells, e.g., CD25+ depleted cells, e.g., by a method described herein.

In one embodiment, the method further comprises transducing a cell from the population of T regulatory-depleted cells, e.g., the population of CD25+ depleted cells, with a vector comprising a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein. In one embodiment, the vector is selected from the group consisting of a DNA, a RNA, a plasmid, a lentivirus vector, adenoviral vector, or a retrovirus vector. In one embodiment, the cell from the population of T regulatory-depleted cells, e.g., the population of CD25+ depleted cells, is transduced with a vector once, e.g., within one day after population of immune effector cells are obtained from a blood sample from a subject, e.g., obtained by apheresis.

In one embodiment, the method further comprises generating a population of RNA-engineered cells transiently expressing exogenous RNA from the population of T regulatory-depleted cells, e.g., the population of CD25+ depleted cells. The method comprises introducing an in vitro transcribed RNA or synthetic RNA into a cell from the population, where the RNA
5 comprises a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein.

In one embodiment, cells transduced with a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, are expanded, e.g., by a method described herein. In one embodiment, the cells are expanded in culture for a period of several
10 hours (e.g., about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 18, 21 hours) to about 14 days (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 days). In one embodiment, the cells are expanded for a period of 3 to 9 days. In one embodiment, the cells are expanded for a period of 4 to 9 days. In one embodiment, the cells are expanded for a period of 8 days or less, e.g., 7, 6, 5, 4, or 3 days. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, are expanded in culture for
15 3 or 4 days, and the resulting cells are more potent than the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, are expanded in culture for 5 days, and the resulting cells are more potent than the same cells expanded in culture for 9 days under the same culture conditions. Potency can be defined, e.g., by various T cell functions, e.g. proliferation, target cell killing, cytokine
20 production, activation, migration, or combinations thereof. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 3 or 4 days show at least a one, two, three or four fold increase in cells doublings upon antigen stimulation as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 5 days show at least a one, two,
25 three or four fold increase in cells doublings upon antigen stimulation as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., the cells expressing a CD19 CAR described herein, are expanded in culture for 3 or 4 days, and the resulting cells exhibit higher proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the
30 same culture conditions. In one embodiment, the cells, e.g., the cells expressing a CD19 CAR described herein, are expanded in culture for 5 days, and the resulting cells exhibit higher proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the

same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 3 or 4 days show at least a one, two, three, four, five, ten fold or more increase in pg/ml of proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 5 days show at least a one, two, three, four, five, ten fold or more increase in pg/ml of proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 3, 4, or 5 days show at least as high cytokine production in pg/ml, or at least a one, two, three, four, five, ten fold or more increase in pg/ml of cytokine production, e.g., IL2, IFN-gamma, GM-CSF, TNF-alpha, IL-1b, IL4, IL5, IL6, IL8, or IL10, levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions.

In one embodiment, beads (e.g., CD3/28-stimulatory beads) are removed from the cells by mechanical disruption. In an embodiment, mechanical disruption comprises passage (e.g., repeat passage) cells through a pipette tip, e.g., a narrow bore pipette tip. In some embodiment, the cells are passed through one or more (e.g., 2, 3, 4, or 5 or more) tubes, e.g., narrow bore tubes. In one embodiment, the tubes are part of a closed cell culture system. In some embodiments, the interior diameter of the pipet tip or tubes is less than about 1 mm, 0.9 mm, 0.8 mm, 0.7 mm, 0.5mm, 0.4 mm, 0.3 mm, or 0.2 mm, and optionally the diameter is greater than 0.1 mm, 0.2 mm, 0.3 mm, or 0.4 mm.

In one embodiment, the cells, e.g., CAR cells described herein, e.g., CD19 CAR cells described herein, expanded for 3, 4, or 5 days, have an increased proportion of Tem cells, Tem cells, or both Tem and Tcm cells, compared to as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., CAR cells described herein, e.g., CD19 CAR cells described herein, expanded for 3, 4, or 5 days, have an increased proportion of Tem cells, Teff cells, or both Tem and Teff cells, compared to as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, a population of cells, e.g., CD19CAR cells described herein, expanded for 3, 4, or 5 days, has a percentage of Tnaive like cells (out of Tnaive like, Teff, and Tcm cells) of

at least about 20%, 25%, 30%, 35%, 40%, 45%, or 50%, and optionally up to about 40% or 50%. In one embodiment, a population of cells, e.g., CD19CAR cells described herein, expanded for 3, 4, or 5 days, has a percentage of Tem cells (out of Tnaive like, Teff, and Tcm cells) of at least about 10%, 15%, or 20%, and optionally up to about 15% or 20%. In one
5 embodiment, a population of cells, e.g., CD19CAR cells described herein, expanded for 3, 4, or 5 days, has a percentage of (Tnaive like + Tem) cells (out of Tnaive like, Teff, and Tcm cells) of at least about 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, or 70%, and optionally up to about 50% or 70%.

In one embodiment, the cells, e.g., CD19 CAR cells described herein, expanded for 3, 4,
10 or 5 days, when administered at a dose of 0.5×10^6 cells in the NALM6 assay of Fig. 31A have an activity greater than or equal to a dose of 1×10^6 , 1.5×10^6 , 2×10^6 , 2.5×10^6 , 3×10^6 , 4×10^6 , or 5×10^6 of the same cells expanded in culture for 9 days under the same culture conditions. In some embodiments activity is measured at 1, 2, 3, 4, 5, or 6 weeks.

In one embodiment, the cells are expanded by culturing the cells in the presence of an
15 agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a costimulatory molecule on the surface of the cells, e.g., as described herein. In one embodiment, the agent is a bead conjugated with anti-CD3 antibody, or a fragment thereof, and/or anti-CD28 antibody, or a fragment thereof.

In one embodiment, the cells are expanded in an appropriate media (e.g., media
20 described herein) that may, optionally, contain one or more (e.g., 2, 3, 4, or 5 or more) factor for proliferation and/or viability, including serum (e.g., fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-12, IL-15, IL-21, TGF β , and TNF- α or any other additives for the growth of cells.

In one embodiment, the cells are expanded in an appropriate media (e.g., media
25 described herein) that includes one or more (e.g., 2, 3, 4, or 5 or more) interleukins that result in at least a 200-fold (e.g., 200-fold, 250-fold, 300-fold, 350-fold) increase in cells over a 14 day expansion period, e.g., as measured by a method described herein such as flow cytometry. In one embodiment, the cells are expanded in the presence of IL-15 and/or IL-7 (e.g., IL-15 and IL-7).

In one embodiment, the cells are cryopreserved after the appropriate expansion period. In one embodiment, the cells are cryopreserved according to a method described herein. In one embodiment, the expanded cells are cryopreserved in an appropriate media, e.g., an infusible media, e.g., as described herein.

5 In one embodiment, the method further comprises contacting the population of immune effector cells with a nucleic acid encoding a telomerase subunit, e.g., hTERT. In an embodiment, the nucleic acid is DNA or RNA.

In another aspect, the disclosure features a reaction mixture comprising a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%,
10 4%, 3%, 2%, 1% of CD25+ cells. In one embodiment, the reaction mixture comprises a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the population of cells contains less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of
15 CD25+ cells and less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the population of cells contains less than 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells.

In one embodiment, the reaction mixture comprises a population of T regulatory-
20 depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of a checkpoint inhibitor expressing cells, e.g., a PD1+ cells, LAG3+ cells, or TIM3+ cells. The reaction mixture may further comprise a buffer or other reagent, e.g., a PBS containing solution.

25 In one embodiment, the reaction mixture comprises a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD14+ cells. The reaction mixture may further comprise a buffer or other reagent, e.g., a PBS containing solution.

30

In one embodiment, the reaction mixture can further comprise an agent that activates and/or expands to cells of the population, e.g., an agent that stimulates a CD3/TCR complex associated signal and/or a ligand that stimulates a costimulatory molecule on the surface of the cells, e.g., as described herein. In one embodiment, the agent is a bead conjugated with anti-
5 CD3 antibody, or a fragment thereof, and/or anti-CD28 antibody, or a fragment thereof.

In one embodiment, the reaction mixture further comprises one or more (e.g., 2, 3, 4, or 5) factor for proliferation and/or viability, including serum (e.g., fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-12, IL-15, IL-21, TGF β , and TNF- α or any other additives for the growth of cells. In one embodiment, the reaction
10 mixture further comprises IL-15 and/or IL-7.

In one embodiment, a plurality of the cells of the population in the reaction mixture comprise a nucleic acid molecule, e.g., a nucleic acid molecule described herein, that comprises a CAR encoding sequence, e.g., a CD19 CAR encoding sequence, e.g., as described herein.

In one embodiment, a plurality of the cells of the population in the reaction mixture
15 comprise a vector comprising a nucleic acid sequence encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein. In one embodiment, the vector is a vector described herein, e.g., a vector selected from the group consisting of a DNA, a RNA, a plasmid, a lentivirus vector, adenoviral vector, or a retrovirus vector.

In one embodiment, the reaction mixture further comprises a cryoprotectant or stabilizer
20 such as, e.g., a saccharide, an oligosaccharide, a polysaccharide and a polyol (e.g., trehalose, mannitol, sorbitol, lactose, sucrose, glucose and dextran), salts and crown ethers. In one embodiment, the cryoprotectant is dextran.

In another aspect, the disclosure features a method of making a population of immune
25 effector cells (e.g., T cells, NK cells) engineered to express a CAR, the method comprising providing a population of immune effector cells (e.g., T cells), wherein a plurality of the immune effector cells comprise a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, and expanding the cells of the population in the presence of one or more (e.g., 2, 3, 4, or 5) interleukin that result in at least a 200-fold (e.g., 200-fold,
30 250-fold, 300-fold, 350-fold) increase in cells over a 14 day expansion period, e.g., as

measured by a method described herein such as flow cytometry. In one embodiment, the cells of the population are expanded in the presence of IL-15 and/or IL-7, e.g., IL-15 and IL-7.

In one embodiment, the cells are expanded in culture for a period of several hours (e.g., about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 18, 21 hours) to about 14 days (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 days). In one embodiment, the cells are expanded for a period of 4 to 9 days. In one embodiment, the cells are expanded for a period of less than 8 days, e.g., 7, 6 or 5 days. In one embodiment, the cells are expanded in culture for 5 days, and the resulting cells are more potent than the same cells expanded in culture for 9 days under the same culture conditions. Potency can be defined, e.g., by various T cell functions, e.g. proliferation, target cell killing, cytokine production, activation, migration, or combinations thereof. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 5 days show at least a one, two, three or four fold increase in cells doublings upon antigen stimulation as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., the cells expressing a CD19 CAR described herein, are expanded in culture for 5 days, and the resulting cells exhibit higher proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 5 days show at least a one, two, three, four, five, ten fold or more increase in pg/ml of proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions.

In one embodiment, the cells are expanded by culturing the cells in the presence of an agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a costimulatory molecule on the surface of the cells, e.g., as described herein. In one embodiment, the agent is a bead conjugated with anti-CD3 antibody, or a fragment thereof, and/or anti-CD28 antibody, or a fragment thereof.

In one embodiment, the provided population of immune effector cells is a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells. In one embodiment, the provided population of immune effector cells is a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%,

30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the provided population of cells contains less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the provided
5 population of cells contains less than 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells.

In one embodiment, the provided population of immune effector cells is a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%,
10 4%, 3%, 2%, 1% of a checkpoint inhibitor expressing cells, e.g., a PD1+ cells, LAG3+ cells, or TIM3+ cells.

In one embodiment, the provided population of immune effector cells is a population of T regulatory-depleted cells containing less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%,
15 4%, 3%, 2%, 1% of CD14+ cells.

In one embodiment, the method further comprises, prior to expansion, removing T regulatory cells, e.g., CD25+ T cells, from the population, to thereby provide a population of T regulatory-depleted cells, e.g., CD25+ depleted cells to be expanded. In one embodiment, the T regulatory cells, e.g., CD25+ cells, are removed by a method described herein.

20 In one embodiment, the method further comprises, prior to expansion, removing T regulatory cells, e.g., CD14+ cells, from the population, to thereby provide a population of CD14+ depleted cells to be expanded. In one embodiment, the T regulatory cells, e.g., CD14+ cells, are removed by a method described herein.

In one embodiment, the method further comprises contacting the population of immune
25 effector cells with a nucleic acid encoding a telomerase subunit, e.g., hTERT. In an embodiment, the nucleic acid is DNA or RNA.

In another aspect, the disclosure features a reaction mixture comprising a population of immune effector cells wherein a plurality of the cells of the population in the reaction mixture comprise a nucleic acid molecule, e.g., a nucleic acid molecule described herein, that comprises

a CAR encoding sequence, e.g., a CD19 CAR encoding sequence, e.g., as described herein, and IL-7 and/or IL-15.

In one embodiment, a plurality of the cells of the population in the reaction mixture comprise a vector comprising a nucleic acid sequence encoding a CAR, e.g., a CAR described
5 herein, e.g., a CD19 CAR described herein. In one embodiment, the vector is a vector described herein, e.g., a vector selected from the group consisting of a DNA, a RNA, a plasmid, a lentivirus vector, adenoviral vector, or a retrovirus vector.

In one embodiment, the reaction mixture can further comprise an agent that activates and/or expands to cells of the population, e.g., an agent that stimulates a CD3/TCR complex
10 associated signal and/or a ligand that stimulates a costimulatory molecule on the surface of the cells, e.g., as described herein. In one embodiment, the agent is a bead conjugated with anti-CD3 antibody, or a fragment thereof, and/or anti-CD28 antibody, or a fragment thereof.

In another aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells) engineered to express a CAR, the method comprising
15 providing a population of immune effector cells (e.g., T cells), wherein a plurality of the immune effector cells comprise a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, and expanding the cells of the population in culture for 5 days, wherein the resulting cells are more potent, as measured by cell proliferation levels upon antigen stimulation, as compared to the same cells expanded in culture for 9 days under the
20 same culture conditions.

In one embodiment, the cells expanded for 5 days show at least a one, two, three or four fold increase in cells doublings upon antigen stimulation as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., the cells expressing a CD19 CAR described herein, are expanded in culture for 5
25 days, and the resulting cells exhibit higher proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells expanded for 5 days show at least a one, two, three, four, five, ten fold or more increase in pg/ml of proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in
30 culture for 9 days under the same culture conditions.

In another aspect, the disclosure features methods comprising administering to a subject a population of immune effector cells made by a method described herein and engineered to express a CAR, e.g. a CAR described herein, e.g., a CD19 CAR described herein.

In one embodiment, method provides an anti-tumor immunity in a subject having
5 cancer, e.g., a hematological cancer such as, e.g., CLL. In one embodiment, the method is a method of treating a subject having cancer, e.g., a hematological cancer described herein, such as, e.g., a leukemia (e.g., CLL, ALL) or a lymphoma (e.g., MCL, HL). In one embodiment, the population of cells are autologous to the subject administered the population. In one embodiment, the population of cells is allogeneic to the subject administered the population. In
10 one embodiment, the subject is a human.

In one embodiment, the disease associated with a tumor antigen, e.g., a tumor antigen described herein, e.g., CD19, is selected from a proliferative disease such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia, or is a non-cancer related indication associated with expression of a tumor
15 antigen described herein. In one embodiment, the disease is a cancer described herein, e.g., a cancer described herein as being associated with a target described herein. In one embodiment, the hematologic cancer is leukemia. In one embodiment, the cancer is selected from the group consisting of one or more acute leukemias including but not limited to B-cell acute lymphoid leukemia ("BALL"), T-cell acute lymphoid leukemia ("TALL"), acute lymphoid leukemia
20 (ALL); one or more chronic leukemias including but not limited to chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL); additional hematologic cancers or hematologic conditions including, but not limited to B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, diffuse large B cell lymphoma, follicular lymphoma, hairy cell leukemia, small cell- or a large cell-follicular lymphoma,
25 malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and "preleukemia" which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells,
30 and to disease associated with expression of a tumor antigen described herein include, but not limited to, atypical and/or non-classical cancers, malignancies, precancerous conditions or

proliferative diseases expressing a tumor antigen as described herein; and any combination thereof. In another embodiment, the disease associated with a tumor antigen described herein is a solid tumor, e.g., a solid tumor described herein, e.g., prostatic, colorectal, pancreatic, cervical, gastric, ovarian, head, or lung cancer.

5 In one embodiment, the population of immune effector cells transduced with a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, are expanded, e.g., by a method described herein. In one embodiment, the cells are expanded for a period of 8 days or less, e.g., 7, 6, 5, 4, or 3 days. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, are expanded in culture for 5 days, and the resulting cells are more
10 potent than the same cells expanded in culture for 9 days under the same culture conditions, e.g., as described herein. In one embodiment, the subject is administered 10^4 to 10^6 immune effector cells per kg body weight of the subject. In one embodiment, the subject receives an initial administration of a population of immune effector cells (e.g., an initial administration of 10^4 to 10^6 immune effector cells per kg body weight of the subject, e.g., 10^4 to 10^5 immune
15 effector cells per kg body weight of the subject), a plurality of which comprise the nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, and one or more (e.g., 2, 3, 4, or 5) subsequent administrations of a population of immune effector cells (e.g., one or more subsequent administration of 10^4 to 10^6 immune effector cells per kg body weight of the subject, e.g., 10^4 to 10^5 immune effector cells per kg body weight of the subject),
20 a plurality of which comprise a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein. In one embodiment, the one or more subsequent administrations are administered less than 15 days, e.g., 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, or 2 days after the previous administration, e.g., less than 4, 3, 2 days after the previous administration. In one embodiment, the subject receives a total of about 10^6 immune effector
25 cells per kg body weight of the subject over the course of at least three administrations of a population of immune effector cells, e.g., the subject receives an initial dose of 1×10^5 immune effector cells, a second administration of 3×10^5 immune effector cells, and a third administration of 6×10^5 immune effector cells, and, e.g., each administration is administered less than 4, 3, 2 days after the previous administration.

30 In one aspect, the disclosure features a population of autologous immune effector cells, a plurality of which are transfected or transduced with a vector comprising a nucleic acid

molecule encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, wherein the population of cells contains less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 50%, 40%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CLL cells. In one embodiment, the population of cells
5 contains less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 15%, 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells. In one embodiment, the population of cells contains less than 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells and less than 10%, 5%, 4%, 3%, 2%, 1% of tumor cells, e.g., CD25 expressing tumor cells, e.g., CLL cells.

10 In another aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells). In an embodiment, the method comprises: providing a population of immune effector cells (e.g., T cells or NK cells), and contacting the population of immune effector cells with a nucleic acid encoding a CAR, and a nucleic acid encoding a telomerase subunit, e.g., hTERT, under conditions that allow for CAR and telomerase
15 expression.

In an embodiment, the nucleic acid encoding the telomerase subunit is RNA. In another embodiment, the nucleic acid encoding the telomerase subunit is DNA. In an embodiment, the nucleic acid encoding the telomerase subunit comprises a promoter capable of driving expression of the telomerase subunit.

20 In a related aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells) the method comprising: providing a population of immune effector cells (e.g., T cells), contacting the population of immune effector cells with a nucleic acid encoding a CAR and an RNA encoding a telomerase subunit, e.g., hTERT, under conditions that allow for CAR and telomerase expression.

25 In an embodiment, the nucleic acid encoding the CAR and the RNA encoding the telomerase subunit are part of the same nucleic acid molecule. In an embodiment the nucleic acid encoding the CAR and the RNA encoding the telomerase subunit are part of separate nucleic acid molecules.

In an embodiment, the method comprises contacting the population of immune effector
30 cells with a nucleic acid encoding the CAR and the RNA encoding the telomerase subunit at

substantially the same time. In an embodiment, the method comprises contacting the population of immune effector cells with a nucleic acid encoding the CAR before contacting the population of immune effector cells with the RNA encoding the telomerase subunit. In an embodiment, the method comprises contacting the population of immune effector cells with a nucleic acid encoding the CAR after contacting the population of immune effector cells with the RNA encoding the telomerase subunit.

In an embodiment, the RNA encoding the telomerase subunit is mRNA. In an embodiment, the RNA encoding the telomerase subunit comprises a poly(A) tail. In an embodiment, the RNA encoding the telomerase subunit comprises a 5' cap structure.

In an embodiment, the method comprises transfecting the immune effector cells with the RNA encoding the telomerase subunit. In an embodiment, the method comprises transducing the immune effector cells with the RNA encoding the telomerase subunit. In an embodiment, the method comprises electroporating the immune effector cells with the RNA encoding the telomerase subunit, under conditions that allow for CAR and telomerase expression.

In another aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells) comprising: providing a population of immune effector cells (e.g., T cells or NK cells) that express a CAR and/or comprise a nucleic acid encoding a CAR; and contacting the population of immune effector cells with a nucleic acid encoding a telomerase subunit, e.g., hTERT, under conditions that allow for hTERT expression.

In another aspect, the disclosure features a method of making a population of immune effector cells (e.g., T cells, NK cells), comprising: providing a population of immune effector cells (e.g., T cells or NK cells) that express a nucleic acid encoding a telomerase subunit, e.g., hTERT, and contacting the population of immune effector cells with a nucleic acid encoding a CAR, under conditions that allow for CAR expression.

In one aspect, this disclosure provides an immune effector cell (e.g., T cell or NK cell) comprising: a nucleic acid encoding a CAR, e.g., a CAR as described herein; and a nucleic acid encoding an exogenous telomerase subunit, e.g., hTERT. In an embodiment, the nucleic acid encoding an exogenous telomerase subunit is RNA, e.g., mRNA.

In one aspect, this disclosure provides an immune effector cell (e.g., T cell or NK cell) comprising: a CAR, e.g., a CAR as described herein; and an exogenous telomerase subunit, e.g., hTERT. In an embodiment, the cell does not comprise DNA, e.g., exogenous DNA, e.g., a vector, encoding the exogenous telomerase subunit. For instance, the cell may have been
5 contacted with mRNA encoding the exogenous telomerase subunit.

Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references (e.g., sequence database reference numbers) mentioned herein are incorporated by reference in their
10 entirety. For example, all GenBank, Unigene, and Entrez sequences referred to herein, e.g., in any Table herein, are incorporated by reference. Unless otherwise specified, the sequence accession numbers specified herein, including in any Table herein, refer to the database entries current as of December 29, 2014. When one gene or protein references a plurality of sequence accession numbers, all of the sequence variants are encompassed.

15 In addition, the materials, methods, and examples are illustrative only and not intended to be limiting. Headings, sub-headings or numbered or lettered elements, e.g., (a), (b), (i) etc., are presented merely for ease of reading. The use of headings or numbered or lettered elements in this document does not require the steps or elements be performed in alphabetical order or that the steps or elements are necessarily discrete from one another. Other features, objects,
20 and advantages of the invention will be apparent from the description and drawings, and from the claims.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-1H show the differential effects of γ_c cytokines and IL-18 on CAR-T cell
25 accumulation. **Figure 1A** is a schematic diagram of C4-27z CAR vector. **Figure 1B** is two representative FACS histogram plots of CAR expression on CD4+ and CD8+ T cells 48 hours after lentiviral transduction. **Figure 1C** is a graph showing the overall accumulation of CAR-T cells in response to various cytokines exposure. T cells were transduced and exposed to various exogenous cytokines with final concentrations of 10ng/mL from the next day (day 0). The
30 numbers of CAR-T cells were calculated based on the number of T cells and the percentages of

CAR expression. The curves are representative of 6 donors. *P <0.05, ***P <0.001. NC, no cytokine. **Figure 1D** is a graph showing CAR expression by T cells 15 days after lentiviral transduction. The bar graph depicts CAR expression levels ($\pm$ SEM, n=6) on the surface of CD3+, CD4+ and CD8+ T cells, with the expression of CAR in NC group normalized as 1. *P <0.05 versus IL-2 group; N.S., no statistical difference. **Figure 1E** is a histogram showing the proliferation of T cells in response to various cytokines. On day 7 after lentivirus transduction, T cells in NC group were labeled with CFSE (2.5 μ M), and then exposed to various cytokines. Seven days later, T cells were analyzed for CFSE dilution by flow cytometry. **Figure 1F** is a graph showing the viability of T cells 15 days after lentiviral transduction. T cells from various cytokine groups are stained with Annexin V and 7-AAD, and then analyzed for the proportions of viable cells (both Annexin V and 7-AAD negative). *P <0.05, **P <0.01 versus IL-2 group (n=6). **Figures 1G** and **1H** show Bcl-xL expression by CAR-T cells. On day 15 days after lentiviral transduction, CAR-T cells are assessed for expression of Bcl-2 protein by flow cytometry. **Figure 1G** is a representative FACS histogram plot of Bcl-xL expression in various cytokine groups. **Figure 1H** is a graph depicting Bcl-xL expression ($\pm$ SEM) in CD4+ and CD8+ CAR-T cells of 6 donors. *P <0.01 versus IL-2 group.

Figures 2A-2I shows the memory T cell subsets of CAR-T cells. **Figure 2A** are representative FACS plots showing the gating strategy of T cell subsets analysis. The T cells are split into four subsets based on CD45RA and CD62L expression, then the expression of CCR7, CD27, CD28 and CD95 are further evaluated for every subset. The CD95 expression is significantly upregulated upon lentiviral transduction. **Figure 2B** shows CD95 expression in CD45RA+CD62L+ subpopulation of T cells before transduction and CAR-T cells 15 days after transduction. **Figures 2C** and **2D** are graphs showing the increase of memory stem T cell (Tscm) proportions in CD4+ (**Figure 2C**) and CD8+ T cells (**Figure 2D**) after lentiviral transduction. Tscm are defined as CD45RA+CD62L+CD95+CCR7+ T cell subsets. **Figure 2E** is a graph showing the correlation between the amount of naïve T (Tn, defined as CD45RA+CD62L+CD95- subpopulation) in T cells pre-transduction and the proportion of Tscm in CAR-T cells after transduction (n=6). Left bars represents the percentages of Tn in CD4+ and CD8+ T cells before transduction and right bars represents the percentages of Tscm in CD4+ and CD8+ CAR-T cells. *P <0.05, **P <0.01. **Figures 2F, 2G, and 2H** are graphs showing the distribution of subsets of CD4+ and CD8+ CAR-T cells. The T subsets in CD4+ and CD8+ CAR-T cells are defined based on the CD95, CD45RA and CD62L expression. The

proportions of Tscm are compared among various cytokine groups, *P <0.05, **P <0.01, versus IL-2 group (n=6). (F) Self-renew and differentiation of different subsets of CAR-T cells. FACS-sorted CAR+ Tscm, Tcm, Tem and Temra cells are cultured exposed to IL-2 (10ng/mL) for 3 days, then analyzed the phenotypes based on CD45RA and CD62L expression (n=3).

- 5 **Figure 2I** is a histogram plot showing the proliferation of various subsets of CAR-T cells in response to IL-2. FACS-sorted CAR+ Tscm, Tcm, Tem and Temra cells were labeled with CFSE (2.5μM), and then cultured exposed to IL-2 (10ng/mL) for 3 days. Three days later, T cells were analyzed for CFSE dilution.

- Figures 3A-3B:** shows the correlation between CD45 RA expression and CFSE intensity. **Figure 3A** demonstrates that CD45RA expression is inversely correlated with CFSE intensity. **Figure 3B** shows that for all cytokine groups (IL-2, IL-7, IL-15, IL-18 and IL-21), CD45RA+ T cells exhibited much lower CFSE levels than CD45RA dim and negative T cells indicating that CD45RA+ T cells had stronger proliferation activity than CD45RA- T cells.

- Figures 4A-4B** show the phenotypes of CAR-T cells resulting from exposure to different cytokines. **Figure 4A** is representative FACS dot-plots showing the expression of CD45RA, CD62L, CCR7, CD27 and CD28 on CAR- and CAR+ T cells exposed to IL-2 for 14 days after lentiviral transduction. **Figure 4B** is a series of graphs showing the quantitation of CD45RA, CD62L, CCR7, CD27, CD28 and IL7Rα expression on the surface of CAR-T cells in indicated cytokine groups. The histograms represent mean value ± SEM of expression levels from 6 independent donors. *P <0.05, **P <0.01 versus IL-2 group.

- Figures 5A-5I** show the Functional analysis of CAR-T cells exposed to different cytokines. **Figure 5A** is representative FACS plots show the staining of intracellular IFN-γ, TNF-α and IL-2 in CAR-T cells. **Figure 5B, 5C, and 5D** are quantitative plots showing the percentages of cytokine-producing CAR-T cells in various cytokine groups (n=6) for production of IFNγ (**Figure 5B**), TNF-α (**Figure 5C**) and IL-2 (**Figure 5D**). Lentiviral transduced T cells are exposed to indicated cytokines for 14 days, and then co-cultured with SKOV3 cells for 5 hours before harvested for flow cytometry analysis. **Figure 5E** is a series of pie charts depicting the proportion of cells producing different numbers of cytokines (IFN-γ, TNF-α and IL-2) after SKOV3 stimulation. *P <0.05. **Figure 5F** are representative FACS plots showing the expression of perforin and granzyme B (GranzB) in CAR-T cells. **Figures 5G and 5H** are quantitative plots showing the percentages of perforin (**Figure 5G**) and Granzyme B

expression (**Figure 5H**) in CAR-T cells in various cytokine groups (n=6). Lentiviral transduced T cells are exposed to indicated cytokines for 14 days, and then co-cultured with SKOV3 cells for 5 hours before harvested for flow cytometry analysis. **Figure 5I** is a graph showing the antigen specific cytotoxic activity of CAR-T cells. Fourteen days after indicated cytokine exposure, the CAR-T cells were assessed for cytolytic ability by using a luciferase-based assay after 18-hour coculture with SKOV3 at the indicated E/T ratios. Untransduced T cells (UNT) served as negative effector controls. Data shown are mean value $\pm$ SEM of six independent cytolytic assays.

Figure 6A-6C: shows the phenotype and function of the CAR-T cells described above in **Figures 5A-5I**. **Figures 6A** and **6B** show that CD62L⁺ CAR-T cells (T_{scm} and T_{cm}) exhibited less cytokine production activity (**Figure 6A** and **6B**) and weaker cytolytic capacity (**Figure 6C**) when compared with CD62L⁻ CAR-T cells (T_{em} and T_{emra}).

Figures 7A-7E show the expansion and phenotype of CAR-T cells exposed to antigen challenge. **Figure 7A** is two graphs showing the overall accumulation and viability of CAR-T cells in the exposure to antigen and indicated cytokines. The T cells exposed to IL-2 are harvested on day 15, and then co-cultured with SKOV3 at E/T ratios of 5:1 and indicated cytokines (10ng/mL) for 7 days, with the supplement of SKOV3 cells on the first and fourth day (The same protocol in **Figures 7C, 7D**, and **7E**). The expansion folds are mean value $\pm$ SEM. T cells are stained with Annexin V and 7-AAD, and then analyzed for the proportions of viable cells at the same day. *P < 0.05 versus IL-2 group. **Figure 7B** depicts two graphs showing the overall accumulation and viability of CAR-T previously exposed to indicated cytokines upon antigen challenge. The T cells exposed to indicated cytokines are harvested on day 15, and then co-cultured with SKOV3 at E/T ratios of 5:1 for 7 days. The expansions of CAR-T cells are calculated and the viability of T cells are evaluated on the seventh day. **Figure 7C** is a graph showing CAR expression by T cells after 7-day's coculture with SKOV3 and indicated cytokines. *P < 0.05 versus IL-2. **Figure 7D** is two graphs showing the quantitation of CD27 and CD28 expression on CAR-T cells after 7-day's coculture with SKOV3 and indicated cytokines. *P < 0.05 versus IL-2. **Figure 7E** is two graphs showing the distribution of memory T subsets of CD4⁺ and CD8⁺ CAR-T cells in various cytokine groups. N.S., no statistical difference.

Figures 8A-8G show the antitumor activity of various CAR-T cells with previous cytokine exposure. **Figure 8A** is an in vivo experiment scheme. **Figure 8B** Tumor growth curves of mice treated with various cytokine exposed C4-27z CAR-T cells, anti-CD19-27z CAR-T cells and untransduced T cells. The data are presented as mean value $\pm$ SEM. The arrow indicates the time of T cell infusion. **Figure 8C** is bioluminescence images show fLuc+ SKOV3 tumors in NSG mice immediately before (day 38) , two weeks (day53) and five weeks (day 74) after first intravenous injection of CAR-T cells. **Figure 8D** is a graph showing the quantitation of circulating human CD4+ and CD8+ T cell counts in mice peripheral blood 15 days after the first dose of CAR-T cell infusion. **Figure 8E** is a graph showing the quantitation of CAR expression on circulating human CD4+ and CD8+ T cells in mice blood. **Figure 8F** is a graph showing the distribution of T-cell subsets of circulating human T cells in mice blood based on CD45RA and CD62L staining. **Figure 8G** is a graph showing the quantitation of CD27 and CD28 expression on circulating human CD4+ and CD8+ T cells in mice blood.

Figures 9A and 9B show FAC plots. **Figure 9A** is a series of FACs plots showing the distribution of CD45, CD3, and CD25 expression in cells from apheresis of a CLL patient. **Figure 9B** is a series of FACS plots (top) showing the CD3 and CD19 populations and histograms (bottom) showing CD14 expression of cells from apheresis, cells selected with anti-CD3/CD28, cells depleted for CD25, and the CD25 enriched cells.

Figure 10 is two FACs plots comparing the distribution of CD4+ and CD8+ T cell populations after CD3/CD28 selection or CD25 depletion.

Figures 11A-11C show the comparison of proliferation capacity between CD3/CD28 selected cells and CD25 depleted cells. **Figure 11A** is a graph showing the total cell number at the indicated days in culture. **Figure 11B** is a graph showing the quantified population doublings at each indicated day in culture. **Figure 11C** shows the percentage of viable cells at the indicated days in culture.

Figures 12A and 12B show the effect of CD25 depletion on lentiviral transduction of CAR19. **Figure 12A** is a series of FACS plots showing the efficiency of CD25 depletion. **Figure 12B** is a series of FACS plots showing the CAR19 expression of untransduced cells, CD3 selected cells, and the CD25 depleted cells.

Figure 13 is a series of FACS plots showing the distribution of CD3, CD19, and CD25 expression in cells from PBMCs from a patient before CD25 depletion or culture with cytokine supplement.

Figure 14 is a series of FACS plots showing the distribution of CD3 and CD19 in
5 unmanipulated PBMCs and CD25-depleted PBMCs after culture with the indicated cytokine supplements, IL-7, IL-15, or IL-7 and IL-15.

Figure 15 is a graph showing the total number of cells after 10 days of culture with the indicated cytokine supplements.

Figure 16 is a graph showing the percentage of green fluorescent protein (GFP)
10 signaling as an indication of lentiviral transfection levels for donor cells transfected on two days, the day of (day 0) and the day after (day 1) the cells were obtained by apheresis (day 0+1); donor cells transfected once, the day of obtaining the cells by apheresis (day 0); donor cells transfected once, the day after the cells were obtained by apheresis (day 1); donor cells transfected once, two day after the cells were obtained by apheresis (day 2); and donor cells
15 transfected once, the day after the cells were obtained by apheresis (day 3).

Figure 17 comprises graphs showing expansion profile in population doublings (Figure 17A) and mean size (fL)(Figure 17B) of PBMCs that have been stimulated with anti-CD3 and CD28 beads, and left either unmanipulated (UTD) or transduced with a CD19 CAR (CD19.BBz), de-beaded, and then harvested at Day 5 and D9.

Figure 18 comprises graphs depicting cytotoxicity as a percent lysis of CD19
20 expressing K562 cells treated with PMBCs that have been stimulated with anti-CD3 and CD28 beads, and left either unmanipulated (UTD) or transduced with a CD19 CAR (CD19.BBz), de-beaded, and then harvested at Day 5 and D9.

Figure 19 comprises graphs depicting proliferation of PBMCs stimulated with anti-
25 CD3 and CD28 beads (3x28 beads), wild type K562 cells, CD19 expressing K562 cells, ALL cells (Nalm6) or CLL cells (PI14). The PBMCs have been left either unmanipulated (UTD) or transduced with a CD19 CAR (CART19), de-beaded, and then harvested at Day 5 and D9.

Figure 20 is a schematic of an exemplary manufacturing scheme.

Figure 21 is a schematic of an exemplary manufacturing scheme.

Figure 22 comprises graphs depicting the level of cell proliferation of two different manufacturing batches of donor cells transfected with the CTL019 CAR, Patient 15 (left panels) and Patient 21 (right panels), expanded over a period of 0 to 9 days.

Figure 23 comprises graphs showing proinflammatory cytokine production, IFN- γ , GM-CSF, TNF- α and IL-4 of two different manufacturing batches of donor cells transfected with either CTL019 CAR, namely Patient 15 cells, or an ss1-mesoCAR, namely Patient 21 cells, and expanded over a period of 0 to 9 days after apheresis.

Figure 24 comprises graphs depicting production levels IFN- γ , TNF- α , IL-6, IL-8, IL-2, IL-1 β , GM-CSF and IL-4 in donor cells stimulated with anti-CAR19-idiotype antibody beads or control beads, transfected with CTL019 CAR and expanded for 5 to 9 days. No cytokine or low cytokine levels (<200 pg/ml) were detected with the control beads.

Figure 25 is a graph depicting cell killing based upon total lysates using a luciferase assay of Nalm6 (ALL) cells of PBMCs left either unmanipulated (UTD) or transduced with a CD19 CAR (CART19), de-beaded, and then harvested at Day 5 and D9. Various ratios of PMBCs to Nalm6 cells (effector (E):Target (T)) were cultured. As shown CART19 cells harvested at day 5 possess a better killing capacity.

Figure 26 is a graph depicting long term in vivo killing capacity of PBMCs left either unmanipulated (UTD) or transduced with a CD19 CAR (CART19), de-beaded, and then harvested at Day 5 and D9. The PBMCs were introduced into non-obese diabetic/severe combined immunodeficiency mice inoculated with Nalm6 cells.

Figures 27A-27D indicate that bead removal at early timepoints does not induce cell loss. Fig. 27A is a Coulter analysis illustrating mean T cell volume and concentration before and after removing α CD3/ α CD28 coated magnetic beads via magnet on day 3. Cells were then frozen after bead removal. Fig. 27B shows that cell number was evaluated before and after removal of magnetic beads. The concentration was similar. Fig. 27C is an overlay showing the mean cell volume of same cells before and after thaw. Cells retain their volume during the freeze thaw process. Results are representative of at least 15 different experiments. Fig. 27D shows a representative Coulter analysis of T cell products before and after α CD3/ α CD28 coated magnetic bead removal demonstrating the efficiency of process.

Figures 28A-28B shows that T cells that are activated, transduced with a CD19-specific CAR bearing BBz signaling and harvested from T cells cultures at time points as early as day 3 following activation have potent, antigen-specific cytotoxic activity in vitro. Figure 28A, upper panel, four-hour killing assay using K562-WT cells. Figure 28B, lower panel, four-hour killing assay using K562-19 cells. Figure 28B upper panels, transduction efficiency using IL7/IL15 (left) or IL2 (right). Figure 28B center panel, population doublings using IL7/IL15 compared to IL2. Figure 28B lower panel, volume using IL7/IL15 compared to IL2.

Figures 29A-29C show that day 3 and day 9 4-1BB ζ CART cells produce similar patterns of cytokines in response to K562-CD19. Day 3 and day 9 CART 19 cells, previously cultured in medium with IL-2 or IL-7/15, were incubated with either K562-CD19 (Fig. 29A and 29B), medium alone for 24h (Fig. 29C). “No antigen” in Fig. 29A and Fig. 29B indicates cells stimulated with wild-type K562 cells without CD19. Cytokines were measured in culture supernatants by cytokine bead array (Luminex). Figure 29C shows representative cytokine production in CART 19 T cells 24h after thaw. The cytokine levels are minimal.

Figures 30A-30B show a progressive transition towards a more differentiated phenotype with an increasing proportion of Tem and Tcm cells.

Figure 31A-31D. Fig. 31A shows the experimental setup for testing the potency of CART cells produced under different conditions. Fig. 31B shows the potency of IL2-treated CART cells in slowing Nalm6 tumor growth in mice. D3, D5, and D9 indicate CART cells that were harvested on day 3, day 5, or day 9. 3e6 and 0.5e6 indicate the number of CAR+ cells administered. Proportions of CAR+ cells are shown in Fig. 31C. Fig. 31D compares the potency of IL2-treated and IL7/IL15-treated CART cells in slowing Nalm6 tumor growth in mice.

Figure 32. In vitro expansion of CAR+ T cells in media contained IL-2 and IL-7/15, following antigen stimulation. Left panel, PBMCs from healthy human donors were stimulated with α CD3/ α CD28 coated Dynal beads on day 0, and lenti-virally transduced with 19-BB ζ on day 1. These cells were expanded for 9 days in medium supplemented either with IL-2 or IL-7/15. T cells were counted by flow cytometry using bead-based counting every other day. Right panel, Mean T cell volume (fl) over the course of expansion was monitored. Data are representative of at least six independent donors.

Figure 33. Day 3 and day 9 CART19 cells cultured in medium with either IL-2 or IL-7/15 proliferate in response to K562-CD19, and not to K562-WT or medium alone. T cells were labeled with CFSE and co-cultured with either K562-CD19, wild-type K562 or medium alone as a negative control, for 120 hours. The number of proliferating T cells was significantly higher in response to K562-CD19 as compared to wild-type K562 or medium alone and was comparable between IL-7/15 and IL-2 groups. Results are representative of two different donors.

DETAILED DESCRIPTION

10 Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains.

The term “a” and “an” refers to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

The term “about” when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of $\pm 20\%$ or in some instances $\pm 10\%$, or in some instances $\pm 5\%$, or in some instances $\pm 1\%$, or in some instances $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods.

The term “Chimeric Antigen Receptor” or alternatively a “CAR” refers to a set of polypeptides, typically two in the simplest embodiments, which when in an immune effector cell, provides the cell with specificity for a target cell, typically a cancer cell, and with intracellular signal generation. In some embodiments, a CAR comprises at least an extracellular antigen binding domain, a transmembrane domain and a cytoplasmic signaling domain (also referred to herein as “an intracellular signaling domain”) comprising a functional signaling domain derived from a stimulatory molecule and/or costimulatory molecule as defined below. In some embodiments, the set of polypeptides are in the same polypeptide chain (e.g., comprise a chimeric fusion protein). In some embodiments, the set of polypeptides are not contiguous with each other, e.g., are in different polypeptide chains. In some embodiments, the set of

polypeptides include a dimerization switch that, upon the presence of a dimerization molecule, can couple the polypeptides to one another, e.g., can couple an antigen binding domain to an intracellular signaling domain. In one embodiment, the stimulatory molecule of the CAR is the zeta chain associated with the T cell receptor complex. In one aspect, the cytoplasmic

5 signaling domain comprises a primary signaling domain (e.g., a primary signaling domain of CD3-zeta). In one embodiment, the cytoplasmic signaling domain further comprises one or more functional signaling domains of at least one costimulatory molecule as defined below. In one embodiment, the costimulatory molecule is a costimulatory molecule described herein, e.g., 4-1BB (i.e., CD137), CD27, ICOS, and/or CD28. In one embodiment, the CAR comprises a

10 chimeric fusion protein comprising an extracellular antigen binding domain, a transmembrane domain and an intracellular signaling domain comprising a functional signaling domain of a stimulatory molecule. In one embodiment, the CAR comprises a chimeric fusion protein comprising an extracellular antigen binding domain, a transmembrane domain and an intracellular signaling domain comprising a functional signaling domain of a co-stimulatory

15 molecule and a functional signaling domain of a stimulatory molecule. In one embodiment, the CAR comprises a chimeric fusion protein comprising an extracellular antigen binding domain, a transmembrane domain and an intracellular signaling domain comprising two functional signaling domains of one or more co-stimulatory molecule(s) and a functional signaling domain of a stimulatory molecule. In one embodiment, the CAR comprises a chimeric fusion protein

20 comprising an extracellular antigen binding domain, a transmembrane domain and an intracellular signaling domain comprising at least two functional signaling domains of one or more co-stimulatory molecule(s) and a functional signaling domain of a stimulatory molecule. In one embodiment, the CAR comprises an optional leader sequence at the amino-terminus (N-terminus) of the CAR fusion protein. In one embodiment, the CAR further comprises a leader

25 sequence at the N-terminus of the extracellular antigen binding domain, wherein the leader sequence is optionally cleaved from the antigen binding domain (e.g., a scFv) during cellular processing and localization of the CAR to the cellular membrane.

A CAR that comprises an antigen binding domain (e.g., a scFv, or TCR) that targets a specific tumor antigen X, such as those described herein, is also referred to as XCAR. For

30 example, a CAR that comprises an antigen binding domain that targets CD19 is referred to as CD19CAR.

The term “signaling domain” refers to the functional portion of a protein which acts by transmitting information within the cell to regulate cellular activity via defined signaling pathways by generating second messengers or functioning as effectors by responding to such messengers.

5 The term “antibody,” as used herein, refers to a protein, or polypeptide sequence derived from an immunoglobulin molecule which specifically binds with an antigen. Antibodies can be polyclonal or monoclonal, multiple or single chain, or intact immunoglobulins, and may be derived from natural sources or from recombinant sources. Antibodies can be tetramers of immunoglobulin molecules.

10 The term “antibody fragment” refers to at least one portion of an antibody, that retains the ability to specifically interact with (e.g., by binding, steric hindrance, stabilizing/destabilizing, spatial distribution) an epitope of an antigen. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab')₂, Fv fragments, scFv antibody fragments, disulfide-linked Fvs (sdFv), a Fd fragment consisting of the VH and CH1 domains,
15 linear antibodies, single domain antibodies such as sdAb (either VL or VH), camelid VHH domains, multi-specific antibodies formed from antibody fragments such as a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region, and an isolated CDR or other epitope binding fragments of an antibody. An antigen binding fragment can also be incorporated into single domain antibodies, maxibodies, minibodies,
20 intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, e.g., Hollinger and Hudson, Nature Biotechnology 23:1126-1136, 2005). Antigen binding fragments can also be grafted into scaffolds based on polypeptides such as a fibronectin type III (Fn3)(see U.S. Patent No.: 6,703,199, which describes fibronectin polypeptide minibodies).

 The term “scFv” refers to a fusion protein comprising at least one antibody fragment
25 comprising a variable region of a light chain and at least one antibody fragment comprising a variable region of a heavy chain, wherein the light and heavy chain variable regions are contiguously linked, e.g., via a synthetic linker, e.g., a short flexible polypeptide linker, and capable of being expressed as a single chain polypeptide, and wherein the scFv retains the specificity of the intact antibody from which it is derived. Unless specified, as used herein an
30 scFv may have the VL and VH variable regions in either order, e.g., with respect to the N-

terminal and C-terminal ends of the polypeptide, the scFv may comprise VL-linker-VH or may comprise VH-linker-VL.

The portion of a CAR comprising an antibody or antibody fragment thereof may exist in a variety of forms where the antigen binding domain is expressed as part of a contiguous polypeptide chain including, for example, a single domain antibody fragment (sdAb), a single chain antibody (scFv) and a humanized antibody (Harlow et al., 1999, In: Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY; Harlow et al., 1989, In: Antibodies: A Laboratory Manual, Cold Spring Harbor, New York; Houston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; Bird et al., 1988, Science 242:423-426). In one embodiment, the antigen binding domain of a CAR comprises an antibody fragment. In a further embodiment, the CAR comprises an antibody fragment that comprises a scFv.

As used herein, the term "binding domain" or "antibody molecule" refers to a protein, e.g., an immunoglobulin chain or fragment thereof, comprising at least one immunoglobulin variable domain sequence. The term "binding domain" or "antibody molecule" encompasses antibodies and antibody fragments. In an embodiment, an antibody molecule is a multispecific antibody molecule, e.g., it comprises a plurality of immunoglobulin variable domain sequences, wherein a first immunoglobulin variable domain sequence of the plurality has binding specificity for a first epitope and a second immunoglobulin variable domain sequence of the plurality has binding specificity for a second epitope. In an embodiment, a multispecific antibody molecule is a bispecific antibody molecule. A bispecific antibody has specificity for no more than two antigens. A bispecific antibody molecule is characterized by a first immunoglobulin variable domain sequence which has binding specificity for a first epitope and a second immunoglobulin variable domain sequence that has binding specificity for a second epitope.

The portion of the CAR of the invention comprising an antibody or antibody fragment thereof may exist in a variety of forms where the antigen binding domain is expressed as part of a contiguous polypeptide chain including, for example, a single domain antibody fragment (sdAb), a single chain antibody (scFv), a humanized antibody, or bispecific antibody (Harlow et al., 1999, In: Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY; Harlow et al., 1989, In: Antibodies: A Laboratory Manual, Cold Spring Harbor, New York; Houston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; Bird et al., 1988,

Science 242:423-426). In one aspect, the antigen binding domain of a CAR composition of the invention comprises an antibody fragment. In a further aspect, the CAR comprises an antibody fragment that comprises a scFv.

5 The term “antibody heavy chain,” refers to the larger of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations, and which normally determines the class to which the antibody belongs.

The term “antibody light chain,” refers to the smaller of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations. Kappa (κ) and lambda (λ) light chains refer to the two major antibody light chain isotypes.

10 The term “complementarity determining region” or “CDR,” as used herein, refers to the sequences of amino acids within antibody variable regions which confer antigen specificity and binding affinity. For example, in general, there are three CDRs in each heavy chain variable region (e.g., HCDR1, HCDR2, and HCDR3) and three CDRs in each light chain variable region (LCDR1, LCDR2, and LCDR3). The precise amino acid sequence boundaries of a
15 given CDR can be determined using any of a number of well-known schemes, including those described by Kabat et al. (1991), “Sequences of Proteins of Immunological Interest,” 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (“Kabat” numbering scheme), Al-Lazikani et al., (1997) JMB 273,927-948 (“Chothia” numbering scheme), or a combination thereof. Under the Kabat numbering scheme, in some embodiments, the CDR
20 amino acid residues in the heavy chain variable domain (VH) are numbered 31-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3); and the CDR amino acid residues in the light chain variable domain (VL) are numbered 24-34 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3). Under the Chothia numbering scheme, in some embodiments, the CDR amino acids in the VH are numbered 26-32 (HCDR1), 52-56 (HCDR2), and 95-102 (HCDR3); and the CDR amino
25 acid residues in the VL are numbered 26-32 (LCDR1), 50-52 (LCDR2), and 91-96 (LCDR3). In a combined Kabat and Chothia numbering scheme, in some embodiments, the CDRs correspond to the amino acid residues that are part of a Kabat CDR, a Chothia CDR, or both. For instance, in some embodiments, the CDRs correspond to amino acid residues 26-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3) in a VH, e.g., a mammalian VH, e.g., a
30 human VH; and amino acid residues 24-34 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3) in a VL, e.g., a mammalian VL, e.g., a human VL.

The term “recombinant antibody” refers to an antibody which is generated using recombinant DNA technology, such as, for example, an antibody expressed by a bacteriophage or yeast expression system. The term should also be construed to mean an antibody which has been generated by the synthesis of a DNA molecule encoding the antibody and which DNA
5 molecule expresses an antibody protein, or an amino acid sequence specifying the antibody, wherein the DNA or amino acid sequence has been obtained using recombinant DNA or amino acid sequence technology which is available and well known in the art.

The term “antigen” or “Ag” refers to a molecule that provokes an immune response. This immune response may involve either antibody production, or the activation of specific
10 immunologically-competent cells, or both. The skilled artisan will understand that any macromolecule, including virtually all proteins or peptides, can serve as an antigen. Furthermore, antigens can be derived from recombinant or genomic DNA. A skilled artisan will understand that any DNA, which comprises a nucleotide sequences or a partial nucleotide sequence encoding a protein that elicits an immune response therefore encodes an “antigen” as
15 that term is used herein. Furthermore, one skilled in the art will understand that an antigen need not be encoded solely by a full length nucleotide sequence of a gene. It is readily apparent that the present invention includes, but is not limited to, the use of partial nucleotide sequences of more than one gene and that these nucleotide sequences are arranged in various combinations to encode polypeptides that elicit the desired immune response. Moreover, a skilled artisan will
20 understand that an antigen need not be encoded by a “gene” at all. It is readily apparent that an antigen can be generated synthesized or can be derived from a biological sample, or might be macromolecule besides a polypeptide. Such a biological sample can include, but is not limited to a tissue sample, a tumor sample, a cell or a fluid with other biological components.

The term “autologous” refers to any material derived from the same individual to whom
25 it is later to be re-introduced into the individual.

The term “allogeneic” refers to any material derived from a different animal of the same species as the individual to whom the material is introduced. Two or more individuals are said to be allogeneic to one another when the genes at one or more loci are not identical. In some aspects, allogeneic material from individuals of the same species may be sufficiently unlike
30 genetically to interact antigenically.

The term “xenogeneic” refers to any material derived from an animal of a different species.

The term “cancer” refers to a disease characterized by the uncontrolled growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body. Examples of various cancers are described herein and include but are not limited to, breast cancer, prostate cancer, ovarian cancer, cervical cancer, skin cancer, pancreatic cancer, colorectal cancer, renal cancer, liver cancer, brain cancer, lymphoma, leukemia, lung cancer and the like. The terms “tumor” and “cancer” are used interchangeably herein, e.g., both terms encompass solid and liquid, e.g., diffuse or circulating, tumors. As used herein, the term “cancer” or “tumor” includes premalignant, as well as malignant cancers and tumors.

“Derived from” as that term is used herein, indicates a relationship between a first and a second molecule. It generally refers to structural similarity between the first molecule and a second molecule and does not connote or include a process or source limitation on a first molecule that is derived from a second molecule. For example, in the case of an intracellular signaling domain that is derived from a CD3zeta molecule, the intracellular signaling domain retains sufficient CD3zeta structure such that it has the required function, namely, the ability to generate a signal under the appropriate conditions. It does not connote or include a limitation to a particular process of producing the intracellular signaling domain, e.g., it does not mean that, to provide the intracellular signaling domain, one must start with a CD3zeta sequence and delete unwanted sequence, or impose mutations, to arrive at the intracellular signaling domain.

The phrase “disease associated with expression of a tumor antigen as described herein” includes, but is not limited to, a disease associated with expression of a tumor antigen as described herein or condition associated with cells which express a tumor antigen as described herein including, e.g., proliferative diseases such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia; or a noncancer related indication associated with cells which express a tumor antigen as described herein. In one embodiment, a cancer associated with expression of a tumor antigen as described herein is a hematological cancer. In one embodiment, a cancer associated with expression of a tumor antigen as described herein is a solid cancer. Further diseases associated with expression of a tumor antigen as described herein include, but not limited to, e.g., atypical

and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases associated with expression of a tumor antigen as described herein. Non-cancer related indications associated with expression of a tumor antigen as described herein include, but are not limited to, e.g., autoimmune disease, (e.g., lupus), inflammatory disorders (allergy and
5 asthma) and transplantation. In some embodiments, the tumor antigen-expressing cells express, or at any time expressed, mRNA encoding the tumor antigen. In an embodiment, the tumor antigen-expressing cells produce the tumor antigen protein (e.g., wild-type or mutant), and the tumor antigen protein may be present at normal levels or reduced levels. In an embodiment, the tumor antigen -expressing cells produced detectable levels of a tumor antigen protein at one
10 point, and subsequently produced substantially no detectable tumor antigen protein.

The phrase “disease associated with expression of CD19” includes, but is not limited to, a disease associated with expression of CD19 or condition associated with cells which express CD19 including, e.g., proliferative diseases such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia; or a
15 noncancer related indication associated with cells which express CD19. In one aspect, a cancer associated with expression of CD19 is a hematological cancer. In one aspect, the hematological cancer is a leukemia or a lymphoma. In one aspect, a cancer associated with expression of CD19 includes cancers and malignancies including, but not limited to, e.g., one or more acute leukemias including but not limited to, e.g., acute myeloid leukemia (AML), B-cell acute
20 Lymphoid Leukemia (BALL), T-cell acute Lymphoid Leukemia (TALL), acute lymphoid leukemia (ALL); one or more chronic leukemias including but not limited to, e.g., chronic myelogenous leukemia (CML), Chronic Lymphoid Leukemia (CLL). Additional cancers or hematologic conditions associated with expression of CD19 comprise, but are not limited to, e.g., B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's
25 lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma (MCL), Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin lymphoma, Hodgkin lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom
30 macroglobulinemia, myeloproliferative neoplasm; a histiocytic disorder (e.g., a mast cell disorder or a blastic plasmacytoid dendritic cell neoplasm); a mast cell disorder, e.g., systemic mastocytosis or mast cell leukemia; B-cell prolymphocytic leukemia, plasma cell myeloma,

and “preleukemia” which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like. Further diseases associated with expression of CD19 include, but not limited to, e.g., atypical and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases associated with expression of CD19. Non-cancer related indications associated with expression of CD19 include, but are not limited to, e.g., autoimmune disease, (e.g., lupus), inflammatory disorders (allergy and asthma) and transplantation. In some embodiments, the tumor antigen-expressing cells express, or at any time expressed, mRNA encoding the tumor antigen. In an embodiment, the tumor antigen-expressing cells produce the tumor antigen protein (e.g., wild-type or mutant), and the tumor antigen protein may be present at normal levels or reduced levels. In an embodiment, the tumor antigen -expressing cells produced detectable levels of a tumor antigen protein at one point, and subsequently produced substantially no detectable tumor antigen protein. In other embodiments, the disease is a CD19-negative cancer, e.g., a CD19-negative relapsed cancer. In some embodiments, the tumor antigen (e.g., CD19)-expressing cell expresses, or at any time expressed, mRNA encoding the tumor antigen. In an embodiment, the tumor antigen (e.g., CD19)-expressing cell produces the tumor antigen protein (e.g., wild-type or mutant), and the tumor antigen protein may be present at normal levels or reduced levels. In an embodiment, the tumor antigen (e.g., CD19)-expressing cell produced detectable levels of a tumor antigen protein at one point, and subsequently produced substantially no detectable tumor antigen protein.

The phrase “disease associated with expression of a B-cell antigen” includes, but is not limited to, a disease associated with expression of one or more of CD19, CD20, CD22 or ROR1, or a condition associated with cells which express, or at any time expressed, one or more of CD19, CD20, CD22 or ROR1, including, e.g., proliferative diseases such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia; or a noncancer related indication associated with cells which express one or more of CD19, CD20, CD22 or ROR1. For the avoidance of doubt, a disease associated with expression of the B-cell antigen may include a condition associated with cells which do not presently express the B-cell antigen, e.g., because the antigen expression has been downregulated, e.g., due to treatment with a molecule targeting the B-cell antigen, e.g., a B-cell targeting CAR, but which at one time expressed the antigen. The phrase “disease associated with expression of a B-cell antigen” includes a disease associated with expression of CD19, as

described herein. In embodiments, the CAR-expressing cells are used to treat a disease associated with a B-cell antigen. In embodiments, a CAR produced by a method herein comprises an antigen binding domain that targets a B-cell antigen.

The term “relapse” as used herein refers to reappearance of a disease (e.g., cancer) after an initial period of responsiveness, e.g., after prior treatment with a therapy, e.g., cancer therapy (e.g., complete response or partial response). The initial period of responsiveness may involve the level of cancer cells falling below a certain threshold, e.g., below 20%, 15%, 10%, 5%, 4%, 3%, 2%, or 1%. The reappearance may involve the level of cancer cells rising above a certain threshold, e.g., above 20%, 15%, 10%, 5%, 4%, 3%, 2%, or 1%. For example, e.g., in the context of B-ALL, the reappearance may involve, e.g., a reappearance of blasts in the blood, bone marrow (> 5%), or any extramedullary site, after a complete response. A complete response, in this context, may involve < 5% BM blast. More generally, in an embodiment, a response (e.g., complete response or partial response) can involve the absence of detectable MRD (minimal residual disease). In an embodiment, the initial period of responsiveness lasts at least 1, 2, 3, 4, 5, or 6 days; at least 1, 2, 3, or 4 weeks; at least 1, 2, 3, 4, 6, 8, 10, or 12 months; or at least 1, 2, 3, 4, or 5 years.

“Refractory” as used herein refers to a disease, e.g., cancer, that does not respond to a treatment. In embodiments, a refractory cancer can be resistant to a treatment before or at the beginning of the treatment. In other embodiments, the refractory cancer can become resistant during a treatment. A refractory cancer is also called a resistant cancer.

The term “conservative sequence modifications” refers to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody or antibody fragment containing the amino acid sequence. Such conservative modifications include amino acid substitutions, additions and deletions. Modifications can be introduced into an antibody or antibody fragment of the invention by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan),

nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues within a CAR described herein can be replaced with other amino acid residues from the same side chain family and the altered CAR can be tested using the functional assays described herein.

The term “stimulation,” refers to a primary response induced by binding of a stimulatory molecule (e.g., a TCR/CD3 complex or CAR) with its cognate ligand (or tumor antigen in the case of a CAR) thereby mediating a signal transduction event, such as, but not limited to, signal transduction via the TCR/CD3 complex or signal transduction via the appropriate NK receptor or signaling domains of the CAR. Stimulation can mediate altered expression of certain molecules.

The term “stimulatory molecule,” refers to a molecule expressed by an immune cell (e.g., T cell, NK cell, B cell) that provides the cytoplasmic signaling sequence(s) that regulate activation of the immune cell in a stimulatory way for at least some aspect of the immune cell signaling pathway. In one aspect, the signal is a primary signal that is initiated by, for instance, binding of a TCR/CD3 complex with an MHC molecule loaded with peptide, and which leads to mediation of a T cell response, including, but not limited to, proliferation, activation, differentiation, and the like. A primary cytoplasmic signaling sequence (also referred to as a “primary signaling domain”) that acts in a stimulatory manner may contain a signaling motif which is known as immunoreceptor tyrosine-based activation motif or ITAM. Examples of an ITAM containing cytoplasmic signaling sequence that is of particular use in the invention includes, but is not limited to, those derived from CD3 zeta, common FcR gamma (FCER1G), Fc gamma RIIa, FcR beta (Fc Epsilon R1b), CD3 gamma, CD3 delta, CD3 epsilon, CD79a, CD79b, DAP10, and DAP12. In a specific CAR of the invention, the intracellular signaling domain in any one or more CARS of the invention comprises an intracellular signaling sequence, e.g., a primary signaling sequence of CD3-zeta. In a specific CAR of the invention, the primary signaling sequence of CD3-zeta is the sequence provided as SEQ ID NO:9 (mutant CD3 zeta), or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like. In a specific CAR of the invention, the primary signaling sequence of CD3-zeta is the sequence as provided in SEQ ID NO:10 (wild-type

human CD3 zeta), or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like.

The term “antigen presenting cell” or “APC” refers to an immune system cell such as an accessory cell (e.g., a B-cell, a dendritic cell, and the like) that displays a foreign antigen
5 complexed with major histocompatibility complexes (MHC's) on its surface. T-cells may recognize these complexes using their T-cell receptors (TCRs). APCs process antigens and present them to T-cells.

An “intracellular signaling domain,” as the term is used herein, refers to an intracellular portion of a molecule. The intracellular signaling domain can generate a signal that promotes
10 an immune effector function of the CAR containing cell, e.g., a CART cell. Examples of immune effector function, e.g., in a CART cell, include cytolytic activity and helper activity, including the secretion of cytokines. In embodiments, the intracellular signaling domain is the portion of a protein which transduces the effector function signal and directs the cell to perform a specialized function. While the entire intracellular signaling domain can be employed, in
15 many cases it is not necessary to use the entire chain. To the extent that a truncated portion of the intracellular signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. The term intracellular signaling domain is thus meant to include any truncated portion of the intracellular signaling domain sufficient to transduce the effector function signal.

In an embodiment, the intracellular signaling domain can comprise a primary
20 intracellular signaling domain. Exemplary primary intracellular signaling domains include those derived from the molecules responsible for primary stimulation, or antigen dependent stimulation. In an embodiment, the intracellular signaling domain can comprise a costimulatory intracellular domain. Exemplary costimulatory intracellular signaling domains include those
25 derived from molecules responsible for costimulatory signals, or antigen independent stimulation. For example, in the case of a CART, a primary intracellular signaling domain can comprise a cytoplasmic sequence of a T cell receptor, and a costimulatory intracellular signaling domain can comprise cytoplasmic sequence from co-receptor or costimulatory molecule.

30 A primary intracellular signaling domain can comprise a signaling motif which is known as an immunoreceptor tyrosine-based activation motif or ITAM. Examples of ITAM

containing primary cytoplasmic signaling sequences include, but are not limited to, those derived from CD3 zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, CD278 ("ICOS"), FcεRI, and CD66d, CD32, DAP10, and DAP12.

The term "zeta" or alternatively "zeta chain", "CD3-zeta" or "TCR-zeta" is defined as the protein provided as GenBank Acc. No. BAG36664.1, or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like, and a "zeta stimulatory domain" or alternatively a "CD3-zeta stimulatory domain" or a "TCR-zeta stimulatory domain" is defined as the amino acid residues from the cytoplasmic domain of the zeta chain that are sufficient to functionally transmit an initial signal necessary for T cell activation. In one aspect the cytoplasmic domain of zeta comprises residues 52 through 164 of GenBank Acc. No. BAG36664.1 or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like, that are functional orthologs thereof. In one aspect, the "zeta stimulatory domain" or a "CD3-zeta stimulatory domain" is the sequence provided as SEQ ID NO:9. In one aspect, the "zeta stimulatory domain" or a "CD3-zeta stimulatory domain" is the sequence provided as SEQ ID NO:10.

The term "costimulatory molecule" refers to the cognate binding partner on a T cell that specifically binds with a costimulatory ligand, thereby mediating a costimulatory response by the T cell, such as, but not limited to, proliferation. Costimulatory molecules are cell surface molecules other than antigen receptors or their ligands that are required for an efficient immune response. Costimulatory molecules include, but are not limited to MHC class I molecule, TNF receptor proteins, Immunoglobulin-like proteins, cytokine receptors, integrins, signalling lymphocytic activation molecules (SLAM proteins), activating NK cell receptors, BTLA, a Toll ligand receptor, OX40, CD2, CD7, CD27, CD28, CD30, CD40, CDS, ICAM-1, LFA-1 (CD11a/CD18), 4-1BB (CD137), B7-H3, CDS, ICAM-1, ICOS (CD278), GITR, BAFFR, LIGHT, HVEM (LIGHTR), KIRDS2, SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46, CD19, CD4, CD8alpha, CD8beta, IL2R beta, IL2R gamma, IL7R alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d, ITGAE, CD103, ITGAL, CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, NKG2D, NKG2C, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRTAM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150,

IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, and a ligand that specifically binds with CD83.

A costimulatory intracellular signaling domain refers to an intracellular portion of a costimulatory molecule. The intracellular signaling domain can comprise the entire
5 intracellular portion, or the entire native intracellular signaling domain, of the molecule from which it is derived, or a functional fragment thereof.

The intracellular signaling domain can comprise the entire intracellular portion, or the entire native intracellular signaling domain, of the molecule from which it is derived, or a functional fragment thereof.

10 The term “4-1BB” refers to a member of the TNFR superfamily with an amino acid sequence provided as GenBank Acc. No. AAA62478.2, or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like; and a “4-1BB costimulatory domain” is defined as amino acid residues 214-255 of GenBank Acc. No. AAA62478.2, or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like.
15 In one aspect, the “4-1BB costimulatory domain” is the sequence provided as SEQ ID NO:7 or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like.

“Immune effector cell,” as that term is used herein, refers to a cell that is involved in an immune response, e.g., in the promotion of an immune effector response. Examples of
20 immune effector cells include T cells, e.g., alpha/beta T cells and gamma/delta T cells, B cells, natural killer (NK) cells, natural killer T (NKT) cells, mast cells, and myeloid-derived phagocytes.

“Immune effector function or immune effector response,” as that term is used herein, refers to function or response, e.g., of an immune effector cell, that enhances or promotes an
25 immune attack of a target cell. E.g., an immune effector function or response refers a property of a T or NK cell that promotes killing or the inhibition of growth or proliferation, of a target cell. In the case of a T cell, primary stimulation and co-stimulation are examples of immune effector function or response.

The term “effector function” refers to a specialized function of a cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines.

The term “encoding” refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (e.g., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene, cDNA, or RNA, encodes a protein if transcription and translation of mRNA corresponding to that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

Unless otherwise specified, a “nucleotide sequence encoding an amino acid sequence” includes all nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. The phrase nucleotide sequence that encodes a protein or a RNA may also include introns to the extent that the nucleotide sequence encoding the protein may in some version contain an intron(s).

The term “endogenous” refers to any material from or produced inside an organism, cell, tissue or system.

The term “exogenous” refers to any material introduced from or produced outside an organism, cell, tissue or system.

The term “expression” refers to the transcription and/or translation of a particular nucleotide sequence driven by a promoter.

The term “transfer vector” refers to a composition of matter which comprises an isolated nucleic acid and which can be used to deliver the isolated nucleic acid to the interior of a cell. Numerous vectors are known in the art including, but not limited to, linear polynucleotides, polynucleotides associated with ionic or amphiphilic compounds, plasmids, and viruses. Thus, the term “transfer vector” includes an autonomously replicating plasmid or a virus. The term should also be construed to further include non-plasmid and non-viral

compounds which facilitate transfer of nucleic acid into cells, such as, for example, a polylysine compound, liposome, and the like. Examples of viral transfer vectors include, but are not limited to, adenoviral vectors, adeno-associated virus vectors, retroviral vectors, lentiviral vectors, and the like.

5 The term “expression vector” refers to a vector comprising a recombinant polynucleotide comprising expression control sequences operatively linked to a nucleotide sequence to be expressed. An expression vector comprises sufficient cis-acting elements for expression; other elements for expression can be supplied by the host cell or in an in vitro expression system. Expression vectors include all those known in the art, including cosmids, 10 plasmids (e.g., naked or contained in liposomes) and viruses (e.g., lentiviruses, retroviruses, adenoviruses, and adeno-associated viruses) that incorporate the recombinant polynucleotide.

 The term “lentivirus” refers to a genus of the Retroviridae family. Lentiviruses are unique among the retroviruses in being able to infect non-dividing cells; they can deliver a significant amount of genetic information into the DNA of the host cell, so they are one of the 15 most efficient methods of a gene delivery vector. HIV, SIV, and FIV are all examples of lentiviruses.

 The term “lentiviral vector” refers to a vector derived from at least a portion of a lentivirus genome, including especially a self-inactivating lentiviral vector as provided in Milone et al., Mol. Ther. 17(8): 1453–1464 (2009). Other examples of lentivirus vectors that 20 may be used in the clinic, include but are not limited to, e.g., the LENTIVECTOR® gene delivery technology from Oxford BioMedica, the LENTIMAX™ vector system from Lentigen and the like. Nonclinical types of lentiviral vectors are also available and would be known to one skilled in the art.

 The term “homologous” or “identity” refers to the subunit sequence identity between 25 two polymeric molecules, e.g., between two nucleic acid molecules, such as, two DNA molecules or two RNA molecules, or between two polypeptide molecules. When a subunit position in both of the two molecules is occupied by the same monomeric subunit; e.g., if a position in each of two DNA molecules is occupied by adenine, then they are homologous or identical at that position. The homology between two sequences is a direct function of the 30 number of matching or homologous positions; e.g., if half (e.g., five positions in a polymer ten subunits in length) of the positions in two sequences are homologous, the two sequences are

50% homologous; if 90% of the positions (e.g., 9 of 10), are matched or homologous, the two sequences are 90% homologous.

“Humanized” forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies and antibody fragments thereof are human immunoglobulins (recipient antibody or antibody fragment) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, a humanized antibody/antibody fragment can comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications can further refine and optimize antibody or antibody fragment performance. In general, the humanized antibody or antibody fragment thereof will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or a significant portion of the FR regions are those of a human immunoglobulin sequence. The humanized antibody or antibody fragment can also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature*, 321: 522-525, 1986; Reichmann et al., *Nature*, 332: 323-329, 1988; Presta, *Curr. Op. Struct. Biol.*, 2: 593-596, 1992.

“Fully human” refers to an immunoglobulin, such as an antibody or antibody fragment, where the whole molecule is of human origin or consists of an amino acid sequence identical to a human form of the antibody or immunoglobulin.

The term “isolated” means altered or removed from the natural state. For example, a nucleic acid or a peptide naturally present in a living animal is not “isolated,” but the same nucleic acid or peptide partially or completely separated from the coexisting materials of its natural state is “isolated.” An isolated nucleic acid or protein can exist in substantially purified form, or can exist in a non-native environment such as, for example, a host cell.

In the context of the present invention, the following abbreviations for the commonly occurring nucleic acid bases are used. “A” refers to adenosine, “C” refers to cytosine, “G” refers to guanosine, “T” refers to thymidine, and “U” refers to uridine.

The term “operably linked” or “transcriptional control” refers to functional linkage
5 between a regulatory sequence and a heterologous nucleic acid sequence resulting in expression of the latter. For example, a first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence.
10 Operably linked DNA sequences can be contiguous with each other and, e.g., where necessary to join two protein coding regions, are in the same reading frame.

The term “parenteral” administration of an immunogenic composition includes, e.g., subcutaneous (s.c.), intravenous (i.v.), intramuscular (i.m.), or intrasternal injection, intratumoral, or infusion techniques.

15 The term “nucleic acid” or “polynucleotide” refers to deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), or a combination of a DNA or RNA thereof, and polymers thereof in either single- or double-stranded form. The term “nucleic acid” includes a gene, cDNA or an mRNA. In one embodiment, the nucleic acid molecule is synthetic (e.g., chemically synthesized) or recombinant. Unless specifically limited, the term encompasses nucleic acids
20 containing analogues or derivatives of natural nucleotides that have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions), alleles, orthologs, SNPs, and complementary sequences as well as the sequence explicitly
25 indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608 (1985); and Rossolini et al., *Mol. Cell. Probes* 8:91-98 (1994)).

30 The terms “peptide,” “polypeptide,” and “protein” are used interchangeably, and refer to a compound comprised of amino acid residues covalently linked by peptide bonds. A protein

or peptide must contain at least two amino acids, and no limitation is placed on the maximum number of amino acids that can comprise a protein's or peptide's sequence. Polypeptides include any peptide or protein comprising two or more amino acids joined to each other by peptide bonds. As used herein, the term refers to both short chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, which generally are referred to in the art as proteins, of which there are many types. "Polypeptides" include, for example, biologically active fragments, substantially homologous polypeptides, oligopeptides, homodimers, heterodimers, variants of polypeptides, modified polypeptides, derivatives, analogs, fusion proteins, among others. A polypeptide includes a natural peptide, a recombinant peptide, or a combination thereof.

The term "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a polynucleotide sequence.

The term "promoter/regulatory sequence" refers to a nucleic acid sequence which is required for expression of a gene product operably linked to the promoter/regulatory sequence. In some instances, this sequence may be the core promoter sequence and in other instances, this sequence may also include an enhancer sequence and other regulatory elements which are required for expression of the gene product. The promoter/regulatory sequence may, for example, be one which expresses the gene product in a tissue specific manner.

The term "constitutive" promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a cell under most or all physiological conditions of the cell.

The term "inducible" promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a cell substantially only when an inducer which corresponds to the promoter is present in the cell.

The term "tissue-specific" promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide encodes or specified by a gene, causes the gene product to be produced in a cell substantially only if the cell is a cell of the tissue type corresponding to the promoter.

The terms “cancer associated antigen” or “tumor antigen” interchangeably refers to a molecule (typically protein, carbohydrate or lipid) that is preferentially expressed on the surface of a cancer cell, either entirely or as a fragment (e.g., MHC/peptide), in comparison to a normal cell, and which is useful for the preferential targeting of a pharmacological agent to the cancer cell. In some embodiments, a tumor antigen is a marker expressed by both normal cells and cancer cells, e.g., a lineage marker, e.g., CD19 on B cells. In certain aspects, the tumor antigens of the present invention are derived from, cancers including but not limited to primary or metastatic melanoma, thymoma, lymphoma, sarcoma, lung cancer, liver cancer, non-Hodgkin lymphoma, Hodgkin lymphoma, leukemias, uterine cancer, cervical cancer, bladder cancer, kidney cancer and adenocarcinomas such as breast cancer, prostate cancer, ovarian cancer, pancreatic cancer, and the like. In some embodiments, a cancer-associated antigen is a cell surface molecule that is overexpressed in a cancer cell in comparison to a normal cell, for instance, 1-fold over expression, 2-fold overexpression, 3-fold overexpression or more in comparison to a normal cell. In some embodiments, a cancer-associated antigen is a cell surface molecule that is inappropriately synthesized in the cancer cell, for instance, a molecule that contains deletions, additions or mutations in comparison to the molecule expressed on a normal cell. In some embodiments, a cancer-associated antigen will be expressed exclusively on the cell surface of a cancer cell, entirely or as a fragment (e.g., MHC/peptide), and not synthesized or expressed on the surface of a normal cell. In some embodiments, the CARs of the present invention includes CARs comprising an antigen binding domain (e.g., antibody or antibody fragment) that binds to a MHC presented peptide. Normally, peptides derived from endogenous proteins fill the pockets of Major histocompatibility complex (MHC) class I molecules, and are recognized by T cell receptors (TCRs) on CD8 + T lymphocytes. The MHC class I complexes are constitutively expressed by all nucleated cells. In cancer, virus-specific and/or tumor-specific peptide/MHC complexes represent a unique class of cell surface targets for immunotherapy. TCR-like antibodies targeting peptides derived from viral or tumor antigens in the context of human leukocyte antigen (HLA)-A1 or HLA-A2 have been described (see, e.g., Sastry et al., J Virol. 2011 85(5):1935-1942; Sergeeva et al., Blood, 2011 117(16):4262-4272; Verma et al., J Immunol 2010 184(4):2156-2165; Willemsen et al., Gene Ther 2001 8(21) :1601-1608 ; Dao et al., Sci Transl Med 2013 5(176) :176ra33 ; Tassev et al., Cancer Gene Ther 2012 19(2):84-100). For example, TCR-like antibody can be identified from screening a library, such as a human scFv phage displayed library.

The term “flexible polypeptide linker” or “linker” as used in the context of a scFv refers to a peptide linker that consists of amino acids such as glycine and/or serine residues used alone or in combination, to link variable heavy and variable light chain regions together. In one embodiment, the flexible polypeptide linker is a Gly/Ser linker and comprises the amino acid sequence (Gly-Gly-Gly-Ser)_n (SEQ ID NO: 15), where n is a positive integer equal to or greater than 1. For example, n=1, n=2, n=3, n=4, n=5, n=6, n=7, n=8, n=9 and n=10. In one embodiment, the flexible polypeptide linkers include, but are not limited to, (Gly₄Ser)₄ (SEQ ID NO:27) or (Gly₄Ser)₃ (SEQ ID NO:28). In another embodiment, the linkers include multiple repeats of (Gly₂Ser), (GlySer) or (Gly₃Ser) (SEQ ID NO:29). Also included within the scope of the invention are linkers described in WO2012/138475, incorporated herein by reference).

As used herein, a 5' cap (also termed an RNA cap, an RNA 7-methylguanosine cap or an RNA m⁷G cap) is a modified guanine nucleotide that has been added to the “front” or 5' end of a eukaryotic messenger RNA shortly after the start of transcription. The 5' cap consists of a terminal group which is linked to the first transcribed nucleotide. Its presence is critical for recognition by the ribosome and protection from RNases. Cap addition is coupled to transcription, and occurs co-transcriptionally, such that each influences the other. Shortly after the start of transcription, the 5' end of the mRNA being synthesized is bound by a cap-synthesizing complex associated with RNA polymerase. This enzymatic complex catalyzes the chemical reactions that are required for mRNA capping. Synthesis proceeds as a multi-step biochemical reaction. The capping moiety can be modified to modulate functionality of mRNA such as its stability or efficiency of translation.

As used herein, “in vitro transcribed RNA” refers to RNA, e.g., mRNA, that has been synthesized in vitro. Generally, the in vitro transcribed RNA is generated from an in vitro transcription vector. The in vitro transcription vector comprises a template that is used to generate the in vitro transcribed RNA.

As used herein, a “poly(A)” is a series of adenosines attached by polyadenylation to the mRNA. In some embodiments of a construct for transient expression, the polyA is between 50 and 5000 (SEQ ID NO: 30), e.g., greater than 64, e.g., greater than 100, e.g., greater than 300 or 400 poly(A) sequences can be modified chemically or enzymatically to modulate mRNA functionality such as localization, stability or efficiency of translation.

As used herein, “polyadenylation” refers to the covalent linkage of a polyadenylyl moiety, or its modified variant, to a messenger RNA molecule. In eukaryotic organisms, most messenger RNA (mRNA) molecules are polyadenylated at the 3' end. The 3' poly(A) tail is a long sequence of adenine nucleotides (often several hundred) added to the pre-mRNA through the action of an enzyme, polyadenylate polymerase. In higher eukaryotes, the poly(A) tail is added onto transcripts that contain a specific sequence, the polyadenylation signal. The poly(A) tail and the protein bound to it aid in protecting mRNA from degradation by exonucleases. Polyadenylation is also important for transcription termination, export of the mRNA from the nucleus, and translation. Polyadenylation occurs in the nucleus immediately after transcription of DNA into RNA, but additionally can also occur later in the cytoplasm. After transcription has been terminated, the mRNA chain is cleaved through the action of an endonuclease complex associated with RNA polymerase. The cleavage site is usually characterized by the presence of the base sequence AAUAAA near the cleavage site. After the mRNA has been cleaved, adenosine residues are added to the free 3' end at the cleavage site.

As used herein, “transient” refers to expression of a non-integrated transgene for a period of hours, days or weeks, wherein the period of time of expression is less than the period of time for expression of the gene if integrated into the genome or contained within a stable plasmid replicon in the host cell.

Apheresis is the process in which whole blood is removed from an individual, separated into select components, and the remainder returned to circulation. Generally, there are two methods for the separation of blood components, centrifugal and non-centrifugal. Leukapheresis results in the active selection and removal of the patient's white blood cells.

As used herein, the terms “treat”, “treatment” and “treating” refer to the reduction or amelioration of the progression, severity and/or duration of a proliferative disorder, or the amelioration of one or more symptoms (e.g., one or more discernible symptoms) of a proliferative disorder resulting from the administration of one or more therapies (e.g., one or more therapeutic agents such as a CAR of the invention). In specific embodiments, the terms “treat”, “treatment” and “treating” refer to the amelioration of at least one measurable physical parameter of a proliferative disorder, such as growth of a tumor, not necessarily discernible by the patient. In other embodiments the terms “treat”, “treatment” and “treating” -refer to the inhibition of the progression of a proliferative disorder, either physically by, e.g., stabilization

of a discernible symptom, physiologically by, e.g., stabilization of a physical parameter, or both. In other embodiments the terms “treat”, “treatment” and “treating” refer to the reduction or stabilization of tumor size or cancerous cell count.

5 The term “signal transduction pathway” refers to the biochemical relationship between a variety of signal transduction molecules that play a role in the transmission of a signal from one portion of a cell to another portion of a cell. The phrase “cell surface receptor” includes molecules and complexes of molecules capable of receiving a signal and transmitting signal across the membrane of a cell.

10 The term “subject” is intended to include living organisms in which an immune response can be elicited (e.g., mammals, human).

The term, a “substantially purified” cell refers to a cell that is essentially free of other cell types. A substantially purified cell also refers to a cell which has been separated from other cell types with which it is normally associated in its naturally occurring state. In some instances, a population of substantially purified cells refers to a homogenous population of cells. In other instances, this term refers simply to cell that have been separated from the cells with which they are naturally associated in their natural state. In some aspects, the cells are cultured in vitro. In other aspects, the cells are not cultured in vitro.

20 In the context of the present invention, “tumor antigen” or “hyperproliferative disorder antigen” or “antigen associated with a hyperproliferative disorder” refers to antigens that are common to specific hyperproliferative disorders. In certain embodiments, the tumor antigen is derived from a cancer including but not limited to primary or metastatic melanoma, thymoma, lymphoma, sarcoma, lung cancer, liver cancer, non-Hodgkin lymphoma, Hodgkin lymphoma, leukemias, uterine cancer, cervical cancer, bladder cancer, kidney cancer and adenocarcinomas such as breast cancer, prostate cancer, ovarian cancer, pancreatic cancer, and the like.

25 The term “transfected” or “transformed” or “transduced” refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A “transfected” or “transformed” or “transduced” cell is one which has been transfected, transformed or transduced with exogenous nucleic acid. The cell includes the primary subject cell and its progeny.

The term “specifically binds,” refers to an antibody, or a ligand, which recognizes and binds with a cognate binding partner protein present in a sample, but which antibody or ligand does not substantially recognize or bind other molecules in the sample.

Ranges: throughout this disclosure, various aspects of the invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. As another example, a range such as 95-99% identity, includes something with 95%, 96%, 97%, 98% or 99% identity, and includes subranges such as 96-99%, 96-98%, 96-97%, 97-99%, 97-98% and 98-99% identity. This applies regardless of the breadth of the range.

Description

Provided herein are methods of manufacturing immune effector cells (e.g., T cells, NK cells) that can be engineered with a CAR, e.g., a CAR described herein, and reaction mixtures and compositions comprising such cells.

In one aspect, the disclosure features an immune effector cell (e.g., T cell, NK cell) engineered to express a CAR, wherein the engineered immune effector cell exhibits an antitumor property. An exemplary antigen is a cancer associated antigen (i.e., tumor antigen) described herein. In one aspect, a cell is transformed with the CAR and the CAR is expressed on the cell surface. In some embodiments, the cell (e.g., T cell, NK cell) is transduced with a viral vector encoding a CAR. In some embodiments, the viral vector is a retroviral vector. In some embodiments, the viral vector is a lentiviral vector. In some such embodiments, the cell may stably express the CAR. In another embodiment, the cell (e.g., T cell, NK cell) is transfected with a nucleic acid, e.g., mRNA, cDNA, DNA, encoding a CAR. In some such embodiments, the cell may transiently express the CAR.

Furthermore, the present invention provides CAR-expressing cell, e.g., CART compositions and their use in medicaments or methods for treating, among other diseases, cancer or any malignancy or autoimmune diseases involving cells or tissues which express a tumor antigen as described herein.

- 5 In one aspect, the CAR of the invention can be used to eradicate a normal cell that express a tumor antigen as described herein, thereby applicable for use as a cellular conditioning therapy prior to cell transplantation.

Sources of Immune effector cells

- 10 In embodiments, prior to expansion and genetic modification or other modification, a source of cells, e.g., T cells or natural killer (NK) cells, can be obtained from a subject. Examples of subjects include humans, monkeys, chimpanzees, dogs, cats, mice, rats, and transgenic species thereof. T cells can be obtained from a number of sources, including peripheral blood mononuclear cells, bone marrow, lymph node tissue, cord blood, thymus
15 tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors.

- In certain aspects of the present disclosure, immune effector cells, e.g., T cells, can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled artisan, such as Ficoll™ separation. In one aspect, cells from the circulating blood of an individual are obtained by apheresis. The apheresis product typically contains
20 lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. In one aspect, the cells collected by apheresis may be washed to remove the plasma fraction and, optionally, to place the cells in an appropriate buffer or media for subsequent processing steps. In one embodiment, the cells are washed with phosphate buffered saline (PBS). In an alternative embodiment, the wash solution lacks
25 calcium and may lack magnesium or may lack many if not all divalent cations.

- Initial activation steps in the absence of calcium can lead to magnified activation. As those of ordinary skill in the art would readily appreciate a washing step may be accomplished by methods known to those in the art, such as by using a semi-automated “flow-through” centrifuge (for example, the Cobe 2991 cell processor, the Baxter CytoMate, or the
30 Haemonetics Cell Saver 5) according to the manufacturer’s instructions. After washing, the

cells may be resuspended in a variety of biocompatible buffers, such as, for example, Ca-free, Mg-free PBS, PlasmaLyte A, or other saline solution with or without buffer. Alternatively, the undesirable components of the apheresis sample may be removed and the cells directly resuspended in culture media.

5 In one aspect, T cells are isolated from peripheral blood lymphocytes by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLLTM gradient or by counterflow centrifugal elutriation.

The methods described herein can include, e.g., selection of a specific subpopulation of immune effector cells, e.g., T cells, that are a T regulatory cell-depleted population, CD25+
10 depleted cells, using, e.g., a negative selection technique, e.g., described herein. In some embodiments, the population of T regulatory-depleted cells contains less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1% of CD25+ cells.

In one embodiment, T regulatory cells, e.g., CD25+ T cells, are removed from the population using an anti-CD25 antibody, or fragment thereof, or a CD25-binding ligand, e.g.
15 IL-2. In one embodiment, the anti-CD25 antibody, or fragment thereof, or CD25-binding ligand is conjugated to a substrate, e.g., a bead, or is otherwise coated on a substrate, e.g., a bead. In one embodiment, the anti-CD25 antibody, or fragment thereof, is conjugated to a substrate as described herein.

In one embodiment, the T regulatory cells, e.g., CD25+ T cells, are removed from the
20 population using CD25 depleting reagent from MiltenyiTM. In one embodiment, the ratio of cells to CD25 depletion reagent is 1e7 cells to 20 uL, or 1e7 cells to 15 uL, or 1e7 cells to 10 uL, or 1e7 cells to 5 uL, or 1e7 cells to 2.5 uL, or 1e7 cells to 1.25 uL. In one embodiment, e.g., for T regulatory cells, e.g., CD25+ depletion, greater than 500 million cells/ml is used. In a further aspect, a concentration of cells of 600, 700, 800, or 900 million cells/ml is used.

25 In one embodiment, the population of immune effector cells to be depleted includes about 6×10^9 CD25+ T cells. In other aspects, the population of immune effector cells to be depleted include about 1×10^9 to 1×10^{10} CD25+ T cell, and any integer value in between. In one embodiment, the resulting population T regulatory-depleted cells has 2×10^9 T regulatory cells, e.g., CD25+ cells, or less (e.g., 1×10^9 , 5×10^8 , 1×10^8 , 5×10^7 , 1×10^7 , or less CD25+
30 cells).

In one embodiment, the T regulatory cells, e.g., CD25+ cells, are removed from the population using the CliniMAC system with a depletion tubing set, such as, e.g., tubing 162-01. In one embodiment, the CliniMAC system is run on a depletion setting such as, e.g., DEPLETION2.1.

5 Without wishing to be bound by a particular theory, decreasing the level of negative regulators of immune cells (e.g., decreasing the number of unwanted immune cells, e.g., T_{REG} cells), in a subject prior to apheresis or during manufacturing of a CAR-expressing cell product significantly reduces the risk of subject relapse. For example, methods of depleting T_{REG} cells are known in the art. Methods of decreasing T_{REG} cells include, but are not limited to,
10 cyclophosphamide, anti-GITR antibody (an anti-GITR antibody described herein), CD25-depletion, mTOR inhibitor, and combinations thereof.

 In some embodiments, the manufacturing methods comprise reducing the number of (e.g., depleting) T_{REG} cells prior to manufacturing of the CAR-expressing cell. For example, manufacturing methods comprise contacting the sample, e.g., the apheresis sample, with an
15 anti-GITR antibody and/or an anti-CD25 antibody (or fragment thereof, or a CD25-binding ligand), e.g., to deplete T_{REG} cells prior to manufacturing of the CAR-expressing cell (e.g., T cell, NK cell) product.

 Without wishing to be bound by a particular theory, decreasing the level of negative regulators of immune cells (e.g., decreasing the number of unwanted immune cells, e.g., T_{REG}
20 cells), in a subject prior to apheresis or during manufacturing of a CAR-expressing cell product can reduce the risk of a subject's relapse. In an embodiment, a subject is pre-treated with one or more therapies that reduce T_{REG} cells prior to collection of cells for CAR-expressing cell product manufacturing, thereby reducing the risk of subject relapse to CAR-expressing cell treatment. In an embodiment, methods of decreasing T_{REG} cells include, but are not limited to,
25 administration to the subject of one or more of cyclophosphamide, anti-GITR antibody, CD25-depletion, or a combination thereof. In an embodiment, methods of decreasing T_{REG} cells include, but are not limited to, administration to the subject of one or more of cyclophosphamide, anti-GITR antibody, CD25-depletion, mTOR inhibitor, or a combination thereof. Administration of one or more of cyclophosphamide, anti-GITR antibody, CD25-
30 depletion, or a combination thereof, can occur before, during or after an infusion of the CAR-expressing cell product. Administration of one or more of cyclophosphamide, anti-GITR

antibody, CD25-depletion, mTOR inhibitor, or a combination thereof, can occur before, during or after an infusion of the CAR-expressing cell product.

In some embodiments, the manufacturing methods comprise reducing the number of (e.g., depleting) T_{REG} cells prior to manufacturing of the CAR-expressing cell. For example, 5 manufacturing methods comprise contacting the sample, e.g., the apheresis sample, with an anti-GITR antibody and/or an anti-CD25 antibody (or fragment thereof, or a CD25-binding ligand), e.g., to deplete T_{REG} cells prior to manufacturing of the CAR-expressing cell (e.g., T cell, NK cell) product.

In an embodiment, a subject is pre-treated with cyclophosphamide prior to collection of 10 cells for CAR-expressing cell product manufacturing, thereby reducing the risk of subject relapse to CAR-expressing cell treatment (e.g., CTL019 treatment). In an embodiment, a subject is pre-treated with an anti-GITR antibody prior to collection of cells for CAR-expressing cell (e.g., T cell or NK cell) product manufacturing, thereby reducing the risk of subject relapse to CAR-expressing cell treatment.

15 In an embodiment, the CAR-expressing cell (e.g., T cell, NK cell) manufacturing process is modified to deplete T_{REG} cells prior to manufacturing of the CAR-expressing cell (e.g., T cell, NK cell) product (e.g., a CTL019 product). In an embodiment, CD25-depletion is used to deplete T_{REG} cells prior to manufacturing of the CAR-expressing cell (e.g., T cell, NK cell) product (e.g., a CTL019 product).

20 In one embodiment, the population of cells to be removed are neither the regulatory T cells or tumor cells, but cells that otherwise negatively affect the expansion and/or function of CART cells, e.g. cells expressing CD14, CD11b, CD33, CD15, or other markers expressed by potentially immune suppressive cells. In one embodiment, such cells are envisioned to be removed concurrently with regulatory T cells and/or tumor cells, or following said depletion, or 25 in another order.

The methods described herein can include more than one selection step, e.g., more than one depletion step. Enrichment of a T cell population by negative selection can be accomplished, e.g., with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method is cell sorting and/or selection via negative magnetic 30 immunoadherence or flow cytometry that uses a cocktail of monoclonal antibodies directed to

cell surface markers present on the cells negatively selected. For example, to enrich for CD4+ cells by negative selection, a monoclonal antibody cocktail can include antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8.

The methods described herein can further include removing cells from the population which express a tumor antigen, e.g., a tumor antigen that does not comprise CD25, e.g., CD19, CD30, CD38, CD123, CD20, CD14 or CD11b, to thereby provide a population of T regulatory-depleted, e.g., CD25+ depleted, and tumor antigen depleted cells that are suitable for expression of a CAR, e.g., a CAR described herein. In one embodiment, tumor antigen expressing cells are removed simultaneously with the T regulatory, e.g., CD25+ cells. For example, an anti-CD25 antibody, or fragment thereof, and an anti-tumor antigen antibody, or fragment thereof, can be attached to the same substrate, e.g., bead, which can be used to remove the cells or an anti-CD25 antibody, or fragment thereof, or the anti-tumor antigen antibody, or fragment thereof, can be attached to separate beads, a mixture of which can be used to remove the cells. In other embodiments, the removal of T regulatory cells, e.g., CD25+ cells, and the removal of the tumor antigen expressing cells is sequential, and can occur, e.g., in either order.

Also provided are methods that include removing cells from the population which express a check point inhibitor, e.g., a check point inhibitor described herein, e.g., one or more of PD1+ cells, LAG3+ cells, and TIM3+ cells, to thereby provide a population of T regulatory-depleted, e.g., CD25+ depleted cells, and check point inhibitor depleted cells, e.g., PD1+, LAG3+ and/or TIM3+ depleted cells. Exemplary check point inhibitors include PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR (e.g., TGFRbeta), e.g., as described herein. In one embodiment, check point inhibitor expressing cells are removed simultaneously with the T regulatory, e.g., CD25+ cells. For example, an anti-CD25 antibody, or fragment thereof, and an anti-check point inhibitor antibody, or fragment thereof, can be attached to the same bead which can be used to remove the cells, or an anti-CD25 antibody, or fragment thereof, and the anti-check point inhibitor antibody, or fragment thereof, can be attached to separate beads, a mixture of which can be used to remove the cells. In other embodiments, the removal of T regulatory cells, e.g.,

CD25+ cells, and the removal of the check point inhibitor expressing cells is sequential, and can occur, e.g., in either order.

Methods described herein can include a positive selection step. For example, T cells can isolated by incubation with anti-CD3/anti-CD28 (e.g., 3x28)-conjugated beads, such as
5 DYNABEADS® M-450 CD3/CD28 T, for a time period sufficient for positive selection of the desired T cells. In one embodiment, the time period is about 30 minutes. In a further embodiment, the time period ranges from 30 minutes to 36 hours or longer and all integer values there between. In a further embodiment, the time period is at least 1, 2, 3, 4, 5, or 6 hours. In yet another embodiment, the time period is 10 to 24 hours, e.g., 24 hours. Longer
10 incubation times may be used to isolate T cells in any situation where there are few T cells as compared to other cell types, such in isolating tumor infiltrating lymphocytes (TIL) from tumor tissue or from immunocompromised individuals. Further, use of longer incubation times can increase the efficiency of capture of CD8+ T cells. Thus, by simply shortening or lengthening the time T cells are allowed to bind to the CD3/CD28 beads and/or by increasing or decreasing
15 the ratio of beads to T cells (as described further herein), subpopulations of T cells can be preferentially selected for or against at culture initiation or at other time points during the process. Additionally, by increasing or decreasing the ratio of anti-CD3 and/or anti-CD28 antibodies on the beads or other surface, subpopulations of T cells can be preferentially selected for or against at culture initiation or at other desired time points.

20 In one embodiment, a T cell population can be selected that expresses one or more of IFN- γ , TNF α , IL-17A, IL-2, IL-3, IL-4, GM-CSF, IL-10, IL-13, granzyme B, and perforin, or other appropriate molecules, e.g., other cytokines. Methods for screening for cell expression can be determined, e.g., by the methods described in PCT Publication No.: WO 2013/126712.

For isolation of a desired population of cells by positive or negative selection, the
25 concentration of cells and surface (e.g., particles such as beads) can be varied. In certain aspects, it may be desirable to significantly decrease the volume in which beads and cells are mixed together (e.g., increase the concentration of cells), to ensure maximum contact of cells and beads. For example, in one aspect, a concentration of 10 billion cells/ml, 9 billion/ml, 8 billion/ml, 7 billion/ml, 6 billion/ml, or 5 billion/ml is used. In one aspect, a concentration of 1
30 billion cells/ml is used. In yet one aspect, a concentration of cells from 75, 80, 85, 90, 95, or

100 million cells/ml is used. In further aspects, concentrations of 125 or 150 million cells/ml can be used.

Using high concentrations can result in increased cell yield, cell activation, and cell expansion. Further, use of high cell concentrations allows more efficient capture of cells that may weakly express target antigens of interest, such as CD28-negative T cells, or from samples where there are many tumor cells present (e.g., leukemic blood, tumor tissue, etc.). Such populations of cells may have therapeutic value and would be desirable to obtain. For example, using high concentration of cells allows more efficient selection of CD8+ T cells that normally have weaker CD28 expression.

In a related aspect, it may be desirable to use lower concentrations of cells. By significantly diluting the mixture of T cells and surface (e.g., particles such as beads), interactions between the particles and cells is minimized. This selects for cells that express high amounts of desired antigens to be bound to the particles. For example, CD4+ T cells express higher levels of CD28 and are more efficiently captured than CD8+ T cells in dilute concentrations. In one aspect, the concentration of cells used is 5×10^6 /ml. In other aspects, the concentration used can be from about 1×10^5 /ml to 1×10^6 /ml, and any integer value in between.

In other aspects, the cells may be incubated on a rotator for varying lengths of time at varying speeds at either 2-10°C or at room temperature.

In one embodiment, a plurality of the immune effector cells of the population do not express diacylglycerol kinase (DGK), e.g., is DGK-deficient. In one embodiment, a plurality of the immune effector cells of the population do not express Ikaros, e.g., is Ikaros-deficient. In one embodiment, a plurality of the immune effector cells of the population do not express DGK and Ikaros, e.g., is both DGK and Ikaros-deficient.

T cells for stimulation can also be frozen after a washing step. Wishing not to be bound by theory, the freeze and subsequent thaw step provides a more uniform product by removing granulocytes and to some extent monocytes in the cell population. After the washing step that removes plasma and platelets, the cells may be suspended in a freezing solution. While many freezing solutions and parameters are known in the art and will be useful in this context, one method involves using PBS containing 20% DMSO and 8% human serum albumin, or culture

media containing 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin and 7.5% DMSO, or 31.25% Plasmalyte-A, 31.25% Dextrose 5%, 0.45% NaCl, 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin, and 7.5% DMSO or other suitable cell freezing media containing for example, Hespan and PlasmaLyte A, the cells then are frozen to -80°C at a rate of 1° per minute and stored in the vapor phase of a liquid nitrogen storage tank. Other methods of controlled freezing may be used as well as uncontrolled freezing immediately at -20° C or in liquid nitrogen.

In certain aspects, cryopreserved cells are thawed and washed as described herein and allowed to rest for one hour at room temperature prior to activation using the methods of the present invention.

Also contemplated in the context of the invention is the collection of blood samples or apheresis product from a subject at a time period prior to when the expanded cells as described herein might be needed. As such, the source of the cells to be expanded can be collected at any time point necessary, and desired cells, such as T cells, isolated and frozen for later use in immune effector cell therapy for any number of diseases or conditions that would benefit from immune effector cell therapy, such as those described herein. In one aspect a blood sample or an apheresis is taken from a generally healthy subject. In certain aspects, a blood sample or an apheresis is taken from a generally healthy subject who is at risk of developing a disease, but who has not yet developed a disease, and the cells of interest are isolated and frozen for later use. In certain aspects, the T cells may be expanded, frozen, and used at a later time. In certain aspects, samples are collected from a patient shortly after diagnosis of a particular disease as described herein but prior to any treatments. In a further aspect, the cells are isolated from a blood sample or an apheresis from a subject prior to any number of relevant treatment modalities, including but not limited to treatment with agents such as natalizumab, efalizumab, antiviral agents, chemotherapy, radiation, immunosuppressive agents, such as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAMPATH, anti-CD3 antibodies, cytoxan, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, and irradiation.

In a further aspect of the present invention, T cells are obtained from a patient directly following treatment that leaves the subject with functional T cells. In this regard, it has been observed that following certain cancer treatments, in particular treatments with drugs that

damage the immune system, shortly after treatment during the period when patients would normally be recovering from the treatment, the quality of T cells obtained may be optimal or improved for their ability to expand ex vivo. Likewise, following ex vivo manipulation using the methods described herein, these cells may be in a preferred state for enhanced engraftment and in vivo expansion. Thus, it is contemplated within the context of the present invention to collect blood cells, including T cells, dendritic cells, or other cells of the hematopoietic lineage, during this recovery phase. Further, in certain aspects, mobilization (for example, mobilization with GM-CSF) and conditioning regimens can be used to create a condition in a subject wherein repopulation, recirculation, regeneration, and/or expansion of particular cell types is favored, especially during a defined window of time following therapy. Illustrative cell types include T cells, B cells, dendritic cells, and other cells of the immune system.

In one embodiment, the immune effector cells expressing a CAR molecule, e.g., a CAR molecule described herein, are obtained from a subject that has received a low, immune enhancing dose of an mTOR inhibitor. In an embodiment, the population of immune effector cells, e.g., T cells, to be engineered to express a CAR, are harvested after a sufficient time, or after sufficient dosing of the low, immune enhancing, dose of an mTOR inhibitor, such that the level of PD1 negative immune effector cells, e.g., T cells, or the ratio of PD1 negative immune effector cells, e.g., T cells/ PD1 positive immune effector cells, e.g., T cells, in the subject or harvested from the subject has been, at least transiently, increased.

In other embodiments, population of immune effector cells, e.g., T cells, which have, or will be engineered to express a CAR, can be treated ex vivo by contact with an amount of an mTOR inhibitor that increases the number of PD1 negative immune effector cells, e.g., T cells or increases the ratio of PD1 negative immune effector cells, e.g., T cells/ PD1 positive immune effector cells, e.g., T cells.

It is recognized that the methods of the application can utilize culture media conditions comprising 5% or less, for example 2%, human AB serum, and employ known culture media conditions and compositions, for example those described in Smith et al., "Ex vivo expansion of human T cells for adoptive immunotherapy using the novel Xeno-free CTS Immune Cell Serum Replacement" *Clinical & Translational Immunology* (2015) 4, e31; doi:10.1038/cti.2014.31.

In one embodiment, the methods of the application can utilize culture media conditions comprising serum-free medium. In one embodiment, the serum free medium is OpTmizer CTS (LifeTech), Immunocult XF (Stemcell technologies), CellGro (CellGenix), TexMacs (Miltenyi), Stemline (Sigma), Xvivo15 (Lonza), PrimeXV (Irvine Scientific), or StemXVivo
5 (RandD systems). The serum-free medium can be supplemented with a serum substitute such as ICSR (immune cell serum replacement) from LifeTech. The level of serum substitute (e.g., ICSR) can be, e.g., up to 5%, e.g., about 1%, 2%, 3%, 4%, or 5%.

In one embodiment, a T cell population is diacylglycerol kinase (DGK)-deficient. DGK-deficient cells include cells that do not express DGK RNA or protein, or have reduced or
10 inhibited DGK activity. DGK-deficient cells can be generated by genetic approaches, e.g., administering RNA-interfering agents, e.g., siRNA, shRNA, miRNA, to reduce or prevent DGK expression. Alternatively, DGK-deficient cells can be generated by treatment with DGK inhibitors described herein.

In one embodiment, a T cell population is Ikaros-deficient. Ikaros-deficient cells
15 include cells that do not express Ikaros RNA or protein, or have reduced or inhibited Ikaros activity, Ikaros-deficient cells can be generated by genetic approaches, e.g., administering RNA-interfering agents, e.g., siRNA, shRNA, miRNA, to reduce or prevent Ikaros expression. Alternatively, Ikaros-deficient cells can be generated by treatment with Ikaros inhibitors, e.g., lenalidomide.

20 In embodiments, a T cell population is DGK-deficient and Ikaros-deficient, e.g., does not express DGK and Ikaros, or has reduced or inhibited DGK and Ikaros activity. Such DGK and Ikaros-deficient cells can be generated by any of the methods described herein.

In an embodiment, the NK cells are obtained from the subject. In another embodiment, the NK cells are an NK cell line, e.g., NK-92 cell line (Conkwest).

25 **Allogeneic CAR**

In embodiments described herein, the immune effector cell can be an allogeneic immune effector cell, e.g., T cell or NK cell. For example, the cell can be an allogeneic T cell, e.g., an allogeneic T cell lacking expression of a functional T cell receptor (TCR) and/or human leukocyte antigen (HLA), e.g., HLA class I and/or HLA class II.

A T cell lacking a functional TCR can be, e.g., engineered such that it does not express any functional TCR on its surface, engineered such that it does not express one or more subunits that comprise a functional TCR (e.g., engineered such that it does not express (or exhibits reduced expression) of TCR alpha, TCR beta, TCR gamma, TCR delta, TCR epsilon, and/or TCR zeta) or engineered such that it produces very little functional TCR on its surface. Alternatively, the T cell can express a substantially impaired TCR, e.g., by expression of mutated or truncated forms of one or more of the subunits of the TCR. The term “substantially impaired TCR” means that this TCR will not elicit an adverse immune reaction in a host.

A T cell described herein can be, e.g., engineered such that it does not express a functional HLA on its surface. For example, a T cell described herein, can be engineered such that cell surface expression HLA, e.g., HLA class I and/or HLA class II, is downregulated. In some embodiments, downregulation of HLA may be accomplished by reducing or eliminating expression of beta-2 microglobulin (B2M).

In some embodiments, the T cell can lack a functional TCR and a functional HLA, e.g., HLA class I and/or HLA class II.

Modified T cells that lack expression of a functional TCR and/or HLA can be obtained by any suitable means, including a knock out or knock down of one or more subunit of TCR or HLA. For example, the T cell can include a knock down of TCR and/or HLA using siRNA, shRNA, clustered regularly interspaced short palindromic repeats (CRISPR) transcription-activator like effector nuclease (TALEN), or zinc finger endonuclease (ZFN).

In some embodiments, the allogeneic cell can be a cell which does not express or expresses at low levels an inhibitory molecule, e.g. by any method described herein. For example, the cell can be a cell that does not express or expresses at low levels an inhibitory molecule, e.g., that can decrease the ability of a CAR-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR (e.g., TGFRbeta). Inhibition of an inhibitory molecule, e.g., by inhibition at the DNA, RNA or protein level, can optimize a CAR-expressing cell performance. In embodiments, an inhibitory nucleic acid, e.g., an inhibitory nucleic acid, e.g., a dsRNA, e.g., an siRNA or shRNA, a

clustered regularly interspaced short palindromic repeats (CRISPR), a transcription-activator like effector nuclease (TALEN), or a zinc finger endonuclease (ZFN), e.g., as described herein, can be used.

5 *siRNA and shRNA to inhibit TCR or HLA*

In some embodiments, TCR expression and/or HLA expression can be inhibited using siRNA or shRNA that targets a nucleic acid encoding a TCR and/or HLA, and/or an inhibitory molecule described herein (e.g., PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1,
10 CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR beta), in a cell, e.g., T cell.

Expression systems for siRNA and shRNAs, and exemplary shRNAs, are described, e.g., in paragraphs 649 and 650 of International Application WO2015/142675, filed March 13,
15 2015, which is incorporated by reference in its entirety.

CRISPR to inhibit TCR or HLA

“CRISPR” or “CRISPR to TCR and/or HLA” or “CRISPR to inhibit TCR and/or HLA” as used herein refers to a set of clustered regularly interspaced short palindromic repeats, or a system comprising such a set of repeats. “Cas”, as used herein, refers to a CRISPR-
20 associated protein. A “CRISPR/Cas” system refers to a system derived from CRISPR and Cas which can be used to silence or mutate a TCR and/or HLA gene, and/or an inhibitory molecule described herein (e.g., PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC
25 class I, MHC class II, GAL9, adenosine, and TGFR beta), in a cell, e.g., T cell.

The CRISPR/Cas system, and uses thereof, are described, e.g., in paragraphs 651-658 of International Application WO2015/142675, filed March 13, 2015, which is incorporated by reference in its entirety.

TALEN to inhibit TCR and/or HLA

“TALEN” or “TALEN to HLA and/or TCR” or “TALEN to inhibit HLA and/or TCR” refers to a transcription activator-like effector nuclease, an artificial nuclease which can be used to edit the HLA and/or TCR gene, and/or an inhibitory molecule described herein (e.g., PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR beta), in a cell, e.g., T cell.

TALENs, and uses thereof, are described, e.g., in paragraphs 659-665 of International Application WO2015/142675, filed March 13, 2015, which is incorporated by reference in its entirety.

Zinc finger nuclease to inhibit HLA and/or TCR

“ZFN” or “Zinc Finger Nuclease” or “ZFN to HLA and/or TCR” or “ZFN to inhibit HLA and/or TCR” refer to a zinc finger nuclease, an artificial nuclease which can be used to edit the HLA and/or TCR gene, and/or an inhibitory molecule described herein (e.g., PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (e.g., CEACAM-1, CEACAM-3 and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR beta), in a cell, e.g., T cell.

ZFNs, and uses thereof, are described, e.g., in paragraphs 666-671 of International Application WO2015/142675, filed March 13, 2015, which is incorporated by reference in its entirety.

Telomerase expression

Telomeres play a crucial role in somatic cell persistence, and their length is maintained by telomerase (TERT). Telomere length in CLL cells may be very short (Roth et al., “Significantly shorter telomeres in T-cells of patients with ZAP-70+/CD38 chronic lymphocytic leukaemia” British Journal of Haematology, 143, 383-386., August 28 2008), and may be even shorter in manufactured CAR-expressing cells, e.g., CART19 cells, limiting their potential to expand after adoptive transfer to a patient. Telomerase expression can rescue CAR-expressing cells from replicative exhaustion.

While not wishing to be bound by any particular theory, in some embodiments, a therapeutic T cell has short term persistence in a patient, due to shortened telomeres in the T cell; accordingly, transfection with a telomerase gene can lengthen the telomeres of the T cell and improve persistence of the T cell in the patient. See Carl June, "Adoptive T cell therapy for cancer in the clinic", *Journal of Clinical Investigation*, 117:1466-1476 (2007). Thus, in an embodiment, an immune effector cell, e.g., a T cell, ectopically expresses a telomerase subunit, e.g., the catalytic subunit of telomerase, e.g., TERT, e.g., hTERT. In some aspects, this disclosure provides a method of producing a CAR-expressing cell, comprising contacting a cell with a nucleic acid encoding a telomerase subunit, e.g., the catalytic subunit of telomerase, e.g., TERT, e.g., hTERT. The cell may be contacted with the nucleic acid before, simultaneous with, or after being contacted with a construct encoding a CAR.

Telomerase expression may be stable (e.g., the nucleic acid may integrate into the cell's genome) or transient (e.g., the nucleic acid does not integrate, and expression declines after a period of time, e.g., several days). Stable expression may be accomplished by transfecting or transducing the cell with DNA encoding the telomerase subunit and a selectable marker, and selecting for stable integrants. Alternatively or in combination, stable expression may be accomplished by site-specific recombination, e.g., using the Cre/Lox or FLP/FRT system.

Transient expression may involve transfection or transduction with a nucleic acid, e.g., DNA or RNA such as mRNA. In some embodiments, transient mRNA transfection avoids the genetic instability sometimes associated with stable transfection with TERT. Transient expression of exogenous telomerase activity is described, e.g., in International Application WO2014/130909, which is incorporated by reference herein in its entirety. In embodiments, mRNA-based transfection of a telomerase subunit is performed according to the messenger RNA Therapeutics™ platform commercialized by Moderna Therapeutics. For instance, the method may be a method described in US Pat. No. 8710200, 8822663, 8680069, 8754062, 8664194, or 8680069.

In an embodiment, hTERT has the amino acid sequence of GenBank Protein ID AAC51724.1 (Meyerson et al., "hEST2, the Putative Human Telomerase Catalytic Subunit Gene, Is Up-Regulated in Tumor Cells and during Immortalization" *Cell* Volume 90, Issue 4, 22 August 1997, Pages 785–795):

MPRAPRCRAVRSLLRSHYREVLPLATFVRRLLGPQGWRLVQRGDP
 AAFRALVAQCLVCPWDARPPPAAPSFRQVSLKELVARVLQRLCERGAKNVLAFGFA
 5 LLDGARGGPPEAFTTSVRSYLPNTVTDALRGSGAWGLLLRRVGDDVLVHLLARCALFV
 LVAPSCAYQVCGPPLYQLGAATQARPPPHASGPRRRLGCERAWNHSVREAGVPLGLPA
 10 PGARRRGGSSASRSLPLPKRPRRGAAPERTPVGQGSWAHPGRTRGPSDRGFCVVSPA
 RPAEEATSLEGALSGTRHSHPSVGRQHHAGPPSTSRPPRPWDTPCPPVYAETKHFLYS
 SGDKEQLRPSFLLSSLRPSLTGARRLVETIFLGSRPWMPGTPRRLPRLPQRYWQMRPL
 15 FLELLGNHAQCPYGVLLKTHCPLRAAVTPAAGVCAREKPGGSVAPEEEDTDPRRLVQ
 LLRQHSSPWQVYGFVRACLRLRLVPPGLWGSRHNERFLRNTKKFISLGKHAKLSLQEL
 20 TWKMSVRGCAWLRRSPGVGCVPAAEHRLREEILAKFLHWLMSVYVVELLRSFYVTET
 TFQKNRLFFYRKS SVWSKLQSIGIRQHLKRVQLRELSAEVRQHREARPALLTSRLRFI
 PKPDGLRP IVNMDYVVGARTFRREKRAERLTSRVKALF SVLNYERARRPGLLGASVLG
 25 LDDIHRAWRTFVLRVRAQDPPELYFVKVDVTGAYDTIPQDRLTEVIASIIKPQNTYC
 VRRYAVVQKAAHGHVRKAFKSHVSTLTDLQPYMRQFVAHLQETSPLRDAVVIEQSSSL
 30 NEASSGLFDVFLRFMCHHAVRIRGKSYVQCQGIPQGSILSTLLCSLCYGDMENKLFAG
 IRRDGLLLRLVDDFLLVTPHLTHAKTFLRTLVRGVPEYGCVVNLRKTVVNFVVEDEAL
 GGTA FVQMPAHGLFPWCGLLLDTRTLEVQSDYSSYARTSIRASLTFNRGFKAGRNMRR
 35 KLFGVLRRLKCHSLFLDLQVNSLQTVCTNIYKILLQAYRFHACVLQLPFHQQVWKNPT
 FFLRVISDTASLCYSILKAKNAGMSLGAKGAAGPLPSEAVQWLCHQAFLCLKLTRHRVT
 YVPLLGSRLTAQTQLSRKLP GTTLTALEAAANPALPSDFKTILD
 40 (SEQ ID NO: 108)

In an embodiment, the hTERT has a sequence at least 80%, 85%, 90%, 95%, 96%,
 97%, 98%, or 99% identical to the sequence of SEQ ID NO: 108. In an embodiment, the
 hTERT has a sequence of SEQ ID NO: 108. In an embodiment, the hTERT comprises a
 45 deletion (e.g., of no more than 5, 10, 15, 20, or 30 amino acids) at the N-terminus, the C-
 terminus, or both. In an embodiment, the hTERT comprises a transgenic amino acid sequence
 (e.g., of no more than 5, 10, 15, 20, or 30 amino acids) at the N-terminus, the C-terminus, or
 both.

In an embodiment, the hTERT is encoded by the nucleic acid sequence of GenBank Accession No. AF018167 (Meyerson et al., "hEST2, the Putative Human Telomerase Catalytic Subunit Gene, Is Up-Regulated in Tumor Cells and during Immortalization" Cell Volume 90, Issue 4, 22 August 1997, Pages 785–795):

```

5      1  caggcagcgt ggtcctgctg cgcacgtggg aagccctggc cccggccacc cccgcgatgc
      61  cgcgcgctcc ccgctgccga gccgtgogct ccttgcctgc cagccactac cgcgaggtgc
     121  tgcgcgtggc cacgttcgtg cggcgccctg gggcccaggg ctggcggctg gtgcagcgcg
     181  gggaccceggc ggctttccgc gcgctggtgg cccagtgcct ggtgtgcgtg ccttgggaag
     241  caeggcgcgc ccccgccgcc cctccttccc gccaggtgtc ctgcctgaag gagctggtgg
10    301  cccgagtgct gcagaggctg tgcgagcgcg gcgcaagaa cgtgctggcc ttcggcttcg
     361  cgctgctgga cggggccccc gggggccccc ccgaggcctt caccaccagc gtgcgcagct
     421  acctgcccaa cacggtgacc gacgcactgc gggggagcgg ggcgtggggg ctgctgttgc
     481  gcgcgctggg cgcgcagctg ctggttcacc tgcgtggcac ctgcgcgctc tttgtgctgg
     541  tggctcccag ctgcgcctac caggtgtgcg ggccgcgct gtaccagctc ggcgctgcc
15    601  ctcaggcccg gcccgcgcca cagctagtgc gaccocgaag gcgtctggga tgcgaacggg
     661  cctggaacca tagcgtcagg gagccggggg tccccctggg cctgccagcc ccgggtgcga
     721  ggaggcgcgg gggcagtgcc agccgaagtc tgccgttgcc caagaggccc aggcgtggcg
     781  ctgcccctga gccggagcgg acgcccgttg gccaggggtc ctggggccac ccgggcagga
     841  cgcgtggacc gagtgaccgt ggtttctgtg ttgtgtcacc tgccagaccc gccgaagaag
20    901  ccacctcttt ggagggtgcg ctctctggca cgcgccactc ccacctatcc gtgggcgcgc
     961  agcaccacgc gggcccccca tccacatcgc ggccaccacg tccctgggac acgcttgtc
    1021  ccccggtgta cgcgcgagac aagcacttcc tctactctc aggcgacaag gacgagctgc
    1081  ggccctcctt cctactcagc tctctgaggg ccagcctgac tggcgctcgg aggcctgtgg
    1141  agaccatctt tctgggttcc aggcctgga tgccagggac tcccgccagg ttgcccgcgc
25    1201  tgcccacagc ctactggcaa atgcggcccc tgtttctgga gctgcttggg aaccacgcgc
     1261  agtgccccta cggggtgctc ctcaagacgc actgcccgct gcgagctgcg gtcacccag
     1321  cagccggtgt ctgtgcccgg gagaagcccc agggctctgt ggcgccccc gaggaggagg
     1381  acacagaccc ccgtcgctcg gtgcagctgc tccgccagca cagcagcccc tggcaggtgt
     1441  acggcttcgt gcgggctgc ctgcgcgggc ttgtgcccc aggcctctgg ggctccaggc
30    1501  acaacgaacg ccgcttcctc aggaacacca agaagttcat ctccctgggg aagcatgcca
     1561  agctctcgct gcaggagctg acgtggaaga tgagcgtgcg gggtgcgct tggctgcgca
     1621  ggagcccagg ggttggtgtg gttccggccg cagagcaccg tctgcgtgag gagatcctgg
     1681  ccaagttcct gcactggctg atgagtggtg acgtcgctga gctgctcagg tcttctttt
     1741  atgtcacgga gaccacgttt caaaagaaca ggctcttttt ctaccggaag agtgtctgga
35    1801  gcaagttgca aagcattgga atcagacagc acttgaagag ggtgcagctg cgggagctgt
     1861  cggaagcaga ggtcaggcag catcggaag ccaggcccg cctgctgacg tccagactcc
     1921  gcttcatccc caagcctgac gggctgcggc cgatttgtga catggactac gtcgtgggag
     1981  ccagaacgtt ccgcagagaa aagaggcccg agcgtctcac ctgaggggtg aaggcactgt
     2041  tcagcgtgct caactacgag cggcgccggc gccccggcct cctgggcgcc tctgtgctgg
40    2101  gcctggacga tatccacagg gcctggcgca ccttcgtgct gcgtgtgcgg gcccaggacc
     2161  cgcgcgctga gctgtacttt gtcaaggtgg atgtgacggg cgcgtacgac accatcccc
     2221  aggacaggct cacggaggtc atcgccagca tcatcaaacc ccagaacacg tactgcgtgc
     2281  gtcggtatgc cgtggtccag aaggccgccc atgggcacgt ccgcaaggcc ttcaagagcc
     2341  acgtctctac cttgacagac ctccagccgt acatgcgaca gttcgtggct cacctgcagg
45    2401  agaccagccc gctgagggat gccgtcgta tcgagcagag ctccctccctg aatgaggcca
     2461  gcagtggcct cttcgacgtc ttctacgct tcatgtgcca ccacgcgctg cgcacagggg
     2521  gcaagtctta cgtccagtgc caggggatcc cgcagggtc catcctctcc acgctgctct
     2581  gcagcctgtg ctacggcgac atggagaaca agctgtttgc ggggattcgg cgggacgggc
     2641  tgctcctgcg tttggtggat gatttcttgt ttgtgacacc tcacctcacc caccgaaaa
50    2701  ccttctcag gacctggtc cgaggtgtcc ctgagtatgg ctgcgtggtg aacttcgga
     2761  agacagtggg gaacttccct gtagaagacg aggcctggg tggcacggct tttgttcaga
     2821  tgccggccca cggcctattc cctgggtgcg gcctgctgct ggatacccg accctggagg
     2881  tgcagagcga ctactccagc tatgcccgga cctccatcag agccagctc accttcaacc
     2941  gcggcttcaa ggctgggagg aacatgcgtc gcaaaactct tggggtcttg cggctgaagt

```

5 3001 gtcacagcct gtttctggat ttgcaggtga acagcctcca gacggtgtgc accaacatct
 3061 acaagatcct cctgctgcag gcgtacaggt ttcaogcatg tgtgctgcag ctcccatttc
 3121 atcagcaagt ttggaagaac cccacatttt tctgcgcggt catctctgac acggcctccc
 3181 tctgctactc catcctgaaa gccagaacg cagggatgtc gctgggggccc aagggcgccg
 3241 cccggccctct gccctccgag gccgtgcagt ggctgtgcc acaagcattc ctgctcaage
 3301 tgactcgaca ccgtgtcacc tacgtgccac tctgggggtc actcaggaca gccagacgc
 3361 agctgagtcg gaagctcccg gggacgacgc tgactgccct ggaggccgca gccaaccccg
 3421 cactgccctc agacttcaag accatcctgg actgatggcc acccgccac agccaggccg
 3481 agagcagaca ccagcagccc tgtcacgccg ggctctacgt cccagggagg gagggcgccg
 10 3541 ccacaccag gcccgacccg ctgggagtc gaggcctgag tgagtgtttg gccgaggcct
 3601 gcatgtccgg ctgaaggctg agtgtccggc tgaggcctga gcgagtgtcc agccaagggc
 3661 tgagtgtcca gcacacctgc cgtcttcaact tccccacagg ctggcgctcg gctccacccc
 3721 agggccagct tttctcacc aggagcccg cttccactcc ccacatagga atagtccatc
 3781 cccagattcg ccattgttca cccctcgccc tgccctcctt tgccctccac ccccaccatc
 15 3841 caggtggaga cccctgagaag gacctggga gctctggga tttggagtga ccaaaggtgt
 3901 gccctgtaca caggcgagga cctgcacct ggatgggggt cctgtgggt caaattgggg
 3961 ggaggtgctg tgggagtaaa atactgaata tatgagtttt tcagttttga aaaaaaaaaa
 4021 aaaaaaa
 (SEQ ID NO: 23)
 20

In an embodiment, the hTERT is encoded by a nucleic acid having a sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the sequence of SEQ ID NO: 23.

In an embodiment, the hTERT is encoded by a nucleic acid of SEQ ID NO: 23.

Chimeric Antigen Receptor (CAR)

25 The present invention provides immune effector cells (e.g., T cells, NK cells) that are engineered to contain one or more CARs that direct the immune effector cells to cancer. This is achieved through an antigen binding domain on the CAR that is specific for a cancer associated antigen. There are two classes of cancer associated antigens (tumor antigens) that can be targeted by the CARs described herein: (1) cancer associated antigens that are expressed
 30 on the surface of cancer cells; and (2) cancer associated antigens that itself is intracellular, however, a fragment of such antigen (peptide) is presented on the surface of the cancer cells by MHC (major histocompatibility complex).

Accordingly, an immune effector cell, e.g., obtained by a method described herein, can be engineered to contain a CAR that target one of the following cancer associated antigens
 35 (tumor antigens): CD19, CD123, CD22, CD30, CD171, CS-1, CLL-1, CD33, EGFRvIII, GD2, GD3, BCMA, Tn Ag, PSMA, ROR1, FLT3, FAP, TAG72, CD38, CD44v6, CEA, EPCAM, B7H3, KIT, IL-13Ra2, Mesothelin, IL-11Ra, PSCA, VEGFR2, LewisY, CD24, PDGFR-beta, PRSS21, SSEA-4, CD20, Folate receptor alpha, ERBB2 (Her2/neu), MUC1, EGFR, NCAM, Prostase, PAP, ELF2M, Ephrin B2, IGF-I receptor, CAIX, LMP2, gp100, bcr-abl, tyrosinase,

EphA2, Fucosyl GM1, sLe, GM3, TGS5, HMWMAA, o-acetyl-GD2, Folate receptor beta, TEM1/CD248, TEM7R, CLDN6, TSHR, GPRC5D, CXORF61, CD97, CD179a, ALK, Plsialic acid, PLAC1, GloboH, NY-BR-1, UPK2, HAVCR1, ADRB3, PANX3, GPR20, LY6K, OR51E2, TARP, WT1, NY-ESO-1, LAGE-1a, legumain, HPV E6,E7, MAGE-A1, MAGE A1, ETV6-AML, sperm protein 17, XAGE1, Tie 2, MAD-CT-1, MAD-CT-2, Fos-related antigen 1, p53, p53 mutant, prostatein, survivin and telomerase, PCTA-1/Galectin 8, MelanA/MART1, Ras mutant, hTERT, sarcoma translocation breakpoints, ML-IAP, ERG (TMPRSS2 ETS fusion gene), NA17, PAX3, Androgen receptor, Cyclin B1, MYCN, RhoC, TRP-2, CYP1B1, BORIS, SART3, PAX5, OY-TES1, LCK, AKAP-4, SSX2, RAGE-1, human telomerase reverse transcriptase, RU1, RU2, intestinal carboxyl esterase, and mut hsp70-2.

Bispecific CARs

In an embodiment a multispecific antibody molecule is a bispecific antibody molecule. A bispecific antibody has specificity for no more than two antigens. A bispecific antibody molecule is characterized by a first immunoglobulin variable domain sequence which has binding specificity for a first epitope and a second immunoglobulin variable domain sequence that has binding specificity for a second epitope. In an embodiment the first and second epitopes are on the same antigen, e.g., the same protein (or subunit of a multimeric protein). In an embodiment the first and second epitopes overlap. In an embodiment the first and second epitopes do not overlap. In an embodiment the first and second epitopes are on different antigens, e.g., different proteins (or different subunits of a multimeric protein). In an embodiment a bispecific antibody molecule comprises a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a first epitope and a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a half antibody having binding specificity for a first epitope and a half antibody having binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a half antibody, or fragment thereof, having binding specificity for a first epitope and a half antibody, or fragment thereof, having binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a scFv, or fragment thereof, have binding specificity for a first epitope and a scFv, or fragment thereof, have binding specificity for a second epitope.

In certain embodiments, the antibody molecule is a multi-specific (e.g., a bispecific or a trispecific) antibody molecule. Protocols for generating bispecific or heterodimeric antibody molecules, and various configurations for bispecific antibody molecules, are described in, e.g., paragraphs 455-458 of WO2015/142675, filed March 13, 2015, which is incorporated by
5 reference in its entirety.

In one aspect, the bispecific antibody molecule is characterized by a first immunoglobulin variable domain sequence, e.g., a scFv, which has binding specificity for CD19, e.g., comprises a scFv as described herein, or comprises the light chain CDRs and/or heavy chain CDRs from a scFv described herein, and a second immunoglobulin variable
10 domain sequence that has binding specificity for a second epitope on a different antigen.

Chimeric TCR

In one aspect, the antibodies and antibody fragments of the present invention (e.g., CD19 antibodies and fragments) can be grafted to one or more constant domain of a T cell receptor ("TCR") chain, for example, a TCR alpha or TCR beta chain, to create a chimeric
15 TCR. Without being bound by theory, it is believed that chimeric TCRs will signal through the TCR complex upon antigen binding. For example, an scFv as disclosed herein, can be grafted to the constant domain, e.g., at least a portion of the extracellular constant domain, the transmembrane domain and the cytoplasmic domain, of a TCR chain, for example, the TCR alpha chain and/or the TCR beta chain. As another example, an antibody fragment, for
20 example a VL domain as described herein, can be grafted to the constant domain of a TCR alpha chain, and an antibody fragment, for example a VH domain as described herein, can be grafted to the constant domain of a TCR beta chain (or alternatively, a VL domain may be grafted to the constant domain of the TCR beta chain and a VH domain may be grafted to a TCR alpha chain). As another example, the CDRs of an antibody or antibody fragment may be
25 grafted into a TCR alpha and/or beta chain to create a chimeric TCR. For example, the LCDRs disclosed herein may be grafted into the variable domain of a TCR alpha chain and the HCDRs disclosed herein may be grafted to the variable domain of a TCR beta chain, or vice versa. Such chimeric TCRs may be produced, e.g., by methods known in the art (For example, Willemsen RA et al, Gene Therapy 2000; 7: 1369-1377; Zhang T et al, Cancer Gene Ther
30 2004; 11: 487-496; Aggen et al, Gene Ther. 2012 Apr;19(4):365-74).

Non-Antibody Scaffolds

In embodiments, the antigen binding domain comprises a non-antibody scaffold, e.g., a fibronectin, ankyrin, domain antibody, lipocalin, small modular immuno-pharmaceutical, maxybody, Protein A, or affilin. The non-antibody scaffold has the ability to bind to target
5 antigen on a cell. In embodiments, the antigen binding domain is a polypeptide or fragment thereof of a naturally occurring protein expressed on a cell. In some embodiments, the antigen binding domain comprises a non-antibody scaffold. A wide variety of non-antibody scaffolds can be employed so long as the resulting polypeptide includes at least one binding region which specifically binds to the target antigen on a target cell.

10 Non-antibody scaffolds include: fibronectin (Novartis, MA), ankyrin (Molecular Partners AG, Zurich, Switzerland), domain antibodies (Domantis, Ltd., Cambridge, MA, and Ablynx nv, Zwijnaarde, Belgium), lipocalin (Pieris Proteolab AG, Freising, Germany), small modular immuno-pharmaceuticals (Trubion Pharmaceuticals Inc., Seattle, WA), maxybodies (Avidia, Inc., Mountain View, CA), Protein A (Affibody AG, Sweden), and affilin (gamma-
15 crystallin or ubiquitin) (Scil Proteins GmbH, Halle, Germany).

In an embodiment the antigen binding domain comprises the extracellular domain, or a counter-ligand binding fragment thereof, of molecule that binds a counterligand on the surface of a target cell.

The immune effector cells can comprise a recombinant DNA construct comprising
20 sequences encoding a CAR, wherein the CAR comprises an antigen binding domain (e.g., antibody or antibody fragment, TCR or TCR fragment) that binds specifically to a tumor antigen, e.g., an tumor antigen described herein, and an intracellular signaling domain. The intracellular signaling domain can comprise a costimulatory signaling domain and/or a primary signaling domain, e.g., a zeta chain. As described elsewhere, the methods described herein can
25 include transducing a cell, e.g., from the population of T regulatory-depleted cells, with a nucleic acid encoding a CAR, e.g., a CAR described herein.

In specific aspects, a CAR comprises a scFv domain, wherein the scFv may be preceded by an optional leader sequence such as provided in SEQ ID NO: 1, and followed by an optional hinge sequence such as provided in SEQ ID NO:2 or SEQ ID NO:36 or SEQ ID NO:38, a
30 transmembrane region such as provided in SEQ ID NO:6, an intracellular signalling domain

that includes SEQ ID NO:7 or SEQ ID NO:16 and a CD3 zeta sequence that includes SEQ ID NO:9 or SEQ ID NO:10, e.g., wherein the domains are contiguous with and in the same reading frame to form a single fusion protein.

In one aspect, an exemplary CAR constructs comprise an optional leader sequence (e.g.,
5 a leader sequence described herein), an extracellular antigen binding domain (e.g., an antigen binding domain described herein), a hinge (e.g., a hinge region described herein), a transmembrane domain (e.g., a transmembrane domain described herein), and an intracellular stimulatory domain (e.g., an intracellular stimulatory domain described herein). In one aspect, an exemplary CAR construct comprises an optional leader sequence (e.g., a leader sequence
10 described herein), an extracellular antigen binding domain (e.g., an antigen binding domain described herein), a hinge (e.g., a hinge region described herein), a transmembrane domain (e.g., a transmembrane domain described herein), an intracellular costimulatory signaling domain (e.g., a costimulatory signaling domain described herein) and/or an intracellular primary signaling domain (e.g., a primary signaling domain described herein).

15 An exemplary leader sequence is provided as SEQ ID NO: 1. An exemplary hinge/spacer sequence is provided as SEQ ID NO: 2 or SEQ ID NO:36 or SEQ ID NO:38. An exemplary transmembrane domain sequence is provided as SEQ ID NO:6. An exemplary sequence of the intracellular signaling domain of the 4-1BB protein is provided as SEQ ID NO: 7. An exemplary sequence of the intracellular signaling domain of CD27 is provided as SEQ
20 ID NO:16. An exemplary CD3zeta domain sequence is provided as SEQ ID NO: 9 or SEQ ID NO:10.

In one aspect, the immune effector cell comprises a recombinant nucleic acid construct comprising a nucleic acid molecule encoding a CAR, wherein the nucleic acid molecule comprises a nucleic acid sequence encoding an antigen binding domain, wherein the sequence
25 is contiguous with and in the same reading frame as the nucleic acid sequence encoding an intracellular signaling domain. An exemplary intracellular signaling domain that can be used in the CAR includes, but is not limited to, one or more intracellular signaling domains of, e.g., CD3-zeta, CD28, CD27, 4-1BB, and the like. In some instances, the CAR can comprise any combination of CD3-zeta, CD28, 4-1BB, and the like.

30 The nucleic acid sequences coding for the desired molecules can be obtained using recombinant methods known in the art, such as, for example by screening libraries from cells

expressing the nucleic acid molecule, by deriving the nucleic acid molecule from a vector known to include the same, or by isolating directly from cells and tissues containing the same, using standard techniques. Alternatively, the nucleic acid of interest can be produced synthetically, rather than cloned.

5 Nucleic acids encoding a CAR can be introduced into the immune effector cells using, e.g., a retroviral or lentiviral vector construct.

Nucleic acids encoding a CAR can also be introduced into the immune effector cell using, e.g., an RNA construct that can be directly transfected into a cell. A method for generating mRNA for use in transfection involves *in vitro* transcription (IVT) of a template
10 with specially designed primers, followed by polyA addition, to produce a construct containing 3' and 5' untranslated sequence ("UTR") (e.g., a 3' and/or 5' UTR described herein), a 5' cap (e.g., a 5' cap described herein) and/or Internal Ribosome Entry Site (IRES) (e.g., an IRES described herein), the nucleic acid to be expressed, and a polyA tail, typically 50-2000 bases in length (e.g., described in the Examples, e.g., SEQ ID NO:35). RNA so produced can efficiently
15 transfect different kinds of cells. In one embodiment, the template includes sequences for the CAR. In an embodiment, an RNA CAR vector is transduced into a cell, e.g., a T cell by electroporation.

Antigen binding domain

In one aspect, a plurality of the immune effector cells, e.g., the population of T
20 regulatory-depleted cells, include a nucleic acid encoding a CAR that comprises a target-specific binding element otherwise referred to as an antigen binding domain. The choice of binding element depends upon the type and number of ligands that define the surface of a target cell. For example, the antigen binding domain may be chosen to recognize a ligand that acts as a cell surface marker on target cells associated with a particular disease state. Thus, examples
25 of cell surface markers that may act as ligands for the antigen binding domain in a CAR described herein include those associated with viral, bacterial and parasitic infections, autoimmune disease and cancer cells.

In one aspect, the portion of the CAR comprising the antigen binding domain comprises an antigen binding domain that targets a tumor antigen, e.g., a tumor antigen described herein.

The antigen binding domain can be any domain that binds to the antigen including but not limited to a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, and a functional fragment thereof, including but not limited to a single-domain antibody such as a heavy chain variable domain (VH), a light chain variable domain (VL) and a variable domain (VHH) of camelid derived nanobody, and to an alternative scaffold known in the art to function as antigen binding domain, such as a recombinant fibronectin domain, a T cell receptor (TCR), or a fragment thereof, e.g., single chain TCR, and the like. In some instances, it is beneficial for the antigen binding domain to be derived from the same species in which the CAR will ultimately be used in. For example, for use in humans, it may be beneficial for the antigen binding domain of the CAR to comprise human or humanized residues for the antigen binding domain of an antibody or antibody fragment.

In an embodiment, the antigen binding domain comprises an anti-CD19 antibody, or fragment thereof, e.g., an scFv. For example, the antigen binding domain comprises a variable heavy chain and a variable light chain listed in Table 1. The linker sequence joining the variable heavy and variable light chains can be, e.g., any of the linker sequences described herein, or alternatively, can be GSTSGSGKPGSGEGSTKG (SEQ ID NO:104).

Table 1: Anti-CD19 antibody binding domains

CD19	huscFv1 (SEQ ID NO: 39)	EIVMTQSPATLSLSPGERATLSCRASQDISKYLNWYQQKPGQAPRLLIYHTSRLHSGIPARFSGSGSGTDYTLTISSLPEDFAVYFCQQGNTLPYTFGGQTKLEIKG GGGSGGGSGGGSGVQLQESGPGLVKPSSETLSLTCTVSGVSLPDYGVSWIRQPPGKLEWIGVIWGSETTYSSSLKSRVTISKDNSKNQVSLKLSSVTAADTAVYYCAKHYYYGGSYAMDYWGQGLVTVSS
CD19	huscFv2 (SEQ ID NO: 40)	Eivmtqspatls slspgeratls crasqdiskylnw yqqkpgqap rlliyhtsr l hsgip arfsgsgsgtdytl t isslp edfavyfcqqgntlpytf gggtkleikggggsgggsg gggsqvlqesgpglvkps etls l tctvsgvslpdygvswir qppgkglewigviwgse t tyyqsslksrvtiskd nsknqvslklssvtaadtavyyca khyyyggsyamydgqgt lvtvss
CD19	huscFv3 (SEQ ID NO: 41)	Qvqlqesgpglvkps etls l tctvsgvslpdygvswir qppgkglewigviwgsettyy ssslksrvtiskd nsknqvslklssvtaadtavyyca khyyyggsyamydgqgtlvtv ssggggsgggsgggsgseivmtqspatls slspgeratls crasqdiskylnw yqqkpgq ap rlliyhtsr l hsgiparfsgsgsgtdytl t isslp edfavyfcqqgntlpytf ggg tkleik
CD19	huscFv4 (SEQ ID NO: 42)	Qvqlqesgpglvkps etls l tctvsgvslpdygvswir qppgkglewigviwgsettyy qsslksrvtiskd nsknqvslklssvtaadtavyyca khyyyggsyamydgqgtlvtv ssggggsgggsgggsgseivmtqspatls slspgeratls crasqdiskylnw yqqkpgq ap rlliyhtsr l hsgiparfsgsgsgtdytl t isslp edfavyfcqqgntlpytf ggg tkleik
CD19	huscFv5 (SEQ ID)	Eivmtqspatls slspgeratls crasqdiskylnw yqqkpgqap rlliyhtsr l hsgip arfsgsgsgtdytl t isslp edfavyfcqqgntlpytf gggtkleikggggsgggsg ggsgggsgqvlqesgpglvkps etls l tctvsgvslpdygvswir qppgkglewigv iwgsettyyssslksrvtiskd nsknqvslklssvtaadtavyyca khyyyggsyamyd

	NO: 43)	wgqgtlvtvss
CD19	huscFv6 (SEQ ID NO: 44)	Eivmtqspatlsispgeratlscrasqdiskylnwyqqkpgqaprrliihtsrhsgip arfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfqggtkleik <u>ggggsgggsg</u> <u>ggsgggsgsgvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigv</u> iwgsettyyqsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdy wgqgtlvtvss
CD19	huscFv7 (SEQ ID NO: 45)	Qvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgsettyy sslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgqgtlvtv ss <u>ggggsgggsgggsgggsgggg</u> seivmtqspatlsispgeratlscrasqdiskylnwyq qkpgqaprrliihtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqgntlpy tfqggtkleik
CD19	huscFv8 (SEQ ID NO: 46)	Qvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgsettyy qsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgqgtlvtv ss <u>ggggsgggsgggsgggsgggg</u> seivmtqspatlsispgeratlscrasqdiskylnwyq qkpgqaprrliihtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqgntlpy tfqggtkleik
CD19	huscFv9 (SEQ ID NO: 47)	Eivmtqspatlsispgeratlscrasqdiskylnwyqqkpgqaprrliihtsrhsgip arfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfqggtkleik <u>ggggsgggsg</u> <u>ggsgggsgsgvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigv</u> iwgsettyynsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdy wgqgtlvtvss
CD19	Hu scFv10 (SEQ ID NO: 48)	Qvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgsettyy nsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgqgtlvtv ss <u>ggggsgggsgggsgggsgggg</u> seivmtqspatlsispgeratlscrasqdiskylnwyq qkpgqaprrliihtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqgntlpy tfqggtkleik
CD19	Hu scFv11 (SEQ ID NO: 49)	Eivmtqspatlsispgeratlscrasqdiskylnwyqqkpgqaprrliihtsrhsgip arfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfqggtkleik <u>ggggsgggsg</u> <u>ggsgvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgse</u> ttyynsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgqgt lvtvss
CD19	Hu scFv12 (SEQ ID NO: 50)	Qvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgsettyy nsslskrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgqgtlvtv ss <u>ggggsgggsgggsgggsgggg</u> seivmtqspatlsispgeratlscrasqdiskylnwyqqkpgq aprrliihtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfqg tkleik
CD19	muCTL0 19 (SEQ ID NO: 51)	Diqmtqtsslsaslgdrvtiscrasqdiskylnwyqqkpdgtvkllyhtsrhsgvp srfsgsgsgtdysltisnleqediatyfcqqgntlpytfgggtkleit <u>ggggsgggsg</u> <u>gggsevklqesgpglvapsqslsvtctvsgvslpdygvswirpprglewlgviwgse</u> ttyynsalksrliikdsksqvflkmnslqtdtdaiyycahyyyggsyamdywgqgt svtvss

Table 2: Additional anti-CD19 antibody binding domains

Antibody	VH Sequence	VL Sequence
SSJ25-C1	QVQLLES GAELVRPGSSVKISCKASGYAFSS YWMNWVKRPGQGLEWIGQIYPGDGDTNYNG KFKGQATLTADKSSSTAYMQLSGLTSEDSAV YSCARKTISSVDFYFDYWQGTTVT (SEQ	ELVLTQSPKFMSTSVGDRVSVTCKASQNVGTNVA WYQQKPGQSPKPLIYSATYRNSGVDRFTGSGSG TDFTLTITNVQSKDLADYFYFCQYNRPYPYTS TKLEIKRRS (SEQ ID NO: 4)

	ID NO: 3)	
--	-----------	--

Any CD19 CAR, e.g., the CD19 antigen binding domain of any known CD19 CAR, can be used in accordance with the present disclosure. For example, LG-740; CD19 CAR described in the US Pat. No. 8,399,645; US Pat. No. 7,446,190; Xu et al., Leuk Lymphoma. 5 2013 54(2):255-260(2012); Cruz et al., Blood 122(17):2965-2973 (2013); Brentjens et al., Blood, 118(18):4817-4828 (2011); Kochenderfer et al., Blood 116(20):4099-102 (2010); Kochenderfer et al., Blood 122 (25):4129-39(2013); and 16th Annu Meet Am Soc Gen Cell Ther (ASGCT) (May 15-18, Salt Lake City) 2013, Abst 10.

Exemplary target antigens that can be targeted using the CAR-expressing cells, include, 10 but are not limited to, CD19, CD123, EGFRvIII, mesothelin, among others, as described in, for example, WO 2014/130635, WO 2014/130657, and WO 2015/090230, each of which is herein incorporated by reference in its entirety.

In one embodiment, the CAR T cell that specifically binds to CD19 has the USAN designation TISAGENLECLEUCEL-T. CTL019 is made by a gene modification of T cells is 15 mediated by stable insertion via transduction with a self-inactivating, replication deficient Lentiviral (LV) vector containing the CTL019 transgene under the control of the EF-1 alpha promoter. CTL019 can be a mixture of transgene positive and negative T cells that are delivered to the subject on the basis of percent transgene positive T cells.

In other embodiments, the CAR-expressing cells can specifically bind to human CD19, 20 *e.g.*, can include a CAR molecule, or an antigen binding domain (*e.g.*, a humanized antigen binding domain) according to Table 3 of WO2014/153270, incorporated herein by reference.

In other embodiments, the CAR-expressing cells can specifically bind to CD123, *e.g.*, can include a CAR molecule (*e.g.*, any of the CAR1-CAR8), or an antigen binding domain according to Tables 1-2 of WO 2014/130635, incorporated herein by reference.

25 In other embodiments, the CAR-expressing cells can specifically bind to EGFRvIII, *e.g.*, can include a CAR molecule, or an antigen binding domain according to Table 2 or SEQ ID NO:11 of WO 2014/130657, incorporated herein by reference.

In other embodiments, the CAR-expressing cells can specifically bind to mesothelin, *e.g.*, can include a CAR molecule, or an antigen binding domain according to Tables 2-3 of WO 2015/090230, incorporated herein by reference.

In one embodiment, the antigen binding domain comprises one, two three (*e.g.*, all three) heavy chain CDRs, HC CDR1, HC CDR2 and HC CDR3, from an antibody listed above, and/or one, two, three (*e.g.*, all three) light chain CDRs, LC CDR1, LC CDR2 and LC CDR3, from an antibody listed above. In one embodiment, the antigen binding domain comprises a heavy chain variable region and/or a variable light chain region of an antibody listed or described above.

In some embodiments, the tumor antigen is a tumor antigen described in International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety. In some embodiments, the tumor antigen is chosen from one or more of: CD19; CD123; CD22; CD30; CD171; CS-1 (also referred to as CD2 subset 1, CRACC, SLAMF7, CD319, and 19A24); C-type lectin-like molecule-1 (CLL-1 or CLECL1); CD33; epidermal growth factor receptor variant III (EGFRvIII); ganglioside G2 (GD2); ganglioside GD3 (aNeu5Ac(2-8)aNeu5Ac(2-3)bDGalp(1-4)bDGlc(1-1)Cer); TNF receptor family member B cell maturation (BCMA); Tn antigen ((Tn Ag) or (GalNAc α -Ser/Thr)); prostate-specific membrane antigen (PSMA); Receptor tyrosine kinase-like orphan receptor 1 (ROR1); Fms-Like Tyrosine Kinase 3 (FLT3); Tumor-associated glycoprotein 72 (TAG72); CD38; CD44v6; Carcinoembryonic antigen (CEA); Epithelial cell adhesion molecule (EPCAM); B7H3 (CD276); KIT (CD117); Interleukin-13 receptor subunit alpha-2 (IL-13Ra2 or CD213A2); Mesothelin; Interleukin 11 receptor alpha (IL-11Ra); prostate stem cell antigen (PSCA); Protease Serine 21 (Testisin or PRSS21); vascular endothelial growth factor receptor 2 (VEGFR2); Lewis(Y) antigen; CD24; Platelet-derived growth factor receptor beta (PDGFR-beta); Stage-specific embryonic antigen-4 (SSEA-4); CD20; Folate receptor alpha; Receptor tyrosine-protein kinase ERBB2 (Her2/neu); Mucin 1, cell surface associated (MUC1); epidermal growth factor receptor (EGFR); neural cell adhesion molecule (NCAM); Prostase; prostatic acid phosphatase (PAP); elongation factor 2 mutated (ELF2M); Ephrin B2; fibroblast activation protein alpha (FAP); insulin-like growth factor 1 receptor (IGF-I receptor), carbonic anhydrase IX (CAIX); Proteasome (Prosome, Macropain) Subunit, Beta Type, 9 (LMP2); glycoprotein 100 (gp100); oncogene fusion protein consisting of breakpoint cluster region

(BCR) and Abelson murine leukemia viral oncogene homolog 1 (Abl) (bcr-abl); tyrosinase; ephrin type-A receptor 2 (EphA2); Fucosyl GM1; sialyl Lewis adhesion molecule (sLe); ganglioside GM3 (aNeu5Ac(2-3) β DGalp(1-4) β DGlc(1-1)Cer); transglutaminase 5 (TGS5); high molecular weight-melanoma-associated antigen (HMWMAA); o-acetyl-GD2 ganglioside (OAcGD2); Folate receptor beta; tumor endothelial marker 1 (TEM1/CD248); tumor endothelial marker 7-related (TEM7R); claudin 6 (CLDN6); thyroid stimulating hormone receptor (TSHR); G protein-coupled receptor class C group 5, member D (GPRC5D); chromosome X open reading frame 61 (CXORF61); CD97; CD179a; anaplastic lymphoma kinase (ALK); Polysialic acid; placenta-specific 1 (PLAC1); hexasaccharide portion of globoH glycosphingolipid (GloboH); mammary gland differentiation antigen (NY-BR-1); uroplakin 2 (UPK2); Hepatitis A virus cellular receptor 1 (HAVCR1); adrenoceptor beta 3 (ADRB3); pannexin 3 (PANX3); G protein-coupled receptor 20 (GPR20); lymphocyte antigen 6 complex, locus K 9 (LY6K); Olfactory receptor 51E2 (OR51E2); TCR Gamma Alternate Reading Frame Protein (TARP); Wilms tumor protein (WT1); Cancer/testis antigen 1 (NY-ESO-1); Cancer/testis antigen 2 (LAGE-1a); Melanoma-associated antigen 1 (MAGE-A1); ETS translocation-variant gene 6, located on chromosome 12p (ETV6-AML); sperm protein 17 (SPA17); X Antigen Family, Member 1A (XAGE1); angiopoietin-binding cell surface receptor 2 (Tie 2); melanoma cancer testis antigen-1 (MAD-CT-1); melanoma cancer testis antigen-2 (MAD-CT-2); Fos-related antigen 1; tumor protein p53 (p53); p53 mutant; prostein; surviving; telomerase; prostate carcinoma tumor antigen-1 (PCTA-1 or Galectin 8), melanoma antigen recognized by T cells 1 (MelanA or MART1); Rat sarcoma (Ras) mutant; human Telomerase reverse transcriptase (hTERT); sarcoma translocation breakpoints; melanoma inhibitor of apoptosis (ML-IAP); ERG (transmembrane protease, serine 2 (TMPRSS2) ETS fusion gene); N-Acetyl glucosaminyl-transferase V (NA17); paired box protein Pax-3 (PAX3); Androgen receptor; Cyclin B1; v-myc avian myelocytomatosis viral oncogene neuroblastoma derived homolog (MYCN); Ras Homolog Family Member C (RhoC); Tyrosinase-related protein 2 (TRP-2); Cytochrome P450 1B1 (CYP1B1); CCCTC-Binding Factor (Zinc Finger Protein)-Like (BORIS or Brother of the Regulator of Imprinted Sites), Squamous Cell Carcinoma Antigen Recognized By T Cells 3 (SART3); Paired box protein Pax-5 (PAX5); proacrosin binding protein sp32 (OY-TES1); lymphocyte-specific protein tyrosine kinase (LCK); A kinase anchor protein 4 (AKAP-4); synovial sarcoma, X breakpoint 2 (SSX2); Receptor for Advanced Glycation Endproducts (RAGE-1); renal ubiquitous 1 (RU1); renal ubiquitous 2 (RU2);

legumain; human papilloma virus E6 (HPV E6); human papilloma virus E7 (HPV E7); intestinal carboxyl esterase; heat shock protein 70-2 mutated (mut hsp70-2); CD79a; CD79b; CD72; Leukocyte-associated immunoglobulin-like receptor 1 (LAIR1); Fc fragment of IgA receptor (FCAR or CD89); Leukocyte immunoglobulin-like receptor subfamily A member 2 (LILRA2); CD300 molecule-like family member f (CD300LF); C-type lectin domain family 12 member A (CLEC12A); bone marrow stromal cell antigen 2 (BST2); EGF-like module-containing mucin-like hormone receptor-like 2 (EMR2); lymphocyte antigen 75 (LY75); Glypican-3 (GPC3); Fc receptor-like 5 (FCRL5); and immunoglobulin lambda-like polypeptide 1 (IGLL1).

10 In one embodiment, the antigen binding domain comprises one, two three (e.g., all three) heavy chain CDRs, HC CDR1, HC CDR2 and HC CDR3, from an antibody listed above, and/or one, two, three (e.g., all three) light chain CDRs, LC CDR1, LC CDR2 and LC CDR3, from an antibody listed above. In one embodiment, the antigen binding domain comprises a heavy chain variable region and/or a variable light chain region of an antibody listed or
15 described above.

 In one aspect, the anti-tumor antigen binding domain is a fragment, e.g., a single chain variable fragment (scFv). In one aspect, the anti-a cancer associate antigen as described herein binding domain is a Fv, a Fab, a (Fab')₂, or a bi-functional (e.g. bi-specific) hybrid antibody (e.g., Lanzavecchia et al., Eur. J. Immunol. 17, 105 (1987)). In one aspect, the antibodies and
20 fragments thereof of the invention binds a cancer associate antigen as described herein protein with wild-type or enhanced affinity.

 In some instances, scFvs can be prepared according to a method known in the art (see, for example, Bird et al., (1988) Science 242:423-426 and Huston et al., (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883). ScFv molecules can be produced by linking VH and VL
25 regions together using flexible polypeptide linkers. The scFv molecules comprise a linker (e.g., a Ser-Gly linker) with an optimized length and/or amino acid composition. The linker length can greatly affect how the variable regions of a scFv fold and interact. In fact, if a short polypeptide linker is employed (e.g., between 5-10 amino acids) intrachain folding is prevented. Interchain folding is also required to bring the two variable regions together to form
30 a functional epitope binding site. For examples of linker orientation and size see, e.g., Hollinger et al. 1993 Proc Natl Acad. Sci. U.S.A. 90:6444-6448, U.S. Patent Application Publication

Nos. 2005/0100543, 2005/0175606, 2007/0014794, and PCT publication Nos. WO2006/020258 and WO2007/024715, which are incorporated herein by reference.

An scFv can comprise a linker of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more amino acid residues between its VL and VH regions. The linker sequence may comprise any naturally occurring amino acid. In some embodiments, the linker sequence comprises amino acids glycine and serine. In another embodiment, the linker sequence comprises sets of glycine and serine repeats such as (Gly₄Ser)_n, where n is a positive integer equal to or greater than 1 (SEQ ID NO:25). In one embodiment, the linker can be (Gly₄Ser)₄ (SEQ ID NO:27) or (Gly₄Ser)₃ (SEQ ID NO:28).
10 Variation in the linker length may retain or enhance activity, giving rise to superior efficacy in activity studies.

In another aspect, the antigen binding domain is a T cell receptor ("TCR"), or a fragment thereof, for example, a single chain TCR (scTCR). Methods to make such TCRs are known in the art. See, e.g., Willemsen RA et al, Gene Therapy 7: 1369–1377 (2000); Zhang T et al, Cancer Gene Ther 11: 487–496 (2004); Aggen et al, Gene Ther. 19(4):365-74 (2012)
15 (references are incorporated herein by its entirety). For example, scTCR can be engineered that contains the V α and V β genes from a T cell clone linked by a linker (e.g., a flexible peptide). This approach is very useful to cancer associated target that itself is intracellular, however, a fragment of such antigen (peptide) is presented on the surface of the cancer cells by MHC.

20

Transmembrane domain

With respect to the transmembrane domain, in various embodiments, a CAR can be designed to comprise a transmembrane domain that is attached to the extracellular domain of the CAR. A transmembrane domain can include one or more additional amino acids adjacent to
25 the transmembrane region, e.g., one or more amino acid associated with the extracellular region of the protein from which the transmembrane was derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 up to 15 amino acids of the extracellular region) and/or one or more additional amino acids associated with the intracellular region of the protein from which the transmembrane protein is derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 up to 15 amino acids of the intracellular region). In one
30 aspect, the transmembrane domain is one that is associated with one of the other domains of the

CAR. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins, e.g., to minimize interactions with other members of the receptor complex. In one aspect, the transmembrane domain is capable of homodimerization
 5 with another CAR on the cell surface of a CAR-expressing cell. In a different aspect, the amino acid sequence of the transmembrane domain may be modified or substituted so as to minimize interactions with the binding domains of the native binding partner present in the same CART.

The transmembrane domain may be derived either from a natural or from a recombinant
 10 source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein. In one aspect the transmembrane domain is capable of signalling to the intracellular domain(s) whenever the CAR has bound to a target. A transmembrane domain of particular use in this invention may include at least the transmembrane region(s) of e.g., the alpha, beta or zeta chain of the T-cell receptor, CD28, CD27, CD3 epsilon, CD45, CD4, CD5,
 15 CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154. In some embodiments, a transmembrane domain may include at least the transmembrane region(s) of, e.g., KIR2DS2, OX40, CD2, CD27, LFA-1 (CD11a, CD18), ICOS (CD278), 4-1BB (CD137), GITR, CD40, BAFFR, HVEM (LIGHTR), SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46, CD160, CD19, IL2R beta, IL2R gamma, IL7R α , ITGA1, VLA1, CD49a,
 20 ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d, ITGAE, CD103, ITGAL, CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, TNFR2, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRTAM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162),
 25 LTBR, PAG/Cbp, NKG2D, NKG2C, or CD19.

In some instances, the transmembrane domain can be attached to the extracellular region of the CAR, e.g., the antigen binding domain of the CAR, via a hinge, e.g., a hinge from a human protein. For example, in one embodiment, the hinge can be a human Ig (immunoglobulin) hinge, e.g., an IgG4 hinge, or a CD8a hinge. In one embodiment, the hinge
 30 or spacer comprises (e.g., consists of) the amino acid sequence of SEQ ID NO:2. In one aspect,

the transmembrane domain comprises (e.g., consists of) a transmembrane domain of SEQ ID NO: 6.

In one aspect, the hinge or spacer comprises an IgG4 hinge. For example, in one embodiment, the hinge or spacer comprises a hinge of the amino acid sequence

5 ESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDSQEDPEVQFNW
YVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEK
TISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK
TTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLGKM
(SEQ ID NO:36). In some embodiments, the hinge or spacer comprises a hinge encoded by a

10 nucleotide sequence of

GAGAGCAAGTACGGCCCTCCCTGCCCCCTTGCCCTGCCCCCGAGTTCTTGGGCGG
ACCCAGCGTGTTCCCTGTTCCCCCAAGCCCAAGGACACCTGATGATCAGCCGGA
CCCCGAGGTGACCTGTGTGGTGGTGGACGTGTCCCAGGAGGACCCCGAGGTCCA
GTTCAACTGGTACGTGGACGGCGTGAGGTGCACAACGCCAAGACCAAGCCCCGG
15 GAGGAGCAGTTCAATAGCACCTACCGGGTGGTGTCCGTGCTGACCGTGCTGCACCA
GGACTGGCTGAACGGCAAGGAATACAAGTGTAAGGTGTCCAACAAGGGCCTGCCC
AGCAGCATCGAGAAAACCATCAGCAAGGCCAAGGGCCAGCCTCGGGAGCCCCAGG
TGTACACCCTGCCCCCTAGCCAAGAGGAGATGACCAAGAACCAGGTGTCCCTGAC
CTGCCTGGTGAAGGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAAC
20 GGCCAGCCCGAGAACAACACTACAAGACCACCCCCCTGTGCTGGACAGCGACGGCA
GCTTCTTCCCTGTACAGCCGGCTGACCGTGGACAAGAGCCGGTGGCAGGAGGGCAA
CGTCTTTAGCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAGA
GCCTGAGCCTGTCCCTGGGCAAGATG (SEQ ID NO:37).

In one aspect, the hinge or spacer comprises an IgD hinge. For example, in one

embodiment, the hinge or spacer comprises a hinge of the amino acid sequence

RWPESPKAQASSVPTAQPQAEGSLAKATTAPATTRNTGRGGEEKKKEKEKEEQEERET
KTPECPSHTQPLGVYLLTPAVQDLWLRDKATFTCFVVGSDLKDAHLTWEVAGKVPTG
GVEEGLLERHSNGSQSQHSRLTLPRSLWNAGTSVTCTLNHPSLPPQRLMALREPAAQA
PVKLSLNLASSDPPEAASWLLCEVSGFSPPNILLMWLEDQREVNTSGFAPARPPPQPG
30 STTFWAWSVLRVPAPPSPQPATYTCVVSHEDSRTLLNASRSLEVS YVTDH (SEQ ID
NO:38). In some embodiments, the hinge or spacer comprises a hinge encoded by a nucleotide

sequence of

AGGTGGCCCGAAAGTCCCAAGGCCAGGCATCTAGTGTTCTACTGCACAGCCCCA
 GGCAGAAGGCAGCCTAGCCAAAGCTACTACTGCACCTGCCACTACGCGCAATACT
 GGCCGTGGCGGGGAGGAGAAGAAAAAGGAGAAAGAGAAAGAAGAACAGGAAGA
 5 GAGGGAGACCAAGACCCCTGAATGTCCATCCCATACCCAGCCGCTGGGCGTCTATC
 TCTTGACTCCCGCAGTACAGGACTTGTGGCTTAGAGATAAGGCCACCTTTACATGT
 TTCGTCGTGGGCTCTGACCTGAAGGATGCCCATTTGACTTGGGAGGTTGCCGGAAA
 GGTACCCACAGGGGGGGTTGAGGAAGGGTTGCTGGAGCGCCATTCCAATGGCTCT
 CAGAGCCAGCACTCAAGACTCACCTTCCGAGATCCCTGTGGAACGCCGGGACCTC
 10 TGTCACATGTACTCTAAATCATCCTAGCCTGCCCCACAGCGTCTGATGGCCCTTAG
 AGAGCCAGCCGCCAGGCACCAAGCTTAGCCTGAATCTGCTCGCCAGTAGTG
 ATCCCCCAGAGGCCGCCAGCTGGCTCTTATGCGAAGTGTCCGGCTTTAGCCCCCCC
 AACATCTTGCTCATGTGGCTGGAGGACCAGCGAGAAGTGAACACCAGCGGCTTCG
 CTCCAGCCCCGGCCCCACCCCAGCCGGGTTCTACCACATTCTGGGCCTGGAGTGTC
 15 TTAAGGGTCCCAGCACCACTAGCCCCCAGCCAGCCACATACACCTGTGTTGTGTC
 CCATGAAGATAGCAGGACCCTGCTAAATGCTTCTAGGAGTCTGGAGGTTTCCTACG
 TGA CTGACCATT (SEQ ID NO:103).

In one aspect, the transmembrane domain may be recombinant, in which case it will
 comprise predominantly hydrophobic residues such as leucine and valine. In one aspect a triplet
 20 of phenylalanine, tryptophan and valine can be found at each end of a recombinant
 transmembrane domain.

Optionally, a short oligo- or polypeptide linker, between 2 and 10 amino acids in length
 may form the linkage between the transmembrane domain and the cytoplasmic region of the
 CAR. A glycine-serine doublet provides a particularly suitable linker. For example, in one
 25 aspect, the linker comprises the amino acid sequence of GGGGSGGGGS (SEQ ID NO: 5). In
 some embodiments, the linker is encoded by a nucleotide sequence of
 GGTGGCGGAGGTTCTGGAGGTGGAGGTTCC (SEQ ID NO: 8).

In one aspect, the hinge or spacer comprises a KIR2DS2 hinge.

Cytoplasmic domain

The cytoplasmic domain or region of the CAR includes an intracellular signaling domain. An intracellular signaling domain is generally responsible for activation of at least one of the normal effector functions of the immune cell in which the CAR has been introduced.

5 Examples of intracellular signaling domains for use in a CAR described herein include the cytoplasmic sequences of the T cell receptor (TCR) and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any recombinant sequence that has the same functional capability.

10 It is known that signals generated through the TCR alone are insufficient for full activation of the T cell and that a secondary and/or costimulatory signal is also required. Thus, T cell activation can be said to be mediated by two distinct classes of cytoplasmic signaling sequences: those that initiate antigen-dependent primary activation through the TCR (primary intracellular signaling domains) and those that act in an antigen-independent manner to provide
15 a secondary or costimulatory signal (secondary cytoplasmic domain, e.g., a costimulatory domain).

 A primary signaling domain regulates primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary intracellular signaling domains that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor
20 tyrosine-based activation motifs or ITAMs.

 Examples of ITAM containing primary intracellular signaling domains that are of particular use in the invention include those of TCR zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, CD278 (also known as "ICOS"), FcεRI, DAP10, DAP12, and CD66d. In one embodiment, a CAR of the invention comprises an
25 intracellular signaling domain, e.g., a primary signaling domain of CD3-zeta, e.g., a CD3-zeta sequence described herein.

 In one embodiment, a primary signaling domain comprises a modified ITAM domain, e.g., a mutated ITAM domain which has altered (e.g., increased or decreased) activity as compared to the native ITAM domain. In one embodiment, a primary signaling domain
30 comprises a modified ITAM-containing primary intracellular signaling domain, e.g., an

optimized and/or truncated ITAM-containing primary intracellular signaling domain. In an embodiment, a primary signaling domain comprises one, two, three, four or more ITAM motifs.

Costimulatory Signaling Domain

5 The intracellular signalling domain of the CAR can comprise the CD3-zeta signaling domain by itself or it can be combined with any other desired intracellular signaling domain(s) useful in the context of a CAR of the invention. For example, the intracellular signaling domain of the CAR can comprise a CD3 zeta chain portion and a costimulatory signaling domain. The costimulatory signaling domain refers to a portion of the CAR comprising the intracellular
10 domain of a costimulatory molecule. In one embodiment, the intracellular domain is designed to comprise the signaling domain of CD3-zeta and the signaling domain of CD28. In one aspect, the intracellular domain is designed to comprise the signaling domain of CD3-zeta and the signaling domain of ICOS.

A costimulatory molecule can be a cell surface molecule other than an antigen receptor
15 or its ligands that is required for an efficient response of lymphocytes to an antigen. Examples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3, and a ligand that specifically binds with CD83, and the like. For example, CD27 costimulation has been demonstrated to enhance expansion, effector function, and survival of human CART cells
20 in vitro and augments human T cell persistence and antitumor activity in vivo (Song et al. Blood. 2012; 119(3):696-706). Further examples of such costimulatory molecules include CDS, ICAM-1, GITR, BAFFR, HVEM (LIGHTR), SLAMF7, NKp80 (KLRF1), NKp30, NKp44, NKp46, CD160, CD19, CD4, CD8alpha, CD8beta, IL2R beta, IL2R gamma, IL7R alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d,
25 ITGAE, CD103, ITGAL, CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRTAM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, NKG2D,
30 NKG2C and PAG/Cbp.

The intracellular signaling sequences within the cytoplasmic portion of the CAR may be linked to each other in a random or specified order. Optionally, a short oligo- or polypeptide linker, for example, between 2 and 10 amino acids (e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids) in length may form the linkage between intracellular signaling sequences. In one embodiment, a glycine-serine doublet can be used as a suitable linker. In one embodiment, a single amino acid, e.g., an alanine, a glycine, can be used as a suitable linker.

In one aspect, the intracellular signaling domain is designed to comprise two or more, e.g., 2, 3, 4, 5, or more, costimulatory signaling domains. In an embodiment, the two or more, e.g., 2, 3, 4, 5, or more, costimulatory signaling domains, are separated by a linker molecule, e.g., a linker molecule described herein. In one embodiment, the intracellular signaling domain comprises two costimulatory signaling domains. In some embodiments, the linker molecule is a glycine residue. In some embodiments, the linker is an alanine residue.

In one aspect, the intracellular signaling domain is designed to comprise the signaling domain of CD3-zeta and the signaling domain of CD28. In one aspect, the intracellular signaling domain is designed to comprise the signaling domain of CD3-zeta and the signaling domain of 4-1BB. In one aspect, the signaling domain of 4-1BB is a signaling domain of SEQ ID NO: 7. In one aspect, the signaling domain of CD3-zeta is a signaling domain of SEQ ID NO: 9.

In one aspect, the intracellular signaling domain is designed to comprise the signaling domain of CD3-zeta and the signaling domain of CD27. In one aspect, the signaling domain of CD27 comprises an amino acid sequence of QRRKYRSNKGESPVPAEPCRYSCPREEEGSTIPIQEDYRKPEPACSP (SEQ ID NO:16). In one aspect, the signalling domain of CD27 is encoded by a nucleic acid sequence of AGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCC GCCCGGGGCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCA GCCTATCGCTCC (SEQ ID NO:14).

In one aspect, the CAR-expressing cell described herein can further comprise a second CAR, e.g., a second CAR that includes a different antigen binding domain, e.g., to the same target or a different target (e.g., a target other than a cancer associated antigen described herein or a different cancer associated antigen described herein, e.g., CD19, CD33, CLL-1, CD34, FLT3, or folate receptor beta). In one embodiment, the second CAR includes an antigen

binding domain to a target expressed the same cancer cell type as the cancer associated antigen. In one embodiment, the CAR-expressing cell comprises a first CAR that targets a first antigen and includes an intracellular signaling domain having a costimulatory signaling domain but not a primary signaling domain, and a second CAR that targets a second, different, antigen and
5 includes an intracellular signaling domain having a primary signaling domain but not a costimulatory signaling domain. While not wishing to be bound by theory, placement of a costimulatory signaling domain, e.g., 4-1BB, CD28, ICOS, CD27 or OX-40, onto the first CAR, and the primary signaling domain, e.g., CD3 zeta, on the second CAR can limit the CAR activity to cells where both targets are expressed. In one embodiment, the CAR expressing
10 cell comprises a first cancer associated antigen CAR that includes an antigen binding domain that binds a target antigen described herein, a transmembrane domain and a costimulatory domain and a second CAR that targets a different target antigen (e.g., an antigen expressed on that same cancer cell type as the first target antigen) and includes an antigen binding domain, a transmembrane domain and a primary signaling domain. In another embodiment, the CAR
15 expressing cell comprises a first CAR that includes an antigen binding domain that binds a target antigen described herein, a transmembrane domain and a primary signaling domain and a second CAR that targets an antigen other than the first target antigen (e.g., an antigen expressed on the same cancer cell type as the first target antigen) and includes an antigen binding domain to the antigen, a transmembrane domain and a costimulatory signaling domain.

20 In another aspect, the disclosure features a population of CAR-expressing cells, e.g., CART cells. In some embodiments, the population of CAR-expressing cells comprises a mixture of cells expressing different CARs.

For example, in one embodiment, the population of CART cells can include a first cell expressing a CAR having an antigen binding domain to a cancer associated antigen described
25 herein, and a second cell expressing a CAR having a different antigen binding domain, e.g., an antigen binding domain to a different a cancer associated antigen described herein, e.g., an antigen binding domain to a cancer associated antigen described herein that differs from the cancer associate antigen bound by the antigen binding domain of the CAR expressed by the first cell.

30 As another example, the population of CAR-expressing cells can include a first cell expressing a CAR that includes an antigen binding domain to a cancer associated antigen

described herein, and a second cell expressing a CAR that includes an antigen binding domain to a target other than a cancer associate antigen as described herein. In one embodiment, the population of CAR-expressing cells includes, e.g., a first cell expressing a CAR that includes a primary intracellular signaling domain, and a second cell expressing a CAR that includes a secondary signaling domain.

In another aspect, the disclosure features a population of cells wherein at least one cell in the population expresses a CAR having an antigen binding domain to a cancer associated antigen described herein, and a second cell expressing another agent, e.g., an agent which enhances the activity of a CAR-expressing cell. For example, in one embodiment, the agent can be an agent which inhibits an inhibitory molecule. Inhibitory molecules, e.g., PD-1, can, in some embodiments, decrease the ability of a CAR-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD-1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (CEACAM-1, CEACAM-3, and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR (e.g., TGFRbeta). In one embodiment, the agent which inhibits an inhibitory molecule comprises a first polypeptide, e.g., an inhibitory molecule, associated with a second polypeptide that provides a positive signal to the cell, e.g., an intracellular signaling domain described herein. In one embodiment, the agent comprises a first polypeptide, e.g., of an inhibitory molecule such as PD-1, PD-L1, CTLA4, TIM3, CEACAM (CEACAM-1, CEACAM-3, and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and TGFR beta, or a fragment of any of these, and a second polypeptide which is an intracellular signaling domain described herein (e.g., comprising a costimulatory domain (e.g., 41BB, CD27, OX40 or CD28, e.g., as described herein) and/or a primary signaling domain (e.g., a CD3 zeta signaling domain described herein). In one embodiment, the agent comprises a first polypeptide of PD-1 or a fragment thereof, and a second polypeptide of an intracellular signaling domain described herein (e.g., a CD28 signaling domain described herein and/or a CD3 zeta signaling domain described herein).

The sequences of anti-CD19 binding domains are provided herein in Table 1. Full CAR constructs can be generated using any of the antigen binding domains described in Table 1 with one or more additional CAR component provided below.

- leader (amino acid sequence) (SEQ ID NO: 1)
MALPVTALLPLALLHAARP
- **leader (nucleic acid sequence) (SEQ ID NO: 12)**
ATGGCCCTGCCTGTGACAGCCCTGCTGCTGCCTCTGGCTCTGCTGCTGCATG
5 CCGCTAGACCC
- CD8 hinge (amino acid sequence) (SEQ ID NO: 2)
TTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACD
- **CD8 hinge (nucleic acid sequence) (SEQ ID NO: 13)**
ACCACGACGCCAGCGCCGCGACCACCAACACCGGCGCCCACCATCGCGTC
10 GCAGCCCCTGTCCCTGCGCCCAGAGGCGTGCCGGCCAGCGGCGGGGGGCGCAGTG
CACACGAGGGGGCTGGACTTCGCCTGTGAT
- CD8 transmembrane (amino acid sequence) (SEQ ID NO: 6)
IYIWAPLAGTCGVLLSLVITLYC
- 15 • **transmembrane (nucleic acid sequence) (SEQ ID NO: 17)**
ATCTACATCTGGGCGCCCTTGGCCGGGACTTGTGGGGTCCTTCTCCTGTCAC
TGGTTATCACCCCTTTACTGC
- 4-1BB Intracellular domain (amino acid sequence) (SEQ ID NO: 7)
20 KRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL
- **4-1BB Intracellular domain (nucleic acid sequence) (SEQ ID NO: 18)**
AAACGGGGCAGAAAGAACTCCTGTATATATTCAAACAACCATTTATGAG
ACCAGTACAACTACTCAAGAGGAAGATGGCTGTAGCTGCCGATTTCCAGAAGAA
GAAGAAGGAGGATGTGAACTG
25
- CD3 zeta domain (amino acid sequence) (SEQ ID NO: 9)
RVKFSRSADAPAYKQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRR
KNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDLGLYQGLSTATKDTYDALHM
QALPPR
- 30 • **CD3 zeta (nucleic acid sequence) (SEQ ID NO: 20)**

AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGCGTACAAGCAGGGCCA
 GAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTG
 GACAAGAGACGTGGCCGGGACCCTGAGATGGGGGGAAAGCCGAGAAGGAAGAAC
 CCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCTACA
 5 GTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTA
 CCAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCC
 TGCCCCCTCGC

- **CD3 zeta domain (amino acid sequence; NCBI Reference Sequence NM_000734.3) (SEQ ID NO:10)**

RVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRR
 KNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHM
 QALPPR

- **CD3 zeta (nucleic acid sequence; NCBI Reference Sequence NM_000734.3); (SEQ ID NO:21)**

agagtgaagttcagcaggagcgagacgccccgcgtaccagcagggccagaaccagctctataacgagctcaatctag
 gacgaagagaggagtacgatgtttggacaagagacgtggccgggaccctgagatgggggaaagccgagaaggaagaaccctca
 ggaaggcctgtacaatgaactgcagaaagataagatggcggaggcctacagtgagattgggatgaaaggcgagcgccggaggggca
 aggggcacgatggcctttaccagggtctcagtacagccaccaaggacacctacgagcccttcacatgcaggccctgccccctgcg

20

IgG4 Hinge (amino acid sequence) (SEQ ID NO:36)

ESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVSQEDPEV
 QFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLP
 SSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE
 25 NNYKTTTPVLDSGDFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLG
 KM

IgG4 Hinge (nucleotide sequence) (SEQ ID NO:37)

GAGAGCAAGTACGGCCCTCCCTGCCCCCTTGCCCTGCCCCCGAGTTCCTG
 30 GGCGGACCCAGCGTGTTCTGTTCCTGTTCCCCCAAGCCCAAGGACACCCTGATGATCAG
 CCGGACCCCCGAGGTGACCTGTGTGGTGGTGGACGTGTCCAGGAGGACCCCGAG

GTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAGACCAAGC
CCCGGGAGGAGCAGTTCAATAGCACCTACCGGGTGGTGTCCGTGCTGACCGTGCTG
CACCAGGACTGGCTGAACGGCAAGGAATACAAGTGTAAGGTGTCCAACAAGGGCC
TGCCCAGCAGCATCGAGAAAACCATCAGCAAGGCCAAGGGCCAGCCTCGGGAGCC
5 CCAGGTGTACACCCTGCCCCCTAGCCAAGAGGAGATGACCAAGAACCAGGTGTCC
CTGACCTGCCTGGTGAAGGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGA
GCAACGGCCAGCCCGAGAACAACCTACAAGACCACCCCCCTGTGCTGGACAGCGA
CGGCAGCTTCTTCCTGTACAGCCGGCTGACCGTGGACAAGAGCCGGTGGCAGGAG
GGCAACGTCTTTAGCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCA
10 GAAGAGCCTGAGCCTGTCCCTGGGCAAGATG

EF1 alpha promoter

CGTGAGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCACAGTCC
CCGAGAAGTTGGGGGGAGGGGTCCGCAATTGAACCGGTGCCTAGAGAAGGTGGCG
15 CGGGGTAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCTTTTTCCCGAGGGTG
GGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTTCGCAACGGG
TTTGCCGCCAGAACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTT
TACGGGTTATGGCCCTTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGAT
TCTTGATCCCGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCGCTT
20 AAGGAGCCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGC
CGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTCTCTA
GCCATTTAAAATTTTTGATGACCTGCTGCGACGCTTTTTTTCTGGCAAGATAGTCTT
GTAAATGCGGGCCAAGATCTGCACACTGGTATTTTCGGTTTTTTGGGGCCGCGGGCGG
CGACGGGGCCCGTGCGTCCCAGCGCACATGTTTCGGCGAGGCGGGGCCTGCGAGCG
25 CGGCCACCGAGAATCGGACGGGGGTAGTCTCAAGCTGGCCGGCCTGCTCTGGTGC
CTGGCCTCGCGCCGCCGTGTATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCCGGTCG
GCACCAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCT
CAAAATGGAGGACGCGGCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAA
GGAAAAGGGCCTTTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGG
30 GCGCCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTCTTTAGGT
TGGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTG
AAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCTTTTTGAGT

TTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTTTTTTCTTCCAT
TTCAGGTGTCGTGA (SEQ ID NO: 11).

Gly/Ser (SEQ ID NO:25)

5 GGGGS

Gly/Ser (SEQ ID NO:26): This sequence may encompass 1-6 "Gly Gly Gly Gly Ser" repeating units

GGGGS GGGGS GGGGS GGGGS GGGGS GGGGS

10 Gly/Ser (SEQ ID NO:27)

GGGGS GGGGS GGGGS GGGGS

Gly/Ser (SEQ ID NO:28)

GGGGS GGGGS GGGGS

15

Gly/Ser (SEQ ID NO:29)

GGGS

PolyA (SEQ ID NO:30): A5000

20 PolyA (SEQ ID NO:31): A100

PolyT (SEQ ID NO:32): T5000

PolyA (SEQ ID NO:33): A5000

PolyA (SEQ ID NO:34): A400

PolyA (SEQ ID NO:35): A2000

25 • Gly/Ser (SEQ ID NO:15): This sequence may encompass 1-10 "Gly Gly Gly Ser" repeating units

GGGSGGGSGG GSGGGSGGGG GGGSGGGSGG GSGGGSGGGG

Exemplary CD19 CAR constructs that can be used in the methods described herein are shown in **Table 3**:

Table 3: CD19 CAR Constructs

Name	SEQ ID	Sequence
CAR 1		
CAR1 scFv domain	39	EIVMTQSPATLSLSPGERATLSCRASQDISKYLNWYQQKPGQAPRLLIYHT SRLHSGIPARFSGSGSGTDYTLTISSLPEDFAVYFCQQGNTLPYTFGQGT KLEIKGGGSGGGGSGGGGSGVQLQESGPGLVKPSSETLSLTCTVSGVSLPD YGVSWIRQPPGKGLEWIGVIWGSETTYSSSLKSRVTISKDNSKNQVSLKL SSVTAADTAVYYCAKHYYYGGSYAMDYWGQGT LVTVSS
103101 CAR1 Soluble scFv - nt	52	atggccctccctgtcaccgcctgctgcttcgctggctcttctgctccacgcgc tcggcccgaaattgtgatgaccagtcaccgcctcttagcctttcaccgggtg agcgcgcaacctgtcttgagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggtcctcgcttctgatctaccacaccagcgggt ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacacc tcactatcagctcactgcagccagaggacttcgctgtctatttctgtcagcaagg aacaccctgccctacacctttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaa gcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagc ggagtgtctctcccgattacggggtgtcttggatcagacagccaccgggaagg tctggaatggattggagtgatttggggtctgagactacttactactcttcatccc tcaagtacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacaccgcctgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacaggtactctggtcacctgt ccagccaccaccatcatcaccatcaccat
103101 CAR1 Soluble scFv - aa	64	<u>MALPVTALLPLALLHAARP</u> eivmtqspatls slspgeratls crasqdiskylnw yqqkpgqaprlliyhtsrlhsgiparfsgsgsgtdytltiisslpedfavyfcqqg ntlpytfgggtkleikgggsgggsgggsgvqlqesgpglvkpssetls lctvs gvslpdygvswirppgkglewigviwgsettyssslksrvtiskdnsknqvslk lssvtaadtavyycahyyyggsyamydywggtlvtvss <u>hhhhhhhh</u>
104875 CAR 1 – Full - nt	90	atggccctccctgtcaccgcctgctgcttcgctggctcttctgctccacgcgc tcggcccgaaattgtgatgaccagtcaccgcctcttagcctttcaccgggtg agcgcgcaacctgtcttgagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggtcctcgcttctgatctaccacaccagcgggt ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacacc tcactatcagctcactgcagccagaggacttcgctgtctatttctgtcagcaagg aacaccctgccctacacctttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaa gcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagc

		ggagtgtctctccccgattacggggtgtcttggatcagacagccaccgggaaggg tctggaatggattggagtgatttggggctctgagactacttactactcttcatccc tcaagtcacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacacccgcgtgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacaggggtactctggtcaccgtgt ccagcaccactaccccagcaccgagggccaccaccccggctcctaccatcgctcc cagcctctgtccctgcgtccggagggcatgtagaccgcagctgggtggggcgtgca taccgggggtcttgacttcgctgcgatatctacatttggggccctctggctggta cttgcggggtcctgctgctttcactcgtgatcactctttactgtaagcgcggtcgg aagaagctgctgtacatctttaagcaacccttcagaggcctgtgcagactactca agaggaggacggctgttcatgccggttcccagaggaggaggaaggcggtgcgaac tgcgcgtaaaattcagccgcagcgcagatgctccagcctacaagcaggggcagaac cagctctacaacgaactcaatcttggtcggagagaggagtacgacgtgctggaciaa gaggagaggacgggacccagaaatggcggggaagccgcgcagaaagaatccccaa agggcctgtacaacgagctccaaaaggataagatggcagaagcctatagcgagatt ggatgaaaggggaacgcagaagaggcaaaggccacgacggactgtaccagggact cagcaccgccaccaaggacacctatgacgctcttcacatgcaggccctgccgctc gg
104875 CAR 1 – Full - aa	77	MALPVTALLPLALLHAARPeivmtqspatlsispgeratlsc <u>rasqdiskylnw</u> yqqkpgqaprllliy <u>htsrllhs</u> giparfsgsgsgtdytlitisslqpedfavyfc <u>ggg</u> <u>ntlpyt</u> fgqgtkleikgggsgggsgggsgvqlqesgpglvkpssetlsltctvs gvslp <u>dygvs</u> wirppgkglewig <u>viwgsettyyssslks</u> rvtiskdnknqvsllk lssvtaadtavyycak <u>hyyyggsyamyd</u> wgggtlvtvsssttpaprpptpaptias qplsrlrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgr kkllyifkqpfmrpvqttgeedgcscrfeeeeeggcelrvkfsrsadapaykqqn qlynelnlgrreeydvldkrrgrdpemggkprknpqeglynelqdkmaeysei gmkgerrrgkghdglyqglstatkdydalhmqalppr
CAR 2		
CAR2 scFv domain	40	eivmtqspatlsispgeratlscrasqdiskylnwyqqkpgqaprllliyhtsrllhs giparfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfgqgtkleikgggsg gggsgggsgvqlqesgpglvkpssetlsltctvsgvslpdygvswirppgkgle wigviwgsettyyqssslksrvtiskdnknqvsllkssvtaadtavyycakhyyyg gsyamydwwgggtlvtvss
103102 CAR2 - Soluble scFv - nt	53	atggccctccctgtcaccgcctgtgtcttccgtggctcttctgctccacgcgc tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaaccctgtcttgcagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggtcctcgcttctgatctaccacaccagccggct ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacaccc

		<p>tcactatcagctcactgcagccagaggacttcgctgtctatcttctgtcagcaaggg aacaccctgccctacacctttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaaa gcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagc ggagtgtctctccccgattacggggtgtcttggatcagacagccaccgggaaggg tctggaatggattggagtgatttggggctctgagactacttactaccaatcatccc tcaagtacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacaccgccgtgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacaggggtactctggtcacctgt ccagccaccaccatcatcaccatcaccat</p>
103102 CAR2 - Soluble scFv - aa	65	<p><u>MALPVTALLLPLALLHAARP</u>eivmtqspatlsispgeratlscrasqdiskylnw yqqkpgqaprllyhtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqg ntlpytfgggtkleikgggsgggsgggsgvqlqesgpglvkpsetlsltctvs gvslpdygvswirppgkglewigviwgsettyyqsslksrvtiskdnknqvsllk lssvtaadtavyycakhyyyggsyamdywggtlvtvss<u>hhhhhhh</u></p>
104876 CAR 2 - Full - nt	91	<p>atggccctccctgtcaccgccttgccttccgctggctcttctgctccacgcgcg tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcacccgggtg agcgcgcaaccctgtcttgcagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggtcctcgccttctgatctaccacaccagccggct ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacaccc tcactatcagctcactgcagccagaggacttcgctgtctatcttctgtcagcaaggg aacaccctgccctacacctttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaaa gcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagc ggagtgtctctccccgattacggggtgtcttggatcagacagccaccgggaaggg tctggaatggattggagtgatttggggctctgagactacttactaccaatcatccc tcaagtacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacaccgccgtgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacaggggtactctggtcacctgt ccagcaccactacccagcaccgagggccacccaccccggtcctaccatcgctcc cagcctctgtccctgcgtccggaggtcatgtagaccgcagctggtggggcgtgca taccgggggtcttgacttcgctgcgatatctacatttgggcccctctggtggta cttgcgggggtcctgctgctttcactcgtgatcactctttactgtaagcgcggtcgg aagaagctgctgtacatctttaagcaacccttcatgaggcctgtgcagactactca agaggaggacggctgttcatgccgggttcccagaggaggaggaaggcggctgcgaac tgcgctgaaattcagcgcagcgcagatgctccagcctacaagcaggggcagaac cagctctacaacgaactcaatcttggtcggagagaggagtacgacgtgctggaaa gcggagaggacgggaccagaaatggcggggaagccgcgcagaaagaatccccaa</p>

		agggcctgtacaacgagctccaaaaggataagatggcagaagcctatagcgagatt ggtatgaaaggggaacgcagaagaggcaaaggccacgacggactgtaccagggact cagcaccgccaccaaggacacctatgacgctcttcacatgcaggccctgccgctc gg
104876 CAR 2 - Full - aa	78	MALPVTALLLPLALLLHAARPeivmtqspatlsispgeratlsc <u>rasqdiskyl</u> nw yqqkpgqaprlly <u>htsr</u> lhsgiparfsgsgsgtdytlitisslqpedfavyfc <u>ggg</u> <u>ntlpyt</u> fgggtkleikggggsgggsgggsgvqlqesgpglvkpssetlsltctvs gvslp <u>dygvs</u> wirppgkglewig <u>viwgsettyyqsslks</u> rvtiskdsknqvslk lssvtaadtavyycak <u>hyyyggsyamdy</u> wgqgtlvtvsssttppaprpptpaptias qplslrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgr klllyifkqpfmrvpqtteedgcscrfeeeeeggcclrvkfsrsadapaykqqn qlynelnlgrreeydvldkrrgrdpemggkprknpqeglynelqdkmaeysei gmkgerrrgkghdglyqglstatkdydalhmqalppr
CAR 3		
CAR3 scFv domain	41	qvqlqesgpglvkpssetlsltctvsgvslpdygvswirppgkglewigviwgset tyyssslksrvtiskdsknqvslklssvtaadtavyycakhyyyggsyamdywgq gtlvtvssggsgggsgggsgggseivmtqspatlsispgeratlscrasqdiskyl nwyqqkpgqaprllyhtsr lhsg iparfsgsgsgtdytlitisslqpedfavyfcq qgntlpytfgggtkleik
103104 CAR 3 - Soluble scFv - nt	54	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacgcttcaagaatcagggcctggctctggtgaagccatctgaga ctctgtccctcacttgaccgtgagcggagtgtccctcccagactacggagtgagc tggattagacagcctcccggaaaggactggagtggatcgagagtgatttggggtag cgaaaccacttactattcatcttcctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgccgctgacaccgcc gtgtattactgtgccaagcattactactatggagggtcctacgccatggactactg gggccagggaactctggtcactgtgtcatctggtggaggaggtagcggaggaggcg ggagcgggtggaggtggctccgaaatcgtgatgaccagagccctgcaaccctgtcc ctttctcccggggaacgggctaccctttcttgtcgggcatcacaagatatctcaaa atacctcaattggtatcaacagaagccgggacaggcccctaggcttcttatctacc acacctctgcctgcatagcgggattcccgcacgcttttagcgggtctggaagcggg accgactacactctgaccatctcatctctccagccgaggacttcgccgtctactt ctgccagcagggtaacaccctgccgtacaccttcggccagggcaccaagcttgaga tcaaacatcaccaccatcatcaccatcac
103104 CAR 3 - Soluble scFv - aa	66	<u>MALPVTALLLPLALLLHAARP</u> qvqlqesgpglvkpssetlsltctvsgvslpdygvs wirppgkglewigviwgsettyyssslksrvtiskdsknqvslklssvtaadta vyycakhyyyggsyamdywgqgtlvtvssggsgggsgggsgggseivmtqspatls ispgeratlscrasqdiskylnwyqqkpgqaprllyhtsr lhsg iparfsgsgsg

		tdytlTisslqpedfavyfcqqgntlpytfgqgkkleik <u>hhhhhhhh</u>
104877 CAR 3 – Full - nt	92	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgcgc tcgcccacaagtcacagcttcaagaatcagggcctggctctggtgaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgtccctccagactacggagtgagc tggattagacagcctcccggaaaggactggagtggatcgagagtatttgggtag cgaaaccacttactattcatcttccctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgcgcgtgacaccgcc gtgtattactgtgccaaagcattactactatggagggtcctacgccatggactactg gggcccaggaactctggctactgtgtcatctggaggaggtagcggaggaggcg ggagcgggtggaggtggctccgaaatcgtgatgaccagagccctgcaaccctgtcc ctttctcccggggaacgggctaccctttctgtcgggcatcacaagatatctaaa atacctcaattggatatcaacagaagccgggacagggccctaggtcttctatctacc acacctctcgctgcatagcgggattcccgcacgctttagcgggtctggaagcggg accgactacactctgaccatctcatctctccagccgaggacttcgccgtctactt ctgccagcagggtaacaccctgccgtacaccttcggccagggcaccaagcttgaga tcaaaaccactactccgctccaaggccaccacccctgcccagaccatcgctct cagccgctttccctgcgtccggaggcatgtagaccgcagctggtggggcctgca taccggggtcttgacttcgctgcgatatctacatttggggccctctggttgga cttgcggggtcctgctgctttcactcgtgatcactctttactgtaagcgggctcg aagaagctgctgtacatctttaagcaacccttcatgaggcctgtgcagactactca agaggaggacggctgttcatgccggttcccagaggaggaggaaggcggtgcgaac tgcgctgaaattcagccgcagcgcagatgctccagcctacaagcaggggcagaac cagctctacaacgaactcaatcttggtcggagagaggagtacgacgtgctggaaa gcggagaggacgggaccagaaatggcggggaagccgcgcagaaagaatccccag aggcctgtacaacgagctccaaaaggataagatggcagaagcctatagcgagatt ggtatgaaaggggaacgcagaagaggcaaaggccacgagactgtaccagggact cagcaccgccaccaaggacacctatgacgctcttcacatgcaggccctgccgcctc gg
104877 CAR 3 – Full - aa	79	MALPVTALLLPLALLLHAARPqvqlqesgpglvkpsetlsltctvsgvslp <u>dygvs</u> wirqppgkglewig <u>viwgsettyysslks</u> rvtiskdnknqvsllkssvtaadta vyycak <u>hyyyggsyamyd</u> wgggtlvtvssggggsgggsggggseivmtqspatls lspgeratlsc <u>rasqdiskyl</u> nwyqqkpgqaprlliy <u>htsrlhs</u> giparfsgsgsg tdytlTisslqpedfavyfc <u>qqgntlpyt</u> fgqgkkleiktttpaprpptpaptias qplslrpeacrpaaggavhtrgldfacdiyiwaplagtcgvlllslvitlyckrgr kklllyifkqpfmrpvqttqeedgcscrffpeeeeggcelrvkfsrsadapaykqqgn qlynelnlgrreeydvldkrrgrdpemggkprknpqeglynelqkdmaeysei gmkgerrrgkghdglyqglstatkdydalhmqalppr
CAR 4		

CAR4 scFv domain	42	qvqlqesgpgglvlpksetlsltctvsgvslpdygvswirppgkglewigviwgset tyyqsslsksrvtiskdnsknqvslklssvtaadtavyycahyyyggsyamdywgq gtlvtvssggsgggsgggsgggseivmtqspatlsispgeratlscrasqdiskyl nwyqqkpgqaprllyhtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcq qgntlpytfgggtkleik
103106 CAR4 – Soluble scFv - nt	55	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacagcttcaagaatcagggcctggctctggtgaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgtccctcccagactacggagtgagc tggttagacagcctcccgaaagggactggagtggatcgagtgatttggggtag cgaaaccacttactatcaatcttccctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgccgctgacaccgcc gtgtattactgtgccaagcattactactatggagggtcctacgccatggactactg gggccaggaactctggtcactgtgtcatctggtggaggaggtagcggaggaggcg ggagcgggtggaggtggctccgaaatcgtgatgaccagagccctgcaaccctgtcc ctttctcccggggaacgggctaccctttcttgtcgggcatcacaagatatctcaa atacctcaattggtatcaacagaagccgggacaggccccctaggcttcttatctacc acacctctcgctgcatagcgggattcccgacgcttttagcgggtctggaagcggg accgactacactctgaccatctcatctctccagccgaggacttcgccgtctactt ctgccagcagggtaacaccctgccgtacaccttcggccagggcaccaagcttgaga tcaaacatcaccaccatcatcaccatcac
103106 CAR4 – Soluble scFv -aa	67	<u>MALPVTALLPLALLHAARP</u> qvqlqesgpgglvlpksetlsltctvsgvslpdygvs wirppgkglewigviwgsettyyqsslsksrvtiskdnsknqvslklssvtaadta vyycakhyyyggsyamdywgqgtlvtvssggsgggsgggsgggseivmtqspatls lspgeratlscrasqdiskylnwyqqkpgqaprllyhtsrhsgiparfsgsgsg tdytlitisslqpedfavyfcqgntlpytfgggtkleik <u>hhhhhhh</u>
104878 CAR 4 – Full - nt	93	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacagcttcaagaatcagggcctggctctggtgaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgtccctcccagactacggagtgagc tggttagacagcctcccgaaagggactggagtggatcgagtgatttggggtag cgaaaccacttactatcaatcttccctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgccgctgacaccgcc gtgtattactgtgccaagcattactactatggagggtcctacgccatggactactg gggccaggaactctggtcactgtgtcatctggtggaggaggtagcggaggaggcg ggagcgggtggaggtggctccgaaatcgtgatgaccagagccctgcaaccctgtcc ctttctcccggggaacgggctaccctttcttgtcgggcatcacaagatatctcaa atacctcaattggtatcaacagaagccgggacaggccccctaggcttcttatctacc acacctctcgctgcatagcgggattcccgacgcttttagcgggtctggaagcggg accgactacactctgaccatctcatctctccagccgaggacttcgccgtctactt

		ctgccagcagggtaaacacctgccgtacaccttcggccagggcaccaagcttgaga tcaaaaccactactcccgtccaaggccaccacccctgccccgaccatcgectct cagccgctttccctgcgtccggagggcatgtagaccgcagctgggtggggccgtgca taccgggggtcttgacttcgcctgcgatatctacatttgggcccctctggctggta cttgcggggtcctgctgctttcactcgtgatcactctttactgtaagcgcggtcgg aagaagctgctgtacatctttaagcaacccttcatgaggcctgtgcagactactca agaggaggacggctgttcatgccggttcccagaggaggaggaaggcggtgcgaac tgcgcggtgaaattcagccgcagcgcagatgctccagcctacaagcaggggcagaac cagctctacaacgaactcaatcttggtcggagagaggagtacgacgtgctggacaa gcggagaggacgggaccagaaatgggcgggaagccgcgcagaaagaatccccag agggcctgtacaacgagctccaaaaggataagatggcagaagcctatagcgagatt ggatatgaaaggggaacgcagaagaggcaaaggccacgacggaactgtaccagggact cagcaccgccaccaaggacacctatgacgctcttcacatgcaggccctgccgcctc gg
104878 CAR 4 – Full - aa	80	MALPVTALLLPLALLLHAARPvqlqesgpglvkpsetlsltctvsgvslp <u>dygvs</u> wirqppgkglewig <u>viwgsettyyqsslks</u> rvtiskdnsknqvsllssvtaadta vyycak <u>hyyyggsyamdy</u> wgggtlvtvssggggsgggsggggseivmtqspatls lspgeratlsc <u>rasqdiskyl</u> nwyqqkpgqaprlliy <u>htsrllhs</u> giparfsgsgsg tdytltisslqpedfavyfc <u>qggntlpyt</u> fgqgtkleiktttpprptpaptias qplslrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgr klllyifkqpfmrpvqttqeedgcscrfeeeeeggcclrvkfsrsadapaykqqn qlynelnlgrreeydvldkrrgrdpemggkprknpqeglynelqkdmaeysei gmkgerrrgkghdglyqglstatkdydalhmqalppr
CAR 5		
CAR5 scFv domain	43	eivmtqspatls lspgeratlscrasqdiskyl nwyqqkpgqaprlliyhtsrllhs giparfsgsgsgtdytltisslqpedfavyfcqggntlpytfgqgtkleikggggs ggggsgggsgggsgvqlqesgpglvkpsetlsltctvsgvslpdygvswirqp gkglewigviwgsettyyssslksrvtiskdnsknqvsllssvtaadtavyycak hyyyggsyamdywgggtlvtvss
99789 CAR5 - Soluble scFv - nt	56	atggccctcccagtgaccgctctgctgctgcctctcgacttcttctccatgccgc tcggcctgagatcgatcatgacccaaagccccgtacctgtccctgtcaccggcg agagggaacccttcatgcagggcagccaggacatttctaagtacctcaactgg tatcagcagaagccagggcaggtcctcgctgctgatctaccacaccagcgcct ccacagcggtatccccgccagatttccgggagcgggtctggaaccgactacacc tcaccatctcttctctgcagcccgaggatttccgcgtctatttctgccagcgggg aatactctgccgtacaccttcgggtcaaggtaccaagctggaaatcaagggaggcgg aggatcaggcgggtggcggaagcggaggaggtggctccggaggaggaggttcccaag tgcagcttcaagaatcaggaccggacttgtgaagccatcagaaccctctccctg

		acttgtaaccgtgtccggtgtgagcctccccgactacggagtctcttggattcgcca gcctccgggaagggctcttgaatggattgggtgatttggggatcagagactactt actactcttcatcacttaagtcacgggtcaccatcagcaaagataatagcaagaac caagtgtcacttaagctgtcatctgtgaccgcccgtgacaccgccgtgtactattg tgccaaacattactattacggagggctcttatgctatggactactggggacagggga ccctgggtgactgtctctagccatcaccatcaccaccatcatcac
99789 CAR5 - Soluble scFv -aa	68	<u>MALPVTALLLPLALLHAARP</u> eivmtqspatlsispgeratlscrasqdiskylnw yqqkpgqaprlliyhtsrhsgiparfsgsgsgtdytlitisslqpedfavyfcqqg ntlpytfgggtkleikgggsgggsgggsgggsgggsgvqlqesgpglvkpsetlsl tctvsgvslpdygvswirppgkglewigviwgsettyysssllksrvtiskdnkn qvsllkssvtaadtavyycahyyyggsyamdyygggtlvtvss hhhhhhhh
104879 CAR 5 – Full - nt	94	atggccctccctgtcaccgcctgtctgcttcgctggctcttctgctccacgcgc tcggcccgaattgtgatgaccagtcacccgccactcttagcctttcacccgggtg agcgcgcaaccctgtcttgcagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggctcctcgcttctgactctaccacaccagccggct ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacacc tcactatcagctcactgcagccagaggacttcgctgtctatcttctgtcagcaagg aacaccctgccctacaccttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagcggcggaggcgggagccagg tccaaactccaagaaagcggaccgggtcttgtgaagccatcagaaactcttctactg acttgtaactgtgagcggagtgtctctccccgattacgggggtgtcttggatcagaca gccaccggggaagggctctggaatggattggagtgatttggggctctgagactactt actactcttcatccctcaagtcacgcgtcaccatctcaaaggacaactctaagaat cagggtgtcactgaaactgtcatctgtgaccgcagccgacaccgccgtgtactattg cgctaagcattactattatggcgggagctacgcaatggattactggggacagggta ctctggtcaccgtgtccagcaccactacccagcaccgagggccacccaccccggt cctaccatcgctcccagcctctgtccctgcgtccggaggcatgtagaccgcagc tgggtggggcgtgcatacccggggtcttgacttcgcctgcgatatctacatttggg ccctctggctggtacttgccgggtcctgctgcttctactcgtgatcactctttac tgtaaagcgggtcggaagaagctgctgtacatctttaagcaaccctcatgaggcc tgtgcagactactcaagaggaggacggctgttcatgccggttccagaggaggagg aaggcgggtgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac aagcaggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaatggcggggaagccgcgca gaaagaatccccaaagaggccctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaggccacgcagg actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgcctcgg

104879 CAR 5 – Full - aa	81	MALPVTALLLPLALLLHAARPeivmtqspatlsispgeratlsc <u>rasqdiskylnw</u> yqqkpgqaprlly <u>htsrllhs</u> giparfsgsgsgtdytlitisslqpedfavyfc <u>ggg</u> <u>ntlpyt</u> fgggtkleikgggsgggsgggsgggsgggsgvqlqesgpglvkpsetls1 tctvsgvslp <u>dygvs</u> swirppgkglewig <u>viwgsettyyssslks</u> rvtiskdnsk qvs1klssvtaadtavyyca <u>hyyyggsyamdy</u> wgggtlvtvsssttpaprpptpa ptiasqplslrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitly ckrgrklllyifkqpfmrpvqttqeedgcscrfeeeeeggcclrvkfsrsadapay kqqgnqlynelnlgrreeydvldkrrgrdpemggkprkrnpqeglynelqkdkmae ayseigmkgerrrgkghdglyqglstatkdydalhmqalppr
CAR 6		
CAR6 scFv domain	44	eivmtqspatlsispgeratlscrasqdiskylnwyqqkpgqaprllyhtsrllhs giparfsgsgsgtdytlitisslqpedfavyfcqqgntlpytfgggtkleikgggsg gggsgggsgggsgvqlqesgpglvkpsetls1tctvsgvslpdygvswwirpp gkglewigviwgsettyyqss1ksrvtiskdnskqvs1klssvtaadtavyyca hyyyggsyamdywgggtlvtvss
99790 CAR6 - Soluble scFv - nt	57	atggccctcccagtgaccgctctgctgctgcctctcgacttcttctccatgccgc tcggcctgagatcgatgacccaaagccccgctaccctgtccctgtcaccggcg agagggcaaccctttcatgcagggccagccaggacatttctaagtacctcaactgg tatcagcagaagccaggcaggctcctcgctgctgatctaccacaccagccgcct ccacagcggatccccgccagattttccgggagcgggtctggaaccgactacaccc tcaccatctcttctctgcagcccgaggatttcgcctgtctatttctgccagcgggg aatactctgccgtacaccttcgggtcaaggtaccaagctggaaatcaagggaggcgg aggatcaggcgggtggcggaagcggaggaggtggctccggaggaggaggttcccaag tgacgcttcaagaatcaggaccggacttgtgaagccatcagaaaccctctccctg acttgtaccgtgtccgggtgtgagcctccccgactacggaggtctcttggaatcgcca gcctccggggaagggtcttgaatggattgggggtgatttggggatcagagactactt actaccagtcacttaagtcacgggtcaccatcagcaaagataatagcaagaac caagtgtcacttaagctgtcatctgtgaccgccgtgacaccgccgtgtactattg tgccaaacattactattacggagggtcttatgctatggactactggggacagggga ccctggtgactgtctctagccatcaccatcaccaccatcatcac
99790 CAR6 - Soluble scFv - aa	69	<u>MALPVTALLLPLALLLHAARP</u> eivmtqspatlsispgeratlscrasqdiskylnw yqqkpgqaprllyhtsrllhsigiparfsgsgsgtdytlitisslqpedfavyfcqqg ntlpytfgggtkleikgggsgggsgggsgggsgggsgvqlqesgpglvkpsetls1 tctvsgvslpdygvswwirppgkglewigviwgsettyyqss1ksrvtiskdnsk qvs1klssvtaadtavyyca <u>hyyyggsyamdy</u> wgggtlvtvss <u>hhhhhhhh</u>
104880 CAR6 – Full - nt	95	atggccctccctgtcaccgcctctgctgcttccgctggctcttctgctccacgccgc tcggcccgaatgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaaccctgtcttgagagcctcccaagacatctcaaaataccttaattgg

		tatcaacagaagcccgacaggctcctcgcttctgatctaccacaccagccggtccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacacctactatcagctcactgcagccagaggacttcgctgtctatcttctgtcagaagggaacacctgccctacacctttggacagggcaccaagctcgagattaaagggtggaggtggcagcggaggaggtgggtccggcgggtggaggaagcggaggcggaggagccaggccaactccaagaaagcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagcggagtgtctctccccgattacggggtgtcttggatcagacagccacggggaagggtctggaatggattggagtgatttggggtctgagactacttactaccaatcatccctcaagtcacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaactgtcatctgtgaccgcagccgacaccgcggtgtactattgcgctaagcattactattatggcgggagctacgcaatggattactggggacagggtactctggtcaccgtgtccagcaccactaccccagcaccgaggccacccaccccggtcctaccatcgctcccagcctctgtccctgcgtccggaggcatgtagaccgcagctgggtggggcgtgcatacccggggtcttgacttcgcctgcgatatctacatttgggcccctctggctggtacttgccgggtcctgctgctttcactcgtgatcactctttactgtaagcgcggtcggaagaagctgctgtacatctttaagcaacccttatgaggccgtgcagactactcaagaggaggacggctgttcatgccggttcccagaggaggaggaggcggctgcgaactgcgcgtgaaattcagccgcagcgcagatgtccagcctacaagcaggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagtagacgtgctggacaagcggagaggacgggacccagaaatggcggggaagccgcgca gaaagaatccccagagggcctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaaggccacgacgg actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggcctgccgcctcgg
104880 CAR6 – Full – aa	82	MALPVTALLLPLALLLHAARPeivmtqspatlsispgeratlsc <u>rasqdiskylnw</u> yqqkpgqaprlliy <u>htsrllhs</u> giparfsgsgsgtdytlitisslqpedfavyfc <u>ggg</u> <u>ntlpyt</u> fgqgtkleikggggsgggsgggsgggsgvqlqesgpglvkpsetls tctvsgvslp <u>dyqvs</u> wirppgkglewig <u>viwgsettyygssls</u> krvtiskdnsk nqvslklssvtaadtavyyca <u>hyyyyggsyamdy</u> wgqgtlvtvssttppaprptpa ptiasqplslrpeacrpaaggavhtrglfacdiyiwaplagtcgvllslvitly ckrgrkkllyifkqpfmrpvqttqeedgcscrfeeeeeggcelrvkfsrsadapay kqqgnqlynelnlgrreeydvldkrrgrdpemggkprrknpqeglynelqkdmae ayseigmkgerrrrgkghdglyqglstatkdydalhmqalppr
CAR 7		
CAR7 scFv domain	45	qvqlqesgpglvkpsetls tctvsgvslpdygvswirppgkglewigviwgset tyyslksrvtiskdnsk nqvslklssvtaadtavyyca <u>hyyyyggsyamdy</u> wgq gtlvtvssggggsgggsgggsgggsgggseivmtqspatlsispgeratlscrasq diskylwnwyqqkpgqaprlliyhtsrllhs giparfsgsgsgtdytlitisslqpedfa

		vyfcqqgntlpytfgggtkleik
100796 CAR7 - Soluble scFv - nt	58	atggcactgcctgtcactgcctcctgctgcctctggccctccttctgcatgccgc caggccccaagtcacagctgcaagagtcaggaccggactggagaagccgtctgaga ctctctcactgacttgtaccgtcagcggcgtgtccctcccagactacggagtgca tggatccgccaacctcccggaagggcttgaatggattgggtgtcatctggggttc tgaaaccacctaactcatcttccctgaagtcagggtgaccatcagcaaggata attccaagaaccaggtcagccttaagctgtcatctgtgaccgtgctgacaccgcc gtgtattactgcgccaaagcactactattacggaggaagctacgctatggactattg gggacagggcactctcgtgactgtgagcagcggcggtggagggtctggagggtggag gatccgggtgggtgggtcaggcggaggaggagcagagattgtgatgactcagtc ccagccaccctttctctttcaccggcgagagcaaccctgagctgtagagccag ccaggacatttctaagtacctaactggatcagcaaaaaccggggcaggccctc gcctcctgatctaccatacctcacgccttactctggatccccgctcggtttagc ggatcaggatctggtaccgactacactctgaccatttccagcctgcagccagaaga tttcgcagtgtatttctgccagcagggcaatacccttccctacaccttcgggtcagg gaaccaagctcgaaatcaagcaccatcaccatcatcaccaccat
100796 CAR7 - Soluble scFv - aa	70	<u>MALPVTALLLPLALLHAARP</u> qvqlqesgpglvkpseltctvsgvslpdygvs wirqppgkglewigviwgsettyyssslksrvtiskdnskqvslklssvtaadta vyycakhyyyggsyamdywgqgtlvtvssggggsgggsgggsgggseivmtqs patlslspgeratlscrasqdiskylnwyqqkpgqaprlliyhtsrhsgiparfs gsgsgtdytltiisslqpedfavvyfcqqgntlpytfgggtkleik <u>hhhhhhh</u>
104881 CAR 7 Full - nt	96	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgcgc tcgcccacaagtcacagcttcaagaatcagggcctggtctggtgaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgccctcccagactacggagtgagc tggattagacagcctcccggaagggactggagtgatcgagtgatttggggtag cgaaaccacttactattcatcttccctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgcgcgtgacaccgcc gtgtattactgtgccaaagcattactactatggagggtcctacgccatggactactg gggccagggaactctggtcactgtgtcatctggtggaggaggtagcggaggaggcg ggagcgggtggagggtggtccggagggtggcggaagcgaaatcgtgatgaccagagc cctgcaaccctgtccctttctcccggaacgggtaccctttcttgcgggcac acaagatatctcaaaatacctcaattggtatcaacagaagccgggacagggcccta ggcttcttatctaccacacctctgcctgcatagcgggattccgcacgctttagc gggtctggaagcgggaccgactacactctgaccatctcatctctccagcccaggga cttcgcgctctacttctgccagcagggtaacaccctgccgtacaccttcggccagg gcaccaagcttgagatcaaaaccactactccgctccaaggccaccaccctgcc ccgaccatcgctctcagccgctttccctgcgtccggaggcatgtagaccgcagc tggtggggcgtgcatacccggttcttgacttcgcctgcgatatctacatttggg

		<p>ccccctctggctggtacttgccgggtcctgctgctttcactcgtgatcactctttac tgaagcgcggctcggaagaagctgctgtacatctttaagcaacccttcagaggcc tgtgcagactactcaagaggaggacggctgttcagtcgggttcccagaggaggagg aaggcggctgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac aagcaggggagagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaaatggcggggaagccgcgca gaaagaatccccagagggtcgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaaggccacgacgg actgtaccagggtactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgcctcgg</p>
104881 CAR 7 Full - aa	83	<p>MALPVTALLLPALLLHAARPqvqlqesgpglvkpsetlsltctvsgvslpdygvs wirqppgkglewigviwgsettyysslksrvtiskdsknqvslklssvtaadta vyycakhyyyggsyamydwgggtlvtvssggggsgggsgggsggggseivmtqs patlsispgeratlscrasqdiskylnwyqqkpgqaprllyhtsrhsgiparfs gsgsgtdytlitisslqpedfavyfcqggntlpytfgggtkleiktttpprptpa ptiasqplslrpeacrpaaggavhtrglldfacdiyiwaplagtcgvllslvitly ckrgrklllyifkqpfmrpvqttqeedgcscrfeeeeeggcclrvkfsrsadapay kqqgnqlynelnlgrreeydvldkrrgrdpemggkprkrnpqeglynelqkdmae ayseigmkgerrrrgkghdglyqglstatkdydalhmqalppr</p>
CAR 8		
CAR8 scFv domain	46	<p>qvqlqesgpglvkpsetlsltctvsgvslpdygvswirqppgkglewigviwgset tyyqsslksrvtiskdsknqvslklssvtaadtavyycahyyyggsyamydwgg gtlvtvssggggsgggsgggsggggseivmtqspatlsispgeratlscrasqd iskylnwyqqkpgqaprllyhtsrhsgiparfsgsgsgtdytlitisslqpedfa vyfcqggntlpytfgggtkleik</p>
100798 CAR8 - Soluble scFv - nt	59	<p>atggcactgcctgtcactgcctcctgctgcctctggccctccttctgcatgccgc caggccccaaagtcagctgcaagagtcaggaccggactggtgaagccgtctgaga ctctctcactgacttgtaccgtcagcggcgtgtccctccccgactacggagtgtca tggatccgccaaacctcccgggaaagggttgatggattggtgtcatctggggttc tgaaaccacactactaccagtcttcctgaagtcagggtgaccatcagcaaggata attccaagaaccaggtcagccttaagctgtcatctgtgaccgctgctgacaccgcc gtgtattactgcgccaaagcactactattacggaggaagctacgctatggactattg gggacagggcactctcgtgactgtgagcagcggcggtggagggtctggagggtggag gatccgggtggtggtgggtcaggcggaggaggagcgagattgtgatgactcagtc ccagccaccctttctctttcaccggcgagagagcaaccctgagctgtagagccag ccaggacatttctaagtacctcaactggtatcagcaaaaaccggggcaggccccctc gctcctgatctaccatacctcacgccttctactctggtatccccgctcggttagc ggatcaggatctggtaccgactacactctgaccatttccagcctgcagccagaaga</p>

		tttcgcagtgatattctgccagcagggcaatacccttcccttacaccttcgggtcagg gaaccaagctcgaaatcaagcaccatcaccatcatcatcaccac
100798 CAR8 - Soluble scFv - aa	71	<u>MALPVTALLLPLALLLHAARP</u> qvqlqesgpglvkpssetlsltctvsgvslpdygvs wirqppgkglewigviwgsettyyqsslksrvtiskdsknqvslklssvtaadta vyycakhyyyggsyamydwgggtlvtvssgggsgggsgggsggggseivmtqs patlsispgeratlscrasqdiskylnwyyqqkpgqaprlliyhtsrhsgiparfs gsgsgtdytlttisslqpedfavyfcqqgntlpvtfgggtkleik <u>hhhhhhhh</u>
104882 CAR 8 – Full - nt	97	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacagcttcaagaatcagggcctggctctggatgaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgtccctcccagactacggagtgagc tggattagacagcctcccgaaagggaactggagtggatcgagagtatttggggtag cgaaaccacttactatcaatcttcctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgcgcgtgacaccgcc gtgtattactgtgccaaagcattactactatggagggtcctacgccatggactactg gggacaggaactctggctcactgtgtcatctggaggaggtagcggaggaggcg ggagcgggtggaggtggctccggaggcgggtgggtcagaaatcgtgatgaccagagc cctgcaaccctgtccctttctccggggaacgggctaccctttcttgcgggcac acaagatatctcaaaatacctcaattggatcaacagaagccgggacagggcccta ggcttcttatctaccacacctctcgctgcatagcgggattcccgacagcttttagc gggtctggaagcgggaccgactacactctgaccatctcatctctccagcccagga cttcgcgcgtctacttctgccagcagggtaacaccctgccgtacaccttcggccagg gcaccaagcttgagatcaaaaccactactccgctccaagggcaccacccctgcc ccgaccatcgctctcagccgctttccctgcgtccggaggcatgtagaccgcagc tgggtgggcccgtgcatacccggggtcttgacttcgcctgcgatatctacatttggg ccctctggctggtacttgccgggtcctgctgctttcactcgtgatcactctttac tgtaagcgcggctcggaagaagctgctgtacatctttaagcaaccctcatgaggcc tgtgcagactactcaagaggaggacggctgtcatgccggttcccagaggaggagg aaggcggctgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac aagcaggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaatgggcggaagccgcgca gaaagaatcccaagagggcctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggatgaaaggggaacgcagaagaggcaaaggccacgacgg actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgcctcgg
104882 CAR 8 – Full - aa	84	<u>MALPVTALLLPLALLLHAARP</u> qvqlqesgpglvkpssetlsltctvsgvslp <u>dygvs</u> wirqppgkglewig <u>viwgsettyyqsslks</u> rvtiskdsknqvslklssvtaadta vyycak <u>hyyyggsyamyd</u> wgggtlvtvssgggsgggsgggsggggseivmtqs patlsispgeratlsc <u>rasqdiskyl</u> nwyyqqkpgqaprlliy <u>htsrhsg</u> iparfs

		gsgsgtdytlttisslqpedfavyfc gggntlpyt fgggtkleiktttpprptpa ptiasqplslrpeacrpaaggavhtrglfacdiyiwaplagtcgvllslvitly ckrgrklllyifkqpfmrpvqttqeedgcscrfeeeeeggcclrvkfsrsadapay kqqgnqlynelnlgrreeydvldkrrgrdpemggkprrknpqeglynelqkdmae ayseigmkgerrrgkghdglyqglstatkdydalhmqlppr
CAR 9		
CAR9 scFv domain	47	eivmtqspatlsislspgeratlscrasqdiskylnwyyqqkpgqaprlliyhtsrllhs giparfsgsgsgtdytlttisslqpedfavyfcqqgntlpytfgggtkleikggggs gggsgggsgggsgggsgvqlqesgpglvkpsetlsltctvsgvslpdygvswirqpp gkglewigviwgsettyynsslskrvtiskdnsknqvs1klssvtaadtavyycak hyyyggsyamdywgggtlvtvss
99789 CAR9 - Soluble scFv - nt	60	atggccctcccagtgaccgctctgctgctgcctctcgcaacttcttctccatgccgc tcggcctgagatcgatgacccaaagccccgctaccctgtccctgtcaccgcggcg agagggcaaccctttcatgcagggccagccaggacatttctaagtacctcaactgg tatcagcagaagccagggcaggctcctcgctgctgatctaccacaccagcgcct ccacagcggatcccccgccagattttccgggagcgggtctggaaccgactacaccc tcaccatctcttctctgcagcccgaggatttcgcccgtctatttctgccagcagggg aatactctgccgtacaccttcgggtcaaggtaccaagctggaaatcaagggaggcgg aggatcaggcgggtggcggaagcggaggagggtggctccggaggaggagggtcccaag tgcagcttcaagaatcaggaccggacttgtgaagccatcagaaacctctcctg acttgtaccgtgtccgggtgtgagcctccccgactacggagtctcttggtatcgcca gcctccggggaagggtcttgaatggattgggggtgatttggggatcagagactactt actacaattcatcacttaagtacgggtcaccatcagcaaagataatagcaagaac caagtgtcacttaagctgtcatctgtgaccgccgctgacaccgccgtgtactattg tgccaaacattactattacggagggtcttatgctatggactactggggacagggga ccctggtgactgtctctagccatcaccatcaccaccatcatcac
99789 CAR9 - Soluble scFv - aa	72	<u>MALPVTALLLPLALLLHAARP</u> eivmtqspatlsislspgeratlscrasqdiskylnw yqqkpgqaprlliyhtsrllhsigiparfsgsgsgtdytlttisslqpedfavyfcqqg ntlpytfgggtkleikgggsgggsgggsgggsgvqlqesgpglvkpsetls1 tctvsgvslpdygvswirqppgkglewigviwgsettyynsslskrvtiskdnskn qvs1klssvtaadtavyycakhyyyggsyamdywgggtlvtvss <u>hhhhhhhh</u>
105974 CAR 9 – Full - nt	98	atggccctccctgtcaccgcctgctgcttccgctggctcttctgctccacgccgc tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaaccctgtcttgcagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggctcctcgcttctgatctaccacaccagcggct ccattctggaatccctgccagggtcagcggtagcggatctgggaccgactacaccc tcactatcagctcactgcagccagaggacttcgctgtctatttctgtcagcaaggg aacaccctgccctacacctttggacagggcaccaagctcgagattaaaggtggagg

		<p>tggcagcggaggaggtgggtccggcgggtggaggaagcggaggcgggtgggagccagg tccaactccaagaaagcggaccgggtcttgtgaagccatcagaaactctttcactg acttgtactgtgagcggagtgtctctccccgattacgggggtgtcttggatcagaca gccaccggggaaggggtctggaatggattggagtgatttggggctctgagactactt actacaactcatccctcaagtcacgcgtcaccatctcaaaggacaactctaagaat caggtgtcactgaaactgtcatctgtgaccgcagccgacaccgcctgtactattg cgctaagcattactattatggcgggagctacgcaatggattactggggacagggtta ctctggtcaccgtgtccagcaccactaccccagcaccgaggccacccaccccggt cctaccatcgctcccagcctctgtccctgcgtccggaggcatgtagaccgcagc tgggtggggcctgcatacccggggtcttgacttgcgctgcgatctacatttggg cccctctggctggtacttgcgggggtcctgctgctttcactcgtgatcactctttac tghtaagcgggtcggaagaagctgctgtacatctttaagcaacccttcatgaggcc tgtgcagactactcaagaggaggacggctgttcatgccgggtccagaggaggagg aaggcggctgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac aagcagggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaatggcggggaagccgcgca gaaagaatccccagaggggcctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaaggccacgacgg actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgctcgg</p>
105974 CAR 9 – Full - aa	85	<p>MALPVTALLPLALLHAARPeivmtqspatlsispgeratlsc<u>rasqdiskylnw</u> yqqkpgqaprlliy<u>htsrllhs</u>giparfsgsgsgtdytlitisslqpedfavyfc<u>ggg</u> <u>ntlpyt</u>fgqgkkleikggggsgggsgggsgggsgvqlqesgpglvkpsetls tctvsgvslp<u>dygvs</u>wirppgkglewig<u>viwgsettyynsslks</u>rvtiskdnsk qvsiklssvtaadtavyycak<u>hyyyggsyamdy</u>wgqgtlvtvssttpprptpa ptiasqplslrpeacrpaaggavhtrglfacdiyiwaplagtcgvllslvitly ckrgrklllyifkqpfmrpvqttqeedgcscrffpeeeeggcelrvkfsrsadapay kqqnqlynelnlgrreeydvldkrgrdpemggkprkrnpqeglynelqkdkmae ayseigmkgerrrrgkghdglyqglstatkdydalhmqalppr</p>
CAR10		
CAR10 scFv domain	48	<p>qvqlqesgpglvkpsetlsltctvsgvslpdygvswirppgkglewigviwgset tyynsslksrvtiskdnskqvsiklssvtaadtavyycakhyyyggsyamdywgq gtlvtvssgggsgggsgggsgggsgggseivmtqspatlsispgeratlscrasqd iskylnwyyqqkpgqaprlliyhtsrllhsgiparfsgsgsgtdytlitisslqpedfa vyfcqqgntlpytfgqgkkleik</p>
100796 CAR10 - Soluble	61	<p>atggcactgcctgtcactgcctcctgctgcctctggccctccttctgcatgccgc caggccccaagtcacagctgcaagagtcaggacccggactggatgaagccgtctgaga ctctctcactgacttgtaccgtcagcggcgtgtccctccccgactacggagtgtca</p>

scFv - nt		<p>tggatccgccaacctcccgggaaagggcttgaatggattggtgtcatctggggttc tgaaccacctaactacaactcttccctgaagtcaggggtgaccatcagcaaggata attccaagaaccaggtcagccttaagctgtcatctgtgaccgctgctgacaccgcc gtgtattactgcgccaagcactactattacggaggaagctacgctatggactattg gggacagggcactctcgtgactgtgagcagcgcggtggagggctctggaggtggag gatccggtgggtgggtcaggcggaggaggagcagagattgtgatgactcagtca ccagccacctttctctttcaccggcgagagcaacctgagctgtagagccag ccaggacatttctaagtacctcaactggtatcagcaaaaaccgggagggccctc gctcctgatctaccatacctcacgccttcactctggtatccccgctcggtttagc ggatcaggatctggtaccgactacactctgaccatttccagcctgcagccagaaga tttcgcagtgtatttctgccagcagggcaataaccttcccttacaccttcggtcagg gaaccaagctcgaaatcaagcaccatcaccatcatcaccaccat</p>
100796 CAR10 - Soluble scFv - aa	73	<p><u>MALPVTALLLPLALLHAARP</u>qvqlqesgpglvkpsetlsltctvsgvslpdygvs wirqppgkglewigviwgsettyynsslksrvtiskdnskqvslklssvtaadta vyycakhyyyggsyamdywgggtlvtvssggggsgggsgggsgggseivmtqs patlsispgeratlscrasqdiskylnwyyqqkpgqaprllyhtsrhsgiparfs gsgsgtdytltiisslqpedfavycqqgntlpytfgggtkleik<u>hhhhhhhh</u></p>
105975 CAR 10 Full - nt	99	<p>atggccctccctgtcaccgcctctgtcctccgctggctcttctgctccacgcgc tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaacctgtcttgcagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggtcctcgccttctgatctaccacaccagccggtc ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacaccc tcactatcagctcactgcagccagaggacttcgctgtctatttctgtcagaaggg aacaccctgcctacacctttggacagggcaccaagctcgagattaaaggtggagg tggcagcggaggaggtgggtccggcggtggaggaagcggaggcgggtgggagccagg tccaactccaagaaagcggaccgggtcttgtgaagccatcagaaactctttcactg acttgtactgtgagcggagtgtctctccccgattacgggggtgtcttggtatcagaca gccaccggggaaggtctggaatggattggagtgatttggggctctgagactactt actacaactcatccctcaagtcacgcgtcaccatctcaaaggacaactctaagaat caggtgtcactgaaactgtcatctgtgaccgcagccgacaccgcggtgtactattg cgctaagcattactattatggcgggagctacgcaatggattactggggacagggta ctctggtcaccgtgtccagcaccactacccagcaccgagccacccaccccggtc cctaccatcgctccagcctctgtccctgcgtccggaggcatgtagaccgcagc tgggtggggccgtgcatacccggggtcttgacttcgcctgcgatctacatttggg ccctctggctggtacttgcggggtcctgctgctttcactcgtgatcactctttac tgtaagcgggtcggaagaagctgctgtacatctttaagcaaccttcatgagggc tgtgcagactactcaagaggaggacggctgttcatgccggttccagaggaggagg aaggcggtgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac</p>

		aagcaggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaatggcggggaagccgcgca gaaagaatccccagaggcctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaaggccacgacgg actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgcctcgg
105975 CAR 10 Full - aa	86	MALPVTALLLP LALLLHAARPEIVMTQSPATLSLSPGERATLSCL <u>RASQDISKYL</u> NW YQQKPGQAPRLLIY <u>HTSRLHS</u> GIPARFSGSGSGTDYTLTISSLPEDFAVYFC <u>QQG</u> <u>NTLPYT</u> FGQGTKLEIKGGGSGGGGSGGGGSGGGGSGVQLQESGPGLVKPSSETLSL TCTVSGVSLP <u>DYGV</u> SWIRQPPGKGLEWIG <u>VIWGSETTYNSSLSK</u> SRVTISKDNSKN QVSLKLSSVTAADTAVYYCAK <u>HYYYGGSYAMDY</u> WGQGLTVTVSSTTTPAPRPPTPA PTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVITLY CKRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAY KQGQNLQYLNELNLGRREEYDVLDRRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAE AYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQALPPR
CAR11		
CAR11 scFv domain	49	eivmtqspatls slspgeratls crasqdiskyl nwyqqkpgqaprlliihtsrlhs giparfsgsgsgtdytl t isslqpedfavyfcqqgntlpytfgggtkleikggggs gggsggggsgvqlqesgpglvkps etlsl tctvsgvslpdygvswirppgkgle wigviwgsettyynsslsk rvtiskdnsk nqvslklssvtaadtavyyca khyyyg gsyamdywgqgtlvtvss
103101 CAR11 - Soluble scFv - nt	62	Atggccctccctgtcaccgcctgctgcttcgctggctcttctgctccacgcgcg tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaaccctgtcttgagagcctcccaagacatctcaaaataccttaattgg tatcaacagaagcccgacaggctcctcgcttctgatctaccacaccagccggct ccattctggaatccctgccaggttcagcggtagcggatctgggaccgactacaccc tactatcagctcactgcagccagaggacttcgctgtctatttctgtcagcaaggg aacaccctgccctacacctttggacagggcaccaagctcgagattaaagggtggagg tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaaa gcggaccgggtcttgtgaagccatcagaaactctttcactgacttgtagtgagc ggagtgtctctccccgattacggggtgtcttgatcagacagccaccggggaaggg tctggaatggattggagtgatttggggtctgagactacttactacaattcatccc tcaagtacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacaccgcgtgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacagggtactctggtcaccgtgt ccagccaccaccatcatcaccatcaccat

103101 CAR11 - Soluble scFv - aa	74	<u>MALPVTALLLPLALLLHAARP</u> eiivmtqspatlsislspgeratlscrasqdiskylnw yqqkpgqaprrllyhtsrhlhsgiparfsgsgsgtdytlitisslqpedfavyfcqqg ntlpytfgqgtkleikgggsgggsgggsgggsgvqlqesgpglvkpssetlsltctvs gvslpdygvswirppgkglewigviwgsettyynsslksrvtiskdnskqnqvsik lssvtaadtavyycakhyyyggsyamdywgqgtlvtvss <u>hhhhhhhh</u>
105976 CAR 11 Full - nt	100	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacagcttcaagaatcagggcctggctgctggaagccatctgaga ctctgtccctcacttgcaccgtgagcggagtgtccctcccagactacggagtgagc tggattagacagcctcccggaaagggactggagtggatcgagtgatttggggtag cgaaaccacttactataactcttcctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgcgcgtgacaccgcc gtgtattactgtgccaagcattactactatggagggtcctacgccatggactactg gggccaggaactctggtcactgtgtcatctggtggaggaggtagcggaggaggcg ggagcgggtggaggtggctccggaggtggcggaagcgaaatcgatgacccagagc cctgcaaccctgtccctttctccggggaacgggctaccctttcttgcgggcac acaagatatctcaaaatacctcaattggtatcaacagaagccgggacagggcccta ggcttcttatctaccacacctctcgctgcatagcgggattcccgcacgcttttagc gggtctggaagcgggaccgactacactctgaccatctcatctctccagcccagga cttcgcgtctacttctgccagcagggtaacaccctgccgtacaccttcggccagg gcaccaagcttgagatcaaaaccactactccgctccaaggccaccaccctgcc ccgaccatcgctctcagccgctttccctgcgtccggagggatgtagaccgcagc tgggtggggcgtgcatacccggttcttgacttcgcctgcgatctacatttggg ccctctggctggtacttgcgggtcctgctgctttcactcgtgatcactctttac tgtaagcgggtcggaagaagctgctgtacatctttaagcaacccttcatgaggcc tgtgcagactactcaagaggaggacggctgttcatgccggttcccagaggaggagg aaggcggctgcgaactgcgcgtgaaattcagccgcagcgcagatgctccagcctac aagcaggggcagaaccagctctacaacgaactcaatcttggtcggagagaggagta cgacgtgctggacaagcggagaggacgggacccagaaatggcggggaagccgcgca gaaagaatccccagagggcctgtacaacgagctccaaaaggataagatggcagaa gcctatagcgagattggtatgaaaggggaacgcagaagaggcaaaggccacgagc actgtaccagggactcagcaccgccaccaaggacacctatgacgctcttcacatgc aggccctgccgcctcgg
105976 CAR 11 Full - aa	87	MALPVTALLLPLALLLHAARPQVQLQESGPGLVKPSSETLSLTCTVSGVSLP <u>DYGV</u> WIRQPPGKGLEWIG <u>VIWGSETTYYNSSLKS</u> RVTISKDNSKNQVSLKLSSVTAADTA VYYCAK <u>HYYYGGSYAMDY</u> WGQGTlvtvssgggsgggsgggsgggsgggseivmtqs PATLSLSPGERATLS <u>CASQDISKYL</u> NWYQQKPGQAPRLIY <u>HTSRLHSG</u> IPARFS GSGSGTDYTLTISSSLQPEDFAVYFC <u>COGNTLPYT</u> FGQGTKLEIKTTTPAPRPPTPA

		PTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVITLY CKRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAY KQGQNQLYNELNLGRREEYDVLDRRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAE AYSEIGMKGERRRGKGDGLYQGLSTATKDTYDALHMQALPPR
CAR12		
CAR12 scFv domain	50	qvqlqesgpglvkpsetlsltctvsgvslpdygvswirqppgkglewlgviwgset tyynsslksrvtiskdnsknqvsllkssvtaadtavyycakhyyyggsyamdywgq gtlvtvssggsgsgsgsgsgsgseivmtqspatlsispgeratlscrasqdiskyl nwyqqkpgqaprlliyhtsrllhsgiparfsgsgsgtdytlttisslqpedfavyfcq qgntlpytfgggtkleik
103104 CAR12 - Soluble scFv - nt	63	atggctctgcccgtgaccgcactcctcctgccactggctctgctgcttcacgccgc tcgcccacaagtcacgcttcaagaatcagggcctggctctggatgaagccatctgaga ctctgtccctcacttgaccgtgagcggagtgtccctcccagactacggagtgagc tggttagacagcctcccggaaaggactggagtggatcggagtgtattgggtag cgaaaccacttactataactcttccctgaagtcacgggtcaccatttcaaaggata actcaaagaatcaagtgagcctcaagctctcatcagtcaccgccgctgacaccgcc gtgtattactgtgccaagcattactactatggagggtcctacgccatggactactg gggccagggaactctggctactgtgtcatctggaggaggtagcggaggaggcg ggagcggtaggagggtggctccgaaatcgtgatgaccagagccctgcaaccctgtcc ctttctcccggggaacgggctaccctttcttgcgggcatcacaagatatctcaa atacctcaattggtatcaacagaagccgggacaggccctaggcttcttattctacc acacctctgcctgcatagcgggattcccgcacgcttttagcgggtctggaagcggg accgactacactctgaccatctcatctctccagcccaggacttcgccgtctactt ctgccagcagggtaacaccctgccgtacaccttcggccagggcaccaagcttgaga tcaaacatcaccaccatcatcaccatcac
103104 CAR12 - Soluble scFv -aa	75	<u>MALPVTALLLPLALLHAARP</u> qvqlqesgpglvkpsetlsltctvsgvslpdygvs wirqppgkglewlgviwgsettyynsslksrvtiskdnsknqvsllkssvtaadta vyycakhyyyggsyamdywgqgtlvtvssggsgsgsgsgsgseivmtqspatls lspgeratlscrasqdiskylnwyqqkpgqaprlliyhtsrllhsgiparfsgsgsg tdytlttisslqpedfavyfcqgqgntlpytfgggtkleik <u>hhhhhhhh</u>
105977 CAR 12 - Full - nt	101	atggccctccctgtcaccgcctcctgctgcttccgctggctcttctgctccacgccgc tcggcccgaaattgtgatgaccagtcacccgccactcttagcctttcaccgggtg agcgcgcaaccctgtcttgagagcctcccaagacatctcaaataccttaattgg tatcaacagaagcccgacaggctcctcgcttctgatctaccacaccagcggct ccattctggaatccctgccagggtcagcggtagcggatctgggaccgactacacc tcactatcagctcactgcagccagaggacttcgctgtctatcttctgtcagcaaggg aacaccctgccctacaccttggacagggcaccaagctcgagattaaaggtggagg

		<p>tggcagcggaggaggtgggtccggcgggtggaggaagccaggtccaactccaagaaa gcggaaccgggtcttgtgaagccatcagaaactctttcactgacttgtactgtgagc ggagtgtctctccccgattacggggtgtcttggatcagacagccaccgggaagg tctggaatggattggagtgttggggctctgagactacttactacaactcatccc tcaagtcacgcgtcaccatctcaaaggacaactctaagaatcaggtgtcactgaaa ctgtcatctgtgaccgcagccgacaccgcctgtactattgcgctaagcattacta ttatggcgggagctacgcaatggattactggggacagggtactctggtcaccgtgt ccagcaccactaccccagcaccgaggccacccaacccggctcctaccatcgctcc cagcctctgtccctgcgtccggaggcatgtagaccgcagctggtggggcctgca taccgggggtcttgacttcgctgcgatatctacatttgggcccctctggtggta cttgcggggctcctgctgctttcactcgtgatcactctttactgtaagcgcggtcgg aagaagctgctgtacatctttaagcaaccctcatgaggcctgtgcagactactca agaggaggacggctgttcatgccggttcccagaggaggaggaaggcggtgcgaac tgcgctgaaattcagccgcagcgcagatgctccagcctacaagcaggggcagaac cagctctacaacgaactcaatcttggtcggagagaggagtacgacgtgctggacia gcggagaggacgggaccagaaatgggcgggaagccgcgcagaaagaatccccaag aggcctgtacaacgagctccaaaaggataagatggcagaagcctatagcgagatt ggtatgaaaggggaacgcagaagaggcaaaggccacgagcactgtaccagggact cagcaccgccaccaaggacacctatgacgctcttcacatgcaggccctgcgcctc gg</p>
105977 CAR 12 – Full - aa	88	<p>MALPVTALLLP LALLLHAARPEIVMTQSPATLSLSPGERATLS <u>C</u><u>R</u><u>A</u><u>S</u><u>Q</u><u>D</u><u>I</u><u>S</u><u>K</u><u>Y</u><u>L</u><u>N</u><u>W</u> YQKPGQAPRLLIY <u>H</u><u>T</u><u>S</u><u>R</u><u>L</u><u>H</u><u>S</u> GIPARFSGSGSGTDYTLTISSLPEDFAVYFC <u>Q</u><u>Q</u><u>G</u> <u>N</u><u>T</u><u>L</u><u>P</u><u>Y</u><u>T</u> FGQTKLEIKGGGSGGGGSGGGGSGVQLQESGPGLVKPSSETLSLTCTVS GVSLP <u>D</u><u>Y</u><u>G</u><u>V</u><u>S</u> WIRQPPGKGLEWIG <u>V</u><u>I</u><u>W</u><u>G</u><u>S</u><u>E</u><u>T</u><u>T</u><u>Y</u><u>N</u><u>S</u><u>S</u><u>L</u><u>K</u><u>S</u> RVTISKDNSKNQVSLK LSSVTAADTAVYYCAK <u>H</u><u>Y</u><u>Y</u><u>G</u><u>G</u><u>S</u><u>Y</u><u>A</u><u>M</u><u>D</u><u>Y</u> WQGTLTVTSSTTPAPRPPTPAPTIAS QPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVITLYCKRGR KKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAYKQGQN QLYNELNLGRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEI GMKGERRRGKGDGLYQGLSTATKDTYDALHMQALPPR</p>
CTL019		
CTL019 – Soluble scFv-Histag - nt	22	<p>atggccctgcccgtaaccgctctgctgctgccccttgcctctgcttcttcatgcagc aaggccggacatccagatgacccaaaccacctcatccctctctgcctctcttggag acagggtgaccatttcttgtcgcgccagccaggacatcagcaagtatctgaactgg tatcagcagaagccggacggaaccgtgaagctcctgatctaccatacctctcgct gcatagcggcgtgccctcacgcttctctggaagcggatcaggaaccgattattctc tactatttcaaatcttgagcaggaagatattgccacctatttctgccagcagggt</p>

		aataccctgccctacaccttcggaggagggaccaagctcgaaatcacccggtggagg aggcagcggcggtggagggctctggtggaggtggttctgaggtgaagctgcaagaat caggccctggacttgtggccccctcacagtccttgagcgtgacttgccacgtgtcc ggagtctccctgccgactacggagtgtcatggatcagacaacctccacggaaagg actggaatggctcggtgtcatctggggtagcgaaactacttactacaattcagccc tcaaaagcaggctgactattatcaaggacaacagcaagtcaccaagctctttcttaag atgaactcactccagactgacgacaccgcaatctactattgtgctaagcactacta ctacggaggatcctacgctatggattactggggacaaggtacttccgtcactgtct cttcacaccatcatcaccatcaccatcac
CTL019 – Soluble scFv-Histag - aa	76	<u>MALPVTALLPLALLHAARP</u> diqmtqttsslsaslgdrvtiscrasqdiskylnw yqqkpdgtvkllyhtsrllhsgvpsrfsrgsgsgtdysltisnleqediatyfcqqg ntlpytfgggtkleitggggsgggsgggggsevglqesgpglvapsqslsvtctvs gvslpdygvswirpprkglewlgviwgsettyynsalksrlltiikdnksqvflk mns1qtdtdtaiyycahyyyggsyamdywggtsvtvss <u>hhhhhhhh</u>
CTL019 Full - nt	102	atggccttaccagtgaaccgccttgcctcctgcgctggccttgcctcctacgcgc caggccggacatccagatgacacagactacatcctccctgtctgcctctctgggag acagagtaccatcagttgcagggaagtcaggacattagtaaatatttaaattgg tatcagcagaaaccagatggaactgttaaactcctgatctaccatacatcaagatt acactcaggagtcccatcaagggtcagtgccagtggtctggaacagattattctc tcaccattagcaacctggagcaagaagatattgccacttacttttgccaacagggg aatacgttccgtacacgttcggaggggggaccaagctggagatcacaggtggcgg tggtcgggcggtggtgggtcgggtggcgccggtctgaggtgaaactgcaggagt caggacctggcctggtggcgccctcacagagcctgtccgtcacatgcactgtctca ggggtctcattaccgcactatggtgtaagctggattcgccagcctccacgaaaggg tctggagtggctgggagtaatatggggtagtgaaccacatactataattcagctc tcaaatccagactgaccatcatcaaggacaactccaagagccaagttttcttaaaa atgaacagtctgcaaactgatgacacagccatttactactgtgccaaacattatta ctacggtggtagctatgctatggactactggggccaaggaaacctcagtcaccgtct cctcaaccacgacgccagcgccgcgaccaccaacaccggcgccaccatcgctcg cagccctgtccctgcgcccagaggcgtgccggccagcgcggggggcgcagtga cacgagggggctggacttcgctgtgatatctacatctgggcgcccttggecgga cttgtggggtccttctcctgtcactgggttatcacccttactgcaaacggggcaga aagaaactcctgtatatattcaacaaccatttatgagaccagtacaaactactca agaggaagatggctgtagctgccgatttccagaagaagaaggaggatgtgaac tgagagtgaagttcagcaggagcgcagacgccccgcgtacaagcagggccagaac cagctctataacgagctcaatctaggacgaagagaggagtacgatgttttggaac gagacgtggccgggaccctgagatggggggaaagccgagaaggaagaacctcagg aaggcctgtacaatgaactgcagaaagataagatggcgaggcctacagtgaatt

		gggatgaaaggcgagcgccggaggggcaaggggcacgatggcctttaccaggggtct cagtacagccaccaaggacacctacgacgcccttcacatgcaggccctgccccctc gc
CTL019 Full - aa	89	MALPVTALLLPLALLLHAARPdiqmtqttsslsaslgdrvtiscrasqdiskylnw yqqkpdgtvkllyhtsrhlsgvpsrfsfgsgsgtdysltisnleqediatyfcqqg ntlpytfgggkkleitgggsgggsggggsevglqesgpglvapsqslsvtctvs gvslpdygvswirqprrkglewlgviwgsettyynsalksrltiikdnksqvflk mns1qtddtaiyycahyyyggsyamdywgqgtsvtvssttppaprpptpaptias qp1slrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgr kkllyifkqpfmrpvqttqeedgcscrffeeeeggcclrvkfsrsadapaykqgn qlynelnlgrreeydvldkrrgrdpemggkprknpqeglynelqkdmaeysei gmkgerrrgkghdglyqglstatkdttydalhmqalppr
CTL019 scFv domain	51	diqmtqttsslsaslgdrvtiscrasqdiskylnwyyqqkpdgtvkllyhtsrhlsg vpsrfsfgsgsgtdysltisnleqediatyfcqqgntlpytfgggkkleitgggsg gggsggggsevglqesgpglvapsqslsvtctvsqslpdygvswirqprrkgle wlgviwgsettyynsalksrltiikdnksqvflkmns1qtddtaiyycahyyyg gsyamdywgqgtsvtvss

Co-expression of CAR with Other Molecules or Agents

Co-expression of a Second CAR

In one aspect, the CAR-expressing cell described herein can further comprise a second

5 CAR, e.g., a second CAR that includes a different antigen binding domain, e.g., to the same target (e.g., CD19) or a different target (e.g., a target other than CD19, e.g., a target described herein). In one embodiment, the CAR-expressing cell comprises a first CAR that targets a first antigen and includes an intracellular signaling domain having a costimulatory signaling domain but not a primary signaling domain, and a second CAR that targets a second, different, antigen

10 and includes an intracellular signaling domain having a primary signaling domain but not a costimulatory signaling domain. Placement of a costimulatory signaling domain, e.g., 4-1BB, CD28, CD27, OX-40 or ICOS, onto the first CAR, and the primary signaling domain, e.g., CD3 zeta, on the second CAR can limit the CAR activity to cells where both targets are expressed. In one embodiment, the CAR expressing cell comprises a first CAR that includes

15 an antigen binding domain, a transmembrane domain and a costimulatory domain and a second CAR that targets another antigen and includes an antigen binding domain, a transmembrane domain and a primary signaling domain. In another embodiment, the CAR expressing cell

comprises a first CAR that includes an antigen binding domain, a transmembrane domain and a primary signaling domain and a second CAR that targets another antigen and includes an antigen binding domain to the antigen, a transmembrane domain and a costimulatory signaling domain.

5 In one embodiment, the CAR-expressing cell comprises an XCAR described herein and an inhibitory CAR. In one embodiment, the inhibitory CAR comprises an antigen binding domain that binds an antigen found on normal cells but not cancer cells, e.g., normal cells that also express X. In one embodiment, the inhibitory CAR comprises the antigen binding domain, a transmembrane domain and an intracellular domain of an inhibitory molecule. For example,
10 the intracellular domain of the inhibitory CAR can be an intracellular domain of PD1, PD-L1, PD-L2, CTLA4, TIM3, CEACAM (CEACAM-1, CEACAM-3, and/or CEACAM-5), LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, and TGFR (e.g., TGFRbeta).

15 In one embodiment, when the CAR-expressing cell comprises two or more different CARs, the antigen binding domains of the different CARs can be such that the antigen binding domains do not interact with one another. For example, a cell expressing a first and second CAR can have an antigen binding domain of the first CAR, e.g., as a fragment, e.g., an scFv, that does not form an association with the antigen binding domain of the second CAR, e.g., the
20 antigen binding domain of the second CAR is a VHH.

In some embodiments, the antigen binding domain comprises a single domain antigen binding (SDAB) molecules include molecules whose complementary determining regions are part of a single domain polypeptide. Examples include, but are not limited to, heavy chain variable domains, binding molecules naturally devoid of light chains, single domains derived
25 from conventional 4-chain antibodies, engineered domains and single domain scaffolds other than those derived from antibodies. SDAB molecules may be any of the art, or any future single domain molecules. SDAB molecules may be derived from any species including, but not limited to mouse, human, camel, llama, lamprey, fish, shark, goat, rabbit, and bovine. This term also includes naturally occurring single domain antibody molecules from species other than
30 Camelidae and sharks.

In one aspect, an SDAB molecule can be derived from a variable region of the immunoglobulin found in fish, such as, for example, that which is derived from the immunoglobulin isotype known as Novel Antigen Receptor (NAR) found in the serum of shark. Methods of producing single domain molecules derived from a variable region of NAR ("IgNARs") are described in WO 03/014161 and Streltsov (2005) Protein Sci. 14:2901-2909.

According to another aspect, an SDAB molecule is a naturally occurring single domain antigen binding molecule known as heavy chain devoid of light chains. Such single domain molecules are disclosed in WO 9404678 and Hamers-Casterman, C. et al. (1993) Nature 363:446-448, for example. For clarity reasons, this variable domain derived from a heavy chain molecule naturally devoid of light chain is known herein as a VHH or nanobody to distinguish it from the conventional VH of four chain immunoglobulins. Such a VHH molecule can be derived from Camelidae species, for example in camel, llama, dromedary, alpaca and guanaco. Other species besides Camelidae may produce heavy chain molecules naturally devoid of light chain; such VHHs are within the scope of the invention.

The SDAB molecules can be recombinant, CDR-grafted, humanized, camelized, de-immunized and/or in vitro generated (e.g., selected by phage display).

It has also been discovered, that cells having a plurality of chimeric membrane embedded receptors comprising an antigen binding domain that interactions between the antigen binding domain of the receptors can be undesirable, e.g., because it inhibits the ability of one or more of the antigen binding domains to bind its cognate antigen. Accordingly, disclosed herein are cells having a first and a second non-naturally occurring chimeric membrane embedded receptor comprising antigen binding domains that minimize such interactions. Also disclosed herein are nucleic acids encoding a first and a second non-naturally occurring chimeric membrane embedded receptor comprising an antigen binding domains that minimize such interactions, as well as methods of making and using such cells and nucleic acids. In an embodiment the antigen binding domain of one of the first and the second non-naturally occurring chimeric membrane embedded receptor, comprises an scFv, and the other comprises a single VH domain, e.g., a camelid, shark, or lamprey single VH domain, or a single VH domain derived from a human or mouse sequence.

In some embodiments, a composition herein comprises a first and second CAR, wherein the antigen binding domain of one of the first and the second CAR does not comprise a variable

light domain and a variable heavy domain. In some embodiments, the antigen binding domain of one of the first and the second CAR is an scFv, and the other is not an scFv. In some embodiments, the antigen binding domain of one of the first and the second CAR comprises a single VH domain, e.g., a camelid, shark, or lamprey single VH domain, or a single VH domain
5 derived from a human or mouse sequence. In some embodiments, the antigen binding domain of one of the first and the second CAR comprises a nanobody. In some embodiments, the antigen binding domain of one of the first and the second CAR comprises a camelid VHH domain.

In some embodiments, the antigen binding domain of one of the first and the second
10 CAR comprises an scFv, and the other comprises a single VH domain, e.g., a camelid, shark, or lamprey single VH domain, or a single VH domain derived from a human or mouse sequence. In some embodiments, the antigen binding domain of one of the first and the second CAR comprises an scFv, and the other comprises a nanobody. In some embodiments, the antigen binding domain of one of the first and the second CAR comprises an scFv, and the other
15 comprises a camelid VHH domain.

In some embodiments, when present on the surface of a cell, binding of the antigen binding domain of the first CAR to its cognate antigen is not substantially reduced by the presence of the second CAR. In some embodiments, binding of the antigen binding domain of the first CAR to its cognate antigen in the presence of the second CAR is at least 85%, 90%,
20 95%, 96%, 97%, 98% or 99%, e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99% of binding of the antigen binding domain of the first CAR to its cognate antigen in the absence of the second CAR.

In some embodiments, when present on the surface of a cell, the antigen binding domains of the first and the second CAR, associate with one another less than if
25 both were scFv antigen binding domains. In some embodiments, the antigen binding domains of the first and the second CAR, associate with one another at least 85%, 90%, 95%, 96%, 97%, 98% or 99% less than, e.g., 85%, 90%, 95%, 96%, 97%, 98% or 99% less than if both were scFv antigen binding domains.

Co-expression of an Agent that Enhances CAR Activity

30 In another aspect, the CAR-expressing cell described herein can further express another agent, e.g., an agent that enhances the activity or fitness of a CAR-expressing cell.

For example, in one embodiment, the agent can be an agent which inhibits a molecule that modulates or regulates, e.g., inhibits, T cell function. In some embodiments, the molecule that modulates or regulates T cell function is an inhibitory molecule. Inhibitory molecules, e.g., PD1, can, in some embodiments, decrease the ability of a CAR-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD1, PD-L1, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, or TGFR beta.

In embodiments, an agent, e.g., an inhibitory nucleic acid, e.g., a dsRNA, e.g., an siRNA or shRNA; or e.g., an inhibitory protein or system, e.g., a clustered regularly interspaced short palindromic repeats (CRISPR), a transcription-activator like effector nuclease (TALEN), or a zinc finger endonuclease (ZFN), e.g., as described herein, can be used to inhibit expression of a molecule that modulates or regulates, e.g., inhibits, T-cell function in the CAR-expressing cell. In an embodiment the agent is an shRNA, e.g., an shRNA described herein. In an embodiment, the agent that modulates or regulates, e.g., inhibits, T-cell function is inhibited within a CAR-expressing cell. For example, a dsRNA molecule that inhibits expression of a molecule that modulates or regulates, e.g., inhibits, T-cell function is linked to the nucleic acid that encodes a component, e.g., all of the components, of the CAR.

In one embodiment, the agent which inhibits an inhibitory molecule comprises a first polypeptide, e.g., an inhibitory molecule, associated with a second polypeptide that provides a positive signal to the cell, e.g., an intracellular signaling domain described herein. In one embodiment, the agent comprises a first polypeptide, e.g., of an inhibitory molecule such as PD1, PD-L1, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GAL9, adenosine, or TGFR beta, or a fragment of any of these (e.g., at least a portion of an extracellular domain of any of these), and a second polypeptide which is an intracellular signaling domain described herein (e.g., comprising a costimulatory domain (e.g., 41BB, CD27 or CD28, e.g., as described herein) and/or a primary signaling domain (e.g., a CD3 zeta signaling domain described herein). In one embodiment, the agent comprises a first polypeptide of PD1 or a fragment thereof (e.g., at least a portion of an extracellular domain of PD1), and a second polypeptide of an intracellular signaling domain described herein (e.g., a

CD28 signaling domain described herein and/or a CD3 zeta signaling domain described herein). PD1 is an inhibitory member of the CD28 family of receptors that also includes CD28, CTLA-4, ICOS, and BTLA. PD-1 is expressed on activated B cells, T cells and myeloid cells (Agata et al. 1996 Int. Immunol 8:765-75). Two ligands for PD1, PD-L1 and PD-L2 have been shown to downregulate T cell activation upon binding to PD1 (Freeman et al. 2000 J Exp Med 192:1027-34; Latchman et al. 2001 Nat Immunol 2:261-8; Carter et al. 2002 Eur J Immunol 32:634-43). PD-L1 is abundant in human cancers (Dong et al. 2003 J Mol Med 81:281-7; Blank et al. 2005 Cancer Immunol. Immunother 54:307-314; Konishi et al. 2004 Clin Cancer Res 10:5094). Immune suppression can be reversed by inhibiting the local interaction of PD1 with PD-L1.

In one embodiment, the agent comprises the extracellular domain (ECD) of an inhibitory molecule, e.g., Programmed Death 1 (PD1), can be fused to a transmembrane domain and intracellular signaling domains such as 41BB and CD3 zeta (also referred to herein as a PD1 CAR). In one embodiment, the PD1 CAR, when used in combinations with an XCAR described herein, improves the persistence of the T cell. In one embodiment, the CAR is a PD1 CAR comprising the extracellular domain of PD1 indicated as underlined in SEQ ID NO: 105. In one embodiment, the PD1 CAR comprises the amino acid sequence of SEQ ID NO:105.

Malpytalllplalllhaarppgwflsdprwnpptfspallvvtgednatftcsfsntsesfvlnwyrmspsnqtdklaaf
pedrsqpgqdcfrvtqlpngrdfhmsvvrarrndsgtylcgaislapkaikeslraelrvterraevptahpspsprpagqfqlvt
 paprpptpaptiasqplslrpeacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgrkklyifkqpfmrpvqttqee
 dgcscrpfecceggcelrvkfsrsadapaykqgqnqlynelnlgreeydvldkrrgrdpemggkprknqpqeglynelqdkkma
 eayseigmkgerrrgkghdglyqglstatkdydalhmqalppr (SEQ ID NO:105).

In one embodiment, the PD1 CAR comprises the amino acid sequence provided below (SEQ ID NO:106).

pgwflsdprwnpptfspallvvtgednatftcsfsntsesfvlnwyrmspsnqtdklaafpedrsqpgqdcfrvtqlp
ngrdfhmsvvrarrndsgtylcgaislapkaikeslraelrvterraevptahpspsprpagqfqlvtttpparpptpaptiasqplslr
 eacrpaaggavhtrgldfacdiyiwaplagtcgvllslvitlyckrgrkklyifkqpfmrpvqttqeedgcscrpfecceggcelrvk
 fsrsadapaykqgqnqlynelnlgreeydvldkrrgrdpemggkprknqpqeglynelqdkkmaeayseigmkgerrrgkghdgl
 yqglstatkdydalhmqalppr (SEQ ID NO:106).

In one embodiment, the agent comprises a nucleic acid sequence encoding the PD1 CAR, e.g., the PD1 CAR described herein. In one embodiment, the nucleic acid sequence for the PD1 CAR is shown below, with the PD1 ECD underlined below in SEQ ID NO: 107

atggccctccctgtcactgcctgtcttccccctgcactcctgtccacgccgctagaccacccggatggttctggactctc
 5 cggatcgcccggtggaatcccccaaccttctaccggcactcttggttgtagctgagggcgataatgcgaccttcacgtgctcgttctccaa
cacctccgaatcattcgtgctgaactggtaccgcatgagcccgtaaaccagaccgacaagctcgccgcgttccgggaagatcggtcgc
aaccgggacagattgtcgggttccgcgtgactcaactgccgaatggcagagacttccacatgagcgtggtccgcgtagggcgaacga
ctccgggacctacctgtcgggagccatctcgctggcgccctaaggcccaaatcaagagagcctgagggccgaactgagagtgaccga
gcgcagagctgaggtgccaaactgcacatccatccccatcgccctggcctgcggggcagtttcagaccctggtcacgaccactccggcg
 10 ccgcgcccaccgactccggccccaactatcgcgagccagccctgtcgtgtagggccggaagcatgccgcctcgccgcggaggtgc
tgtgcataccggggattggacttcgcatgcgacatctacatttgggctcctctcgccggaacttgtggcgtgctccttctgtccctgtcat
cacctgtactgcaagcggggtcggaaaaagcttctgtacattttcaagcagcccttcatgagggccgtgcaaaccacccaggaggagg
acggttgctcctgccggttccccgaagaggaagaaggaggttcgagctgcgcgtgaagtctcccggagcgccgacgccccgcct
ataagcaggggcagaaccagctgtacaacgaactgaacctgggacggcggaagagtacgatgtgctggacaagcggcgcgccgc
 15 ggaccccgaaatggcggaagcctagaagaagaacctcaggaaggcctgtataacgagctgcagaaggacaagatggccgag
gcctactccgaaattgggatgaaggagagcggcgaggggaaaggggcacgacggcctgtaccaaggactgtccaccgccacca
aggacacatacatgatgcctgcacatgcaggccctccccctcgc (SEQ ID NO: 107).

In another example, in one embodiment, the agent which enhances the activity of a CAR-expressing cell can be a costimulatory molecule or costimulatory molecule ligand.

20 Examples of costimulatory molecules include MHC class I molecule, BTLA and a Toll ligand receptor, as well as OX40, CD27, CD28, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), and 4-1BB (CD137). Further examples of such costimulatory molecules include CDS, ICAM-1, GITR, BAFFR, HVEM (LIGHTR), SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46, CD160, CD19, CD4, CD8alpha, CD8beta, IL2R beta, IL2R gamma, IL7R

25 alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d, ITGAE, CD103, ITGAL, CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, NKG2D, NKG2C, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRTAM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108),

30 SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, and a ligand that specifically binds with CD83., e.g., as

described herein. Examples of costimulatory molecule ligands include CD80, CD86, CD40L, ICOSL, CD70, OX40L, 4-1BBL, GITRL, and LIGHT. In embodiments, the costimulatory molecule ligand is a ligand for a costimulatory molecule different from the costimulatory molecule domain of the CAR. In embodiments, the costimulatory molecule ligand is a ligand for a costimulatory molecule that is the same as the costimulatory molecule domain of the CAR. In an embodiment, the costimulatory molecule ligand is 4-1BBL. In an embodiment, the costimulatory ligand is CD80 or CD86. In an embodiment, the costimulatory molecule ligand is CD70. In embodiments, a CAR-expressing immune effector cell described herein can be further engineered to express one or more additional costimulatory molecules or costimulatory molecule ligands.

Co-expression of CAR with a Chemokine Receptor

In embodiments, the CAR-expressing cell described herein, e.g., CD19 CAR-expressing cell, further comprises a chemokine receptor molecule. Transgenic expression of chemokine receptors CCR2b or CXCR2 in T cells enhances trafficking to CCL2- or CXCL1-secreting solid tumors including melanoma and neuroblastoma (Craddock et al., *J Immunother.* 2010 Oct; 33(8):780-8 and Kershaw et al., *Hum Gene Ther.* 2002 Nov 1; 13(16):1971-80). Thus, without wishing to be bound by theory, it is believed that chemokine receptors expressed in CAR-expressing cells that recognize chemokines secreted by tumors, e.g., solid tumors, can improve homing of the CAR-expressing cell to the tumor, facilitate the infiltration of the CAR-expressing cell to the tumor, and enhances antitumor efficacy of the CAR-expressing cell. The chemokine receptor molecule can comprise a naturally occurring or recombinant chemokine receptor or a chemokine-binding fragment thereof. A chemokine receptor molecule suitable for expression in a CAR-expressing cell (e.g., CAR-Tx) described herein include a CXC chemokine receptor (e.g., CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, CXCR6, or CXCR7), a CC chemokine receptor (e.g., CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10, or CCR11), a CX3C chemokine receptor (e.g., CX3CR1), a XC chemokine receptor (e.g., XCR1), or a chemokine-binding fragment thereof. In one embodiment, the chemokine receptor molecule to be expressed with a CAR described herein is selected based on the chemokine(s) secreted by the tumor. In one embodiment, the CAR-expressing cell described herein further comprises, e.g., expresses, a CCR2b receptor or a CXCR2 receptor. In an embodiment, the CAR described herein and the chemokine receptor molecule are on the same vector or are on two different vectors. In embodiments where the CAR described herein

and the chemokine receptor molecule are on the same vector, the CAR and the chemokine receptor molecule are each under control of two different promoters or are under the control of the same promoter.

5 **Nucleic Acid Constructs Encoding a CAR**

The present invention also provides an immune effector cell, e.g., made by a method described herein, that includes a nucleic acid molecules encoding one or more CAR constructs described herein. In one aspect, the nucleic acid molecule is provided as a messenger RNA transcript. In one aspect, the nucleic acid molecule is provided as a DNA construct.

10 The nucleic acid molecules described herein can be a DNA molecule, an RNA molecule, or a combination thereof. In one embodiment, the nucleic acid molecule is an mRNA encoding a CAR polypeptide as described herein. In other embodiments, the nucleic acid molecule is a vector that includes any of the aforesaid nucleic acid molecules.

In one aspect, the antigen binding domain of a CAR of the invention (e.g., a scFv) is
15 encoded by a nucleic acid molecule whose sequence has been codon optimized for expression in a mammalian cell. In one aspect, entire CAR construct of the invention is encoded by a nucleic acid molecule whose entire sequence has been codon optimized for expression in a mammalian cell. Codon optimization refers to the discovery that the frequency of occurrence of synonymous codons (i.e., codons that code for the same amino acid) in coding DNA is
20 biased in different species. Such codon degeneracy allows an identical polypeptide to be encoded by a variety of nucleotide sequences. A variety of codon optimization methods is known in the art, and include, e.g., methods disclosed in at least US Patent Numbers 5,786,464 and 6,114,148.

Accordingly, in one aspect, an immune effector cell, e.g., made by a method described
25 herein, includes a nucleic acid molecule encoding a chimeric antigen receptor (CAR), wherein the CAR comprises an antigen binding domain that binds to a tumor antigen described herein, a transmembrane domain (e.g., a transmembrane domain described herein), and an intracellular signaling domain (e.g., an intracellular signaling domain described herein) comprising a stimulatory domain, e.g., a costimulatory signaling domain (e.g., a costimulatory signaling

domain described herein) and/or a primary signaling domain (e.g., a primary signaling domain described herein, e.g., a zeta chain described herein).

The present invention also provides vectors in which a nucleic acid molecule encoding a CAR, e.g., a nucleic acid molecule described herein, is inserted. Vectors derived from retroviruses such as the lentivirus are suitable tools to achieve long-term gene transfer since they allow long-term, stable integration of a transgene and its propagation in daughter cells. Lentiviral vectors have the added advantage over vectors derived from onco-retroviruses such as murine leukemia viruses in that they can transduce non-proliferating cells, such as hepatocytes. They also have the added advantage of low immunogenicity. A retroviral vector may also be, e.g., a gammaretroviral vector. A gammaretroviral vector may include, e.g., a promoter, a packaging signal (ψ), a primer binding site (PBS), one or more (e.g., two) long terminal repeats (LTR), and a transgene of interest, e.g., a gene encoding a CAR. A gammaretroviral vector may lack viral structural genes such as gag, pol, and env. Exemplary gammaretroviral vectors include Murine Leukemia Virus (MLV), Spleen-Focus Forming Virus (SFFV), and Myeloproliferative Sarcoma Virus (MPSV), and vectors derived therefrom. Other gammaretroviral vectors are described, e.g., in Tobias Maetzig et al., "Gammaretroviral Vectors: Biology, Technology and Application" *Viruses*. 2011 Jun; 3(6): 677–713.

In another embodiment, the vector comprising the nucleic acid encoding the desired CAR is an adenoviral vector (A5/35). In another embodiment, the expression of nucleic acids encoding CARs can be accomplished using of transposons such as sleeping beauty, crisper, CAS9, and zinc finger nucleases. See below June et al. 2009 *Nature Reviews Immunology* 9.10: 704-716, is incorporated herein by reference.

In brief summary, the expression of natural or synthetic nucleic acids encoding CARs is typically achieved by operably linking a nucleic acid encoding the CAR polypeptide or portions thereof to a promoter, and incorporating the construct into an expression vector. The vectors can be suitable for replication and integration eukaryotes. Typical cloning vectors contain transcription and translation terminators, initiation sequences, and promoters useful for regulation of the expression of the desired nucleic acid sequence.

The nucleic acid can be cloned into a number of types of vectors. For example, the nucleic acid can be cloned into a vector including, but not limited to a plasmid, a phagemid, a

phage derivative, an animal virus, and a cosmid. Vectors of particular interest include expression vectors, replication vectors, probe generation vectors, and sequencing vectors.

Further, the expression vector may be provided to a cell in the form of a viral vector. Viral vector technology is well known in the art and is described, for example, in Sambrook et al., 2012, MOLECULAR CLONING: A LABORATORY MANUAL, volumes 1 -4, Cold Spring Harbor Press, NY), and in other virology and molecular biology manuals. Viruses, which are useful as vectors include, but are not limited to, retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, and lentiviruses. In general, a suitable vector contains an origin of replication functional in at least one organism, a promoter sequence, convenient restriction endonuclease sites, and one or more selectable markers, (e.g., WO 01/96584; WO 01/29058; and U.S. Pat. No. 6,326,193).

A number of viral based systems have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene delivery systems. A selected gene can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to cells of the subject either in vivo or ex vivo. A number of retroviral systems are known in the art. In some embodiments, adenovirus vectors are used. A number of adenovirus vectors are known in the art. In one embodiment, lentivirus vectors are used.

Additional promoter elements, e.g., enhancers, regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the thymidine kinase (tk) promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription. Exemplary promoters include the CMV IE gene, EF-1 α , ubiquitin C, or phosphoglycerokinase (PGK) promoters.

An example of a promoter that is capable of expressing a CAR encoding nucleic acid molecule in a mammalian T cell is the EF1 α promoter. The native EF1 α promoter drives expression of the alpha subunit of the elongation factor-1 complex, which is responsible for the

enzymatic delivery of aminoacyl tRNAs to the ribosome. The EF1a promoter has been extensively used in mammalian expression plasmids and has been shown to be effective in driving CAR expression from nucleic acid molecules cloned into a lentiviral vector. See, e.g., Milone et al., *Mol. Ther.* 17(8): 1453–1464 (2009). In one aspect, the EF1a promoter
 5 comprises the sequence provided in the Examples.

Another example of a promoter is the immediate early cytomegalovirus (CMV) promoter sequence. This promoter sequence is a strong constitutive promoter sequence capable of driving high levels of expression of any polynucleotide sequence operatively linked thereto. However, other constitutive promoter sequences may also be used, including, but not limited to
 10 the simian virus 40 (SV40) early promoter, mouse mammary tumor virus (MMTV), human immunodeficiency virus (HIV) long terminal repeat (LTR) promoter, MoMuLV promoter, an avian leukemia virus promoter, an Epstein-Barr virus immediate early promoter, a Rous sarcoma virus promoter, as well as human gene promoters such as, but not limited to, the actin promoter, the myosin promoter, the elongation factor-1 α promoter, the hemoglobin promoter,
 15 and the creatine kinase promoter. Further, the invention should not be limited to the use of constitutive promoters. Inducible promoters are also contemplated as part of the invention. The use of an inducible promoter provides a molecular switch capable of turning on expression of the polynucleotide sequence which it is operatively linked when such expression is desired, or turning off the expression when expression is not desired. Examples of inducible promoters
 20 include, but are not limited to a metallothionine promoter, a glucocorticoid promoter, a progesterone promoter, and a tetracycline promoter.

Another example of a promoter is the phosphoglycerate kinase (PGK) promoter. In embodiments, a truncated PGK promoter (e.g., a PGK promoter with one or more, e.g., 1, 2, 5, 10, 100, 200, 300, or 400, nucleotide deletions when compared to the wild-type PGK promoter
 25 sequence) may be desired.

The nucleotide sequences of exemplary PGK promoters are provided below.

WT PGK Promoter:

ACCCCTCTCTCCAGCCACTAAGCCAGTTGCTCCCTCGGCTGACGGCTGCACG
 CGAGGCCTCCGAACGTCTTACGCCCTTGTGGCGCGCCCGTCCTTGTCCCGGGTGTGA
 30 TGGCGGGGTGTGGGGCGGAGGGCGTGGCGGGGAAGGGCCGGCGACGAGAGCCGC

GCGGGACGACTCGTCGGCGATAACCGGTGTCGGGTAGCGCCAGCCGCGCGACGGT
AACGAGGGACCGCGACAGGCAGACGCTCCCATGATCACTCTGCACGCCGAAGGCA
AATAGTGCAGGCCGTGCGGCGCTTGGCGTTCCTTGGAAGGGCTGAATCCCCGCCTC
GTCCTTCGCAGCGGCCCCCGGGTGTTCCTATCGCCGCTTCTAGGCCCACTGCGAC
5 GCTTGCCTGCACTTCTTACACGCTCTGGGTCCCAGCCGCGGCGACGCAAAGGGCCT
TGGTGCGGGTCTCGTCGGCGCAGGGACGCGTTTGGGTCCCGACGGAACCTTTCCG
CGTTGGGGTTGGGGCACCATAAGCT (SEQ ID NO: 109)

Exemplary truncated PGK Promoters:

PGK100:

10 ACCCTCTCTCCAGCCACTAAGCCAGTTGCTCCCTCGGCTGACGGCTGCACG
CGAGGCCTCCGAACGTCTTACGCTTGTGGCGCGCCCGTCCTTGTCCCGGGTGTGA
TGGCGGGGTG (SEQ ID NO: 110)

PGK200:

15 ACCCTCTCTCCAGCCACTAAGCCAGTTGCTCCCTCGGCTGACGGCTGCACG
CGAGGCCTCCGAACGTCTTACGCTTGTGGCGCGCCCGTCCTTGTCCCGGGTGTGA
TGGCGGGGTGTGGGGCGGAGGGCGTGGCGGGGAAGGGCCGGCGACGAGAGCCGC
GCGGGACGACTCGTCGGCGATAACCGGTGTCGGGTAGCGCCAGCCGCGCGACGGT
AACG (SEQ ID NO:111)

PGK300:

20 ACCCTCTCTCCAGCCACTAAGCCAGTTGCTCCCTCGGCTGACGGCTGCACG
CGAGGCCTCCGAACGTCTTACGCTTGTGGCGCGCCCGTCCTTGTCCCGGGTGTGA
TGGCGGGGTGTGGGGCGGAGGGCGTGGCGGGGAAGGGCCGGCGACGAGAGCCGC
GCGGGACGACTCGTCGGCGATAACCGGTGTCGGGTAGCGCCAGCCGCGCGACGGT
AACGAGGGACCGCGACAGGCAGACGCTCCCATGATCACTCTGCACGCCGAAGGCA
25 AATAGTGCAGGCCGTGCGGCGCTTGGCGTTCCTTGGAAGGGCTGAATCCCCG (SEQ
ID NO:112)

PGK400:

ACCCTCTCTCCAGCCACTAAGCCAGTTGCTCCCTCGGCTGACGGCTGCACG
CGAGGCCTCCGAACGTCTTACGCTTGTGGCGCGCCCGTCCTTGTCCCGGGTGTGA

TGGCGGGGTGTGGGGCGGAGGGCGTGGCGGGGAAGGGCCGGCGACGAGAGCCGC
 GCGGGACGACTCGTCGGCGATAACCGGTGTCGGGTAGCGCCAGCCGCGCGACGGT
 AACGAGGGACCGCGACAGGCAGACGCTCCCATGATCACTCTGCACGCCGAAGGCA
 AATAGTGCAGGCCGTGCGGCGCTTGGCGTTCCTTGGAAGGGCTGAATCCCCGCCTC
 5 GTCCTTCGCAGCGGCCCCCGGGTGTTCCTTCGCGCTTCTAGGCCCACTGCGAC
 GCTTGCCTGCACTTCTTACACGCTCTGGGTCCCAGCCG (SEQ ID NO:113)

A vector may also include, e.g., a signal sequence to facilitate secretion, a
 polyadenylation signal and transcription terminator (e.g., from Bovine Growth Hormone
 10 (BGH) gene), an element allowing episomal replication and replication in prokaryotes (e.g.
 SV40 origin and ColE1 or others known in the art) and/or elements to allow selection (e.g.,
 ampicillin resistance gene and/or zeocin marker).

In order to assess the expression of a CAR polypeptide or portions thereof, the
 expression vector to be introduced into a cell can also contain either a selectable marker gene or
 15 a reporter gene or both to facilitate identification and selection of expressing cells from the
 population of cells sought to be transfected or infected through viral vectors. In other aspects,
 the selectable marker may be carried on a separate piece of DNA and used in a co- transfection
 procedure. Both selectable markers and reporter genes may be flanked with appropriate
 regulatory sequences to enable expression in the host cells. Useful selectable markers include,
 20 for example, antibiotic-resistance genes, such as neo and the like.

Reporter genes are used for identifying potentially transfected cells and for evaluating
 the functionality of regulatory sequences. In general, a reporter gene is a gene that is not
 present in or expressed by the recipient organism or tissue and that encodes a polypeptide
 whose expression is manifested by some easily detectable property, e.g., enzymatic activity.
 25 Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced
 into the recipient cells. Suitable reporter genes may include genes encoding luciferase, beta-
 galactosidase, chloramphenicol acetyl transferase, secreted alkaline phosphatase, or the green
 fluorescent protein gene (e.g., Ui-Tei et al., 2000 FEBS Letters 479: 79-82). Suitable
 expression systems are well known and may be prepared using known techniques or obtained
 30 commercially. In general, the construct with the minimal 5' flanking region showing the highest
 level of expression of reporter gene is identified as the promoter. Such promoter regions may

be linked to a reporter gene and used to evaluate agents for the ability to modulate promoter-driven transcription.

In embodiments, the vector may comprise two or more nucleic acid sequences encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR, and a second CAR, e.g., an inhibitory
5 CAR or a CAR that specifically binds to an antigen other than CD19. In such embodiments, the two or more nucleic acid sequences encoding the CAR are encoded by a single nucleic molecule in the same frame and as a single polypeptide chain. In this aspect, the two or more CARs, can, e.g., be separated by one or more peptide cleavage sites. (e.g., an auto-cleavage site or a substrate for an intracellular protease). Examples of peptide cleavage sites include T2A,
10 P2A, E2A, or F2A sites.

Methods of introducing and expressing genes into a cell are known in the art. In the context of an expression vector, the vector can be readily introduced into a host cell, e.g., mammalian, bacterial, yeast, or insect cell by any method, e.g., one known in the art. For example, the expression vector can be transferred into a host cell by physical, chemical, or
15 biological means.

Physical methods for introducing a polynucleotide into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Methods for producing cells comprising vectors and/or exogenous nucleic acids are well-known in the art. See, for example, Sambrook et al., 2012, MOLECULAR CLONING: A
20 LABORATORY MANUAL, volumes 1 -4, Cold Spring Harbor Press, NY). A suitable method for the introduction of a polynucleotide into a host cell is calcium phosphate transfection.

Biological methods for introducing a polynucleotide of interest into a host cell include the use of DNA and RNA vectors. Viral vectors, and especially retroviral vectors, have become the most widely used method for inserting genes into mammalian, e.g., human cells. Other viral
25 vectors can be derived from lentivirus, poxviruses, herpes simplex virus I, adenoviruses and adeno-associated viruses, and the like. See, for example, U.S. Pat. Nos. 5,350,674 and 5,585,362.

Chemical means for introducing a polynucleotide into a host cell include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and
30 lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes.

An exemplary colloidal system for use as a delivery vehicle in vitro and in vivo is a liposome (e.g., an artificial membrane vesicle). Other methods of state-of-the-art targeted delivery of nucleic acids are available, such as delivery of polynucleotides with targeted nanoparticles or other suitable sub-micron sized delivery system.

5 In the case where a non-viral delivery system is utilized, an exemplary delivery vehicle is a liposome. The use of lipid formulations is contemplated for the introduction of the nucleic acids into a host cell (in vitro, ex vivo or in vivo). In another aspect, the nucleic acid may be associated with a lipid. The nucleic acid associated with a lipid may be encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a
10 liposome via a linking molecule that is associated with both the liposome and the oligonucleotide, entrapped in a liposome, complexed with a liposome, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid. Lipid, lipid/DNA or lipid/expression vector associated compositions are not limited to any particular
15 structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a “collapsed” structure. They may also simply be interspersed in a solution, possibly forming aggregates that are not uniform in size or shape. Lipids are fatty substances which may be naturally occurring or synthetic lipids. For example, lipids include the fatty droplets that naturally occur in the cytoplasm as well as the class of compounds which contain long-chain
20 aliphatic hydrocarbons and their derivatives, such as fatty acids, alcohols, amines, amino alcohols, and aldehydes.

Lipids suitable for use can be obtained from commercial sources. For example, dimyristyl phosphatidylcholine (“DMPC”) can be obtained from Sigma, St. Louis, MO; dicetyl phosphate (“DCP”) can be obtained from K & K Laboratories (Plainview, NY); cholesterol
25 (“Choi”) can be obtained from Calbiochem-Behring; dimyristyl phosphatidylglycerol (“DMPG”) and other lipids may be obtained from Avanti Polar Lipids, Inc. (Birmingham, AL.). Stock solutions of lipids in chloroform or chloroform/methanol can be stored at about - 20°C. Chloroform is used as the only solvent since it is more readily evaporated than methanol. “Liposome” is a generic term encompassing a variety of single and multilamellar lipid vehicles
30 formed by the generation of enclosed lipid bilayers or aggregates. Liposomes can be characterized as having vesicular structures with a phospholipid bilayer membrane and an inner

aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh et al., 1991
5 Glycobiology 5: 505-10). However, compositions that have different structures in solution than the normal vesicular structure are also encompassed. For example, the lipids may assume a micellar structure or merely exist as nonuniform aggregates of lipid molecules. Also contemplated are lipofectamine-nucleic acid complexes.

Regardless of the method used to introduce exogenous nucleic acids into a host cell or
10 otherwise expose a cell to the inhibitor of the present invention, in order to confirm the presence of the recombinant nucleic acid sequence in the host cell, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; "biochemical" assays, such as detecting the presence or absence of a particular peptide, e.g., by
15 immunological means (ELISAs and Western blots) or by assays described herein to identify agents falling within the scope of the invention.

Natural Killer Cell Receptor (NKR) CARs

In an embodiment, the CAR molecule described herein comprises one or more components of a natural killer cell receptor (NKR), thereby forming an NKR-CAR. The NKR
20 component can be a transmembrane domain, a hinge domain, or a cytoplasmic domain from any of the following natural killer cell receptors: killer cell immunoglobulin-like receptor (KIR), e.g., KIR2DL1, KIR2DL2/L3, KIR2DL4, KIR2DL5A, KIR2DL5B, KIR2DS1, KIR2DS2, KIR2DS3, KIR2DS4, KIR2DS5, KIR3DL1/S1, KIR3DL2, KIR3DL3, KIR2DP1, and KIR3DP1; natural cytotoxicity receptor (NCR), e.g., NKp30, NKp44, NKp46; signaling
25 lymphocyte activation molecule (SLAM) family of immune cell receptors, e.g., CD48, CD229, 2B4, CD84, NTB-A, CRACC, BLAME, and CD2F-10; Fc receptor (FcR), e.g., CD16, and CD64; and Ly49 receptors, e.g., LY49A, LY49C. The NKR-CAR molecules described herein may interact with an adaptor molecule or intracellular signaling domain, e.g., DAP12. Exemplary configurations and sequences of CAR molecules comprising NKR components are
30 described in International Publication No. WO2014/145252, the contents of which are hereby incorporated by reference.

Split CAR

In some embodiments, the CAR-expressing cell uses a split CAR. The split CAR approach is described in more detail in publications WO2014/055442 and WO2014/055657. Briefly, a split CAR system comprises a cell expressing a first CAR having a first antigen
5 binding domain and a costimulatory domain (e.g., 41BB), and the cell also expresses a second CAR having a second antigen binding domain and an intracellular signaling domain (e.g., CD3 zeta). When the cell encounters the first antigen, the costimulatory domain is activated, and the cell proliferates. When the cell encounters the second antigen, the intracellular signaling domain is activated and cell-killing activity begins. Thus, the CAR-expressing cell is only fully
10 activated in the presence of both antigens.

Strategies for Regulating Chimeric Antigen Receptors

In some embodiments, a regulatable CAR (RCAR) where the CAR activity can be controlled is desirable to optimize the safety and efficacy of a CAR therapy. There are many ways CAR activities can be regulated. For example, inducible apoptosis using, e.g., a caspase
15 fused to a dimerization domain (see, e.g., Di Stasa et al., N Engl. J. Med. 2011 Nov. 3; 365(18):1673-1683), can be used as a safety switch in the CAR therapy of the instant invention. In one embodiment, the cells (e.g., T cells or NK cells) expressing a CAR of the present invention further comprise an inducible apoptosis switch, wherein a human caspase (e.g., caspase 9) or a modified version is fused to a modification of the human FKB protein that
20 allows conditional dimerization. In the presence of a small molecule, such as a rapalog (e.g., AP 1903, AP20187), the inducible caspase (e.g., caspase 9) is activated and leads to the rapid apoptosis and death of the cells (e.g., T cells or NK cells) expressing a CAR of the present invention. Examples of a caspase-based inducible apoptosis switch (or one or more aspects of such a switch) have been described in, e.g., US2004040047; US20110286980;
25 US20140255360; WO1997031899; WO2014151960; WO2014164348; WO2014197638; WO2014197638; all of which are incorporated by reference herein.

In another example, CAR-expressing cells can also express an inducible Caspase-9 (iCaspase-9) molecule that, upon administration of a dimerizer drug (e.g., rimiducid (also called AP1903 (Bellicum Pharmaceuticals) or AP20187 (Ariad)) leads to activation of the
30 Caspase-9 and apoptosis of the cells. The iCaspase-9 molecule contains a chemical inducer of dimerization (CID) binding domain that mediates dimerization in the presence of a CID. This

results in inducible and selective depletion of CAR-expressing cells. In some cases, the iCaspase-9 molecule is encoded by a nucleic acid molecule separate from the CAR-encoding vector(s). In some cases, the iCaspase-9 molecule is encoded by the same nucleic acid molecule as the CAR-encoding vector. The iCaspase-9 can provide a safety switch to avoid any toxicity of CAR-expressing cells. See, e.g., Song et al. *Cancer Gene Ther.* 2008; 15(10):667-75; Clinical Trial Id. No. NCT02107963; and Di Stasi et al. *N. Engl. J. Med.* 2011; 365:1673-83.

Alternative strategies for regulating the CAR therapy of the instant invention include utilizing small molecules or antibodies that deactivate or turn off CAR activity, e.g., by deleting CAR-expressing cells, e.g., by inducing antibody dependent cell-mediated cytotoxicity (ADCC). For example, CAR-expressing cells described herein may also express an antigen that is recognized by molecules capable of inducing cell death, e.g., ADCC or complement-induced cell death. For example, CAR expressing cells described herein may also express a receptor capable of being targeted by an antibody or antibody fragment. Examples of such receptors include EpCAM, VEGFR, integrins (e.g., integrins $\alpha\beta3$, $\alpha4$, $\alpha1\beta3$, $\alpha4\beta7$, $\alpha5\beta1$, $\alpha\nu\beta3$, $\alpha\nu$), members of the TNF receptor superfamily (e.g., TRAIL-R1, TRAIL-R2), PDGF Receptor, interferon receptor, folate receptor, GPNMB, ICAM-1, HLA-DR, CEA, CA-125, MUC1, TAG-72, IL-6 receptor, 5T4, GD2, GD3, CD2, CD3, CD4, CD5, CD11, CD11a/LFA-1, CD15, CD18/ITGB2, CD19, CD20, CD22, CD23/IgE Receptor, CD25, CD28, CD30, CD33, CD38, CD40, CD41, CD44, CD51, CD52, CD62L, CD74, CD80, CD125, CD147/basigin, CD152/CTLA-4, CD154/CD40L, CD195/CCR5, CD319/SLAMF7, and EGFR, and truncated versions thereof (e.g., versions preserving one or more extracellular epitopes but lacking one or more regions within the cytoplasmic domain).

For example, a CAR-expressing cell described herein may also express a truncated epidermal growth factor receptor (EGFR) which lacks signaling capacity but retains the epitope that is recognized by molecules capable of inducing ADCC, e.g., cetuximab (ERBITUX®), such that administration of cetuximab induces ADCC and subsequent depletion of the CAR-expressing cells (see, e.g., WO2011/056894, and Jonnalagadda et al., *Gene Ther.* 2013; 20(8):853-860). Another strategy includes expressing a highly compact marker/suicide gene that combines target epitopes from both CD32 and CD20 antigens in the CAR-expressing cells described herein, which binds rituximab, resulting in selective depletion of the CAR-expressing

cells, e.g., by ADCC (see, e.g., Philip et al., Blood. 2014; 124(8)1277-1287). Other methods for depleting CAR-expressing cells described herein include administration of CAMPATH, a monoclonal anti-CD52 antibody that selectively binds and targets mature lymphocytes, e.g., CAR-expressing cells, for destruction, e.g., by inducing ADCC. In other embodiments, the CAR-expressing cell can be selectively targeted using a CAR ligand, e.g., an anti-idiotypic antibody. In some embodiments, the anti-idiotypic antibody can cause effector cell activity, e.g., ADCC or ADC activities, thereby reducing the number of CAR-expressing cells. In other embodiments, the CAR ligand, e.g., the anti-idiotypic antibody, can be coupled to an agent that induces cell killing, e.g., a toxin, thereby reducing the number of CAR-expressing cells.

10 Alternatively, the CAR molecules themselves can be configured such that the activity can be regulated, e.g., turned on and off, as described below.

In other embodiments, a CAR-expressing cell described herein may also express a target protein recognized by the T cell depleting agent. In one embodiment, the target protein is CD20 and the T cell depleting agent is an anti-CD20 antibody, e.g., rituximab. In such embodiment, the T cell depleting agent is administered once it is desirable to reduce or eliminate the CAR-expressing cell, e.g., to mitigate the CAR induced toxicity. In other embodiments, the T cell depleting agent is an anti-CD52 antibody, e.g., alemtuzumab, as described in the Examples herein.

15

In other embodiments, an RCAR comprises a set of polypeptides, typically two in the simplest embodiments, in which the components of a standard CAR described herein, e.g., an antigen binding domain and an intracellular signalling domain, are partitioned on separate polypeptides or members. In some embodiments, the set of polypeptides include a dimerization switch that, upon the presence of a dimerization molecule, can couple the polypeptides to one another, e.g., can couple an antigen binding domain to an intracellular signalling domain. In one embodiment, a CAR of the present invention utilizes a dimerization switch as those described in, e.g., WO2014127261, which is incorporated by reference herein. Additional description and exemplary configurations of such regulatable CARs are provided herein and in, e.g., paragraphs 527-551 of International Publication No. WO 2015/090229 filed March 13, 2015, which is incorporated by reference in its entirety. In some embodiments, an RCAR involves a switch domain, e.g., a FKBP switch domain, as set out SEQ ID NO: 114, or comprise a fragment of FKBP having the ability to bind with FRB, e.g., as set out in SEQ ID

20

25

30

NO: 115. In some embodiments, the RCAR involves a switch domain comprising a FRB sequence, e.g., as set out in SEQ ID NO: 116, or a mutant FRB sequence, e.g., as set out in any of SEQ ID Nos. 117-122.

DVPDYASLGGPSSPKKKRKVSRGVQVETISPGDGRTPFKRGQTCVVHYTGMLE
5 DGKKFDSSSRDRNKPFFKMLGKQEVIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHP
GIIPPHATLVFDVELLKLETSY (SEQ ID NO: 114)

VQVETISPGDGRTPFKRGQTCVVHYTGMLEDGKKFDSSSRDRNKPFFKMLGKQE
VIRGWEEGVAQMSVGQRAKLTISPDYAYGATGHPGIIPPHATLVFDVELLKLETS (SEQ
ID NO: 115)

10 ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQ
AYGRDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRISK (SEQ ID NO: 116)

Table 4. Exemplary mutant FRB having increased affinity for a dimerization molecule.

FRB mutant	Amino Acid Sequence	SEQ ID NO:
E2032I mutant	ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRISKTS	117
E2032L mutant	ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDLTQAWDLYYHVFRISKTS	118
T2098L mutant	ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDLLQAWDLYYHVFRISKTS	119
E2032, T2098 mutant	ILWHEMWHEGL X EASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDL X QAWDLYYHVFRISKTS	120
E2032I, T2098L mutant	ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDLLQAWDLYYHVFRISKTS	121
E2032L, T2098L mutant	ILWHEMWHEGLEEASRLYFGERNVKGMEFEVLEPLHAMMERGPQTLKETSFNQAYG RDLMEAQEWCRKYMKSGNVKDLLQAWDLYYHVFRISKTS	122

15 RNA Transfection

Disclosed herein are methods for producing an in vitro transcribed RNA CAR. RNA CAR and methods of using the same are described, e.g., in paragraphs 553-570 of in International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

An immune effector cell can include a CAR encoded by a messenger RNA (mRNA). In one aspect, the mRNA encoding a CAR described herein is introduced into an immune effector cell, e.g., made by a method described herein, for production of a CAR-expressing cell.

In one embodiment, the *in vitro* transcribed RNA CAR can be introduced to a cell as a form of transient transfection. The RNA is produced by *in vitro* transcription using a polymerase chain reaction (PCR)-generated template. DNA of interest from any source can be directly converted by PCR into a template for *in vitro* mRNA synthesis using appropriate primers and RNA polymerase. The source of the DNA can be, for example, genomic DNA, plasmid DNA, phage DNA, cDNA, synthetic DNA sequence or any other appropriate source of DNA. The desired template for *in vitro* transcription is a CAR described herein. For example, the template for the RNA CAR comprises an extracellular region comprising a single chain variable domain of an antibody to a tumor associated antigen described herein; a hinge region (e.g., a hinge region described herein), a transmembrane domain (e.g., a transmembrane domain described herein such as a transmembrane domain of CD8a); and a cytoplasmic region that includes an intracellular signaling domain, e.g., an intracellular signaling domain described herein, e.g., comprising the signaling domain of CD3-zeta and the signaling domain of 4-1BB.

In one embodiment, the DNA to be used for PCR contains an open reading frame. The DNA can be from a naturally occurring DNA sequence from the genome of an organism. In one embodiment, the nucleic acid can include some or all of the 5' and/or 3' untranslated regions (UTRs). The nucleic acid can include exons and introns. In one embodiment, the DNA to be used for PCR is a human nucleic acid sequence. In another embodiment, the DNA to be used for PCR is a human nucleic acid sequence including the 5' and 3' UTRs. The DNA can alternatively be an artificial DNA sequence that is not normally expressed in a naturally occurring organism. An exemplary artificial DNA sequence is one that contains portions of genes that are ligated together to form an open reading frame that encodes a fusion protein. The portions of DNA that are ligated together can be from a single organism or from more than one organism.

PCR is used to generate a template for *in vitro* transcription of mRNA which is used for transfection. Methods for performing PCR are well known in the art. Primers for use in PCR are designed to have regions that are substantially complementary to regions of the DNA to be used as a template for the PCR. "Substantially complementary," as used herein, refers to

sequences of nucleotides where a majority or all of the bases in the primer sequence are complementary, or one or more bases are non-complementary, or mismatched. Substantially complementary sequences are able to anneal or hybridize with the intended DNA target under annealing conditions used for PCR. The primers can be designed to be substantially
5 complementary to any portion of the DNA template. For example, the primers can be designed to amplify the portion of a nucleic acid that is normally transcribed in cells (the open reading frame), including 5' and 3' UTRs. The primers can also be designed to amplify a portion of a nucleic acid that encodes a particular domain of interest. In one embodiment, the primers are designed to amplify the coding region of a human cDNA, including all or portions of the 5' and
10 3' UTRs. Primers useful for PCR can be generated by synthetic methods that are well known in the art. "Forward primers" are primers that contain a region of nucleotides that are substantially complementary to nucleotides on the DNA template that are upstream of the DNA sequence that is to be amplified. "Upstream" is used herein to refer to a location 5' to the DNA sequence to be amplified relative to the coding strand. "Reverse primers" are primers that contain a
15 region of nucleotides that are substantially complementary to a double-stranded DNA template that are downstream of the DNA sequence that is to be amplified. "Downstream" is used herein to refer to a location 3' to the DNA sequence to be amplified relative to the coding strand.

Any DNA polymerase useful for PCR can be used in the methods disclosed herein. The reagents and polymerase are commercially available from a number of sources.

20 Chemical structures with the ability to promote stability and/or translation efficiency may also be used. The RNA in embodiments has 5' and 3' UTRs. In one embodiment, the 5' UTR is between one and 3000 nucleotides in length. The length of 5' and 3' UTR sequences to be added to the coding region can be altered by different methods, including, but not limited to, designing primers for PCR that anneal to different regions of the UTRs. Using this approach,
25 one of ordinary skill in the art can modify the 5' and 3' UTR lengths required to achieve optimal translation efficiency following transfection of the transcribed RNA.

The 5' and 3' UTRs can be the naturally occurring, endogenous 5' and 3' UTRs for the nucleic acid of interest. Alternatively, UTR sequences that are not endogenous to the nucleic acid of interest can be added by incorporating the UTR sequences into the forward and reverse
30 primers or by any other modifications of the template. The use of UTR sequences that are not endogenous to the nucleic acid of interest can be useful for modifying the stability and/or

translation efficiency of the RNA. For example, it is known that AU-rich elements in 3' UTR sequences can decrease the stability of mRNA. Therefore, 3' UTRs can be selected or designed to increase the stability of the transcribed RNA based on properties of UTRs that are well known in the art.

5 In one embodiment, the 5' UTR can contain the Kozak sequence of the endogenous nucleic acid. Alternatively, when a 5' UTR that is not endogenous to the nucleic acid of interest is being added by PCR as described above, a consensus Kozak sequence can be redesigned by adding the 5' UTR sequence. Kozak sequences can increase the efficiency of translation of some RNA transcripts, but does not appear to be required for all RNAs to enable efficient
10 translation. The requirement for Kozak sequences for many mRNAs is known in the art. In other embodiments the 5' UTR can be 5'UTR of an RNA virus whose RNA genome is stable in cells. In other embodiments various nucleotide analogues can be used in the 3' or 5' UTR to impede exonuclease degradation of the mRNA.

To enable synthesis of RNA from a DNA template without the need for gene cloning, a
15 promoter of transcription should be attached to the DNA template upstream of the sequence to be transcribed. When a sequence that functions as a promoter for an RNA polymerase is added to the 5' end of the forward primer, the RNA polymerase promoter becomes incorporated into the PCR product upstream of the open reading frame that is to be transcribed. In one embodiment, the promoter is a T7 polymerase promoter, as described elsewhere herein. Other
20 useful promoters include, but are not limited to, T3 and SP6 RNA polymerase promoters. Consensus nucleotide sequences for T7, T3 and SP6 promoters are known in the art.

In an embodiment, the mRNA has both a cap on the 5' end and a 3' poly(A) tail which determine ribosome binding, initiation of translation and stability mRNA in the cell. On a circular DNA template, for instance, plasmid DNA, RNA polymerase produces a long
25 concatameric product which is not suitable for expression in eukaryotic cells. The transcription of plasmid DNA linearized at the end of the 3' UTR results in normal sized mRNA which is not effective in eukaryotic transfection even if it is polyadenylated after transcription.

On a linear DNA template, phage T7 RNA polymerase can extend the 3' end of the transcript beyond the last base of the template (Schenborn and Mierendorf, *Nuc Acids Res.*,
30 13:6223-36 (1985); Nacheva and Berzal-Herranz, *Eur. J. Biochem.*, 270:1485-65 (2003).

The conventional method of integration of polyA/T stretches into a DNA template is molecular cloning. However polyA/T sequence integrated into plasmid DNA can cause plasmid instability, which is why plasmid DNA templates obtained from bacterial cells are often highly contaminated with deletions and other aberrations. This makes cloning procedures not only laborious and time consuming but often not reliable. That is why a method which allows construction of DNA templates with polyA/T 3' stretch without cloning highly desirable.

The polyA/T segment of the transcriptional DNA template can be produced during PCR by using a reverse primer containing a polyT tail, such as 100T tail (SEQ ID NO: 31) (size can be 50-5000 T (SEQ ID NO: 32)), or after PCR by any other method, including, but not limited to, DNA ligation or in vitro recombination. Poly(A) tails also provide stability to RNAs and reduce their degradation. Generally, the length of a poly(A) tail positively correlates with the stability of the transcribed RNA. In one embodiment, the poly(A) tail is between 100 and 5000 adenosines (e.g., SEQ ID NO: 33).

Poly(A) tails of RNAs can be further extended following in vitro transcription with the use of a poly(A) polymerase, such as E. coli polyA polymerase (E-PAP). In one embodiment, increasing the length of a poly(A) tail from 100 nucleotides to between 300 and 400 nucleotides (SEQ ID NO: 34) results in about a two-fold increase in the translation efficiency of the RNA. Additionally, the attachment of different chemical groups to the 3' end can increase mRNA stability. Such attachment can contain modified/artificial nucleotides, aptamers and other compounds. For example, ATP analogs can be incorporated into the poly(A) tail using poly(A) polymerase. ATP analogs can further increase the stability of the RNA.

5' caps on also provide stability to RNA molecules. In an embodiment, RNAs produced by the methods disclosed herein include a 5' cap. The 5' cap is provided using techniques known in the art and described herein (Cougot, et al., Trends in Biochem. Sci., 29:436-444 (2001); Stepinski, et al., RNA, 7:1468-95 (2001); Elango, et al., Biochim. Biophys. Res. Commun., 330:958-966 (2005)).

The RNAs produced by the methods disclosed herein can also contain an internal ribosome entry site (IRES) sequence. The IRES sequence may be any viral, chromosomal or artificially designed sequence which initiates cap-independent ribosome binding to mRNA and facilitates the initiation of translation. Any solutes suitable for cell electroporation, which can

contain factors facilitating cellular permeability and viability such as sugars, peptides, lipids, proteins, antioxidants, and surfactants can be included.

RNA can be introduced into target cells using any of a number of different methods, for instance, commercially available methods which include, but are not limited to, electroporation
5 (Amaxa Nucleofector-II (Amaxa Biosystems, Cologne, Germany)), (ECM 830 (BTX) (Harvard Instruments, Boston, Mass.) or the Gene Pulser II (BioRad, Denver, Colo.), Multiporator (Eppendorf, Hamburg Germany), cationic liposome mediated transfection using lipofection, polymer encapsulation, peptide mediated transfection, or biolistic particle delivery systems such as “gene guns” (see, for example, Nishikawa, et al. Hum Gene Ther., 12(8):861-70
10 (2001).

Non-viral delivery methods

In some aspects, non-viral methods can be used to deliver a nucleic acid encoding a CAR described herein into a cell or tissue or a subject.

In some embodiments, the non-viral method includes the use of a transposon (also
15 called a transposable element). In some embodiments, a transposon is a piece of DNA that can insert itself at a location in a genome, for example, a piece of DNA that is capable of self-replicating and inserting its copy into a genome, or a piece of DNA that can be spliced out of a longer nucleic acid and inserted into another place in a genome. For example, a transposon comprises a DNA sequence made up of inverted repeats flanking genes for transposition.

Exemplary methods of nucleic acid delivery using a transposon include a Sleeping
20 Beauty transposon system (SBTS) and a piggyBac (PB) transposon system. See, e.g., Aronovich et al. Hum. Mol. Genet. 20.R1(2011):R14-20; Singh et al. Cancer Res. 15(2008):2961–2971; Huang et al. Mol. Ther. 16(2008):580–589; Grabundzija et al. Mol. Ther. 18(2010):1200–1209; Kebriaei et al. Blood. 122.21(2013):166; Williams. Molecular Therapy 25 16.9(2008):1515–16; Bell et al. Nat. Protoc. 2.12(2007):3153-65; and Ding et al. Cell. 122.3(2005):473-83, all of which are incorporated herein by reference.

The SBTS includes two components: 1) a transposon containing a transgene and 2) a source of transposase enzyme. The transposase can transpose the transposon from a carrier plasmid (or other donor DNA) to a target DNA, such as a host cell chromosome/genome. For
30 example, the transposase binds to the carrier plasmid/donor DNA, cuts the transposon

(including transgene(s)) out of the plasmid, and inserts it into the genome of the host cell. See, e.g., Aronovich et al. *supra*.

Exemplary transposons include a pT2-based transposon. See, e.g., Grabundzija et al. *Nucleic Acids Res.* 41.3(2013):1829-47; and Singh et al. *Cancer Res.* 68.8(2008): 2961–2971, all of which are incorporated herein by reference. Exemplary transposases include a Tc1/mariner-type transposase, e.g., the SB10 transposase or the SB11 transposase (a hyperactive transposase which can be expressed, e.g., from a cytomegalovirus promoter). See, e.g., Aronovich et al.; Kebriaei et al.; and Grabundzija et al., all of which are incorporated herein by reference.

10 Use of the SBTS permits efficient integration and expression of a transgene, e.g., a nucleic acid encoding a CAR described herein. Provided herein are methods of generating a cell, e.g., T cell or NK cell, that stably expresses a CAR described herein, e.g., using a transposon system such as SBTS.

15 In accordance with methods described herein, in some embodiments, one or more nucleic acids, e.g., plasmids, containing the SBTS components are delivered to a cell (e.g., T or NK cell). For example, the nucleic acid(s) are delivered by standard methods of nucleic acid (e.g., plasmid DNA) delivery, e.g., methods described herein, e.g., electroporation, transfection, or lipofection. In some embodiments, the nucleic acid contains a transposon comprising a transgene, e.g., a nucleic acid encoding a CAR described herein. In some embodiments, the nucleic acid contains a transposon comprising a transgene (e.g., a nucleic acid encoding a CAR described herein) as well as a nucleic acid sequence encoding a transposase enzyme. In other 20 embodiments, a system with two nucleic acids is provided, e.g., a dual-plasmid system, e.g., where a first plasmid contains a transposon comprising a transgene, and a second plasmid contains a nucleic acid sequence encoding a transposase enzyme. For example, the first and the 25 second nucleic acids are co-delivered into a host cell.

30 In some embodiments, cells, e.g., T or NK cells, are generated that express a CAR described herein by using a combination of gene insertion using the SBTS and genetic editing using a nuclease (e.g., Zinc finger nucleases (ZFNs), Transcription Activator-Like Effector Nucleases (TALENs), the CRISPR/Cas system, or engineered meganuclease re-engineered homing endonucleases).

In some embodiments, use of a non-viral method of delivery permits reprogramming of cells, e.g., T or NK cells, and direct infusion of the cells into a subject. Advantages of non-viral vectors include but are not limited to the ease and relatively low cost of producing sufficient amounts required to meet a patient population, stability during storage, and lack of immunogenicity.

Methods of Manufacture/Production

In some embodiments, the methods disclosed herein further include administering a T cell depleting agent after treatment with the cell (e.g., an immune effector cell as described herein), thereby reducing (e.g., depleting) the CAR-expressing cells (e.g., the CD19CAR-expressing cells). Such T cell depleting agents can be used to effectively deplete CAR-expressing cells (e.g., CD19CAR-expressing cells) to mitigate toxicity. In some embodiments, the CAR-expressing cells were manufactured according to a method herein, e.g., assayed (e.g., before or after transfection or transduction) according to a method herein.

In some embodiments, the T cell depleting agent is administered one, two, three, four, or five weeks after administration of the cell, e.g., the population of immune effector cells, described herein.

In one embodiment, the T cell depleting agent is an agent that depletes CAR-expressing cells, e.g., by inducing antibody dependent cell-mediated cytotoxicity (ADCC) and/or complement-induced cell death. For example, CAR-expressing cells described herein may also express an antigen (e.g., a target antigen) that is recognized by molecules capable of inducing cell death, e.g., ADCC or complement-induced cell death. For example, CAR expressing cells described herein may also express a target protein (e.g., a receptor) capable of being targeted by an antibody or antibody fragment. Examples of such target proteins include, but are not limited to, EpCAM, VEGFR, integrins (e.g., integrins $\alpha\beta3$, $\alpha4$, $\alpha13/4\beta3$, $\alpha4\beta7$, $\alpha5\beta1$, $\alpha\beta3$, $\alpha\gamma$), members of the TNF receptor superfamily (e.g., TRAIL-R1, TRAIL-R2), PDGF Receptor, interferon receptor, folate receptor, GPNMB, ICAM-1, HLA-DR, CEA, CA-125, MUC1, TAG-72, IL-6 receptor, 5T4, GD2, GD3, CD2, CD3, CD4, CD5, CD11, CD11a/LFA-1, CD15, CD18/ITGB2, CD19, CD20, CD22, CD23/IgE Receptor, CD25, CD28, CD30, CD33, CD38, CD40, CD41, CD44, CD51, CD52, CD62L, CD74, CD80, CD125, CD147/basigin, CD152/CTLA-4, CD154/CD40L, CD195/CCR5, CD319/SLAMF7, and EGFR, and truncated

versions thereof (e.g., versions preserving one or more extracellular epitopes but lacking one or more regions within the cytoplasmic domain).

In some embodiments, the CAR expressing cell co-expresses the CAR and the target protein, e.g., naturally expresses the target protein or is engineered to express the target protein.

5 For example, the cell, e.g., the population of immune effector cells, can include a nucleic acid (e.g., vector) comprising the CAR nucleic acid (e.g., a CAR nucleic acid as described herein) and a nucleic acid encoding the target protein.

In one embodiment, the T cell depleting agent is a CD52 inhibitor, e.g., an anti-CD52 antibody molecule, e.g., alemtuzumab.

10 In other embodiments, the cell, e.g., the population of immune effector cells, expresses a CAR molecule as described herein (e.g., CD19CAR) and the target protein recognized by the T cell depleting agent. In one embodiment, the target protein is CD20. In embodiments where the target protein is CD20, the T cell depleting agent is an anti-CD20 antibody, e.g., rituximab.

In further embodiments of any of the aforesaid methods, the methods further include
15 transplanting a cell, e.g., a hematopoietic stem cell, or a bone marrow, into the mammal.

In another aspect, the invention features a method of conditioning a mammal prior to cell transplantation. The method includes administering to the mammal an effective amount of the cell comprising a CAR nucleic acid or polypeptide, e.g., a CD19 CAR nucleic acid or polypeptide. In some embodiments, the cell transplantation is a stem cell transplantation, e.g.,
20 a hematopoietic stem cell transplantation, or a bone marrow transplantation. In other embodiments, conditioning a subject prior to cell transplantation includes reducing the number of target-expressing cells in a subject, e.g., CD19-expressing normal cells or CD19-expressing cancer cells.

25 **Activation and Expansion of Immune Effector Cells (e.g., T cells)**

Immune effector cells such as T cells may be activated and expanded generally using methods as described, for example, in U.S. Patents 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041; and U.S. Patent Application
30 Publication No. 20060121005.

Generally, a population of immune effector cells, e.g., T regulatory cell depleted cells, may be expanded by contact with a surface having attached thereto an agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a costimulatory molecule on the surface of the T cells. In particular, T cell populations may be stimulated as described
5 herein, such as by contact with an anti-CD3 antibody, or antigen-binding fragment thereof, or an anti-CD2 antibody immobilized on a surface, or by contact with a protein kinase C activator (e.g., bryostatin) in conjunction with a calcium ionophore. For co-stimulation of an accessory molecule on the surface of the T cells, a ligand that binds the accessory molecule is used. For example, a population of T cells can be contacted with an anti-CD3 antibody and an anti-CD28
10 antibody, under conditions appropriate for stimulating proliferation of the T cells. To stimulate proliferation of either CD4+ T cells or CD8+ T cells, an anti-CD3 antibody and an anti-CD28 antibody can be used. Examples of an anti-CD28 antibody include 9.3, B-T3, XR-CD28 (Diacclone, Besançon, France) can be used as can other methods commonly known in the art (Berg et al., Transplant Proc. 30(8):3975-3977, 1998; Haanen et al., J. Exp. Med.
15 190(9):1319-1328, 1999; Garland et al., J. Immunol Meth. 227(1-2):53-63, 1999).

In certain aspects, the primary stimulatory signal and the costimulatory signal for the T cell may be provided by different protocols. For example, the agents providing each signal may be in solution or coupled to a surface. When coupled to a surface, the agents may be coupled to the same surface (i.e., in “cis” formation) or to separate surfaces (i.e., in “trans” formation).
20 Alternatively, one agent may be coupled to a surface and the other agent in solution. In one aspect, the agent providing the costimulatory signal is bound to a cell surface and the agent providing the primary activation signal is in solution or coupled to a surface. In certain aspects, both agents can be in solution. In one aspect, the agents may be in soluble form, and then cross-linked to a surface, such as a cell expressing Fc receptors or an antibody or other binding agent
25 which will bind to the agents. In this regard, see for example, U.S. Patent Application Publication Nos. 20040101519 and 20060034810 for artificial antigen presenting cells (aAPCs) that are contemplated for use in activating and expanding T cells in the present invention.

In one aspect, the two agents are immobilized on beads, either on the same bead, i.e., “cis,” or to separate beads, i.e., “trans.” By way of example, the agent providing the primary
30 activation signal is an anti-CD3 antibody or an antigen-binding fragment thereof and the agent providing the costimulatory signal is an anti-CD28 antibody or antigen-binding fragment

thereof; and both agents are co-immobilized to the same bead in equivalent molecular amounts. In one aspect, a 1:1 ratio of each antibody bound to the beads for CD4+ T cell expansion and T cell growth is used. In certain aspects of the present invention, a ratio of anti CD3:CD28 antibodies bound to the beads is used such that an increase in T cell expansion is observed as compared to the expansion observed using a ratio of 1:1. In one particular aspect an increase of from about 1 to about 3 fold is observed as compared to the expansion observed using a ratio of 1:1. In one aspect, the ratio of CD3:CD28 antibody bound to the beads ranges from 100:1 to 1:100 and all integer values there between. In one aspect, more anti-CD28 antibody is bound to the particles than anti-CD3 antibody, i.e., the ratio of CD3:CD28 is less than one. In certain aspects, the ratio of anti CD28 antibody to anti CD3 antibody bound to the beads is greater than 2:1. In one particular aspect, a 1:100 CD3:CD28 ratio of antibody bound to beads is used. In one aspect, a 1:75 CD3:CD28 ratio of antibody bound to beads is used. In a further aspect, a 1:50 CD3:CD28 ratio of antibody bound to beads is used. In one aspect, a 1:30 CD3:CD28 ratio of antibody bound to beads is used. In one aspect, a 1:10 CD3:CD28 ratio of antibody bound to beads is used. In one aspect, a 1:3 CD3:CD28 ratio of antibody bound to the beads is used. In yet one aspect, a 3:1 CD3:CD28 ratio of antibody bound to the beads is used.

Ratios of particles to cells from 1:500 to 500:1 and any integer values in between may be used to stimulate T cells or other target cells. As those of ordinary skill in the art can readily appreciate, the ratio of particles to cells may depend on particle size relative to the target cell. For example, small sized beads could only bind a few cells, while larger beads could bind many. In certain aspects the ratio of cells to particles ranges from 1:100 to 100:1 and any integer values in-between and in further aspects the ratio comprises 1:9 to 9:1 and any integer values in between, can also be used to stimulate T cells. The ratio of anti-CD3- and anti-CD28-coupled particles to T cells that result in T cell stimulation can vary as noted above, however certain suitable values include 1:100, 1:50, 1:40, 1:30, 1:20, 1:10, 1:9, 1:8, 1:7, 1:6, 1:5, 1:4, 1:3, 1:2, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, and 15:1 with one suitable ratio being at least 1:1 particles per T cell. In one aspect, a ratio of particles to cells of 1:1 or less is used. In one particular aspect, a suitable particle: cell ratio is 1:5. In further aspects, the ratio of particles to cells can be varied depending on the day of stimulation. For example, in one aspect, the ratio of particles to cells is from 1:1 to 10:1 on the first day and additional particles are added to the cells every day or every other day thereafter for up to 10 days, at final ratios of from 1:1 to 1:10 (based on cell counts on the day of addition). In one particular aspect, the ratio of particles to

cells is 1:1 on the first day of stimulation and adjusted to 1:5 on the third and fifth days of stimulation. In one aspect, particles are added on a daily or every other day basis to a final ratio of 1:1 on the first day, and 1:5 on the third and fifth days of stimulation. In one aspect, the ratio of particles to cells is 2:1 on the first day of stimulation and adjusted to 1:10 on the third and fifth days of stimulation. In one aspect, particles are added on a daily or every other day basis to a final ratio of 1:1 on the first day, and 1:10 on the third and fifth days of stimulation. One of skill in the art will appreciate that a variety of other ratios may be suitable for use in the present invention. In particular, ratios will vary depending on particle size and on cell size and type. In one aspect, the most typical ratios for use are in the neighborhood of 1:1, 2:1 and 3:1 on the first day.

In further aspects, the cells, such as T cells, are combined with agent-coated beads, the beads and the cells are subsequently separated, and then the cells are cultured. In an alternative aspect, prior to culture, the agent-coated beads and cells are not separated but are cultured together. In a further aspect, the beads and cells are first concentrated by application of a force, such as a magnetic force, resulting in increased ligation of cell surface markers, thereby inducing cell stimulation.

By way of example, cell surface proteins may be ligated by allowing paramagnetic beads to which anti-CD3 and anti-CD28 are attached (3x28 beads) to contact the T cells. In one aspect the cells (for example, 10^4 to 10^9 T cells) and beads (for example, DYNABEADS® M-450 CD3/CD28 T paramagnetic beads at a ratio of 1:1) are combined in a buffer, for example PBS (without divalent cations such as, calcium and magnesium). Again, those of ordinary skill in the art can readily appreciate any cell concentration may be used. For example, the target cell may be very rare in the sample and comprise only 0.01% of the sample or the entire sample (i.e., 100%) may comprise the target cell of interest. Accordingly, any cell number is within the context of the present invention. In certain aspects, it may be desirable to significantly decrease the volume in which particles and cells are mixed together (i.e., increase the concentration of cells), to ensure maximum contact of cells and particles. For example, in one aspect, a concentration of about 10 billion cells/ml, 9 billion/ml, 8 billion/ml, 7 billion/ml, 6 billion/ml, 5 billion/ml, or 2 billion cells/ml is used. In one aspect, greater than 100 million cells/ml is used. In a further aspect, a concentration of cells of 10, 15, 20, 25, 30, 35, 40, 45, or 50 million cells/ml is used. In yet one aspect, a concentration of cells from 75, 80, 85, 90, 95, or 100

million cells/ml is used. In further aspects, concentrations of 125 or 150 million cells/ml can be used. Using high concentrations can result in increased cell yield, cell activation, and cell expansion. Further, use of high cell concentrations allows more efficient capture of cells that may weakly express target antigens of interest, such as CD28-negative T cells. Such
5 populations of cells may have therapeutic value and would be desirable to obtain in certain aspects. For example, using high concentration of cells allows more efficient selection of CD8+ T cells that normally have weaker CD28 expression.

In one embodiment, cells transduced with a nucleic acid encoding a CAR, e.g., a CAR described herein, e.g., a CD19 CAR described herein, are expanded, e.g., by a method
10 described herein. In one embodiment, the cells are expanded in culture for a period of several hours (e.g., about 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 18, 21 hours) to about 14 days (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 days). In one embodiment, the cells are expanded for a period of 4 to 9 days. In one embodiment, the cells are expanded for a period of 8 days or less, e.g., 7, 6 or 5 days. In one embodiment, the cells are expanded in culture for 5 days, and the resulting
15 cells are more potent than the same cells expanded in culture for 9 days under the same culture conditions. Potency can be defined, e.g., by various T cell functions, e.g. proliferation, target cell killing, cytokine production, activation, migration, or combinations thereof. In one embodiment, the cells, e.g., a CD19 CAR cell described herein, expanded for 5 days show at least a one, two, three or four fold increase in cells doublings upon antigen stimulation as
20 compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., the cells expressing a CD19 CAR described herein, are expanded in culture for 5 days, and the resulting cells exhibit higher proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions. In one embodiment, the cells, e.g., a
25 CD19 CAR cell described herein, expanded for 5 days show at least a one, two, three, four, five, ten fold or more increase in pg/ml of proinflammatory cytokine production, e.g., IFN- γ and/or GM-CSF levels, as compared to the same cells expanded in culture for 9 days under the same culture conditions.

Several cycles of stimulation may also be desired such that culture time of T cells can
30 be 60 days or more. Conditions appropriate for T cell culture include an appropriate media (e.g., Minimal Essential Media or RPMI Media 1640 or, X-vivo 15, (Lonza)) that may contain

factors necessary for proliferation and viability, including serum (e.g., fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-12, IL-15, TGF β , and TNF- α or any other additives for the growth of cells known to the skilled artisan. Other additives for the growth of cells include, but are not limited to, surfactant, plasmanate, and reducing agents such as N-acetyl-cysteine and 2-mercaptoethanol. Media can include RPMI 1640, AIM-V, DMEM, MEM, α -MEM, F-12, X-Vivo 15, and X-Vivo 20, Optimizer, with added amino acids, sodium pyruvate, and vitamins, either serum-free or supplemented with an appropriate amount of serum (or plasma) or a defined set of hormones, and/or an amount of cytokine(s) sufficient for the growth and expansion of T cells. Antibiotics, e.g., penicillin and streptomycin, are included only in experimental cultures, not in cultures of cells that are to be infused into a subject. The target cells are maintained under conditions necessary to support growth, for example, an appropriate temperature (e.g., 37° C) and atmosphere (e.g., air plus 5% CO₂).

In one embodiment, the cells are expanded in an appropriate media (e.g., media described herein) that includes one or more interleukin that result in at least a 200-fold (e.g., 200-fold, 250-fold, 300-fold, 350-fold) increase in cells over a 14 day expansion period, e.g., as measured by a method described herein such as flow cytometry. In one embodiment, the cells are expanded in the presence IL-15 and/or IL-7 (e.g., IL-15 and IL-7).

In embodiments, methods described herein, e.g., CAR-expressing cell manufacturing methods, comprise removing T regulatory cells, e.g., CD25+ T cells, from a cell population, e.g., using an anti-CD25 antibody, or fragment thereof, or a CD25-binding ligand, IL-2. Methods of removing T regulatory cells, e.g., CD25+ T cells, from a cell population are described herein. In embodiments, the methods, e.g., manufacturing methods, further comprise contacting a cell population (e.g., a cell population in which T regulatory cells, such as CD25+ T cells, have been depleted; or a cell population that has previously contacted an anti-CD25 antibody, fragment thereof, or CD25-binding ligand) with IL-15 and/or IL-7. For example, the cell population (e.g., that has previously contacted an anti-CD25 antibody, fragment thereof, or CD25-binding ligand) is expanded in the presence of IL-15 and/or IL-7.

In some embodiments a CAR-expressing cell described herein is contacted with a composition comprising an interleukin-15 (IL-15) polypeptide, a interleukin-15 receptor alpha (IL-15Ra) polypeptide, or a combination of both a IL-15 polypeptide and a IL-15Ra

polypeptide e.g., hetIL-15, during the manufacturing of the CAR-expressing cell, e.g., ex vivo. In embodiments, a CAR-expressing cell described herein is contacted with a composition comprising a IL-15 polypeptide during the manufacturing of the CAR-expressing cell, e.g., ex vivo. In embodiments, a CAR-expressing cell described herein is contacted with a composition
5 comprising a combination of both a IL-15 polypeptide and a IL-15 Ra polypeptide during the manufacturing of the CAR-expressing cell, e.g., ex vivo. In embodiments, a CAR-expressing cell described herein is contacted with a composition comprising hetIL-15 during the manufacturing of the CAR-expressing cell, e.g., ex vivo.

In one embodiment the CAR-expressing cell described herein is contacted with a
10 composition comprising hetIL-15 during ex vivo expansion. In an embodiment, the CAR-expressing cell described herein is contacted with a composition comprising an IL-15 polypeptide during ex vivo expansion. In an embodiment, the CAR-expressing cell described herein is contacted with a composition comprising both an IL-15 polypeptide and an IL-15Ra polypeptide during ex vivo expansion. In one embodiment the contacting results in the survival
15 and proliferation of a lymphocyte subpopulation, e.g., CD8+ T cells.

T cells that have been exposed to varied stimulation times may exhibit different characteristics. For example, typical blood or apheresed peripheral blood mononuclear cell products have a helper T cell population (TH, CD4+) that is greater than the cytotoxic or suppressor T cell population (TC, CD8+). Ex vivo expansion of T cells by stimulating CD3 and
20 CD28 receptors produces a population of T cells that prior to about days 8-9 consists predominately of TH cells, while after about days 8-9, the population of T cells comprises an increasingly greater population of TC cells. Accordingly, depending on the purpose of treatment, infusing a subject with a T cell population comprising predominately of TH cells may be advantageous. Similarly, if an antigen-specific subset of TC cells has been isolated it
25 may be beneficial to expand this subset to a greater degree.

Further, in addition to CD4 and CD8 markers, other phenotypic markers vary significantly, but in large part, reproducibly during the course of the cell expansion process. Thus, such reproducibility enables the ability to tailor an activated T cell product for specific purposes.

30 Once a CAR described herein is constructed, various assays can be used to evaluate the activity of the molecule, such as but not limited to, the ability to expand T cells following

antigen stimulation, sustain T cell expansion in the absence of re-stimulation, and anti-cancer activities in appropriate in vitro and animal models. Assays to evaluate the effects of a CAR of the present invention are described in further detail below

Western blot analysis of CAR expression in primary T cells can be used to detect the
5 presence of monomers and dimers, e.g., as described in paragraph 695 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

In vitro expansion of CAR⁺ T cells following antigen stimulation can be measured by flow cytometry. For example, a mixture of CD4⁺ and CD8⁺ T cells are stimulated with
10 α CD3/ α CD28 aAPCs followed by transduction with lentiviral vectors expressing GFP under the control of the promoters to be analyzed. Exemplary promoters include the CMV IE gene, EF-1 α , ubiquitin C, or phosphoglycerokinase (PGK) promoters. GFP fluorescence is evaluated on day 6 of culture in the CD4⁺ and/or CD8⁺ T cell subsets by flow cytometry. See, e.g., Milone *et al.*, Molecular Therapy 17(8): 1453-1464 (2009). Alternatively, a mixture of CD4⁺
15 and CD8⁺ T cells are stimulated with α CD3/ α CD28 coated magnetic beads on day 0, and transduced with CAR on day 1 using a bicistronic lentiviral vector expressing CAR along with eGFP using a 2A ribosomal skipping sequence. Cultures are re-stimulated with either a cancer associated antigen as described herein⁺ K562 cells (K562-expressing a cancer associated antigen as described herein), wild-type K562 cells (K562 wild type) or K562 cells expressing
20 hCD32 and 4-1BBL in the presence of antiCD3 and anti-CD28 antibody (K562-BBL-3/28) following washing. Exogenous IL-2 is added to the cultures every other day at 100 IU/ml. GFP⁺ T cells are enumerated by flow cytometry using bead-based counting. See, e.g., Milone *et al.*, Molecular Therapy 17(8): 1453-1464 (2009).

Sustained CAR⁺ T cell expansion in the absence of re-stimulation can also be measured.
25 See, e.g., Milone *et al.*, Molecular Therapy 17(8): 1453-1464 (2009). Briefly, mean T cell volume (fl) is measured on day 8 of culture using a Coulter Multisizer III particle counter, a Nexcelom Cellometer Vision or Millipore Scepter, following stimulation with α CD3/ α CD28 coated magnetic beads on day 0, and transduction with the indicated CAR on day 1.

Animal models can also be used to measure a CAR-expressing cell activity, e.g., as
30 described in paragraph 698 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

Dose dependent CAR treatment response can be evaluated, e.g., as described in paragraph 699 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

5 Assessment of cell proliferation and cytokine production has been previously described, as described in paragraph 700 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

Cytotoxicity can be assessed by a standard ⁵¹Cr-release assay, e.g., as described in paragraph 701 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference in its entirety.

10 Cytotoxicity can also be assessed by measuring changes in adherent cell's electrical impedance, e.g., using an xCELLigence real time cell analyzer (RTCA). In some embodiments, cytotoxicity is measured at multiple time points.

Imaging technologies can be used to evaluate specific trafficking and proliferation of CARs in tumor-bearing animal models, e.g., as described in paragraph 702 of International Application WO2015/142675, filed March 13, 2015, which is herein incorporated by reference
15 in its entirety.

Other assays, including those described in the Example section herein as well as those that are known in the art can also be used to evaluate the CARs described herein.

Alternatively, or in combination to the methods disclosed herein, methods and
20 compositions for one or more of: detection and/or quantification of CAR-expressing cells (e.g., in vitro or in vivo (e.g., clinical monitoring)); immune cell expansion and/or activation; and/or CAR-specific selection, that involve the use of a CAR ligand, are disclosed. In one exemplary embodiment, the CAR ligand is an antibody that binds to the CAR molecule, e.g., binds to the extracellular antigen binding domain of CAR (e.g., an antibody that binds to the antigen
25 binding domain, e.g., an anti-idiotypic antibody; or an antibody that binds to a constant region of the extracellular binding domain). In other embodiments, the CAR ligand is a CAR antigen molecule (e.g., a CAR antigen molecule as described herein).

In one aspect, a method for detecting and/or quantifying CAR-expressing cells is disclosed. For example, the CAR ligand can be used to detect and/or quantify CAR-expressing

cells in vitro or in vivo (e.g., clinical monitoring of CAR-expressing cells in a patient, or dosing a patient). The method includes:

providing the CAR ligand (optionally, a labelled CAR ligand, e.g., a CAR ligand that includes a tag, a bead, a radioactive or fluorescent label);

5 acquiring the CAR-expressing cell (e.g., acquiring a sample containing CAR-expressing cells, such as a manufacturing sample or a clinical sample);

contacting the CAR-expressing cell with the CAR ligand under conditions where binding occurs, thereby detecting the level (e.g., amount) of the CAR-expressing cells present. Binding of the CAR-expressing cell with the CAR ligand can be detected using standard

10 techniques such as FACS, ELISA and the like.

In another aspect, a method of expanding and/or activating cells (e.g., immune effector cells) is disclosed. The method includes:

providing a CAR-expressing cell (e.g., a first CAR-expressing cell or a transiently expressing CAR cell);

15 contacting said CAR-expressing cell with a CAR ligand, e.g., a CAR ligand as described herein), under conditions where immune cell expansion and/or proliferation occurs, thereby producing the activated and/or expanded cell population.

In certain embodiments, the CAR ligand is present on a substrate (e.g., is immobilized or attached to a substrate, e.g., a non-naturally occurring substrate). In some embodiments, the
20 substrate is a non-cellular substrate. The non-cellular substrate can be a solid support chosen from, e.g., a plate (e.g., a microtiter plate), a membrane (e.g., a nitrocellulose membrane), a matrix, a chip or a bead. In embodiments, the CAR ligand is present in the substrate (e.g., on the substrate surface). The CAR ligand can be immobilized, attached, or associated covalently or non-covalently (e.g., cross-linked) to the substrate. In one embodiment, the CAR ligand is
25 attached (e.g., covalently attached) to a bead. In the aforesaid embodiments, the immune cell population can be expanded in vitro or ex vivo. The method can further include culturing the population of immune cells in the presence of the ligand of the CAR molecule, e.g., using any of the methods described herein.

In other embodiments, the method of expanding and/or activating the cells further
30 comprises addition of a second stimulatory molecule, e.g., CD28. For example, the CAR

ligand and the second stimulatory molecule can be immobilized to a substrate, e.g., one or more beads, thereby providing increased cell expansion and/or activation.

In yet another aspect, a method for selecting or enriching for a CAR expressing cell is provided. The method includes contacting the CAR expressing cell with a CAR ligand as
5 described herein; and selecting the cell on the basis of binding of the CAR ligand.

In yet other embodiments, a method for depleting, reducing and/or killing a CAR expressing cell is provided. The method includes contacting the CAR expressing cell with a CAR ligand as described herein; and targeting the cell on the basis of binding of the CAR ligand, thereby reducing the number, and/or killing, the CAR-expressing cell. In one
10 embodiment, the CAR ligand is coupled to a toxic agent (e.g., a toxin or a cell ablative drug). In another embodiment, the anti-idiotypic antibody can cause effector cell activity, e.g., ADCC or ADC activities.

Exemplary anti-CAR antibodies that can be used in the methods disclosed herein are described, e.g., in WO 2014/190273 and by Jena et al., "Chimeric Antigen Receptor (CAR)-
15 Specific Monoclonal Antibody to Detect CD19-Specific T cells in Clinical Trials", PLOS March 2013 8:3 e57838, the contents of which are incorporated by reference.

In some aspects and embodiments, the compositions and methods herein are optimized for a specific subset of T cells, e.g., as described in US Serial No. PCT/US2015/043219 filed July 31, 2015, the contents of which are incorporated herein by reference in their entirety. In
20 some embodiments, the optimized subsets of T cells display an enhanced persistence compared to a control T cell, e.g., a T cell of a different type (e.g., CD8+ or CD4+) expressing the same construct.

In some embodiments, a CD4+ T cell comprises a CAR described herein, which CAR comprises an intracellular signaling domain suitable for (e.g., optimized for, e.g., leading
25 to enhanced persistence in) a CD4+ T cell, e.g., an ICOS domain. In some embodiments, a CD8+ T cell comprises a CAR described herein, which CAR comprises an intracellular signaling domain suitable for (e.g., optimized for, e.g., leading to enhanced persistence of) a CD8+ T cell, e.g., a 4-1BB domain, a CD28 domain, or another costimulatory domain other than an ICOS domain. In some embodiments, the CAR described herein comprises an antigen
30 binding domain described herein, e.g., a CAR comprising an antigen binding domain.

In an aspect, described herein is a method of treating a subject, e.g., a subject having cancer. The method includes administering to said subject, an effective amount of:

- 1) a CD4+ T cell comprising a CAR (the CARCD4+) comprising:
 - an antigen binding domain, e.g., an antigen binding domain described herein;
 - 5 a transmembrane domain; and
 - an intracellular signaling domain, e.g., a first costimulatory domain, e.g., an ICOS domain; and
 - 2) a CD8+ T cell comprising a CAR (the CARCD8+) comprising:
 - an antigen binding domain, e.g., an antigen binding domain described herein;
 - 10 a transmembrane domain; and
 - an intracellular signaling domain, e.g., a second costimulatory domain, e.g., a 4-1BB domain, a CD28 domain, or another costimulatory domain other than an ICOS domain;wherein the CARCD4+ and the CARCD8+ differ from one another.
- Optionally, the method further includes administering:
- 3) a second CD8+ T cell comprising a CAR (the second CARCD8+) comprising:
 - an antigen binding domain, e.g., an antigen binding domain described herein;
 - 15 a transmembrane domain; and
 - an intracellular signaling domain, wherein the second CARCD8+ comprises an intracellular signaling domain, e.g., a costimulatory signaling domain, not present on the
 - 20 CARCD8+, and, optionally, does not comprise an ICOS signaling domain.

Biopolymer delivery methods

In some embodiments, one or more CAR-expressing cells as disclosed herein can be administered or delivered to the subject via a biopolymer scaffold, e.g., a biopolymer implant.

25 Biopolymer scaffolds can support or enhance the delivery, expansion, and/or dispersion of the CAR-expressing cells described herein. A biopolymer scaffold comprises a biocompatible (e.g., does not substantially induce an inflammatory or immune response) and/or a biodegradable polymer that can be naturally occurring or synthetic. Exemplary biopolymers are described, e.g., in paragraphs 1004-1006 of International Application WO2015/142675,

30 filed March 13, 2015, which is herein incorporated by reference in its entirety.

Pharmaceutical compositions and treatments

In some aspects, the disclosure provides a method of treating a patient, comprising administering CAR-expressing cells produced as described herein, optionally in combination
5 with one or more other therapies. In some aspects, the disclosure provides a method of treating a patient, comprising administering a reaction mixture comprising CAR-expressing cells as described herein, optionally in combination with one or more other therapies. In some aspects, the disclosure provides a method of shipping or receiving a reaction mixture comprising CAR-expressing cells as described herein. In some aspects, the disclosure provides a method of
10 treating a patient, comprising receiving a CAR-expressing cell that was produced as described herein, and further comprising administering the CAR-expressing cell to the patient, optionally in combination with one or more other therapies. In some aspects, the disclosure provides a method of treating a patient, comprising producing a CAR-expressing cell as described herein, and further comprising administering the CAR-expressing cell to the patient, optionally in
15 combination with one or more other therapies. The other therapy may be, e.g., a cancer therapy such as chemotherapy.

In an embodiment, cells expressing a CAR described herein are administered to a subject in combination with a molecule that decreases the Treg cell population. Methods that decrease the number of (e.g., deplete) Treg cells are known in the art and include, e.g., CD25
20 depletion, cyclophosphamide administration, modulating GITR function. Without wishing to be bound by theory, it is believed that reducing the number of Treg cells in a subject prior to apheresis or prior to administration of a CAR-expressing cell described herein reduces the number of unwanted immune cells (e.g., Tregs) in the tumor microenvironment and reduces the subject's risk of relapse.

In one embodiment, a therapy described herein, e.g., a CAR-expressing cell, is administered to a subject in combination with a molecule targeting GITR and/or modulating GITR functions, such as a GITR agonist and/or a GITR antibody that depletes regulatory T cells (Tregs). In embodiments, cells expressing a CAR described herein are administered to a subject in combination with cyclophosphamide. In one embodiment, the GITR binding molecules and/or
30 molecules modulating GITR functions (e.g., GITR agonist and/or Treg depleting GITR antibodies) are administered prior to the CAR-expressing cell. For example, in one embodiment, a GITR agonist can be administered prior to apheresis of the cells. In

embodiments, cyclophosphamide is administered to the subject prior to administration (e.g., infusion or re-infusion) of the CAR-expressing cell or prior to apheresis of the cells. In embodiments, cyclophosphamide and an anti-GITR antibody are administered to the subject prior to administration (e.g., infusion or re-infusion) of the CAR-expressing cell or prior to apheresis of the cells. In one embodiment, the subject has cancer (e.g., a solid cancer or a hematological cancer such as ALL or CLL). In one embodiment, the subject has CLL. In embodiments, the subject has ALL. In embodiments, the subject has a solid cancer, e.g., a solid cancer described herein. Exemplary GITR agonists include, e.g., GITR fusion proteins and anti-GITR antibodies (e.g., bivalent anti-GITR antibodies) such as, e.g., a GITR fusion protein described in U.S. Patent No.: 6,111,090, European Patent No.: 090505B1, U.S. Patent No.: 8,586,023, PCT Publication Nos.: WO 2010/003118 and 2011/090754, or an anti-GITR antibody described, e.g., in U.S. Patent No.: 7,025,962, European Patent No.: 1947183B1, U.S. Patent No.: 7,812,135, U.S. Patent No.: 8,388,967, U.S. Patent No.: 8,591,886, European Patent No.: EP 1866339, PCT Publication No.: WO 2011/028683, PCT Publication No.: WO 2013/039954, PCT Publication No.: WO2005/007190, PCT Publication No.: WO 2007/133822, PCT Publication No.: WO2005/055808, PCT Publication No.: WO 99/40196, PCT Publication No.: WO 2001/03720, PCT Publication No.: WO99/20758, PCT Publication No.: WO2006/083289, PCT Publication No.: WO 2005/115451, U.S. Patent No.: 7,618,632, and PCT Publication No.: WO 2011/051726.

In one embodiment, a CAR expressing cell described herein is administered to a subject in combination with a GITR agonist, e.g., a GITR agonist described herein. In one embodiment, the GITR agonist is administered prior to the CAR-expressing cell. For example, in one embodiment, the GITR agonist can be administered prior to apheresis of the cells. In one embodiment, the subject has CLL.

The methods described herein can further include formulating a CAR-expressing cell in a pharmaceutical composition. Pharmaceutical compositions may comprise a CAR-expressing cell, e.g., a plurality of CAR-expressing cells, as described herein, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or

glutathione; adjuvants (e.g., aluminum hydroxide); and preservatives. Compositions can be formulated, e.g., for intravenous administration.

In one embodiment, the pharmaceutical composition is substantially free of, e.g., there are no detectable levels of a contaminant, e.g., selected from the group consisting of endotoxin, mycoplasma, replication competent lentivirus (RCL), p24, VSV-G nucleic acid, HIV gag, residual anti-CD3/anti-CD28 coated beads, mouse antibodies, pooled human serum, bovine serum albumin, bovine serum, culture media components, vector packaging cell or plasmid components, a bacterium and a fungus. In one embodiment, the bacterium is at least one selected from the group consisting of *Alcaligenes faecalis*, *Candida albicans*, *Escherichia coli*, *Haemophilus influenza*, *Neisseria meningitidis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Streptococcus pneumonia*, and *Streptococcus pyogenes* group A.

When “an immunologically effective amount,” “an anti-cancer effective amount,” “a cancer-inhibiting effective amount,” or “therapeutic amount” is indicated, the precise amount of the compositions to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject). It can generally be stated that a pharmaceutical composition comprising the immune effector cells (e.g., T cells, NK cells) described herein may be administered at a dosage of 10^4 to 10^9 cells/kg body weight, in some instances 10^5 to 10^6 cells/kg body weight, including all integer values within those ranges. T cell compositions may also be administered multiple times at these dosages. The cells can be administered by using infusion techniques that are commonly known in immunotherapy (see, e.g., Rosenberg et al., *New Eng. J. of Med.* 319:1676, 1988).

In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises about 1×10^6 , 1.1×10^6 , 2×10^6 , 3.6×10^6 , 5×10^6 , 1×10^7 , 1.8×10^7 , 2×10^7 , 5×10^7 , 1×10^8 , 2×10^8 , or 5×10^8 cells/kg. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises at least about 1×10^6 , 1.1×10^6 , 2×10^6 , 3.6×10^6 , 5×10^6 , 1×10^7 , 1.8×10^7 , 2×10^7 , 5×10^7 , 1×10^8 , 2×10^8 , or 5×10^8 cells/kg. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises up to about 1×10^6 , 1.1×10^6 , 2×10^6 , 3.6×10^6 , 5×10^6 , 1×10^7 , 1.8×10^7 , 2×10^7 , 5×10^7 , 1×10^8 , 2×10^8 , or 5×10^8 cells/kg. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises about $1.1 \times 10^6 - 1.8 \times 10^7$ cells/kg. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises about 1×10^7 , 2×10^7 , 5

x 10^7 , 1×10^8 , 2×10^8 , 5×10^8 , 1×10^9 , 2×10^9 , or 5×10^9 cells. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises at least about 1×10^7 , 2×10^7 , 5×10^7 , 1×10^8 , 2×10^8 , 5×10^8 , 1×10^9 , 2×10^9 , or 5×10^9 cells. In some embodiments, a dose of CAR cells (e.g., CD19 CAR cells) comprises up to about 1×10^7 , 2×10^7 , 5×10^7 , 1×10^8 , 2×10^8 , 5×10^8 , 1×10^9 , 2×10^9 , or 5×10^9 cells.

In certain aspects, it may be desired to administer activated immune effector cells (e.g., T cells, NK cells) to a subject and then subsequently redraw blood (or have an apheresis performed), activate immune effector cells (e.g., T cells, NK cells) therefrom, and reinfuse the patient with these activated and expanded immune effector cells (e.g., T cells, NK cells). This process can be carried out multiple times every few weeks. In certain aspects, immune effector cells (e.g., T cells, NK cells) can be activated from blood draws of from 10cc to 400cc. In certain aspects, immune effector cells (e.g., T cells, NK cells) are activated from blood draws of 20cc, 30cc, 40cc, 50cc, 60cc, 70cc, 80cc, 90cc, or 100cc.

The administration of the subject compositions may be carried out in any convenient manner. The compositions described herein may be administered to a patient trans arterially, subcutaneously, intradermally, intratumorally, intranodally, intramedullary, intramuscularly, by intravenous (i.v.) injection, or intraperitoneally, e.g., by intradermal or subcutaneous injection. The compositions of immune effector cells (e.g., T cells, NK cells) may be injected directly into a tumor, lymph node, or site of infection.

20

EXAMPLES

The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

Example 1: Optimizing CART production with exogenous cytokines

Cytokines have important functions related to T cell expansion, differentiation, survival and homeostasis. One of the most important cytokine families for clinical use is the common γ -chain (γ_c) family cytokines, which includes interleukin (IL)-2, IL-4, IL-7, IL-9, IL-15 and IL-21 (Liao et al., 2013, *Immunity*, 38:13-25. IL-2 has been widely studied as an immunotherapeutic

agent for cancer. The supplement of IL-2 enhanced the antitumor ability of anti-CD19 CAR-T cells in the clinical trials (Xu et al., 2013, *Lymphoma*, 54:255-60). However, the administration of IL-2 is limited by side effects and a propensity for expansion of regulatory T cells and the effect of activated induced cell death (AICD) (Malek et al., 2010, *Immunity*, 33:153-65; and Lenardo et al., 1999, *Annu Rev Immunol*, 17:221-53). IL-7, IL-15, and IL-21 each can enhance the effectiveness of adoptive immunotherapies and seems to be less toxicity compared with IL-2 (Alves et al., 2007, *Immunol Lett*, 108:113-20). Despite extensive preclinical and clinical studies on the role of the above cytokines, multi-parameter comparative studies on the roles of various exogenous γ_c cytokines on CAR-T cell adoptive therapy are lacking.

Besides γ -chain cytokines, IL-18 is another immunostimulatory cytokine regulating immune responses, which enhances the production of IFN- γ by T cells and augments the cytolytic activity of CTLs (Srivastava et al., 2010, *Curr Med Chem*, 17:3353-7). Administration of IL-18 is safe and well tolerated, even when the dose reaching as high as 1000 μ g/kg (Robertson et al., 2006, *Clin Cancer Res*, 12:4265-73). Therefore, IL-18 could be another candidate used to boost the antitumor of CAR-T cells.

In this example, the effect of administration of different exogenous cytokines was examined for expansion, phenotype, *in vitro* effector functions, and *in vivo* anti-tumor efficacy of T cells and folate receptor alpha (FR α) CART cells.

The following materials and methods were used in the experiments described in this example.

CAR construction and lentivirus preparation

The pELNS-C4-27z CAR vector was constructed as described previously (manuscript under review). Briefly, the pHEN2 plasmid containing the anti-FR α C4/AFRA4 scFv was used as a template for PCR amplification of C4 fragment using the primers of 5'-
ataggatcccagctggtggagtctgggggaggc-3' (SEQ ID NO: 19) and 5'-
atagctagcacctaggacggcagcttggtccc-3' (SEQ ID NO: 24) (BamHI and NheI were underlined). The PCR product and the third generation self-inactivating lentiviral expression vectors pELNS were digested with BamHI and NheI. The digested PCR products were then inserted into the pELNS vector containing CD27-CD3z T-cell signaling domain in which transgene expression is driven by the elongation factor-1 α (EF-1 α) promoter.

High-titer replication-defective lentivirus was generated by transfection of human embryonic kidney cell line 293T (293T) cells with four plasmids (pVSV-G, pRSV.REV, pMDLg/p.RRE and pELNS-C4-27z CAR) by using Express In (Open Biosystems) as described previously. Supernatants were collected at 24h and 48h after transfection and concentrated by ultracentrifugation. The virus titers were determined based on the transduction efficiency of lentivirus to SupT1 cells by using limiting dilution method.

T cells and cell lines

Peripheral blood lymphocytes were obtained from healthy donors after informed consent under a protocol approved by University Institutional Review Board at the University of Pennsylvania. The primary T cells were purchased from the Human Immunology Core after purified by negative selection. T cells were cultured in complete media (RPMI 1640 supplemented with 10% FBS, 100U/mL penicillin, 100µg/mL streptomycin sulfate) and stimulated with anti-CD3 and anti-CD28 mAbs-coated beads (Invitrogen) at a ratio of 1:1 following the instruction. Twenty-four hours after activation, cells were transduced with lentivirus at MOI of 5. Indicated cytokines were added to the transduced T cells from the next day with a final concentration of 10ng/mL. The cytokines were replaced every 3 days.

The 293T cell used for lentivirus packaging and the SupT1 cell used for lentiviral titration were obtained from ATCC. The established ovarian cancer cell lines SKOV3 (FRα+) and C30 (FRα-) was used as target cell for cytokine-secreting and cytotoxicity assay. For bioluminescence assays, SKOV3 was transduced with lentivirus to express firefly luciferase (fLuc).

Flow cytometric analysis and cell sorting

Flow cytometry was performed on a BD FACSCanto. Anti-human CD45 (HI30), CD3 (HIT3a), CD8 (HIT8a), CD45RA (HI100), CD62L (DREG-56), CCR7 (G043H7), IL-7Rα (A019D5), CD27 (M-T271), CD28 (CD28.2), CD95 (DX2), TNF-α (MAb11), IFN-γ (4S.B3), IL-2 (MQ1-17H12), perforin (B-D48), granzym-B (GB11) were obtained from Biolegend. Biotin-SP-conjugated rabbit anti-human IgG (H+L) was purchased from Jackson ImmunoResearch and APC conjugated streptavidin was purchased from Biolegend. Anti-human Bcl-xl (7B2.5) was purchased from SouthernBiotech. Apoptosis kit and TruCount tubes were obtained from BD Bioscience. For peripheral blood T cell count, blood was obtained via retro-

orbital bleeding and stained for the presence of human CD45, CD3, CD4 and CD8 T cells. Human CD45⁺-gated, CD3⁺, CD4⁺ and CD8⁺ subsets were quantified with the TruCount tubes following the manufacturer's instructions.

5 *In vivo study of adoptive cell therapy*

Female non-obese diabetic/severe combined immunodeficiency/ γ -chain^{-/-}(NSG) mice 8 to 12 weeks of age were obtained from the Stem Cell and Xenograft Core of the Abramson Cancer Center, University of Pennsylvania. The mice were inoculated subcutaneously with 3×10^6 fLuc⁺ SKOV3 cells on the flank on day 0. Four or Five mice were randomized per group before treatment. After tumors became palpable, human primary T cells were activated and transduced as described previously. T cells were expanded in the presence of IL-2 (5ng/mL) for about 2 weeks. When the tumor burden was ~ 250 - 300 mm^3 , the mice were injected with 5×10^6 CAR-T cells or 100 μ l saline intravenously and then received daily intraperitoneal injection of 5 μ g of IL-2, IL-7, IL-15, IL-18, IL-21 or phosphate buffer solution (PBS) for 7 days. Tumor dimensions were measured with calipers and tumor volumes were calculated with the following formula: tumor volume = (length $\times$ width²)/2. The number and phenotype of transferred T cells in recipient mouse blood was determined by flow cytometry after retro-orbital bleeding. The mice were euthanized when the tumor volumes were more than 2000 mm^3 and tumors were resected immediately for further analysis.

20

Statistical analysis

Statistical analysis was performed with Prism 5 (GraphPad software) and IBM SPSS Statistics 20.0 software. The data were shown as mean $\pm$ SEM unless clarified. Paired sample t-tests or nonparametric Wilcoxon rank tests were used for comparison of two groups and repeated measures ANOVA or Friedman test were used to test statistical significance of differences among three or more groups. Findings were considered as statistically significant when P-values were less than 0.05.

25

RESULTS

30 **1. Construction and expression of anti-FR α C4 CAR**

The pELNS-C4-27z CAR comprised of the anti-FR α C4 scFv linked to a CD8 α hinge and transmembrane region, followed by a CD3 ζ signaling moiety in tandem with the CD27

intracellular signaling motif (Figure 1A). Primary human T cells were efficiently transduced with C4 CAR lentiviral vectors with transduction efficiencies of 43%~65% when detected at 48h after transduction (Figure 1B). CAR expression levels were comparable between CD4+ and CD8+ T cells ($52.6 \pm 10.2\%$ vs. $49.5 \pm 17.1\%$, $P=0.713$).

5 **2. Influence of cytokines on expansion of CAR transduced T (CAR-T) cell**

The expansion and accumulation of CAR-T cells in the presence of various γ c cytokines and IL-18 was investigated. Three weeks after exposure to the different cytokines in culture, CAR-T cells that had been cultured in the presence of IL2, IL-7 or IL-5 had expanded 1000-2000 fold. CAR-T cells that had been cultured in the presence of IL-18, IL-21 or NC (control,
10 no cytokine) demonstrated a less than 200 fold expansion (Figure 1C).

CAR expression levels were compared 15 days after transduction in different T cell populations: CD3+, CD4+, and CD8+ T cells. CAR expression levels of each group of CAR-T cells grown in the presence of the different cytokines were similar in CD3+, CD4+, and CD8+ T cell populations. CAR-T cells that had been cultured in the presence of IL-21 showed higher
15 levels of CAR expression than the CAR-T cells exposed to other cytokines (including IL-2 and the control cells that were not grown in the presence of cytokines (NC) in the CD3+ and CD4+ T cell populations (Figure 1D).

The reasons contributing to the higher accumulation of CAR-T cells were analyzed, specifically, proliferation and apoptosis of the T cells was assessed. The proliferative response
20 was measured by monitoring cell division of CFSE labeled T cells cultured for 7 days. As shown in Figure 1E, T cells cultured with IL-2 and IL-15 showed the highest proliferative ability, followed by IL-7; while IL-21 and IL-18 were less potent mitogenic stimulants. Apoptosis of the T cells cultured in the different cytokines was tested using Annexin-V staining. The results indicated that T cells cultured in IL-2, IL-7 and IL-15 underwent less
25 apoptosis when compared with NC, IL-18 and IL-21 groups (Figure 1F). Next, the expression of Bcl-xL, a key negative regulator of lymphocyte apoptosis, was examined by FACs analysis. Consistent with the apoptosis results in Figure 1F, there is a trend that IL-2, IL-7 and IL-15 exposure results in the up-regulation of Bcl-xL expression both in CD4+ and CD8+ T cells, while IL-21 did not increase Bcl-xL expression (Figure 1G and 1H). These results indicate that
30 increased accumulation of T cells expanded in the presence of cytokines, e.g., IL-2, IL-7, or IL-15, may be caused by both an increase in proliferation and a decrease in apoptosis by activation of the Bcl-xL anti-apoptotic pathway.

3. Influence of cytokines on the phenotypes of CAR-T cells

Next, the phenotype of the CAR-T cells expanded in the presence of exogenous cytokines was examined. The fresh T cells from healthy donors were generally divided into four subsets based on CD45RA and CD62L expression: 1) naïve T cell (CD45RA+CD62L+, referred to as Tn), 2) central memory T cell (CD45RA-CD62L+, referred to as Tcm), 3) effector memory T cell (CD45RA-CD62L-, referred to as Tem) and 4) CD45RA positive effector T cell (CD45RA+CD62L-, referred to as Temra). Then the expression of CCR7, CD27, CD28, and CD95 are further evaluated for each subset (Figure 2A). The latter three T cell subsets were positive for CD95 while only small part of Tn expressed CD95 ($3.6 \pm 1.4\%$ in CD4+ and $3.7 \pm 1.3\%$ in CD8+ T cells). This small population also co-expressed CD27, CD28 and CCR7, and was considered as memory stem T cells (Tscm). However, after stimulation with anti-CD3/CD28 beads before and after lentiviral transduction with CAR, CD95 was greatly up-regulated to nearly 100% in this population (Figure 2B). The percentages of CD45RA+CD62L+CD95+ T cells were greatly expanded after anti-CD3/CD28 bead stimulation in both CD4+ and CD8+ T and CAR-T cells when compared with the fresh T cells (Figures 2C and 2D). This population highly expressed CD27, CD28 and CCR7 simultaneously (Figure 2A), indicating it could be defined as Tscm. Furthermore, CD8+ CAR-T cells had a higher percentage of Tscm cells, which may be related to the higher proportion of Tn in initial CD8+ T cells (Figure 2F and 2G).

Fourteen days after co-culture with various cytokines, the proportion of T cell subsets of CAR-T cells were investigated by measuring the expression of CD45RA, CD62L and CD95. As shown in Figure 2E, of the CD4+ CAR-T cells, a significantly higher percentage of Tscm cells existed in the IL-7 group compared with the IL-2 group, while the NC and IL-18 groups presented lower percentages of Tscm but higher percentages of Tcm. The distribution of T cell subsets in the IL-15 group was similar with the IL-2 group, while the IL-21 group presented a higher percentage of Tcm, while percentage of Tscm was comparable with the IL-2 group. The CD8+ CAR-T cells demonstrated a similar trend as that of the CD4+ CAR-T cells on the differentiation and distribution of the four T cell subsets for each cytokine-administered group, with higher proportions of Tscm compared with CD4+ CAR-T cells in the corresponding group of CD8+ CAR-T cells.

The abilities of various CAR-T cell subpopulations to self-renew and to differentiate into other cell types were further studied. The four subsets of CAR-T cells were sorted based on CAR, CD45RA and CD62L expression and cultured separately in medium containing IL-2 for 3 days. As shown in Figure 2H, the Tscm subset was able to differentiate into all the other three subsets, and Tcm and Temra subsets were able to differentiate into Tem. These results indicate that CD62L⁺ and CD45RA⁺ T cells were able to differentiate into CD62L⁻ and CD45RA⁻ T cells, respectively. The proliferation capacity of the four subsets was assessed by CFSE dilution and then compared. The results showed the Tscm presented stronger proliferation ability than other subsets (Figure 2I). Furthermore, CD45RA expression inversely correlated with CFSE intensity while CD62L and CCR7 expression directly correlated with proliferation. In all cytokine groups, CD45RA⁺ T cells exhibited much lower CFSE levels than CD45RA^{dim} and negative T cells (Figure 3), indicating that CD45RA⁺ T cells had stronger proliferation activity than CD45RA⁻ T cells. Thus, the increased accumulation of T cells grown in the presence of IL-2, IL-7 and IL-15 may be related to the increased proportion of CD45RA⁺ T cells (which have increased proliferation capacity) (Figure 4B).

With regard to the phenotype of the CAR-T cells, CAR-T cells presented lower expression of CD45RA, CD62L, CD27 and CD28, but higher expression of CCR7 on the surface of T cells (Figure 4A). The influence of cytokines on the phenotype of CAR-T cells were further assessed based on the expression of the following surface markers: CD27, CD28, CD62L, CCR7 and IL7R α . CAR-T cells grown in the presence of IL-18 showed quite similar expression pattern with those grown without cytokine supplement. IL-2 dramatically down-regulated the expressions of CD27, CD28 CD62L, CCR7 and IL7R α when compared with NC control. Of the other γ c cytokines, compared with IL-2 exposed CAR-T cells, IL-7 exposed CAR-T cells presented higher CD62L, CD27 and CD28 expression but significantly decreased CCR7 expression; IL-15 group CAR-T cells presented higher CD27 and CD28 expression; and IL-21 exposed CAR-T cells presented higher CD62L, CCR7, CD27 and CD28 expression, indicating that IL-2 exposure induced the expansion of a subset of T cells with a much more mature T cell phenotype than all other groups (Figure 4B).

4. Influence of cytokines on the effector function of CAR-T cells

To investigate the influence of cytokines on CAR-T cell effector function, the cytokine production capability of CAR-T cells after stimulation with FR α -expressing SKOV3 cells was

assessed. Following 5 hours stimulation, TNF- α , IFN- γ and IL-2 were detectable in the cytoplasm of CAR-T cells, with 41.5-54.0% of the CAR-T cells produced TNF- α , 12.4-15.3% of the CAR-T cells produced IFN γ , and 4.3-6.5% of CAR-T cells produced and IL-2 (Figure 5A&B). IL-2, IL-7 and IL-15 exposure during expansion promoted CAR-T cells to produce
5 TNF- α , while the numbers of IFN- γ and IL-2 producing CAR-T cells were comparable among all the cytokine groups (Figures 5B, 5C, and 5D). Next, the fractions of responding CAR-T cells and their polyfunctionality were compared (Figure 5E). In comparison to exposure to IL-2 during expansion, exposure to IL-18, IL-21 or no cytokine exposure during expansion induced less cytokine-producing CAR-T cells, and less CAR-T cells possessed the ability to produce
10 multiple cytokines when stimulated by target cells. These results are consistent with the phenotype that the CAR-T cells in IL-18, IL-21 and NC groups were less differentiated than those in the IL-2 exposed group.

Then, the effect of cytokine exposure during expansion on the expression of the cytolytic molecules perforin and granzyme-B after antigen stimulation was determined (Figure
15 5I). Similar with TNF- α production, the CAR-T cells exposed to IL-2, IL-7, and IL-15 demonstrated increased perforin expression compared with CAR-T cells exposed to NC, IL-18 and IL-21. However, although CAR-T cells exposed to IL-21 produce less TNF- α and perforin, they produced the highest level of granzyme-B. The next highest levels of granzyme-B production were observed in CAR-T cells exposed to IL-2 and IL-15 during expansion. CAR-T
20 cells in IL-18 group presented the least amount of both perforin and granzyme-B expression after antigen stimulation (Figure 5G and 5H).

Finally, the tumor lysis activity by CAR-T cells exposed to various cytokines during exposure was quantified by luciferase assay. As shown in Figure 5F, CAR-T cells co-cultured with IL-2 and IL-15 lysed the SKOV3 more efficiently than those with NC and IL-18 (both
25 $P < 0.05$).

The association between phenotype of the CAR-T cells and their function was further confirmed. The T cells 14 days were sorted after lentiviral transduction based on CAR and CD62L expression. The CD62L+ CAR-T cells (Tscm and Tcm) exhibited less cytokine production activity and weaker cytolytic capacity when compared with CD62L- CAR-T cells
30 (Tem and Temra) (Figure 6). In this perspective, CAR-T cells exposed to IL-2 and IL-15 produced more cytokines and presented stronger tumor lysis activity, which might be partially attributed to the higher proportions of Tem and Temra in these groups.

5. Expansion and phenotype of CAR-T cells after antigen challenge

To investigate the influence of cytokines on CAR-T cell expansion under the challenge of specific antigen, the CAR-T cells exposed to IL-2 for two weeks were co-cultured with SKOV3 (FR α +) or C30 (FR α -) cells in the presence of indicated cytokines for 7 days. Similar to the antigen-free circumstance, CAR-T cells exposed to IL-2, IL-7 and IL-15 presented higher fold expansion than CAR-T cells exposed to other cytokines. The CAR-T cells exposed to IL-21 during expansion were more likely to undergo apoptosis (Figure 7A). However, when the CAR-T cells exposed to the indicated cytokines for two weeks were co-cultured with SKOV3 or C30 cells without further cytokine supplement for 7 days, the accumulation of CAR-T cells were comparable among all groups, with those having been exposed to IL-15 and IL-18 undergoing the least amount of apoptosis (Figure 7B). The T cells with the highest CAR expression was the group exposed to IL-21 during expansion (Figure 7C). The phenotypes of CAR-T cells were also analyzed. CD27 and CD28 were highly expressed in CD8+ and CD4+ CAR-T cells in all cytokine groups, respectively. Similar to antigen-free expansion, IL-21 exposure increased the expression of CD27 and CD28 in CAR-T cells (Figure 7D). As to the four subsets of memory T cells, the results were different from antigen-free study: Tscm was rare and Tem accounted for more than 50% in no cytokine, IL-18 and IL-21 all groups. Cytokines had no significant impact on the composition of memory T subsets and IL-7 exposure did not favor the increase of Tscm (Figure 7E).

6. Anti-tumor efficacy of various cytokines in animal models

To evaluate the effects of various cytokines during *ex vivo* expansion of CAR-T cells on the efficacy of CAR-T cells *in vivo*, the persistence of CAR-T cells and outcome was investigated by using a NSG mouse xenograft model of ovarian cancer. Mice bearing subcutaneous SKOV3 tumors were intravenously injected with two doses of 5 $\times$ 10⁶ C4-27z CAR-T cells which had been exposed to the indicated cytokines *ex vivo* for 2 weeks previously (Figure 8A). All mice receiving C4-27z CAR-T cell infusion presented less tumor burden when compared with those injected with untransduced T cells and anti-CD19 CAR-T cells (Figure 8B&8C). Of the various cytokine groups, mice receiving CAR-T cells with previous IL-2 exposure showed the highest tumor burden, consistent with the least amount of circulating human T cell in these mice. The tumors in NC, IL-7, IL-15, IL-18 and IL-21 groups were all

significantly suppressed or even disappeared, without any statistical difference on tumor size. The persistence of transferred T cells in the peripheral blood was determined 15 and 32 days after adoptive transfer. Mice receiving IL-7 and IL-21 treated CAR-T cells seemed to have higher amount of human T cells than other groups in the peripheral blood on day +15, while mice receiving IL-2 treated CAR-T cells had the lowest number of human T cells (Figure 8D). As to the percentages of different CAR-T cell populations, NC, IL-15, IL-18 and IL-21 exposed groups all presented higher CD4+ CAR-T cells when compared with IL-2 group, while the percentages of CD8+ CAR-T cells were comparable among all the groups (Figure 8E). Of the T cell phenotypes, CD62L, CD27 and CD28 were expressed only on about 5-10% of T cells and were comparable among all groups, except that CD8+ T cells in IL-21 group expressed higher CD28 than those in IL-2 and NC group (both $P < 0.05$) (Figure 8F&8G). Interestingly, NC and IL-18 had a much higher amount of CD45RA+CD62L+ T cells in the blood, which was opposite to the result obtained from the in vitro experiment (Figure 8F, 2E&2F). On day +32, the circulating human T cells in all CAR-T cell groups expanded significantly except the IL-2 group, with an average T cell account of 14907/ μ l to 19651/ μ l (and only 242/ μ l in the IL-2 group). Two mice died although the tumors were regressed.

DISCUSSION

IL-2 is the most frequently used cytokine for generating lymphocytes for adoptive immunotherapy. It promotes T cell survival and expansion, enhances tumor-killing ability of T cells. However, the action of IL-2 is limited as it results in activation induced cell death (AICD) of T-cell and the development of regulatory T-cell (Malek et al., *Immunity*, 2010, 33:153-65; and Lenardo et al., *Annu Rev Immunol*, 1999, 17:221-53). In this example, IL-2 significantly increased the accumulation of CAR-T cells and their cytotoxicity ability, but IL-2 exposed CAR-T cells presented inferior antitumor immunity in vivo following adoptive transfer. This finding demonstrates an inverse relationship between in vitro tumor-lysis and in vivo tumor eradication. IL-2 exposed CAR-T cells displayed a relative mature phenotype with low expression of CD62L, CCR7, CD27 and CD28, which are less persistent in vivo (Yang et al., *Cancer Immunol Immunother*, 2013, 62:727-36). Recent studies have indicated that adoptive transfer of less differentiated T cells correlates with superior tumor regression, which supports the finding that IL-2 exposed CAR-T cells are less effective than other group (Gattinoni et al., *Nat Med*, 2011, 17:1290-7; and Markley et al., *Blood*, 2010, 115:3508-19).

IL-15 presented similar performance of stimulating CAR-T cell expansion and tumor-lysis function as IL-2, but induced a less differentiated phenotype (higher expression of CD27 and CD28). Therefore, IL-15 supports the persistence of CAR-T cells in vivo and shows better antitumor immunity in animal models.

5 Compared with IL-2 and IL-15, IL-7 showed similar capability to promote CAR-T cell expansion, but induced higher level of CD62L expression and exhibited the highest proportion of CAR-Tscm cells in an antigen-free circumstance. Therefore, compared to CAR-T cells exposed to IL-2, ex vivo exposure of IL-7 without antigen challenge enhanced the antitumor efficacy of the CAR-T cells. IL-7 exposed CAR-T cells did not result in better in vivo
10 antitumor efficacy than IL-2, and efficacy was inferior to IL-15 due to the less expansion of CAR-T cells under antigen challenge.

IL-21 exerted few effects on CAR-T cell accumulation as it could not enhance anti-apoptosis ability, e.g., by promoting Bcl-xL expression. However, IL-21 induced the expansion of less differentiated CAR-T cells, with a phenotype of high expression of CD62L, CCR7,
15 CD27 and CD28, even under the circumstance of antigen challenge. Therefore, IL-21 exposed CAR-T cells showed best persistence in animal models and IL-21 injection in vivo, and also presented a better efficacy in promoting tumor eradication than other cytokine groups except IL-15. These results are consistent with previous finding that less differentiated CAR-T cells correlates with superior tumor regression.

20 IL-18 is proinflammatory cytokine belonging to the IL-1 family, which regulates both innate and adaptive immune responses by activating monocytes, NK cells, and T cells and production of IFN- γ as well as other cytokines in vivo (Srivastava et al., *Curr Med Chem*, 2010, 17:3353-7). The results presented herein indicates that IL-18 has little impact on CAR-T cell's expansion, phenotype and function in ex vivo experiments, as most of the results in IL-18
25 groups are similar and comparable with NC group. IL-18 promoted little proliferation of T cells and maintained more T cell survival under antigen challenge compared to the control (NC) group. In vivo studies show that IL-18 has no significant impact on CAR-T cell efficacy when compared with mice without cytokine supplement.

In summary, the findings of these experiments indicate that IL-2 supplement ex vivo for
30 CAR-T cell expansion is not an optimal strategy although it is widely used. As to IL-18, IL-21 or no cytokine supplement, although they may induced relative effective CAR-T cells, they do not promote CAR-T cell expansion effectively enough, such that enough CAR-T cells could be

prepared for clinical use in a limited expansion time. Therefore, IL-15 and IL-7 may be better agents for CAR-T cell expansion. Furthermore, the combination of IL-7 and IL-15 supplement instructs the generation of Tscm, which is beneficial to produce more “young” CAR-T cells. As to in vivo cytokine injection, all γc cytokines supplement enhance antitumor efficacy, as many of them favor the expansion of CAR-T cells, with IL-15 presenting best effect. Mice receiving IL-15 exposed CAR-T cells by injection had increased efficacy, due in part to the increased expansion ability and increased persistence of the CAR-T cells during tumor treatment. Thus, the results of these experiments indicate that IL-7 and IL-15 show promise to promote CAR-T cell expansion and induce T cell phenotypes that are most efficacious for therapeutic treatment.

Example 2: Effect of CD25 depletion on cell growth and transduction efficiency

The interleukin-2 α -chain, also known as CD25, is expressed by regulatory T cells (Tregs) but has also been observed on chronic B cell leukemia (CLL) cells (in greater than 85% of CLL patients). Tregs have immune suppressing functions and can impede the efficacy of immunotherapy, e.g., by inhibiting T cell proliferation. Current isolation or enrichment of T cells from CLL patients by apheresis usually contains a significantly increased proportion of Tregs as well as CLL cells. The depletion of Tregs and CLL cells in the starting material by CD25 depletion methods may significantly improve the purity of effector T cells, and thereby increase the potency of CAR19 expressing T cells, e.g., CART19 cells.

Figure 9A shows flow cytometry analysis plots of the cells from an apheresis of a CLL patient. Cells were first sorted for CD45 expression, and the CD45-expressing (CD45+) subset was then further analyzed for CD25 and CD3 expression. As shown in the panel on the right of Figure 9A, CLL patients exhibit a high percentage of CD25 expressing cells.

Optimizing CD25 depletion

A validation experiment was performed to identify the optimal conditions for CD25 depletion from the aphereses from two patients using CD25 Reagent from Miltenyi in a CliniMACS System. CD25 depletion reagent was used at 100%, 70%, and 30% of the manufacturer's recommended amount to identify whether the same depletion efficiency could be obtained by using less reagent. Two different tubing sets from Miltenyi were also tested. The depletion was performed in accordance with the manufacturer's directions. The results from the experiments are shown in the table below. For control, selection using anti-CD3/CD28 immunomagnetic beads was performed.

Table 5. Experimental results from CD25 depletion.

CD25 depletion arms		100%	70%	30%
Miltenyi tubing set	161-01			
CliniMACS program	ENRICHMENT1.1			
Patient cells	UPCC04409-15			
%CD45+CD25+ cells	83.56%			
%CD45+CD3+ cells	8.66%			
%CD45+CD3+CD25- cells	5.70%			
#CD25+ cells to target		2.E+09	2.E+09	2.E+09
#apheresed cells for CD25 depletion		2.39E+09	3.41E+09	7.97.E+09
CD25 bead volume used (mL)		2.5	2.5	2.5
Cell# in CD25-depleted fraction		1.05E+09	1.86E+09	3.36E+09
Cell# in CD25-enriched fraction		2.05E+08	2.58E+08	5.19E+08
Expected CD25- T-cell yield		1.36E+08	1.95E+08	4.54E+08
%T cells in depleted fraction		6.26%	4.08%	2.50%
Observed yield CD25- T cells		6.57E+07	7.55E+07	8.40E+07
Yield of CD3+CD25- as % of expected		48%	39%	18%
%B cells in depleted fraction		90.50%	91.6%	95.30%
Viability CD25+ fraction		94.4%	96.2%	91.1%
Viability CD25- fraction		95.8%	95.0%	99.0%

The expected CD25- (CD25-negative) T cell yield represents the calculated CD25- T cell yield calculated by assuming 100% efficiency in the respective manipulations. The observed yield of CD25- T cells represents the number of CD25- T cells after the respective manipulations. As shown in Table 5, using less reagent than recommended by the manufacturer did not result in the same efficiency in CD25 depletion. Using different tubing resulted in an increase in T cell enrichment by one log.

Figure 9B shows representative flow cytometry analysis plots (top panels) demonstrating the efficiency of CD25 depletion compared to the total cells from the apheresis (top left panel of Figure 9B), control CD3/CD28 selected cells (top middle left panel of Figure 9B), CD25 depleted cells (top middle right panel of Figure 9B), and CD25 enriched cells (top right panel of Figure 9B). The histograms (bottom panels) show the monocyte content of the cell population, as determined by CD14 expression of the CD3-CD19- subset. These results indicate efficient CD25 depletion and that CD25 depletion also resulted in significant monocyte content (61.1% CD14-expressing cells compared to less than 2% in the total cells from apheresis, control, and the CD25 enriched cells).

Effect of CD25 depletion on T cell population and proliferation

Next, the quality of the T cell product after CD25 depletion was assessed by determining the proportion of CD4+ and CD8+ T cells and proliferation capacity.

To determine the proportion of specific T cells populations, cells were analyzed by flow cytometry nine days after selection by anti-CD3/CD28 or CD25 depletion as described above.

5 The results show that CD3/CD28-selected T cells had a greater proportion of CD4+ T cells compared to CD25 depleted cells (84.6% compared to 46.8% CD4+ T cells, boxes at the top left of the panels in Figure 10). Conversely, CD25 depleted cells had a greater proportion of CD8+ T cells compared to the CD3/CD28-selected cells (47.2% compared to 11.5% CD8+ T cells; boxes at the bottom right of the panels in Figure 10). Therefore, CD25 depletion results
10 in T cells with a greater proportion of CD8+ T effector cells.

Proliferation capacity and cell viability was also assessed in control (CD3/CD28 selected cells) and CD25 depleted cells. 1.6×10^7 cells from control and CD25 depleted cells were plated and the cell number and viability was determined over 10-13 days. Figure 11A shows the total cell number over time and Figure 11B shows the calculated population
15 doublings (calculated from the total number of cells). The results indicate that the CD25 depleted cells demonstrated similar growth characteristics to the control cells. Figure 11C shows the percentage of viable cells, and the results show that viability was also similar between control and CD25 depleted cells.

Effect of CD25 depletion on lentiviral transduction efficiency

20 The effect of CD25 depletion on lentiviral transduction efficiency was assessed by determining the expression of CAR after transduction. A patient apheresis was depleted with CD25 cells as described above. The efficiency of the CD25 depletion is demonstrated in the flow cytometry analysis plots comparing the CD25-expressing population before (apheresis sample) and after CD25 depletion (CD25-depleted fraction) (Figure 12A). After CD25
25 depletion, the CD25 depleted fraction contained about 59.2% of CD25 negative cells and only 10.3% CD25 positive cells.

The CD25 depleted fraction was transduced with a lentiviral construct encoding CAR19. After 11 days of culture, CAR expression was assessed by flow cytometry. Cells that were untransduced and transduced CD3 selected cells were used as controls. As shown in
30 Figure 12B, CAR19 expression was significantly higher in CD25 depleted cells compared to CD3 selected cells (51.4% compared to 12.8%). This result demonstrates that CD25 depleted

cells have improved lentiviral transduction efficiency, which may be important for improved therapeutic effect in CART therapy.

Example 3: Using cytokines with CD25-depleted cells

5 In this example, the effect of CD25 depletion with cytokine supplement during expansion in culture was examined. Peripheral blood mononuclear cells (PBMCs) were isolated from a patient and were either left unmanipulated or were depleted of CD25-expressing cells as described in Example 2. T cell enrichment was achieved by stimulation with anti-CD3 and CD28 coated beads. The T cells were immediately cultured in media supplemented with 10ng/ml IL-7, 10ng/ml IL-15, or the combination of
 10 10ng/ml IL-7 and 10ng/ml IL-15. At day 3, medium was changed with the same cytokines added. At day 5, the medium containing 100 IU IL-2/ml was added, and the cells were grown for a total of 10 days.

Flow cytometric analysis shows the change in distribution of CD3 and CD19 cells in CD25 depleted cells compared to unmanipulated PBMC (standard CD3/CD28 selection) after
 15 culture in the presence of IL7, IL-15, or IL7 and IL15. The distribution of CD3, CD19, and CD25 expressing cells in the starting population (e.g., before CD25 depletion and before culture with cytokine supplementation) is shown in Figure 13. The starting population had a high proportion of CD3-CD19+ cells (top panel, Figure 11) and a high proportion of CD25-expressing cells (bottom panels, Figure 13). After manipulation (CD25 depletion) and culture
 20 with cytokines, the distribution changed as shown in Figure 14. CD25 depleted cells overall showed greater reduction in CD19-expressing cells compared to the unmanipulated cells.

Proliferation capacity was also assessed for the same cell samples by determining the total number of cells in culture at day 10 after stimulation with anti-CD3 and anti-CD28 coated beads. The cell numbers for each cell sample are shown below and in Figure 15.

25 **Table 6. In vitro expansion**

Cells	Cytokines added	# Cells in culture
Unmanipulated	IL-7	1.24×10^6
	IL-15	0.92×10^6
	IL-7 + IL-15	0.52×10^6
CD25-depleted	IL-7	0.93×10^6
	IL-15	1.95×10^6
	IL-7 + IL-15	3.03×10^6

These results show that supplementation of IL-15 during culture of CD25 depleted T cells resulted in increased expansion compared to unmanipulated cells. Addition of IL-7 and IL-15 in the media during culture resulted in significant increase in expansion compared to unmanipulated cells, and compared to adding the cytokines IL-7 or IL-15 independently. Thus, the combination IL-7 and IL-15 supplement resulted in T cells with the most increased proliferation capacity.

Example 4: Three-day manufacturing process

With the use of gene transfer technologies, T cells can be genetically modified to stably express antibody binding domains on their surface that endow the T cells with specificities that are independent of the constraints imposed by the major histocompatibility complex (MHC). CAR therapies can utilize synthetic proteins expressed on T-cells that fuse an antigen recognition fragment of an antibody (an scFv, or single-chain variable region fragment) with an intracellular domain of the CD3-zeta chain. Upon interaction with a target cell expressing the scFv's cognate antigen, CARs expressed on T cell cells can trigger T-cell activation leading to target cell killing (also referred to as target cell lysis). When combined with additional costimulatory signals such as the cytoplasmic domain of CD137 or CD28, these receptors can also stimulate T cell proliferation and increase CAR-modified T cell (CART) persistence in vivo.

The mechanisms underlying CART persistence are being explored. Signaling domains in CARs appear to be important factors determining persistence; however, a number of published studies in mice and non-human primates (Klebanoff, C. A., et al. (2005). "Central memory self/tumor-reactive CD8+ T cells confer superior antitumor immunity compared with effector memory T cells." *Proc Natl Acad Sci U S A* 102(27): 9571-9576, Berger, C., et al. (2008). "Adoptive transfer of effector CD8+ T cells derived from central memory cells establishes persistent T cell memory in primates." *J Clin Invest* 118(1): 294-305, Hinrichs, C. S., et al. (2009). "Adoptively transferred effector cells derived from naive rather than central memory CD8+ T cells mediate superior antitumor immunity." *Proc Natl Acad Sci U S A* 106(41): 17469-17474, Wang, X., et al. (2011). "Engraftment of human central memory-derived effector CD8+ T cells in immunodeficient mice." *Blood* 117(6): 1888-1898) and data from human clinical trial correlative studies suggest that the phenotype of T cells in the

adoptively transferred product is also an important determinant of T cell persistence following adoptive transfer (Xu, Y., et al. (2014). "Closely related T-memory stem cells correlate with in vivo expansion of CAR.CD19-T cells and are preserved by IL-7 and IL-15." Blood 123(24): 3750-3759). While not wishing to be bound by theory, it is hypothesized that the T cells
5 expanded during ex vivo culture under the commonly employed system of anti-CD3 and anti-CD28 agonist antibodies combined with exogenous IL-2 biases the expanded T cells towards more differentiated cells with reduced persistence. It was hypothesized that replacing IL-2 with IL-7 and IL-15 would bias ex vivo culture over time towards cells with a more optimal Tscm/cm phenotype. It was also hypothesized that products with less ex vivo culture might
10 have a greater proportion of cells with the optimal phenotype for engraftment and persistence following adoptive transfer.

This Example both characterizes the phenotype of cells over time in culture and evaluates the functionality of cells at different time points following activation and transduction.

15 Bead removal from T cells following activation with CD3/28-stimulatory beads can be accomplished with high efficiency by mechanical disruption as shown in **Figures 27A-27D**. Bead detachment was accomplished in these preclinical studies by repeat passage (extrusion) of the cell suspension through a narrow bore pipet tip. In a closed culture system, extrusion to create gentle shearing could be performed using a closed system that incorporates a set of
20 narrow bore tube in a closed system prior to bead removal.

As shown in **Figure 28**, T cells that are activated, transduced with a CD19-specific CAR bearing BBz signaling and harvested from T cells cultures at time points as early as day 3 following activation have potent, antigen-specific cytotoxic activity in vitro. These early harvest cells are also able to produce cytokines with a similar pattern and quantity to those
25 cultured for longer periods of 5 or 9 days ex vivo (**Figure 29A**) with similar basal secretion (**Figure 29B**).

During ex vivo culture, the population of T cells undergoes a progressive transition towards a more differentiated phenotype with an increasing proportion of T effector memory (Tem) and Tcm cells as shown in **Figure 30A**. This increase in proportion of Tem and Tcm
30 cells results from an increase in more differentiated T cell subsets with minimal expansion and/or decrease in the naïve-like/Tscm cells (**Figure 30B**).

Fig. 31A shows the experimental setup for testing the potency of CART cells produced under different conditions. **Fig. 31B** shows the potency of IL2-treated CART cells in slowing Nalm6 tumor growth in mice. Day 3 cells demonstrate enhanced potency at all doses. T cells activated and transduced with a CD19-specific CAR bearing the 4-1BBz signaling domain show more potent in vivo anti-leukemic activity when harvested at the early time points. This increased potency is demonstrated by the lower doses of T cells required to eradicate leukemia with day 3 harvested CART19 cells compared with day 5 or day 9 harvested cells. Proportions of CAR+ cells are shown in **Fig. 31C**.

Cells treated with IL-2 and IL7/15 were harvested in two different days (3, 9). A relatively low dose of 0.7×10^6 CAR+ cells was used. T cells activated, transduced with a CD19-specific CAR bearing the 4-1BBz signaling domain and cultured in IL-7/IL-15 showed more potent activity at limiting doses than cells expanded in IL-2 at early (day 3) and late (day 9) time points (**Figure 31D**). Day 3 cells treated with IL7/15 demonstrate enhanced potency at early time points.

EQUIVALENTS

The disclosures of each and every patent, patent application, and publication cited herein are hereby incorporated herein by reference in their entirety. While this invention has been disclosed with reference to specific aspects, it is apparent that other aspects and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The appended claims are intended to be construed to include all such aspects and equivalent variations.