

METHODS OF USING RAR γ AGONISTS FOR CANCER TREATMENT

FIELD

[0001] The present disclosure relates to methods of treating cancer with RAR γ agonists.

CROSS REFERENCE TO RELATED APPLICATIONS

[0002] The present application claims the benefit of U.S. Provisional Patent Application 63/282,547 filed November 23, 2022, the entire contents of which is incorporated by reference herein.

BACKGROUND

[0003] Retinoids are a class of small molecules that interact with the nuclear receptors retinoic acid receptor (RAR) and retinoid X receptor (RXR). Retinoids that interact with RXR are termed rexinoids and, in some contexts, the term retinoid is used to indicate a compound that interacts with RAR. Both RAR and RXR have three subtypes, α , β , and γ . Retinoids can act as either agonists or antagonists of these receptors. Some retinoids interact with both RAR and RXR and with each of the subtypes, while others will preferentially or specifically interact with a particular receptor type or subtype(s). Retinoids regulate development and the differentiation of multiple types of cells. Retinoids have pleotropic effects depending, in part, on the particular receptor types and subtypes that a particular retinoid activates or inhibits. Retinoids exhibit a degree of structural diversity. It is not possible to reliably predict which receptor type and subtype a particular retinoid will interact with or whether it will act as an agonist or antagonist.

[0004] Further uncertainty about what can be expected from a particular subtype of retinoid arises from a degree of inconsistency in the scientific literature as to what effects may be observed. In tissue culture studies of a leukemic cell line Perri et al. (*Exp Cell Res.* 327(2):183–191, 2014) found that a RAR α agonist and a RAR γ antagonist each inhibited proliferation, and that a combination of the RAR γ antagonist and a dual antagonist of BCL-x_L and MCL-1 was profoundly inhibitory. In contrast, Meister et al. (*Anticancer Res.* 18(3A):1777-1786, 1998) found RAR γ agonists had the greatest antiproliferative effect against a neuroblastoma cell line. In both studies apoptosis was seen as the primary mechanism of inhibiting tumor cell proliferation. Cheung et al. (*Biochem Biophys Res Commun.* 229(1):349-54, 1996) studied the effects of RAR subtype specific agents by transfecting a neuroblastoma cell line with each individual RAR subtype and concluded the RAR β expression was necessary for growth inhibition. Raffo et al. (*Anticancer Res.* 20(3A):1535-43, 2000) observed some apoptotic activity against breast cancer cell lines by agonists of all three subtypes, but the greatest effect from RAR α agonists. As compared to these *in vitro* experiments, what effects any of these


compounds may have *in vivo*, where the whole panoply of pleiotropic effects would be in play, is even less clear.

[0005] Treatments for cancer are ever evolving, gaining in specificity and sophistication. Early non-surgical cancer treatments generally targeted rapidly dividing cells which were more sensitive to radiological and chemical assault. Over time, more specific and less generally toxic treatments have been developed. Some treatments appear to have broad applicability, for example immune checkpoint inhibitors. Others are targeted to cancers that express a particular antigen or other biomarker involved in the regulation of proliferation or differentiation; including many monoclonal antibodies and kinase inhibitors. Yet as the variety of cancer treatments has grown, it has become ever harder to determine which candidate treatments might be productively pursued and for what indications.

SUMMARY


[0006] Disclosed herein are methods of expanding tumor-infiltrating lymphocytes (TIL) *in vitro* and associated methods treating cancer by administering an effective amount of a RAR γ selective agonist and/or expanded TIL to a subject in need thereof.

[0007] One aspect is a method of treating cancer comprising administering, to a patient in need thereof, an effective amount RAR γ -selective agonist having the structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified. However, in some embodiments, the E configuration of the =N-OH group is preferred. R⁷ to R¹² are independently: C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom (for example F, Cl, or Br), a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃. Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl. R¹⁶ is H or F.


[0008] In some embodiments, the RAR γ -selective agonist has the structure



[0009] One aspect is a method of generating, differentiating, or expanding the number of TIL by contacting them with a RARy selective agonist. In some embodiments, contacting comprises culturing the TIL *in vitro* in a media supplemented with the RARy selective agonist. In some embodiments, contacting comprises administering the RARy selective agonist to a subject having a tumor.


[0010] With respect to the above aspects, in some embodiments, the RARy selective agonist is selective in that it has no or negligible agonistic activity for RAR α at clinically relevant concentrations. In some embodiments the RARy selective agonist is selective in that it has no or negligible agonistic activity for both RAR α and RAR β at clinically relevant concentrations.

[0011] In some embodiments, the RARy selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified. However, in some embodiments, the E configuration of the =N-OH group is preferred. R⁷ to R¹² are independently: C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom (for example F, Cl, or Br), a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃. Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl. R¹⁶ is H or F.

[0012] In some embodiments, the RARy selective agonist is 4-((1E,3E)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1E), having the structure




or a pharmaceutically acceptable salt thereof. In some embodiments, the RARy selective agonist is 4-((1E,3Z)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1Z), having the structure




or a pharmaceutically acceptable salt thereof.


[0013] In some embodiments, the RAR γ selective agonist is a compound of structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified. However, in some embodiments, the E configuration of the =N-OH group is preferred. In some embodiments, the RAR γ agonist is 3-fluoro-4-((1E,3E)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2E), having the structure




or a pharmaceutically acceptable salt thereof. In some embodiments, the RAR γ selective agonist is 3-fluoro-4-((1E,3Z)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2Z), having the structure



or a pharmaceutically acceptable salt thereof.

[0014] In some embodiments, the RAR γ selective agonist is a compound of structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; each R¹ and R² are independently H or C₁₋₆ alkyl; R³ and R^{3'} are independently H, or halogen (for example, Cl, F, or Br); (R⁴)₂ comprises R⁴ and R^{4'} which are independently H, halogen (for example, Cl, F, or Br), C₁₋₆ alkyl or C₁₋₆ alkoxy; X is O, S, CH₂, C(R⁵)₂, or NR⁶, wherein each R⁵ and R⁶ are independently H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; and the COOR group is in the meta or para position and the two R⁴ groups occupy the remaining positions on the ring. In some embodiments, the para-position of the COOR group is preferred. In some embodiments, the E configuration of the =N-OH group is preferred. In some embodiments, both R¹ are CH₃. In some embodiments, both R² are H. In some embodiments, X is C(R⁵)₂. In some embodiments, both R⁵ are CH₃. In some embodiments, one R³ is H and the other R³ is F. In some embodiments, both R⁴ are H. In some embodiments, all R³ and R⁴ are H. In some embodiments, R is H. In some embodiments, R is methyl or ethyl. In some embodiments, the COOR group is in the para position. In some embodiments, the COOR group is in the meta position. In some embodiments, the =N-OH group is in the E configuration. In some embodiments the =N-OH group is in the Z configuration. In particular embodiments, R is H and the carboxylic acid group is in the para position, both R¹ are CH₃, both R² are H, X is C(CH₃)₂, all R³ and R⁴ are H, and the =N-OH group is in the E configuration (GA3Ep). Some embodiments specifically include one or more specific substituents at one or more of the variable positions. Some embodiments specifically exclude one or more specific substituents at one or more of the variable positions.

[0015] In some embodiments, the RAR γ selective agonist is CD437, CD2325, CD666, trifarotene, or BMS961.

[0016] One aspect is a method of treating cancer with RAR γ agonist-expanded TIL. In some embodiments, the RAR γ agonist is administered to a subject having a cancer to be treated.

[0017] In alternative embodiments, T lymphocytes are expanded in *in vitro* culture by exposing the culture to the RAR γ agonist, and the RAR γ agonist-expanded T lymphocytes are administered to a subject having a cancer to be treated, as adoptive T cell therapy. That is, the culture media is supplemented with the RAR γ agonist. The concentration of RAR γ agonist in the culture media can be in a range of 1 pM to 1 mM. In some embodiments, the concentration of RAR γ agonist in the culture media is 0.5 nM. In some embodiments, the T lymphocytes are TIL isolated from a tumor explant from a subject. In some embodiments, the

T lymphocytes are peripheral blood mononuclear cells (PBMC). In some embodiments, the T lymphocyte culture further comprises lethally irradiated tumor cells. In some embodiments, the tumor cells to be lethally irradiated are obtained from the subject to be treated. In some embodiments, the culture media is supplemented with interleukin 2 (IL-2). In some embodiments, the expanded T lymphocytes are administered to the subject by infusion, for example, by intravenous or intratumoral infusion.

[0018] In some embodiments, the subject is a mammal. In some embodiments, the subject is a human. In some embodiments, the RAR γ agonist is GA1E. In some embodiments, the RAR γ agonist is GA2E.

[0019] With respect to the above aspects, in some embodiments, the cancer is breast cancer. In some embodiments, the breast cancer is triple negative breast cancer. In some embodiments, the cancer is Her2⁺ breast cancer. In some embodiments, the cancer is lung cancer, for example, non-small cell lung cancer. In some embodiments, the cancer is a solid tumor. In some embodiments, the cancer is a hematologic cancer. In some embodiments, the cancer is a carcinoma, a sarcoma, a melanoma, a glioblastoma, a leukemia, a lymphoma, a myeloma, or a plasma cell cancer.


[0020] In some embodiments, the effective amount of the RAR γ agonist is about 0.01 to about 300 mg/m²/day; however, doses below or above this exemplary range are within the scope of the present disclosure. The daily dose can be about 0.5 to about 100 mg/m²/day, about 1 to about 90 mg/m²/day, about 5 to about 80 mg/m²/day; or at least about 0.02, 0.03, 0.05, 0.07, 0.1, 0.2, 0.3, 0.5, 0.7, 1, 2, 3, 5, 7, 10, 15, 20, 25, 30, 50, 70, or 100 mg/m²/day; or not more than about 0.1, 0.2, 0.3, 0.5, 0.7, 1, 2, 3, 5, 7, 10, 15, 20, 25, 30, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, 225, 250, 275, or 300 mg/m²/day; or a range defined by any two of the foregoing values. In some embodiments, the effective amount for a human is about 0.006 to about 200 mg/day or about 1 to about 100 mg/day.

[0021] In some embodiments, the effective amount of the RAR γ agonist is about 0.27 μ g/kg/day to about 8 mg/kg/day; however, doses below or above this exemplary range are within the scope of the present disclosure. The daily dose can be about 0.013 to about 2.7 mg/kg/day, about 0.025 to about 2.5 mg/kg/day, about 0.130 to about 22 mg/kg/day; or at least about 0.0005, 0.0008, 0.001, 0.0013, 0.0020, 0.0025, 0.005, 0.0008, 0.0010, 0.0013, 0.0020, 0.0027, 0.005, 0.008, 0.010, 0.013, 0.020, 0.027, 0.05, 0.08, 0.1, 0.13, 0.2, 0.5, 0.8, 1.0, 1.3, 1.8, 2.0, or 2.7 mg/kg/day; or not more than about 0.0027, 0.005, 0.008, 0.010, 0.013, 0.020, 0.027, 0.05, 0.08, 0.1, 0.13, 0.2, 0.5, 0.8, 1.0, 1.3, 1.8, 2.0, or 2.7, 3.3, 4.0, 4.7, 5.4, 6.1, 6.8, 7.4, or 8.0 mg/kg/day, or a range defined by any two of the foregoing values.

[0022] With respect to any aspect comprising administering the RAR γ selective agonist to a subject, in some embodiments, the administering occurs periodically throughout an interval of treatment. That is, the RAR γ agonist is administered at regularly occurring time points within the interval of treatment. In some embodiments, periodically is twice daily, once daily, every other day, every third day, or twice weekly. In some embodiments, the administering occurs in repeated cycles throughout an interval of treatment. In some embodiments, a cycle comprises 1) administering the RAR γ selective agonist periodically over a first span of time and 2) suspending administration of the RAR γ selective agonist over a second span of time, after which a new cycle may be initiated. In some embodiments, the first span of time (during which the RAR γ agonist is administered) is 10-15 days, or any integer number of days therein. In some embodiments, the second span of time (during which administration of the RAR γ agonist is suspended) is two weeks to one month or any integer number of days therein. In various embodiments, the interval of treatment extends from a first administration of the RAR γ agonist until, a complete response is achieved, stable disease is obtained, or disease progression again occurs after stable disease or regression.

[0023] With respect to any aspect comprising administering the RAR γ selective agonist to a subject, some embodiments further comprise administering an inhibitor of regulatory T cells (Tregs). In some embodiments, the inhibitor of Tregs is a Treg-depleting antibody. In various embodiments, the Treg-depleting antibody is anti-CD25 antibody, an anti-glucocorticoid-induced tumor necrosis factor-related protein (GITR) antibody, an anti-FoxP3 antibody, an anti-CCR $_4$ antibody, or an anti-folate receptor 4 antibody. In some embodiments, the inhibitor of Tregs comprises an RAR α antagonist.

[0024] With respect to any aspect comprising administering the RAR γ selective agonist to a subject, some embodiments further comprise administering an RXR agonist having the structure



where R is H or C $_{1-6}$ alkyl, or a pharmaceutically acceptable salt thereof. In some embodiments, the RXR agonist is 3,7-dimethyl-6(S),7(S)-methano,7-[1,1,4,4-tetramethyl-1,2,3,4-tetrahydro-naphth-7-yl]2(E), 4(E) heptadienoic acid (IRX4204).

[0025] With respect to any aspect comprising administering the RAR γ -selective agonist to a subject, some embodiments further comprise administering a CAR-T cell. In some

embodiments, the RAR γ -selective agonist is a compound having a structure of GAG1 as defined above.

[0026] With respect to any aspect comprising administering the RAR γ -selective agonist to a subject, some embodiments further comprise administering an anti-PD-1 or anti-PD-L1 antibody.


[0027] With respect to any aspect comprising generating or expanding TIL by contacting them with a RAR γ selective agonist, some embodiments further comprise administering or supplementing culture media, as appropriate, with an anti-PD-1 or anti-PD-L1 antibody.

[0028] One aspect is a method of potentiating chimeric antigen receptor-T (CAR-T) cancer immunotherapy comprising administering an effective amount RAR γ -selective agonist having the structure of GAG1, as defined above, to a cancer patient who is receiving, has received, or is scheduled to receive, CAR-T cells.

[0029] One aspect is a compound having a structure of GAG1 as defined above. One embodiment is 4-((1*E*,3*E*)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1E). Some embodiments are a pharmaceutically-acceptable salt of GA1E. Some embodiments are a C₁₋₆ ester of GA1E. Some embodiments are a pharmaceutical composition comprising GA1E, or a salt or ester thereof.


[0030] One aspect is a method of general synthesis of GAG1 according to the synthetic scheme in Figure 1. A complete synthesis begins with Compound 1 and proceeds through intermediates, Compounds 2 through 5, to produce the species of GAG1, the isomers GA1E and GA1Z. Some embodiments are a method of synthesizing GAG1 species comprising producing intermediate Compound 4 by reacting Compound 3 with *n*-BuLi and *N*-methoxyl-*N*-methylacetamide. Some embodiments are a method of synthesizing GAG1 species comprising producing intermediate Compound 5 by reacting Compound 4 with methyl 3-fluoro-4-formylbenzoate. Some embodiments are a method of synthesizing GAG1 species by reacting Compound 5 with hydroxylamine hydrochloride in the presence of an organic or inorganic base. In some embodiments, the base is pyridine.

[0031] One aspect is intermediate Compound 4, having the structure



A further aspect is a method of synthesizing Compound 4 by reacting Compound 3 with *n*-BuLi and *N*-methoxyl-*N*-methylacetamide.

[0032] One aspect is intermediate Compound 5, having the structure



A further aspect is a method of synthesizing Compound 5 by reacting Compound 4 with methyl 3-fluoro-4-formylbenzoate in the presence of a base, such as NaOH.

BRIEF DESCRIPTION OF DRAWINGS

[0033] Figure 1 presents a scheme for the synthesis of the species of GAG1, GA1E and GA1Z.

[0034] Figure 2 presents a scheme for the synthesis of the species of GAG1, GAG1-1 and GAG1-2.

[0035] Figures 3A-B depict the results of RAR α and RAR γ transactivation assays of GA1E (Figure 3A) and GA2E (Figure 3B).

[0036] Figures 4A-B depict the results of treating a triple negative breast cancer in a syngeneic, immunocompetent mouse model with the RAR γ selective agonists tazarotenic acid and GA2E, versus vehicle control. Figure 4A shows tumor volume over the course of the experiment in each of the three arms of the study, and thus inhibition of tumor growth. Figure 4B shows body weight over the course of the experiment in each of the three arms of the study.

[0037] Figures 5A-B depict the results of flow cytometric analysis of tumors excised at the end of the study described in Example 3, for each of the treatments, GA2E, tazarotenic acid, and vehicle control. Results are shown both as percentage (upper panels) and absolute number of cells/tumor (lower panels). Figure 5A shows the number of live cells in the tumors and their percentage of the total cells (Live), the number of leukocytes based on CD45 staining and their percentage of the live cells (CD45), the total number of T cells based on CD3 staining and their percentage of the leukocytes (Tot T), and CD4⁺ T cell based on CD4 staining and their percentage of total T cells (CD4T). Figure 5B shows the number of memory T helper cells based on staining for CD4 and intracellular IFN γ and their percentage of total CD4⁺ cells (TH IFN γ), the number of Th17 cells based on staining for CD4 and intracellular IL-17 and their percentage of total CD4⁺ cells (TH IL17), the number of T regulatory cells based on staining for CD4, CD25, and FOXP3 and their percentage of total CD4⁺ cells (Treg), CD8⁺ T cell based

on CD8 staining and their percentage of total T cells (CD8T), and memory cytolytic T cells based on staining for CD8 and intracellular IFN γ and their percentage of CD8 $^{+}$ cells (TC IFN γ).

[0038] Figures 6A-B depict weight change in Balb/c mice implanted with EMT6 triple negative breast cancer cells in control mice (◆) and GA2E treated mice at 10 mg/kg (■), 25 mg/kg (▲), or 50 mg/kg (X) from the first day of treatment (Day 0) through the end of the study, reported as mass (Figure 6A) or percent change (Figure 6B).

[0039] Figure 7 depicts tumor growth, in mm 3 , in Balb/c mice implanted with EMT6 triple negative breast cancer cells in control mice (◆) and GA2E treated mice at 10 mg/kg (■), 25 mg/kg (▲), or 50 mg/kg (X), from the first day of treatment (Day 0) through the end of the study.

[0040] Figures 8A-C depict the production of IFN γ (in pg/ml) by recall antigen-stimulated PBMC treated with vehicle control or a RAR γ selective agonist; Figure 8A GA2E; Figure 8B GA3Ep; Figure 8C tazarotenic acid.

[0041] Figures 9A-C depict PBMC proliferation of recall antigen-stimulated CD8 $^{+}$ PBMC treated with vehicle control or a RAR γ selective agonist; Figure 9A GA2E; Figure 9B GA3Ep; Figure 9C tazarotenic acid.

[0042] Figures 10A-B depict weight change in NSG-B2M mice implanted with a Her2 $^{+}$ breast cancer cell line (JIMT-1) and human PBMC in control (◆) and GA2E treated (■) mice, from the first day of treatment (Day 0) through the end of the study, reported as mass (Figure 10A) or percent change (Figure 10B).

[0043] Figure 11 depicts tumor growth, in mm 3 , in NSG-B2M mice implanted with a Her2 $^{+}$ breast cancer cell line (JIMT-1) and human PBMC in control (◆) and GA2E treated (■) mice, from the first day of treatment (Day 0) through the end of the study.

[0044] Figure 12A-E depicts flow cytometric T subset analysis of CD8 $^{+}$ TIL from NSG-B2M mice implanted with a Her2 $^{+}$ breast cancer cell line (JIMT-1) and human PBMC in control (◆) and GA2E treated (■) mice on Day 26 (termination) of the study. The subsets were total CD8 $^{+}$ T cells (Figure 12A, TC), naive CD8 $^{+}$ T cells (Figure 12B, CD8 Naïve), central memory CD8 $^{+}$ T cells (Figure 12C, CD8 TCM), effector memory CD8 $^{+}$ T cells (Figure 12D, CD8 TEM), and terminally differentiated effector CD8 $^{+}$ T cells (Figure 12E, CD8 TEFF). Significance of the difference between treated and control: * indicates $P < 0.05$; *** indicates $P < 0.001$; no asterisks indicates $P > 0.05$.

[0045] Figure 13A-C depicts flow cytometric biomarker analysis of CD8 $^{+}$ TIL from NSG-B2M mice implanted with a Her2 $^{+}$ breast cancer cell line (JIMT-1) and human PBMC in control (◆) and GA2E treated (■) mice on Day 26 (termination) of the study. The subsets were PD-1 $^{+}$ CD8 $^{+}$ T cells (Figure 13A, TC PD1), CD138 $^{+}$ CD8 $^{+}$ T cells (Figure 13B, TC CD18), and CD54 $^{+}$

CD8⁺ T cells (Figure 13C, TC CD54). Significance of the difference between treated and control: ** indicates $P < 0.005$; *** indicates $P < 0.001$; **** indicates $P < 0.0001$.

[0046] Figure 14A-E depicts flow cytometric T subset analysis of CD4⁺ TIL from NSG-B2M mice implanted with a Her2⁺ breast cancer cell line (JIMT-1) and human PBMC in control (♦) and GA2E treated (■) mice on Day 26 (termination) of the study. The subsets were total CD4⁺ T cells (Figure 14A, TH), naïve CD4⁺ T cells (Figure 14B, CD4 Naïve), central memory CD4⁺ T cells (Figure 14C, CD4 TCM), effector memory CD4⁺ T cells (Figure 14D, CD4 TEM), and terminally differentiated effector CD4⁺ T cells (Figure 14E, CD4 TEFF). Significance of the difference between treated and control: * indicates $P < 0.05$; *** indicates $P < 0.001$; no asterisks indicates $P > 0.05$.

[0047] Figure 15A-D depicts flow cytometric biomarker analysis of CD4⁺ TIL from NSG-B2M mice implanted with a Her2⁺ breast cancer cell line (JIMT-1) and human PBMC in control (♦) and GA2E treated (■) mice on Day 26 (termination) of the study. The subsets were PD-1⁺ CD8⁺ T cells (Figure 15A, TH PD1), CD18β⁺ CD4⁺ T cells (Figure 15B, TH CD18), CD54⁺ CD4⁺ T cells (Figure 15C, TH CD54), and Fox3P⁺ CD4⁺ T cells (Figure 15D, Tregs). Significance of the difference between treated and control: * indicates $P < 0.05$; no asterisks indicates $P > 0.05$.

[0048] Figure 16 depicts dose-response curves for cell viability for EMT6 cells grown in the presence of three RARγ agonists GA1E (■), GA2E (●), and GA3Ep (▲).

[0049] Figure 17 depicts dose-response curves for cell viability for Lewis Lung carcinoma (LLC) cells grown in the presence of three RARγ agonists GA1E (■), GA2E (●), and GA3Ep (▲).

[0050] Figures 18A-B depict weight change in B57BL/6 mice implanted with LLC cells in control mice (♦) and GA2E treated mice at 10 mg/kg (■) or 25 mg/kg (▲), from the first day of treatment (Day 0) through the end of the study, reported as mass (Figure 18A) or percent change (Figure 18B).

[0051] Figure 19 depicts tumor growth, in mm³, in B57BL/6 mice implanted with LLC cells in control mice (♦) and GA2E treated mice at 10 mg/kg (■) or 25 mg/kg (▲), from the first day of treatment (Day 0) through the end of the study.

[0052] Figure 20 depicts the results of flow cytometric analysis of LLC tumors excised at the end of the study described in Example 10, for each of the treatments, vehicle control and GA2E at dosages of 10 and 25 mg/kg in columns 1, 2, and 3, respectively, in each panel. Results are shown both as percentage of total T cells (upper panels) and as absolute number of cells pre mg/tumor (lower panels). Depicted are the number of live cells in the tumors and

their percentage of the total cells (Live), the number of leukocytes based on CD45 staining and their percentage of the live cells (CD45), and the total number of T cells based on CD3 staining and their percentage of the leukocytes (Tot T).

[0053] Figure 21 depicts the results of flow cytometric analysis of LLC tumors excised at the end of the study described in Example 10, for each of the treatments, vehicle control and GA2E at dosages of 10 and 25 mg/kg in columns 1, 2, and 3, respectively, in each panel. Results are shown both as percentage of total T cells (upper panels) and as absolute number of cells per mg/tumor (lower panels). Depicted are the number of CD4⁺ T cell based on CD4 staining and their percentage of total T cells (CD4⁺), T helper cells based on positive staining for CD4 and intracellular IFN- γ (Thelper1), and T regulatory cells based on positive staining for CD4, CD25, and FOXP3.

[0054] Figure 22 depicts the results of flow cytometric analysis of LLC tumors excised at the end of the study described in Example 10, for each of the treatments, vehicle control and GA2E at dosages of 10 and 25 mg/kg in columns 1, 2, and 3, respectively, in each panel. Results are shown both as percentage of total T cells (upper panels) and as absolute number of cells per mg/tumor (lower panels). Depicted are the number of CD8⁺ T cell based on CD8 staining and their percentage of total T cells (CD8⁺), cytotoxic T cells based on positive staining for CD8 and intracellular IFN- γ (Cytotoxic CD8), naïve CD8⁺ T cells based on positive staining for CD8 and CD62L, and negative staining for CD44 (CD8 naïve), CD8⁺ central memory T cells (CD8 Tcm) based on positive staining for CD8, CD62L, and CD44, and CD8⁺ effector memory T cells (CD8 Tem) based on positive staining for CD8 and CD44 and negative staining for CD62L.


DESCRIPTION

[0055] Disclosed herein are methods of expanding tumor-infiltrating lymphocytes (TIL) and associated methods of treating cancer by administering an effective amount of a RAR γ selective agonist to a subject in need thereof.

[0056] As used herein “RAR γ selective agonist” refers to RAR γ agonists that have no or negligible agonistic activity for RAR α at some clinically relevant concentrations. In some embodiments, the RAR γ selective agonist is selective in that it has no or negligible agonistic activity for both RAR α and RAR β at some clinically relevant concentrations. “Clinically relevant concentrations” refers to the concentration in the blood (typically determined by assaying plasma or serum) of a subject receiving efficacious doses of the RAR γ selective agonist, or the concentration used in *in vitro* culture for generating, differentiating, or expanding TIL, as relevant to the method in which the RAR γ selective agonist is being used. In some

embodiments, the EC_{50} for activation of RAR γ is at least 100-fold less, or at least 50-fold less, than for the activation of RAR α , or RAR α and RAR β .


[0057] One aspect is a method of treating cancer comprising administering, to a patient in need thereof, an effective amount RAR γ -selective agonist having a structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified. However, in some embodiments, the E configuration of the =N-OH group is preferred. R⁷ to R¹² are independently: C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom (for example F, Cl, or Br), a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃. Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl. R¹⁶ is H or F. In some embodiments, R is methyl or ethyl. In some embodiments, R is H. In some embodiments, R¹⁶ is F.

[0058] Synthetic schemes for species of GAG1 are presented in Figures 1 and 2.


[0059] In some embodiments, in the RAR γ -selective agonist having a structure of GAG1, Y is O:



[0060] In some embodiments, in the RAR γ -selective agonist having a structure of GAG1, the =N-OH group is in the E configuration:



[0061] In some embodiments, the RAR γ -selective agonist having a structure of GAG1 has the structure



[0062] As used herein, alkyl means a moiety composed of carbon and hydrogen containing no double or triple bonds. Alkyl may be linear alkyl, branched alkyl, cycloalkyl, or a combination thereof. C₁₋₆ alkyl, means an alkyl group having 1 to 6 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, pentyl isomers, hexyl isomers, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and combinations thereof having anywhere from 1-6 carbon atoms. These considerations apply equally to the alkyl portion or an alkoxy group.

[0063] With respect to the herein disclosed structures comprising aromatic rings substituted with R groups, when the R group is an alkoxy group, one valence of the oxygen atom is attached to the aromatic ring and the other valence is attached to the alkyl portion of the alkoxy group.

[0064] One aspect disclosed herein is a method of generating, differentiating, or expanding TIL by contacting them with a RAR γ selective agonist. RAR γ agonists stimulate T cell effector function, but not Treg cells, and thus can promote an anti-cancer immune response. In particular, the RAR γ selective agonists described herein can promote expansion of CD8⁺ TIL. IL-2 has been used previously for expanding TIL. *In vitro* culture can result in changes to the T cell receptor repertoire and the *in vitro*-expanded TIL are typically infused with further IL-2 as a growth factor. However, IL-2 is also associated with a variety of toxicities, some associated with capillary leak syndrome presented by edema, hypotension and reduced urine output within hours of infusion, but also fevers, rigors, myalgia and nausea. An alternative agent to expand TIL without producing such substantial toxicity is a long-felt, unmet need.

[0065] As used herein, the term "TIL" refers primarily to the lymphocytes that can be found within a tumor. Such TIL can be isolated from explanted tumor tissue and expanded *in vitro*. Yet TIL infiltrate the tumor from outside the tumor thus can also be found outside the tumor. Therefore, it is also possible to culture PBMC *in vitro* in the presence of tumor antigen (for example, lethally irradiated tumor cells) to generate or expand T cells capable of infiltrating

into tumors. Methods of generating, differentiating, or expanding TIL encompass such use of PBMC.

[0066] In some embodiments, contacting TIL with a RAR γ selective agonist comprises culturing the TIL *in vitro* in a media supplemented with the RAR γ selective agonist. Procedures for culturing TIL are known to those of skill in the art. To summarize a basic procedure, resected tumor tissue is minced into approximate 1-3 mm³ fragments and placed in culture (for example in 24- or 48-well plates with 2 or 1 mL of culture media, respectively) and the TIL allowed to extravasate from the tissue. Alternatively, the tumor tissue may be subjected to enzymatic digestion and/or mechanical disaggregation to obtain a single cell suspension. For expansion, the TIL can be cultured *in vitro* for 3-6 weeks in an appropriate medium, replacing half the media every 2-3 days and splitting at 80% confluence. Tumor cells disappear from the culture over the first 1-3 weeks. To more rapidly expand the TIL, anti-CD3 antibody and 100 to 200-fold excess of irradiated feeder cells (autologous or allogeneic) can be added to the culture. The TIL can be transferred to a bioreactor to support high cell density and achieve a population of from 10⁹ to 2x10¹¹ cells. The RAR γ selective agonist is used continually in the culture.

[0067] In some embodiments, in which TIL are being generated or expanded from PBMC, the PBMC are obtained from the subject to be treated (autologous). In other embodiments, the PBMC are obtained from an HLA-matched donor. In some embodiments, tumor cells are obtained from the subject to be treated, while in other embodiments they are obtained from a different individual. In some embodiments, the PBMC and the tumor cells are obtained from the same individual. In some embodiments, that individual is the subject to be treated. In some embodiments, the PBMC and the tumor cells are obtained from different individuals, for example, the PBMC can be autologous but cultured with tumor cells from an HLA-matched or allogeneic donor. In some embodiments, the PBMC (or TIL) are cultured with lethally-irradiated tumor cells, but in alternative embodiments the culture contains a tumor cell lysate, or purified or synthetic tumor antigens.

[0068] In some embodiments, contacting comprises administering the RAR γ selective agonist to a subject having a tumor.

[0069] One aspect disclosed herein is a method of treating cancer comprising providing RAR γ selective agonist-expanded TIL to a patient in need thereof. In one embodiment, the RAR γ selective agonist is administered to a subject having a cancer to be treated. In an alternative embodiment, TIL are expanded in culture by exposing the culture to the RAR γ selective agonist, and the RAR γ selective agonist-expanded TIL are administered to a subject having a cancer to be treated as adoptive T cell therapy. In some embodiments, the subject

is a mammal. In some embodiments the subject is a human. In some embodiments, the RARy selective agonist is GAG1. In some embodiments, the RARy selective agonist is GAG2. In some embodiments, the RARy selective agonist is GAG3p. In some instances, the =N-OH group of GAG1, GAG2, or GAG3p is in the E configuration. In still further embodiments, the RARy selective agonist is any individual species or group of species encompassed by structures GAG1, GAG2, or GAG3. Some embodiments specifically, exclude any individual species or group of species encompassed by structures GAG1, GAG2, or GAG3.

[0070] With respect to any relevant aspect, in some embodiments, the cancer is a solid tumor, a carcinoma, a sarcoma, or a hematologic cancer. With respect to any relevant aspect, in some embodiments, the cancer is a melanoma, a glioblastoma, renal cell carcinoma, lung cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, gall bladder cancer, laryngeal cancer, liver cancer, thyroid cancer, stomach cancer, salivary gland cancer, prostate cancer, pancreatic cancer ovarian cancer, a leukemia, a lymphoma, a myeloma, or a plasma cell cancer. With respect to any relevant aspect, in some embodiments, the cancer is breast cancer. In some embodiments, the breast cancer is triple negative breast cancer. In some embodiments, the cancer is Her2⁺ breast cancer. In some embodiments, the cancer is lung cancer, for example, non-small cell lung cancer. In some embodiments, the subject in need thereof is human.

[0071] The growth of many breast cancers is stimulated by estrogen, progesterone and (particularly when the cancer expresses excess human epidermal growth factor receptor 2 (Her2)) epidermal growth factor. Agents that antagonize these effects have proven to be effective treatments for breast cancer. However, about 10-20% of breast cancers do not express estrogen receptor, progesterone receptor, or excess Her2. These cancers are therefore termed “triple negative breast cancer”. Triple negative breast cancer is typically more aggressive and has a poorer prognosis than other types of breast cancer, in part due to there being fewer treatment options. Triple negative breast cancer is also more likely to metastasize and more likely to recur after treatment. Breast cancers are graded on their resemblance to normal healthy breast cells on a 3 point scale, with higher grades indicating less resemblance to normal cells. Triple negative breast cancers are often grade 3. Triple negative breast cancers are usually “basal-like” meaning they resemble the basal cells lining the breast ducts. Basal-like cancers tend to be more aggressive and higher grade. The onset of triple negative breast cancer tends to occur at an earlier age, for example, under 50 as compared to other types of breast cancer which are more commonly diagnosed in persons over 60 years of age. About 70% of cancers having a BRCA mutation are triple negative. Altogether, these factors make triple negative breast cancer a particularly difficult disease to treat.

[0072] Current treatments for triple negative breast cancer include neoadjuvant chemotherapy (chemotherapy prior to surgical removal of the tumor); inhibitors of poly ADP-ribose polymerase (PARP), such as olaparib; and immunotherapy for example by PD-1 blockade, such as by atezolizumab, in combination with albumin-bound paclitaxel.

[0073] In some embodiments, in which the RAR γ selective agonist is administered to the subject, the effective amount is about 0.01 to about 300 mg/m²/day; however, doses below or above this exemplary range are within the scope of the present disclosure. The daily dose can be about 0.5 to about 100 mg/m²/day, about 1 to about 90 mg/m²/day, about 5 to about 80 mg/m²/day; or at least about 0.02, 0.03, 0.05, 0.07, 0.1, 0.2, 0.3, 0.5, 0.7, 1, 2, 3, 5, 7, 10, 15, 20, 25, 30, 50, 70, or 100 mg/m²/day; or not more than about 0.1, 0.2, 0.3, 0.5, 0.7, 1, 2, 3, 5, 7, 10, 15, 20, 25, 30, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, 225, 250, 275, or 300 mg/m²/day; or a range defined by any two of the foregoing values. These dosages may be converted into approximately equivalent human dosages in mg/kg/day by dividing by 37.

[0074] In some embodiments, the effective amount of the RAR γ agonist is about 0.27 μ g/kg/day to about 8 mg/kg/day; however, doses below or above this exemplary range are within the scope of the present disclosure. The daily dose can be about 0.013 to about 2.7 mg/kg/day, about 0.025 to about 2.5 mg/kg/day, about 0.130 to about 22 mg/kg/day; or at least about 0.0005, 0.0008, 0.001, 0.0013, 0.0020, 0.0025, 0.005, 0.0008, 0.0010, 0.0013, 0.0020, 0.0027, 0.005, 0.008, 0.010, 0.013, 0.020, 0.027, 0.05, 0.08, 0.1, 0.13, 0.2, 0.5, 0.8, 1.0, 1.3, 1.8, 2.0, or 2.7 mg/kg/day; or not more than about 0.0027, 0.005, 0.008, 0.010, 0.013, 0.020, 0.027, 0.05, 0.08, 0.1, 0.13, 0.2, 0.5, 0.8, 1.0, 1.3, 1.8, 2.0, or 2.7, 3.3, 4.0, 4.7, 5.4, 6.1, 6.8, 7.4, or 8.0 mg/kg/day, or a range defined by any two of the foregoing values. These dosages may be converted into approximately equivalent human dosages in mg/m²/day by multiplying by 37.

[0075] In some embodiments, the effective amount for a human is about 0.006 to about 200 mg/day or about 1 to about 100 mg/day. In some embodiments, the daily dose is given in a single administration. In other embodiments, the daily dose is split into multiple administrations, for example two administrations, 9-15, 10-14, 11-13, or 12 hours apart.

[0076] It is not necessary that the RAR γ selective agonist be present at a therapeutic level in the subject throughout treatment. This is because the primary therapeutic effect of the RAR γ selective agonist is indirect, promoting an immune response, the expansion of TIL, which persists in the absence of the RAR γ selective agonist. Because it takes time for the immune response to develop, the beneficial effect may not be observed until administration of the RAR γ selective agonist is paused. Indeed, in some embodiments, it may be beneficial to have a drug holiday during which administration of the RAR γ selective agonist is suspended. To the

extent that administration of the RARy selective agonist is associated with a toxicity or other undesirable side-effect, a suspension of administration allows time for the subject to recover from the ill effects, rather than for them to continue to worsen. As used herein "suspended" refers to a cessation of drug administration with the intent and expectation that it will resume at a later time and is distinct from discontinuation in which there no plan to resume administration of the drug. Resumption of administration of can be beneficial as immune responses tend to wane over time. Thus, with to any aspect comprising administering the RARy selective agonist to a subject, in some embodiments, the administering occurs periodically throughout an interval of treatment. That is, the RARy agonist is administered at regularly occurring time points within the interval of treatment. In some embodiments, periodically is twice daily, once daily, every other day, every third day, or twice weekly. In some embodiments, the administering occurs in repeated cycles throughout an interval of treatment. In some embodiments, a cycle comprises 1) administering the RARy selective agonist periodically over a first span of time and 2) suspending administration of the RARy selective agonist over a second span of time, after which a new cycle may be initiated. In some embodiments, the first span of time (during which the RARy agonist is administered) is 10-15 days, or any integer number of days therein. In some embodiments, the second span of time (during which administration of the RARy agonist is suspended) is two weeks to one month or any integer number of days therein. In various embodiments, the interval of treatment extends from a first administration of the RARy agonist until a complete response is achieved, stable disease is obtained, or disease progression occurs.

[0077] The RARy selective agonist may be administered to a subject by any suitable route of administration. In some embodiments, the RARy a selective agonist is administered orally. In some embodiments, the RARy selective agonist is administered by injection or infusion, for example, intravenously, subcutaneously, or intratumorally. In some embodiments, the RARy selective agonist is administered intranasally. In some embodiments, the RARy selective agonist is administered by nasal or oral (pulmonary) inhalation.

[0078] *Ex vivo* expanded TIL are generally administered by infusion, for example intravenous or intratumoral infusion. However, in some embodiments, *ex vivo* expanded TIL are administered in a single bolus or split into multiple boluses, intravenously, or intratumorally. In some embodiments, the RARy selective agonist is included in the bolus or infusion with the TIL. In various embodiments, infusion may extend for half an hour, for an hour, for several hours, for a day, or for several days.

[0079] In some embodiments, about 10^9 to $>10^{11}$ TIL are administered, for example, about 1×10^{10} to 2×10^{11} . The RARy selective agonist is used continually in the culture and may be included with the infused, expanded TIL. The RARy agonist may also be administered to the

patient subsequent to TIL infusion for a set period of time or until tumor is eliminated or no further benefit from the TIL is observed.

[0080] In some embodiments, the RAR γ selective agonist is used as monotherapy. In other embodiments, it is used in combination with one of the current therapies, either before, during, or after, the other treatment.


[0081] RAR γ selective agonists promote expansion of TIL that are CD8 $^{+}$ effector cells. Consistent with what is known about CD8 $^{+}$ effector cells generally, these cells express PD-1, which can act as a negative regulator of CD8 $^{+}$ effector cell activity. Thus, in some embodiments, treatment with a RAR γ selective agonist is combined with PD-1 blockade. Treatment with RAR γ selective agonists can also increase the presence of PD-1 $^{+}$ T cells among CD8 $^{+}$ TIL. For this reason as well, in some embodiments, treatment with a RAR γ selective agonist is combined with PD-1 blockade. Several products for PD-1 blockade, also called immune checkpoint inhibition, are undergoing clinical evaluation. These include the anti-PD-1 antibodies pembrolizumab, nivolumab, cemiplimab, dostarlimab, tislelizumab, spartalizumab, camrelizumab, sintilimab, toripalimab, JTX-4014, INCMGA00012, AMP-514, and budigalimab, and the anti-PD-L1 antibodies atezolizumab, avelumab, durvalumab, envafolelimab, CK-301, CS-1001, KN035SHR-1316, CBT-502, BGB-A333, and BMS-936559. Several non-antibody inhibitors of PD-L1 are also in development, including AUNP12 (a 29mer peptide), CA-170 (a small organic molecule), and BMS-986189 (a macrocyclic peptide). AMP-224 is a fusion protein of PD-L2 (also known as B7-DC) with an antibody Fc region, which is being developed as an anti-PD-1 checkpoint inhibitor. Such antibodies constitute means for PD-1 blockade.

[0082] A growing mode of cancer treatment is chimeric antigen receptor T cell (CAR-T) therapy. CAR-T cells are effectively artificial TIL. Accordingly, in some embodiments, treatment with a RAR γ selective agonist, as described herein, is combined with CAR-T therapy. RAR γ -selective agonists can potentiate chimeric antigen receptor-modified immune cells (CAR-MIC) therapy, as described in US2019001563A1 and US20180338940A1 (each of which is incorporated by reference for all that they teach about potentiation of CAR-MIC with RAR γ agonists). Thus, some embodiments are methods of potentiating CAR-MIC cancer immunotherapy comprising administering a RAR γ -selective agonist having the structure of GAG1 to a cancer patient who is receiving, has received, or is scheduled to receive, CAR-MIC. In some instances, the CAR-MIC is a CAR-T cell.

[0083] With respect to any aspect comprising administering the RAR γ selective agonist to a subject, some embodiments further comprise administering an inhibitor of regulatory T cells (Tregs). In some embodiments, the inhibitor of Tregs is a Treg-depleting antibody. In various embodiments, the Treg-depleting antibody is anti-CD25 antibody, an anti-GITR antibody, an

anti-FoxP3 antibody, an anti-CCR₄ antibody, or an anti-folate receptor 4 antibody. In some embodiments, the inhibitor of Tregs comprises a RAR α antagonist.

[0084] With respect to any aspect comprising administering the RAR γ selective agonist to a subject, some embodiments further comprise administering an RXR agonist having the structure



where R is H or C₁₋₆ alkyl, or a pharmaceutically acceptable salt thereof. In some embodiments, R is methyl or ethyl. In some embodiments, R is H; that is the RXR agonist is 3,7-dimethyl-6(S),7(S)-methano-7-[1,1,4,4-tetramethyl-1,2,3,4-tetrahydro-naphth-7-yl]2(E),4(E) heptadienoic acid, also known as IRX4204. Compounds with the structure of XAG have anticancer activity as described in US 2008/0300312A1 and US 2020/0390736A1 (which are incorporated by reference for all that they teach about the treatment of cancer with these RXR agonists). Treatment with RAR γ selective agonists can have undesirable side effects, such as weight loss that can become excessive, especially at higher dosages, as seen in some of the Examples below. Compounds with the structure of XAG also have an anticachectic effect, as described in US 20070185055A1 (which is incorporated by reference for all that it teaches about the treatment of cachexia with RXR agonists), which could counteract or mitigate the side effects of the RAR γ agonist contributing to potentially excessive weight loss. Accordingly, for either or both of these reasons, in some embodiments treatment with a RAR γ selective agonist, as described herein, is combined with treatment with an appropriate RXR agonist, such as IRX4204.

[0085] The average surface area of a human body is generally accepted to be 1.9 m² for an adult male, 1.6 m² for an adult female, and 1.33 m² for a 12-13 year old child. These values can be used to calculate dose ranges for doses of the RAR γ selective agonist. The total daily dosage of RAR γ selective agonist active agents can be administered as a single dose or as two doses administered with a 24 hour period spaced 8 to 16, or 10 to 14, hours apart. For repeated administrations over several days or longer, the treatment can be repeated until a desired suppression of disease or disease symptoms occurs. However, other dosage regimens may be useful and are within the scope of the disclosure. The desired dosage can be delivered by a single bolus administration of the composition, by multiple bolus administrations of the composition, or by continuous infusion administration of the composition.

[0086] The RARy selective agonist can be administered to a mammal using standard administration techniques, including parenteral, oral, intravenous, intraperitoneal, subcutaneous, pulmonary, transdermal, intramuscular, intranasal, buccal, sublingual, or suppository administration. The term "parenteral," as used herein, includes intravenous, intramuscular, subcutaneous, rectal, vaginal, and intraperitoneal administration. The RARy selective agonist preferably is suitable for oral administration, for example as a pill, tablet or capsule.

[0087] Administration may be continuous or intermittent. The dosage may also be determined by the timing and frequency of administration. Thus, the RARy selective agonist disclosed herein can be given on a daily, weekly, biweekly, or monthly basis for a period of time, followed by an optional drug holiday (drug free period) and that this drug administration/drug holiday cycle can be repeated as necessary. For example, the RARy selective agonist could be administered daily for a week, and then not administered for the rest of a month, and then administered for another week, etc. In certain embodiments, the total daily dosage of RARy agonist can be administered as a single dose or as two doses administered with a 24-hour period spaced 8 to 16, or 10 to 14, hours apart.

[0088] The effectiveness of cancer therapy is typically measured in terms of "response." The techniques to monitor responses can be similar to the tests used to diagnose cancer such as, but not limited to:

- A lump or tumor involving some lymph nodes can be felt and measured externally by physical examination.
- Some internal cancer tumors will show up on an x-ray or CT scan and can be measured with a ruler.
- Blood tests, including those that measure organ function can be performed.
- A tumor marker test can be done for certain cancers.

[0089] Regardless of the test used, whether blood test, cell count, or tumor marker test, it is repeated at specific intervals so that the results can be compared to earlier tests of the same type.

[0090] Response to cancer treatment is defined several ways:

- Complete response - all of the cancer or tumor disappears; there is no evidence of disease. Expression level of tumor marker (if applicable) may fall within the normal range.
- Partial response - the cancer has shrunk by a percentage but disease remains. Levels of a tumor marker (if applicable) may have fallen (or increased, based on the tumor marker, as an indication of decreased tumor burden) but evidence of disease remains.

- Stable disease - the cancer has neither grown nor shrunk; the amount of disease has not changed. A tumor marker (if applicable) has not changed significantly.
- Disease progression - the cancer has grown; there is more disease now than before treatment. A tumor marker test (if applicable) shows that a tumor marker has risen.

[0091] Other measures of the efficacy of cancer treatment include intervals of overall survival (that is time to death from any cause, measured from diagnosis or from initiation of the treatment being evaluated)), cancer-free survival (that is, the length of time after a complete response cancer remains undetectable), and progression-free survival (that is, the length of time after disease stabilization or partial response that resumed tumor growth is not detectable).


[0092] There are two standard methods for the evaluation of solid cancer treatment response with regard to tumor size (tumor burden), the WHO and RECIST standards. These methods measure a solid tumor to compare a current tumor with past measurements or to compare changes with future measurements and to make changes in a treatment regimen. In the WHO method, the solid tumor's long and short axes are measured with the product of these two measurements is then calculated; if there are multiple solid tumors, the sum of all the products is calculated. In the RECIST method, only the long axis is measured. If there are multiple solid tumors, the sum of all the long axes measurements is calculated. However, with lymph nodes, the short axis is measured instead of the long axis.

[0093] The terms "treating" or "treatment" broadly include any kind of treatment activity, including the mitigation, or prevention of disease, or an aspect thereof, in man or other animals, or any activity that otherwise affects the structure or any function of the body of man or other animals. Treatment activity includes the administration of the medicaments, dosage forms, and pharmaceutical compositions described herein to a patient, especially according to the various methods of treatment and methods of generating, differentiating or expanding TIL disclosed herein, whether by a healthcare professional, the patient his/herself, or any other person. Treatment activities include the orders, instructions, and advice of healthcare professionals such as physicians, physician's assistants, nurse practitioners, and the like, that are then acted upon by any other person including other healthcare professionals or the patient him/herself. This includes, for example, direction to the patient to undergo, or to a clinical laboratory to perform, a diagnostic procedure, such as for cancer diagnosis and staging, so that ultimately the patient may receive the benefit appropriate treatment. In some embodiments, the orders, instructions, and advice aspect of treatment activity can also include encouraging, inducing, or mandating that a particular medicament, or combination thereof, be chosen for treatment of a condition - and the medicament is actually used - by approving insurance coverage for the medicament, denying coverage for an alternative medicament,

including the medicament on, or excluding an alternative medicament, from a drug formulary, or offering a financial incentive to use the medicament, as might be done by an insurance company or a pharmacy benefits management company, and the like. In some embodiments, treatment activity can also include encouraging, inducing, or mandating that a particular medicament be chosen for treatment of a condition - and the medicament is actually used - by a policy or practice standard as might be established by a hospital, clinic, health maintenance organization, medical practice or physicians group, and the like. All such orders, instructions, and advice are to be seen as conditioning receipt of the benefit of the treatment on compliance with the instruction. In some instances, a financial benefit is also received by the patient for compliance with such orders, instructions, or advice. In some instances, a financial benefit is also received by the healthcare professional for compliance with such orders, instructions, or advice.


RARy Selective Agonists

[0094] One aspect comprises an RARy selective agonist, wherein the RARy selective agonist is a compound having a structure of




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl. However, in some embodiments, the E configuration of the =N-OH group is preferred. R⁷ to R¹² are independently: C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom (for example F, Cl, or Br), a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃. Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl. In some embodiments, R is methyl or ethyl. In some embodiments, R is H. Some embodiments are a pharmaceutical composition or formulation comprising a compound having the structure of GAG1 or a pharmaceutically acceptable salt thereof. With respect to any method aspect, in some embodiments, the RARy selective agonist is a compound having the structure of GAG1 or a pharmaceutically acceptable salt thereof.


[0095] In some embodiments, in the RARy-selective agonist having a structure of GAG1, Y is O:




[0096] In some embodiments, in the RAR γ -selective agonist having a structure of GAG1, the =N-OH group is in the E configuration:




[0097] With respect to any aspect, in some embodiments, the RAR γ selective agonist is 4-((1E,3E)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1E), having the structure



[0098] In some embodiments, the RAR γ selective agonist is 4-((1E,3Z)-3-(3-(*tert*-butyl)-2-ethyl=benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1Z), having the structure




[0099] With respect to any method of use aspect, or any method of generating, differentiating, or expanding TIL, in some embodiments, the RAR γ selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified. However, in some embodiments, the E configuration of the =N-OH group is preferred. In some embodiments, R is methyl or ethyl. In some embodiments, R is H. In some embodiments, R is methyl or ethyl.


[0100] In some embodiments, the RARy selective agonist is 3-fluoro-4-((1E,3E)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2E), having the structure



In some embodiments, the RARy selective agonist is 3-fluoro-4-((1E,3Z)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2Z), having the structure




[0101] With respect to any method of use aspect, or any method of generating, differentiating, or expanding TIL, in some embodiments, the RARy selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; each R¹ and R² are independently H or C₁₋₆ alkyl; R³ and R^{3'} are independently H, or halogen (for example, Cl, F,

or Br); $(R^4)_2$ comprises R^4 and $R^{4'}$ which are independently H, halogen (for example, Cl, F, or Br), C_{1-6} alkyl or C_{1-6} alkoxy; X is O, S, CH_2 , $C(R^5)_2$, or NR^6 , wherein each R^5 and R^6 are independently H or C_{1-6} alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; and the COOR group is in the meta or para position and the two R^4 groups occupy the remaining positions on the ring. In some embodiments, the para-position of the COOR group is preferred. In some embodiments, the E configuration of the =N-OH group is preferred. In some embodiments, both R^1 are CH_3 . In some embodiments, both R^2 are H. In some embodiments, X is $C(R^5)_2$. In some embodiments, both R^5 are CH_3 . In some embodiments, one R^3 is H and the other R^3 is F. In some embodiments, both R^4 are H. In some embodiments, all R^3 and R^4 are H. In some embodiments, R is H. In some embodiments, R is methyl or ethyl. In some embodiments, the para- position of the COOR group is preferred. In some embodiments, the E configuration of the =N-OH group is preferred. In some embodiments, both R^1 are CH_3 . In some embodiments, both R^1 are H. In some embodiments, both R^2 are H. In some embodiments, X is $C(R^5)_2$. In some embodiments, X is CH_2 . In some embodiments, both R^5 are CH_3 . In some embodiments, both R^3 are H, while in other embodiments, one R^3 is H and one R^3 is F. In some embodiments, both R^4 are H. In some embodiments, all R^3 and R^4 are H. In some embodiments, the COOR group is in the para position. In some embodiments, the COOR group is in the meta position. In some embodiments, the =N-OH group is in the E configuration. In some embodiments the =N-OH group is in the Z configuration. In particular embodiments, R is H and the carboxylic acid group is in the para position, both R^1 are CH_3 , both R^2 are H, X is $C(CH_3)_2$, all R^3 and R^4 are H, and the =N-OH group is in the E configuration (GA3Ep). Some embodiments specifically include one or more specific substituents at one or more of the variable positions. Some embodiments specifically include one or more of the alternatives, while other embodiments specifically exclude one or more of these alternatives.


[0102] With respect to GAG3, in some embodiments the RAR γ selective agonist has the structure




and other embodiments have the structure




[0103] With respect to GAG3, in some embodiments the RARy selective agonist has the structure



[0104] In some embodiments, the RARy selective agonist is a compound with the structure



and in other embodiments, the RARy selective agonist is a compound with the structure




[0105] With respect to any of the preceding RARy selective agonists of this section comprising a COOR group, in some embodiments R is H. In embodiments in which R is C₁₋₆ alkyl, the alkyl part of the ester has anywhere from 1 to 6 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, pentyl isomers, hexyl isomers, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and combinations thereof having anywhere from 1-6 carbon atoms, etc. The various subsets (including individual species) and combinations of these esters are contemplated as further distinct embodiments, including straight-chain, branched, and/or cyclic moieties of any length or set of lengths within the C₁₋₆ alkyl genus.

[0106] With respect to any of the preceding RARy selective agonists of this section comprising a =N-OH group, some embodiments comprise either configuration, while other

embodiments comprise only the E configuration, and still other embodiments comprise only the Z configuration, of this group.

[0107] With respect to any method of use aspect, in some embodiments the RAR γ selective agonist is tazarotenic acid (selective for RAR γ over RAR α , but not over RAR β), CD437, CD2325, CD666, trifarotene, or BMS961.



[0108] In some embodiments, the RAR γ agonist is a RAR γ -selective agonist in that it has no or only negligible agonistic activity with RAR α and RAR β at clinically relevant concentrations. In some embodiments, the RAR γ agonist is a RAR γ -selective agonist in that it has no or only negligible agonistic activity with RAR α at clinically relevant concentrations. In various embodiments, a RAR γ selective agonist has no or only negligible agonistic activity with RAR α or RAR β at some clinically relevant concentrations when the agonist's KD for RAR γ is greater than 10-, 20-, 50-, or 100-fold lower than for the other RAR(s). KD is typically determined in a binding assay. In various embodiments, a RAR γ selective agonist has no or only negligible agonistic activity with RAR α or RAR β at some clinically relevant concentrations when the agonist's EC₅₀ with RAR γ is greater than 10-, 20-, 50-, or 100-fold lower than for the other RAR(s). EC₅₀ is typically determined in an activation assay, for example, a transactivation assay. In some embodiments where a specific criteria of selectivity is not stated, the RAR γ selective agonist meets at least one of these criteria, for example the least stringent.

[0109] These RAR γ agonists constitute means for activating RAR γ , or means for generating, differentiating, or expanding TIL. Some embodiments specifically include one or

more of the disclosed genera, sub-genera, or species of these RARy agonists. Some embodiments specifically exclude one or more of the disclosed genera, sub-genera, or species of these RARy agonists.

EXAMPLES

[0110] The following non-limiting examples are provided for illustrative purposes only in order to facilitate a more complete understanding of representative embodiments now contemplated. These examples should not be construed to limit any of the embodiments described in the present specification.

Example 1

Synthesis of GAG1

[0111] The synthetic scheme for the synthesis of species of GAG1, GA1E and GA1Z, are presented in Figure 1.

[0112] *Ethyl-2-oxo-ethyl-5-bromo-salicylate-2-yl-butyrates*

[0113] To a solution of ethyl-5-bromo-salicylate (Compound 1, 24.5 g, 100 mmol) in acetone (400 mL), were added ethyl-2-bromo butyrate (23.4 g, 120 mmol) and potassium carbonate (20.7 g, 150 mmol) and the mixture stirred for 24 h at ambient temperature. Solid material was removed by filtration and the solvent was removed by distillation to yield the title compound (not shown in Figure 1) as a yellow oil. Yield = 23.2 g.

[0114] *5-Bromo-2-ethyl-benzofuran(2H)-3-one (Compound 2)*

[0115] To a cold (0°C) solution of ethyl-2-oxo-ethyl-5-bromo Salicylate-2-yl)-butyrate (23.2 g, 70.3 mmol) in dry toluene (300 mL) was added 20% sodium ethoxide in ethanol (27 g, 80 mmol) in small portions (5 min). The cooling bath was removed and the reaction was stirred at ambient temperature for 12 h. The reaction mixture was washed with water (25 mL) and brine (50 mL). The organic layer was dried and the solvent was removed by distillation under vacuum. The crude product was purified by silica gel column chromatography (10% ethyl acetate in hexane) to afford the title compound (Compound 2) as a white solid. Yield = 16.5 g

[0116] ¹HNMR (CDCl₃): δ 1.09 (triplet, 3H), 1.80 – 1.95 (m, 1H), 2.05 – 2.20 (m, 1H), 4.61 (t, 2), 7.12 (d, 1H), 7.70 (dd, 1H), 7.83 (d, 1H).

[0117] *5-Bromo-3-*t*-butyl-2-ethyl-benzofuran (Compound 3)*

[0118] To a cold (-78°C) suspension of cerium chloride (25.41 g, 103 mmol) in THF (250 mL) was added *t*-BuMgCl in ether (2M, 41.2 mL, 82.4 mmol), and the resulting mixture stirred for 30 min. Then 5-bromo-2-ethyl-benzofuran(2H)-3-one (Compound 2) (16.5 g, 68.66 mmol) in THF (100 mL) was added and the mixture stirred at ambient temperature for 30 min. The reaction was quenched by adding MeOH (-78°C), diluted with ethyl acetate (100 mL), washed with aq. NH₄Cl, water and brine (20 mL each). The organic layer was dried and the solvent was removed by distillation. The resulting residue was a mixture of the tertiary alcohol and unreacted compound 2, the mixture was used in the next step without further purification.

[0119] A mixture of the crude tertiary alcohol (obtained above), dichloromethane (300 mL) and *p*-TSA (200 mg) was stirred for 12 h at ambient temperature. The mixture was washed with aq. NaHCO₃, water and brine (100 mL each). The organic layer was dried and the solvent was removed by distillation. The residual crude product was purified by silicagel chromatography (3% ethyl acetate/ hexane) to afford the title compound (Compound 3) as colorless oil. Yield = 2.8 g

[0120] ¹HNMR (CDCl₃): δ 1.35 (triplet, 3H), 1.54 (s, 9H), 2.97 (q, 2H), 7.28 (d, 1H), 7.33 (dd, 1H), 7.88 (d, 1H).

[0121] *1-(3-*tert*-butyl-2-ethyl-benzofuran-5-yl)-ethanone (Compound 4)*

[0122] To a cold (-78°C.) solution of 5-bromo-3-*t*-butyl-2-ethyl benzofuran (Compound 3) (2.8 g, 10 mmol) in THF (60 mL) was added *n*-BuLi in hexane (2.5 M solution, 4.8 mL, 12 mmol). The mixture was gradually warmed to approximately 10°C over 50 min, then cooled again to -78°C and *N*-methoxyl-*N*-methylacetamide (1.23 g, 12 mmol) was added via syringe (neat). Cooling was removed and the reaction was stirred at ambient temperature for 12 hr. The reaction was quenched by adding aq. NH₄Cl, and the mixture was stirred for 15 min. The mixture was diluted with ethyl acetate (60 mL) washed with water and brine (10 mL each). The organic layer was dried and the solvent was removed by distillation. The product (Compound 4) was purified by silica gel chromatography (7% ethyl acetate in hexane), yield = 2.1 g.

[0123] ¹HNMR (CDCl₃): δ 1.33 (triplet, 3H), 1.55 (s, 9H), 2.68 (s, 3H), 2.97 (q, 2H), 7.87 (d, 1H), 8.41 (d, 2H).

[0124] *4-[(*E*)-3-(3-*tert*-butyl-2-ethyl-benzofuran-5-yl)-3-oxo-prop-1-enyl]-3-fluoro-benzoic acid (Compound 5)*

[0125] Methyl 3-fluoro-4-formylbenzoate (1.56 g, 8.6 mmol) was added to a solution of 1-(3-*tert*-butyl-2-ethyl-benzofuran-5-yl)-ethanone (Compound 4) (2.1 g 8.6 mmol) in 10 mL of 1 N sodium hydroxide and 20 mL of methanol. After stirring at room temperature for 18 h, the

reaction mixture was acidified with 1N HCl and extracted with ethyl acetate (3x10 mL). The combined organic layer was washed with brine (1x10 mL), dried (MgSO₄) and concentrated at reduced pressure. The product (Compound 5) was purified by silicagel chromatography (50% ethyl acetate in hexane), yield = 1.6 gram.

[0126] ¹HNMR (CDCl₃): δ 1.39 (triplet, 3H), 1.60 (s, 9H), 3.02 (q, 2H), 7.32 (s, 1H), 7.51 (d, 1H), 7.83 (t, 1H), 7.85 (d, 1H), 7.92 (d, 1H), 7.99 (d, 1H), 8.03 (d, 1H), 8.52 (d, 1H).

[0127] 4-[(E,3E)-3-(3-*tert*-butyl-2-ethyl-benzofuran-5-yl)-3-hydroxyimino-prop-1-enyl]-3-fluoro-benzoic acid (GA1E)

[0128] To a solution of 4-[(E)-3-(3-*tert*-butyl-2-ethyl-benzofuran-5-yl)-3-oxo-prop-1-enyl]-3-fluoro-benzoic acid, (Compound 5) (1.60 g, 4.06 mmol) in 10 mL of ethanol was added hydroxylamine hydrochloride (653 mg, 9.40 mmol) and pyridine (1.86 g, 23.5 mmol). The reaction mixture was then heated at reflux for 6 h. After cooling to room temperature, the solvent was removed *in vacuo* and the residue was taken up in water. The aqueous layer was adjusted to pH 4 - 5 with 1 N HCl and the product was isolated by filtration. Further purification was done by silica gel chromatography (10% methanol/1% triethylamine/dichloromethane), yield = 660 mg of pure E-isomer (GA1E) and 98 mg of impure Z-isomer (GA1Z) due to incomplete separation.

[0129] GA1E: ¹HNMR (DMSO-D₆): δ 1.31 (triplet, 3H), 1.50 (s, 9H), 2.97 (q, 2H), 6.95 (d, 1H), 7.38 (d, 1H), 7.57 (d, 1H), 7.71 (d, 1H), 7.82 (d, 1H), 7.86 (s, 1H), 7.90 (d, 1H), 7.97 (t, 1H).

Example 2

Specificity of Binding and Activity of GA1E and GA2E

[0130] Transactivation assays were conducted to test the specificity of binding activity of GA1E and GA2E for RAR α and RAR γ . Retinoic acid receptor transactivation activity and binding efficiencies were determined essentially as described in U.S. Pat. Nos.: 5,298,429 and 5,071,773, incorporated by reference herein for all they teach regarding transactivation assays. Transactivation assays employed expression plasmids encoding the full-length receptors RAR α and RAR γ . Reporter plasmids containing the herpes virus thymidine kinase promoter and the appropriate retinoic acid receptor response element (RAREs) were positioned upstream of an open coding region encoding firefly luciferase.

[0131] The ability of GA1E and GA2E to activate RAR α and RAR γ in a transactivation assay was tested over a concentration range of 0.5 to 625 nM (GA2E) or .05 to 3320 nM (GA1E). GA1E induced less than 10% transactivation for RAR α across the whole concentration range, but over 50% transactivation for RAR γ at concentrations $\geq$ 25nM and over 70% at

concentrations ≥ 125 nM (Figure 3A). GA2E induced more than 85% transactivation at all tested concentrations and less than 10% transactivation of RAR α at concentrations of ≤ 25 nM (Figure 3B).

Example 3

Treatment of Triple Negative Breast Cancer in Mice with RAR γ Agonists

[0132] The RAR γ agonists tazarotenic acid and GA2E were evaluated in the EMT6 syngeneic mouse model of breast cancer. 5×10^5 EMT6 cells were injected into the mammary fat pad of immunocompetent female Balb/c mice. When the tumors reached an average volume of 50-150 mm³, mice were matched by tumor size into control or treatment groups (n=10-11).

[0133] The mice were treated with 21 daily oral doses 10 mg/kg of 1 mg/ml tazarotenic acid or GA2E, or vehicle control, starting on Day 0. Tumor volume was measured, and the mice weighed, on days 0, 1, 3, 6, 8, 10, 13, 15, 17, 20, and 21.

[0134] Both tazarotenic acid and GA2E inhibited tumor growth, with GA2E having the greater efficacy (Figure 4A). The inhibition of tumor growth did not become apparent until more than 10 days after dosing had begun. At the end of the study, tazarotenic acid had inhibited tumor growth by 44% and GA2E by 63%. While the vehicle control group had a small weight gain, both treatment groups experienced similar but modest weight loss, between 15 and 20% (Figure 4B), indicating the drug was acceptably well-tolerated for a cancer therapeutic.

Example 4

Increased TIL in Tumors from Triple Negative Breast Cancer in Mice with an RAR γ Selective Agonist

[0135] At the end of the study of Example 3, the tumors were excised, the cells disaggregated and dispersed without use of proteases, stained with various agents, and subjected to flow cytometry. Figure 5A-B present the results of the flow cytometry both as a percentage of cells and as the number of cells per tumor. Generally, only a minority of cells in the tumor were alive, as revealed by staining with the dye FVS700, which penetrates and stains necrotic cells (Figure 5A, "Live" panel). Of these live cells, a substantial proportion were leukocytes (Figure 5A, "CD45" panels). Staining with anti-CD3 (Figure 5A, "Tot T" panels) revealed a mean of only about 5% and 10% of the leukocytes were T cells for the vehicle and tazarotenic acid treated mice, respectively, but in the tumors from the mice treated with GA2E a mean of about 30% of the leukocytes were T cells; a statistically significant difference. The majority of the T cells were CD4⁺ (Figure 5A, "CD4" panels), expected to be predominantly helper cells. Of the CD4⁺ cells, only small proportions were T memory cells (expected to be predominantly TH1 memory cells) based on intracellular staining for IFN γ (Figure 5B, "TH

IFGy" panels), TH17 cells based in intracellular staining for IL-17 (Figure 5B, "TH IL17" panels), or Treg cell based on staining for CD25 and FoxP3 (Figure 5B, "Treg" panels). About 25-30% of the T cells were CD8⁺ (Figure 5B, "CD8 T" panels), expected to be predominantly cytotoxic cells. The proportion of CD8⁺ T cells that were memory cells, based on intracellular staining for IFN γ (Figure 5B, "TC IFN γ " panels), was numerically increased in the tumors from the mice treated with tazarotenic acid and GA2E, though the difference from the vehicle control animals did not achieve statistical significance. Thus GA2E treatment increased the number of TIL.

Example 5

Treatment of Triple Negative Breast Cancer in Mice with Various Dosages of an RAR γ Agonist

[0136] The EMT6 triple negative breast cancer model, essentially as described above, was used to assess the effects of increased dosages of GA2E. The mice were orally administered 10 (n=10), 25 (n=10), or 50 (n=11) mg/kg of GA2E, or vehicle control (n=10) daily for 17 days. However, all of the mice in the 50 mg/kg group were terminated for excessive weight loss not later than Day 12, as was one mouse in the 25 mg/ml group. Two mice in the vehicle control group were terminated at Day 12 with tumors >1500 mm³.

[0137] Overall, the 10 and 25 mg/kg were acceptably well-tolerated for a cancer therapeutic, with only a short excursion to more than 10% weight loss for the 25 mg/kg dosage (Figures 6A-B). The 25 mg/kg dosage was somewhat more effective at inhibiting tumor growth than the 10 mg/kg dosage. The tumor inhibition by the 50 mg/kg dosage was not clearly different than that of the other dosages in the limited time those animals were in the study (Figure 7). Scaling to human according to body surface area, 10 and 25 mg/kg in the mouse correspond to approximately 0.8 and 2.0 mg/kg in humans, respectively.

Example 6

RAR γ Selective Agonists Promote Activity and Proliferation of CD8⁺ Memory T Cells

[0138] Peripheral blood mononuclear cells (PBMC) from a cytomegalovirus (CMV)-positive donor were labeled with carboxyfluorescein succinimidyl ester (CFSE) and stimulated with CMV recall antigens at 1 mg/ml. The PBMC were then treated with one of three different RAR γ selective agonists or vehicle (DMSO) control and incubated at 37°C, 5% CO₂ for four days. Culture supernatant was harvested and assayed for IFN γ production and the PBMC evaluated for proliferation by flow cytometry for CFSE dilution.

[0139] CFSE is a fluorescent reactive cell membrane permeable dye. After it is taken up, CFSE covalently modifies proteins and is retained in cells for long periods of time. The CFSE-modified proteins are generally evenly divided between daughter cells at each division so that

fluorescent intensity of the cells falls by half at each division. Thus diminution of the dye's fluorescent signal can be used as a measure of proliferation.

[0140] Treatment with RAR γ selective agonists increased IFN γ production substantially over the vehicle control at concentrations of 0.1 to 1000 nM for each of GA2E, GA3Ep, and tazarotenic acid (Figures 8A-C). Proliferation of CD8 $^{+}$ cells was also increased (Figure 9A-C). Although the increase over the background level of proliferation is small, it is still quite meaningful as only a small proportion of the CD8 $^{+}$ cells would be expected to be specific for CMV.

Example 7

RAR γ Selective Agonist Promotes Generation of Human CD8 $^{+}$ Effector TIL and Tumor Shrinkage (NSG-B2m Model)

[0141] The NSG-B2M mouse model is a severely immunocompromised animal, combining the lesions non-obese diabetic (NOD), severe combined immunodeficiency (SCID), IL-2 receptor γ null, and β 2-microglobulin null. Human tumor cells readily engraft in this model and cells from the human immune system can also be engrafted in order to study various modes of cancer immunotherapy. Thus, one can model a human immune response to a human tumor in this system. Female NSG-B2M mice were injected in the mammary fat pad with 5×10^6 JIMT-1 cells (human breast carcinoma cell line) suspended in 0.1 mL of phosphate-buffered saline (PBS) with an equal volume of Matrigel®. Once tumors reached a volume of 100-200 mm 3 , the mice were implanted with 1×10^7 human PBMC by subcutaneous injection into the right flank (pre-randomization). PBMC were obtained from two donors and distributed evenly between groups. When tumors reached an average tumor volume of 200-250 mm 3 (about 3 days post-PBMC injection) animals were matched by tumor volume into treatment and control groups, and dosing initiated. This first day of dosing was defined as Day 0. The mice were administered a daily oral dose of 10 mL/kg of 10% dimethyl sulfoxide (DMSO) in PBS as vehicle control or 10 mg/kg of GA2E dissolved in 10% DMSO in PBS as the treatment.

[0142] Tumor volume was measured twice weekly and on the day a study endpoint was reached. The mice were also weighed twice weekly and on the day a study endpoint was reached. Animals exhibiting a weight loss from Day 0 of >10% were provided a food supplement *ad libitum*. Animals exhibiting a weight loss of >20% in any 7-day period or >30% from Day 0 were considered moribund and euthanized.

[0143] Study endpoint was defined as when the mean tumor volume of the control group (uncensored) reaches 1500 mm 3 . If this occurs before Day 28, treatment groups and individual mice were permitted to be dosed and measured up to Day 28. If the mean tumor volume of the control group (uncensored) did not reach 1500 mm 3 by Day 28, then the endpoint for all

animals was the day when the mean tumor volume of the control group (uncensored) reached 1500mm³ up to a maximum of Day 60.

[0144] The GA2E-treated mice lost substantial weight, >10% by Day 12 of the study (Figure 10A-B) and the animals were placed on drug holiday on Day 13 through the end of the study. No impact on tumor growth was observed while the mice were receiving GA2E, however, there was tumor shrinkage subsequent to Day 19 of the study (Figure 11); and several of the mice showed shrinkage subsequent to day 17 (data not shown). These data suggest that the anti-tumor effects of RAR γ agonists is not a direct effect on the tumor cells themselves, but are consistent with RAR γ agonists promoting anti-tumor immunity.

[0145] The mice were sacrificed on Day 26 of the study and the tumors collected and analyzed by flow cytometry. Statistical analysis was by ordinary one-way ANOVA with Tukey's test for significance comparison between groups. Gates were set to identify single cells by forward light scatter, live cells as single cells based on side scatter and staining with the vital stain Fixable Viability Dye eFluor™ 780, white blood cells from live cells based on side scatter and staining with anti-CD45, lymphocytes from white blood cells based on side and forward light scatter, and total T lymphocytes (tot T) from lymphocytes based on side scatter and staining with anti-CD3. Total T lymphocytes were divided into CD4⁺ (TH) and CD8⁺ (TC) lymphocytes based on staining with anti-CD4 and CD8 antibodies and subsets assessed. The following observations were made on a per-mg of tumor tissue basis.

[0146] The concentration of total CD8⁺ lymphocytes was increased in the GA2-treated tumors as compared to controls (Figure 12A, P<0.001). There was no significant change in the concentration of naïve (defined as CCR7⁺ CD45AR⁺), T central memory (CCR7⁺ CD45AR⁻), or T effector memory (CCR7⁻ CD45AR⁻) subsets of CD8⁺ cells in the tumors (Figure 12B, C, and D, respectively). However, there was a substantial increase the concentration of terminally differentiated CD8⁺ T effector cells, defined as CCR7⁻ CD45AR⁺ (Figure 12E, P<0.0001). CCR7 promotes homing to the lymph nodes, thus effector cells, found in the periphery are CCR7⁻. That terminally differentiated CD8⁺ T effector cells, expected to be cytolytic, are increased amongst TIL in shrinking tumors, is consistent with the interpretation that anti-cancer activity of RAR γ agonists, such as GA2E, arises through an immunologic mechanism rather than a direct effect. The concentration of CD8⁺ TIL expressing the T cell markers PD-1, CD18 β , and CD54 was also increased in the GA2E-treated tumors as compared to controls (Figure 13A, P<0.0001; Figure 13B, P<0.001; and Figure 13C, P<0.01; respectively). CD18 and CD54 are both involved in cell-cell adhesion and immune surveillance. Their increased presence in TIL suggests improved efficiency in forming the immunologic synapse and antigen recognition by the CD8⁺ T effector cells. PD-1 is involved in immune homeostasis, facilitating down-regulation of immune responses, for example, through apoptosis of the effector cells

expressing PD-1. The increase in PD-1⁺ cells reflects an upregulated response. Antagonists of the PD-1/PD-L axis, such as anti-PD-1 and anti-PD-L1 antibodies (immune checkpoint inhibitors) may thus further augment the immune response promoted by RAR γ agonists, such as GA2E.

[0147] The concentration of total CD4⁺ lymphocytes was decreased (Figure 14A; $P < 0.05$), as were all of the tested subsets: naïve (defined as CCR7⁺ CD45AR⁺), T central memory (CCR7⁺ CD45AR⁻), T effector memory (CCR7⁻ CD45AR⁻) terminally differentiated CD4⁺ T effector cells (CCR7⁻ CD45AR⁺) (Figure 14B, C, and D, $P < 0.05$ each, and Figure 14D, $P > 0.05$, respectively). The concentration of CD4⁺ TIL expressing the T cell markers PD-1, CD18 β , CD54, and FoxP3 was also increased in the GA2-treated tumors as compared to controls (Figure 15A, $P < 0.05$; Figure 15B, $P < 0.05$; Figure 15C, $P < 0.05$; and Figure 15D, $P > 0.05$; respectively).

[0148] The effect of treatment with the RAR γ agonist GA2E, in this model of human anti-tumor immune response, was to substantially increase the concentration of CD8⁺ TIL in the tumor, especially of terminally differentiated effector cells. In contrast, the concentration of CD4⁺ TIL in the tumor was decreased. Similarly, whereas the concentration PD-1⁺, CD8 β ⁺ and CD54⁺ CD8⁺ TIL was increased, it was decreased CD4⁺ TIL. All of these data are consistent with promotion of a CD8⁺ effector cell-based TIL antitumor response accompanied by a diminished CD4⁺ response. Applicants are not aware of another class of small molecule agents that specifically increases CD8⁺ effector cells in the TIL population. The desirability of such an effect for cancer treatment is manifest.

[0149] The tumor cells were also assessed by flow cytometry for expression of PD-L1, CD18 β , CD58 (LFA-3), and Her2. Treatment with GA2E did not significantly alter the proportion of cells expressing these markers, consistent with RAR γ agonists not having a direct effect on the tumor cells.

Example 8

Effect of RAR γ Agonists on *In Vitro* Culture of Triple Negative Breast Cancer Cell Line

[0150] EMT6 cells were cultured *in vitro* for 5 days the presence of a multiple concentrations of the RAR γ agonists GA1E, GA2E, and GA3Ep and viability assessed by the Cell Titer Glo assay® (Promega). The compounds had no effect at concentrations below 1 μ M, but rapidly became toxic as concentrations exceeded 10 μ M (Figure 16). The IC₅₀ for GA1E, GA2E, and GA3Ep were 52 μ M, 19 μ M, and 101 μ M for GA1E, GA2E, and GA3Ep, respectively. These data confirm that the anti-cancer effects of the RAR γ agonists observed *in vivo* arise not from a direct effect on the tumor cells but, most likely, from promotion of a TIL-based antitumor immune response. That is, at the concentrations used, the RAR γ agonists did not directly

inhibit tumor growth, and therefore their tumor inhibitory effect arises from their promotion of TIL.

Example 9

Effect of RAR γ Agonists on *In Vitro* Culture of Lewis Lung Cancer Cell Line

[0151] Lewis lung carcinoma (LLC) cells were cultured *in vitro* the presence of a multiple concentrations of the RAR γ agonists GA1E, GA2E, and GA3Ep and viability assessed by the Cell Titer Glo assay®. The compounds had no effect at concentrations below 1 μ M, but rapidly became toxic as concentrations exceeded 10 μ M (Figure 17). The IC₅₀ for GA1E, GA2E, and GA3Ep were 96 μ M, 46 μ M, and 45 μ M for GA1E, GA2E, and GA3Ep, respectively. These data confirm that the anti-cancer effects of the RAR γ agonists observed *in vivo* arise not from a direct effect on the tumor cells but, most likely, from promotion of a TIL-based antitumor immune response. That is, at the concentrations used, the RAR γ agonists did not directly inhibit tumor growth, and therefore their tumor inhibitory effect arises from their promotion of TIL.

Example 10

RAR γ Selective Agonist Promotes Generation of Human CD8⁺ Effector TIL and Tumor Shrinkage (LLC Model)

[0152] The LLC model is based on an epidermoid carcinoma of the lung that spontaneously arose in a C57BL/6 mouse some 70 years ago, Syngeneic models have been useful in predicting clinical benefit. C57BL/6 mice, 6-8 weeks old, were injected subcutaneously in the left flank with 0.1 mL containing 3x10⁵ LLC cells. When tumors reached an average tumor volume of 50-150 mm³, the mice were matched by tumor size and sorted into control and treatment groups, and dosing initiated (Day 0). Three groups of 10 mice each were established: vehicle control, 10 mg/kg GA2E, and 25 mg/kg GA2E. The mice received one oral dose per day for 13 days. Tumor volumes and weight were measured on days 0, 2, 4, 6, 9, 11, and 13. This study was ended on Day 13, as the average tumor volume was >1500 mm³, and the tumors analyzed by flow cytometry. Two animals in the 10 mg/kg group and one animal in the 25 mg/kg group were terminated early due to ulceration of the tumor.

[0153] The 10 mg/kg dosage of GA2E was well tolerated, with minimal weight loss over the course of the study. The 25 mg/kg dosage of GA2E was less well tolerated, being associated with steady weight loss over the course of the study, exceeding 10% by about Day 10 (Figure 18A-B). Both dosages of drug inhibited tumor growth, though there was not a significant difference in the tumor inhibition of effect of the two dosages (Figure 19).

[0154] The mice were sacrificed on Day 13 of the study and the tumors collected and analyzed by flow cytometry. Statistical analysis was by ordinary one-way ANOVA with Tukey's

test for significance comparison between groups. Gates were set to identify single cells by forward light scatter, live cells as single cells based on side scatter and staining with the vital stain Fixable Viability Dye eFluor™ 780, white blood cells from live cells based on side scatter and staining with anti-CD45, lymphocytes from white blood cells based on side and forward light scatter, and total T lymphocytes (tot T) from lymphocytes based on side scatter and staining with anti-CD3. Total T lymphocytes were divided into CD4⁺ (TH) and CD8⁺ (TC) lymphocytes based on staining with anti-CD4 and CD8 antibodies and subsets assessed (Figure 20). The following observations were made on a per-mg of tumor tissue basis.

[0155] The concentration of total CD8⁺ lymphocytes was increased in the GA2E-treated tumors as compared to controls (Figure 22, panel CD8⁺; $P < 0.05$ for the 25 mg/kg dosage). The concentration of cytotoxic cells was also increased (Figure 22, panel Cytotoxic CD8; $P < 0.05$ for the 25 mg/kg dosage) as was the concentration of effector memory CD8⁺ T cells in the tumors (Figure 22, panel CD8 Tem; $P < 0.01$ for the 25 mg/kg dosage). The concentration of naïve and central memory CD8⁺ T cells in the tumors did not change significantly with the treatment (Figure 22, panels CD8 naïve and CD8 Tcm, respectively). The concentration of total CD4⁺ T cells, T helper cells and T regulatory cells in the tumor did not change significantly with the treatment (Figure 21). However, as a percentage of total T cell the proportion of Treg cells was significantly reduced (Figure 21, panel Tregs; $P < 0.001$). Accordingly, upon treatment with GA2E, the number of TIL present in the tumors, with phenotypes that should be more effective in killing tumor cells, was expanded.

[0156] The Examples demonstrate that RAR γ agonists inhibit tumor growth in multiple cancer models including breast cancer (Her2⁺ and triple negative) and lung cancer cell lines and promote TIL with mouse and human lymphocytes. As the RAR γ agonists did not appear to have a tumor growth inhibitory effect *in vitro*, it is apparent that the tumor growth inhibitory effect observed *in vivo* is due to increased production and promotion anti-tumor activity of TIL.

Example 11

Pharmacological activation of RAR γ signaling using RAR γ agonists has a cooperative effect with immune checkpoint inhibitor antibody in rejection of B 16 melanoma cells

[0157] The anti-tumor effects of an immune checkpoint inhibitor antibody treatment (such as anti-CTLA-4 antibody) combined with 10 nM of a RAR γ agonist disclosed herein are examined in C57BL/6 mice engrafted with B16F10 tumor cells. Mice treated with vehicles only do not show a survival advantage over untreated control mice. Mice treated with both the immune checkpoint inhibitor antibody and RAR γ agonist have improved survival at 50 days indicating that these two agents cooperate to eliminate the B16 melanoma cells. Surviving mice that undergo combination treatment are resistant to re-challenge with live tumor cells

indicating the effective formation of B16-specific memory cells. Importantly, the anti-melanoma effect is obtained with this combination of drugs without signs of acute or delayed toxicity.

[0158] In closing, it is to be understood that although aspects of the present specification are highlighted by referring to specific embodiments, one skilled in the art will readily appreciate that these disclosed embodiments are only illustrative of the principles of the subject matter disclosed herein. Therefore, it should be understood that the disclosed subject matter is in no way limited to a particular methodology, protocol, and/or reagent, etc., described herein. As such, various modifications or changes to or alternative configurations of the disclosed subject matter can be made in accordance with the teachings herein without departing from the spirit of the present specification. Lastly, the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which is defined solely by the claims. Accordingly, the present invention is not limited to that precisely as shown and described.

[0159] Certain embodiments of the present invention are described herein, including the best mode known to the inventors for carrying out the invention. Of course, variations on these described embodiments will become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventor expects skilled artisans to employ such variations as appropriate, and the inventors intend for the present invention to be practiced otherwise than specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described embodiments in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

[0160] Groupings of alternative embodiments, elements, or steps of the present invention are not to be construed as limitations. Each group member may be referred to and claimed individually or in any combination with other group members disclosed herein. It is anticipated that one or more members of a group may be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

[0161] Unless otherwise indicated, all numbers expressing a characteristic, item, quantity, parameter, property, term, and so forth used in the present specification and claims are to be understood as being modified in all instances by the term "about." As used herein, the term "about" means that the characteristic, item, quantity, parameter, property, or term so qualified encompasses a range of plus or minus ten percent above and below the value of the stated

characteristic, item, quantity, parameter, property, or term. Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical indication should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and values setting forth the broad scope of the invention are approximations, the numerical ranges and values set forth in the specific examples are reported as precisely as possible. Any numerical range or value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Recitation of numerical ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate numerical value falling within the range. Unless otherwise indicated herein, each individual value of a numerical range is incorporated into the present specification as if it were individually recited herein.

[0162] The terms “a,” “an,” “the” and similar referents used in the context of describing the present invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein is intended merely to better illuminate the present invention and does not pose a limitation on the scope of the invention otherwise claimed. No language in the present specification should be construed as indicating any non-claimed element essential to the practice of the invention.


[0163] Specific embodiments disclosed herein may be further limited in the claims using consisting of or consisting essentially of language. When used in the claims, whether as filed or added per amendment, the transition term “consisting of” excludes any element, step, or ingredient not specified in the claims. The transition term “consisting essentially of” limits the scope of a claim to the specified materials or steps and those that do not materially affect the basic and novel characteristic(s). Embodiments of the present invention so claimed are inherently or expressly described and enabled herein.

[0164] All patents, patent publications, and other publications referenced and identified in the present specification are individually and expressly incorporated herein by reference in their entirety for the purpose of describing and disclosing, for example, the compositions and methodologies described in such publications that might be used in connection with the present invention. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission

that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.


CLAIMS

1. A method of treating cancer comprising administering to a patient in need thereof an effective amount RAR γ -selective agonist having the structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; R⁷ to R¹² are independently, C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom, a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃; Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl; and R¹⁶ is H or F.

2. The method of claim 1, wherein the RAR γ -selective agonist has the structure




3. The method of claim 1 or 2, wherein the cancer is a solid tumor.
4. The method of claim 3, wherein the cancer is breast cancer.
5. The method of claim 4, wherein the breast cancer is triple negative breast cancer.
6. The method of claim 4, wherein the breast cancer is Her2⁺ breast cancer.
7. The method of claim 3, wherein the cancer is lung cancer.
8. The method of claim any one of claims 1-7, wherein the effective amount is about 1 to about 100 mg/day.
9. The method of any one of claims 1-8, further comprising administration of a CAR-T cell.
10. The method of any one of claims 1-8, further comprising administration of an immune checkpoint inhibitor.

11. A method of generating or expanding tumor infiltrating lymphocytes (TIL) by contacting the TIL with a RARy selective agonist.
12. The method of claim 11, wherein contacting comprises administering the RARy selective agonist to a subject having cancer.
13. The method of claim 11, wherein contacting comprises culturing the TIL *in vitro* in a media supplemented with the RARy selective agonist.
14. The method of claim 13, comprising isolating TIL from a tumor explant.
15. The method of claim 13, comprising culturing peripheral blood mononuclear cells (PBMC) *in vitro*, in the presence of irradiated tumor cells, in a media supplemented with the RARy selective agonist.
16. The method of claim 15, wherein the PBMC and the irradiated tumor cells are obtained from the same individual.
17. The method of any one of claims 13-16, wherein the media is further supplemented with IL-2.
18. The method of any one of claims 13-17, further comprising supplementing the culture media with an anti-PD-1 or anti-PD-L1 antibody.
19. A method of treating cancer, comprising infusing the TIL generated or expanded according to any one of claims 11-18 to a patient in need thereof.
20. A method of treating cancer, comprising providing RARy selective agonist-expanded TIL to a patient in need thereof.
21. The method of claim 20, wherein providing comprises administering TIL cultured *in vitro* in a media supplemented with the RARy selective agonist.
22. The method of claim 20 or 21, wherein providing comprises administering the RARy selective agonist to the patient.
23. The method of any one of claims 20-22, wherein the cancer is a solid tumor.
24. The method of claim 23 wherein the cancer is breast cancer.
25. The method of claim 24, wherein the breast cancer is triple negative breast cancer.
26. The method of claim 24, wherein the breast cancer is Her2⁺ breast cancer.
27. The method of claim 24, wherein the cancer is lung cancer.
28. The method of claim any one of claims 11-27, wherein the RARy selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; each R¹ and R² are independently H or C₁₋₆ alkyl; R³, R^{3'}, R⁴, and R^{4'} are independently, H or F; X is O, S, CH₂, C(R⁵)₂, or NR⁶, wherein each R⁵ and R⁶ are independently H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified.

29. The method of claim 28, wherein the RARy selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof.

30. The method of claim 28, wherein the RARy selective agonist is a compound of structure




or a pharmaceutically acceptable salt thereof.

31. The method of claim any one of claims 19-30, wherein the RARy selective agonist is a compound of structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; R⁷ to R¹² are independently, C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom, a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃; Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl; and R¹⁶ is H or F.

32. The method of claim 31, wherein the RAR_Y-selective agonist has the structure



33. The method of any one of claims 1-10, or 19-32, comprising administering the RAR_Y selective agonist to a subject, wherein the administering occurs periodically throughout an interval of treatment.

34. The method of any one of claims 1-10, or 19-32, comprising administering the RAR_Y selective agonist to a subject, wherein the administering occurs in repeated cycles throughout an interval of treatment.

35. The method of claim 34, wherein one cycle of treatment comprises 1) administering the RAR_Y selective agonist periodically over a first span of time and 2) suspending administration of the RAR_Y selective agonist over a second span of time, after which a new cycle may be initiated.

36. The method of claim 35, wherein the first span of time is 10-15 days or any integer number of days therein.

37. The method of claim 35 or 36, wherein the second span of time is two weeks to one month or any integer number of days therein.


38. The method of any one of claims 33-37, wherein periodically is daily.

39. The method of any one of claims 33-37, wherein periodically is twice daily.

40. The method of any one of claims 33-37, wherein periodically is every other day.


41. The method of any one of claims 33-40, wherein the interval of treatment extends from a first administration until a complete response is achieved.

42. The method of any one of claims 33-40, wherein the interval of treatment extends from a first administration until the cancer again progresses after stable disease or regression.
43. The method of any one of claims 1-10, or 19-42, further comprising administering an inhibitor of Treg cells.
44. The method of claim 43, wherein the inhibitor of Treg cells comprises a RAR α antagonist.
45. The method of claim 43, wherein the inhibitor of Treg cells comprises a Treg-depleting antibody.
46. The method of claim 45, wherein the Treg-depleting antibody comprises an anti-CD25 antibody, an anti-GITR antibody, an anti-FoxP3 antibody, an anti-CCR₄ antibody, or an anti-folate receptor 4 antibody.
47. The method of any one of claims 1-8, or 19-46 further comprising administration of an anti-PD-1 or anti-PD-L1 antibody.
48. The method of any one of claims 1-10, or 19-46, further comprising administering an RXR agonist having the structure



where R is H or C₁₋₆ alkyl, or a pharmaceutically acceptable salt thereof.


49. The method of claim 48, wherein the RXR agonist is 3,7-dimethyl-6(S),7(S)-methano,7-[1,1,4,4-tetramethyl-1,2,3,4-tetrahydronaphth-7-yl]2(E), 4(E) heptadienoic acid (IRX4204).
50. A method of potentiating chimeric antigen receptor-T (CAR-T) cancer immunotherapy comprising administering an effective amount RAR γ -selective agonist having the structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; R⁷ to R¹² are independently, C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom, a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃; Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl; and R¹⁶ is H or F,

to a cancer patient who is receiving, has received, or is scheduled to receive, CAR-T cells.


51. A method of potentiating immune checkpoint inhibitor cancer immunotherapy comprising administering an effective amount RAR γ -selective agonist having the structure



or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; R⁷ to R¹² are independently, C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom, a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃; Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl; and R¹⁶ is H or F,

to a cancer patient who is receiving, has received, or is scheduled to receive, an immune checkpoint inhibitor.

52. A RAR γ -selective agonist of the structure:




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; R⁷ to R¹² are independently: C₁₋₆ alkyl, a hydrogen atom, an alkoxy group, a halogen atom, a nitro group, a hydroxy group, OCF₃, or COR¹³, where R¹³ is C₁₋₆ alkyl or CF₃. Y is oxygen, sulfur, or NR¹⁴, where R¹⁴ is C₁₋₆ alkyl.

53. The RAR γ -selective agonist of claim 52, wherein R is H, methyl, or ethyl.

54. The RAR γ -selective agonist of claim 52, wherein Y is O.


55. The RAR γ -selective agonist of claim 54 having the following structure:



(GAG1-1).


56. The RAR γ -selective agonist of claim 52, wherein the =N-OH group is in the E configuration.

57. The RAR γ -selective agonist of claim 56 having the following structure:




(GAG1-2).

58. The RAR γ -selective agonist of claim 52, wherein the RAR γ selective agonist is 4-((1*E*,3*E*)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1E), having the structure




(GA1E).

59. The RAR γ -selective agonist of claim 52, wherein the RAR γ selective agonist is 4-((1*E*,3*Z*)-3-(3-(*tert*-butyl)-2-ethyl-benzofuran-5-yl)-3-(hydroxyimino)prop-1-en-1-yl)-3-fluorobenzoic acid (GA1Z), having the structure



(GA1Z).

60. A RARy-selective agonist having the structure




or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl, and the crossed double bond to the =N-OH group indicates that stereochemistry is not specified.


61. The RARy-selective agonist of claim 60, wherein R is H, methyl, or ethyl.

62. The RARy-selective agonist of claim 60, wherein the =N-OH group is in the E configuration.


63. The RARy-selective agonist of claim 60, wherein the RARy selective agonist is 3-fluoro-4-((1E,3E)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2E), having the structure



64. The RARy-selective agonist of claim 60, wherein the RARy selective agonist is 3-fluoro-4-((1E,3Z)-3-(hydroxyimino)-3-(5,5,8,8-tetramethyl-5,6,7,8-tetrahydronaphthalen-2-yl)prop-1-en-1-yl)benzoic acid (GA2Z), having the structure






65. A RARy-selective agonist having the structure





or a pharmaceutically acceptable salt thereof, wherein R is H or C₁₋₆ alkyl; each R¹ and R² are independently H or C₁₋₆ alkyl; R³ and R^{3'} are independently H, or halogen; (R⁴)₂ comprises R⁴ and R^{4'} which are independently H, halogen, C₁₋₆ alkyl or C₁₋₆ alkoxy; X is O, S, CH₂, C(R⁵)₂, or NR⁶, wherein each R⁵ and R⁶ are independently H or C₁₋₆ alkyl; the crossed double bond to the =N-OH group indicates that stereochemistry is not specified; and the COOR group is in the meta or para position and the two R⁴ groups occupy the remaining positions on the ring.

66. The RARy-selective agonist of claim 65, having the structure



67. The RARy-selective agonist of claim 65, having the structure





GA3Em.