


INDOZOLONE DERIVATIVES AS 11B-HSD1 INHIBITORS

The present invention is concerned with novel indazolone derivatives which are useful as 11b-HSD1 inhibitors.

The invention is concerned particularly with compounds of formula (I)



5 wherein

R^1 is hydrogen, lower-alkyl, aryl, or aryl-lower-alkyl;

R^2 is aryl, aryl-lower-alkyl, heteroaryl, heteroaryl-lower-alkyl, cycloalkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN,
10 halogen, lower-alkoxy and $C(O)NR^8R^9$;

R^3 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;

R^4 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;

15 R^5 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;

R^6 is hydrogen or lower-alkyl;

R^7 is aryl, heteroaryl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN,
20 halogen, lower-alkoxy and cycloalkyl;

R^8 and R^9 , independently from each other, are selected from the group consisting of hydrogen and lower-alkyl;

CS / 15.7.05

and pharmaceutically acceptable salts thereof.

Glucocorticoids (cortisol in humans, corticosterone in mice and rats) are an important class of adrenocorticosteroids that regulate many metabolic and homeostatic processes and form a key component of the response to stress. Glucocorticoids act via
5 intracellular glucocorticoid receptors and, in some tissues, mineralocorticoid receptors; both being nuclear transcription factors. Glucocorticoid action on target tissues depends not only on circulating steroid concentrations and the cellular expression of receptors, but also on intracellular enzymes that critically determine to which extent glucocorticoids gain access to receptors in an active form. 11beta-hydroxysteroid dehydrogenases (11beta-
10 HSD's) catalyze the interconversion of the principal active 11-hydroxy-glucocorticoid (Cortisol in men) and their inactive 11-keto metabolites (cortisone in men).

The enzyme 11beta-hydroxysteroid dehydrogenase type 1 (11beta-HSD1) inter-converts inactive into active glucocorticoids, thereby playing a major role in local modulation of cellular agonist concentration and thus activation of corticosteroid
15 receptors in target tissues. In a recent study made by F. Hoffmann-La Roche differences in gene expression in lean and obese men were analyzed using gene array technology in order to identify specific changes in gene expression that might be associated with insulin resistance or altered metabolism. This study revealed that the mRNA for 11beta-HSD1 is approximately two-fold up regulated in adipose tissue in obese individuals. Moreover,
20 overexpressing 11beta-HSD1 in adipocytes of mice led to visceral obesity and to a syndrome-X like phenotype (Masuzaki H. et al., Science. 2001 Dec 7; 294(5549):2166-70.). Taken together, these data very strongly support an important role of 11beta-HSD1 in the induction of obesity and the impairment of glucose homeostasis and lipid parameters. Thus, selective inhibition of this enzyme could lower blood glucose levels in Type 2
25 diabetic patients, normalize elevated lipid parameters and/or reduce weight in obese subjects.

The first pharmacological indication that 11beta-HSD1 inhibition in men might have beneficial effects were obtained by using carbenoxolone, an anti-ulcer drug which inhibits both 11beta-HSD1 and the related enzyme 11beta-HSD2. Treatment with
30 carbenoxolone led to an increase in insulin sensitivity indicating that that inhibition of 11beta-HSD1 may reduce cellular cortisol levels and therefore minimizing some of its deleterious effects. (Walker et al. 1995; J. Clin. Endocrinol. Metab. 80, 31155-3159).

11beta-HSD1 is expressed in many tissues including liver, adipose tissue, vascular smooth muscles, pancreas and brain. Its activity is dependent on NADP(H) and it has a

relatively low affinity for its substrate (compared to 11beta-HSD2). 11 beta-HSD1 in tissue homogenates and when purified is bidirectional, exhibiting both 11beta-dehydrogenase and 11beta-reductase reactions, with greater stability of the dehydrogenase activity (P.M. Stewart and Z.S. Krozowski, Vitam. Horm. 57 (1999), pp. 249–324). However, when the
5 enzyme activity is tested in intact cells, the 11beta-reductase activity predominates, which regenerates active glucocorticoids from inert 11-keto forms. Such glucocorticoid regeneration will increase effective intracellular glucocorticoid levels and thereby amplifying glucocorticoid activity. It is this elevated cellular cortisol concentration that might lead to increased hepatic glucose production, adipocyte differentiation and insulin
10 resistance.

Inhibition of 11beta-HSD1 should not only reduce the typical Syndrome-X / Diabetes associated symptoms, but it should also be save and without major side effect. Studies with the unspecific inhibitor carbenoxolone highlight the importance of developing specific 11beta-HSD1 inhibitors. The inhibition of the 11beta-HSD2 enzyme is
15 badly tolerated and results in increased blood pressure. In contrast inhibition of 11beta-HSD1 should be well tolerated since 11beta-HSD1 knockout mice were found be healthy and to resist hyperglycemia provoked by obesity or stress (Kotelevtsev Y. et al., Proc Natl Acad Sci U S A. 1997 Dec 23;94(26):14924-9). Similar upon starvation these mice had attenuated activation of key hepatic enzymes that are involved in gluconeogenesis. In
20 addition, these mice had improved lipid and lipoprotein profiles suggesting that inhibition of HSD1 might be highly efficacious and safe. Recent reports indicate that 11beta-HSD1 inhibitors might also be beneficial to reduce high blood pressure (Masuzaki H. et al., J Clin Invest. 2003 July;112(1):83-90; Rauz S. et al., QJM. 2003 July;96(7):481-90) to improve cognition (Sandeep TC. et al., Proc Natl Acad Sci U S A. 2004 Apr. 27;101(17):6734-9) or
25 to improve Alzheimer associated deficits. Taken together 11beta-HSD1 inhibition might be a save and efficacious approach to treat symptoms of diabetes, obesity and other diseases.

The compounds of formula I and their pharmaceutically acceptable salts and esters are novel and have valuable pharmacological properties. In particular they are 11b-HSD1
30 inhibitors and they display selectivity against the related 11beta-HSD2 enzyme. Therefore the compounds which are specific 11beta-HSD1 inhibitors represent an approach to e.g. lower blood glucose levels and normalize lipid parameters in Type 2 diabetic patients by modulating the local concentration of the active glucocorticoid cortisol in target tissue (liver, adipose tissue).

35 The compounds of the present invention can be used in the prophylaxis and/or treatment of metabolic disorders, obesity, dyslipidemias, hypertension and/or diabetes,

- 4 -

particularly diabetes Type II. The compounds of this invention can further be used in the prophylaxis and/or treatment of high ocular eye pressure, cognition, Alzheimer and/or neurodegeneration.

Objects of the present invention are the compounds of formula I and their
5 aforementioned salts per se and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts, the use of the said compounds and salts for the prophylaxis and/or therapy of illnesses, especially in the treatment or prophylaxis of metabolic disorders, obesity, dyslipidemiae,
10 hypertension and/or diabetes, particularly diabetes Type II, and the use of the said compounds and salts for the production of medicaments for the treatment or prophylaxis of metabolic disorders, obesity, dyslipidemiae, hypertension and/or diabetes, particularly diabetes Type II.

The compounds of the present invention can further be combined with PPAR
15 (alpha, gamma, delta) agonists, DHEA (dehydroepiandrosterone), DPPIV inhibitors, insulin and/or lipase inhibitors, particularly orlistat.

Unless otherwise indicated, the following definitions are set forth to illustrate and define the meaning and scope of the various terms used to describe the invention herein.

In this specification the term "lower" is used to mean a group consisting of one to seven, preferably of one to four carbon atom(s).

5 The term "halogen" refers to fluorine, chlorine, bromine and iodine, with fluorine, chlorine and bromine being preferred.

 The term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon
10 atoms. Lower-alkyl groups as described below also are preferred alkyl groups.

 The term "lower-alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent alkyl radical of one to seven carbon atoms, preferably one to four carbon atoms. This term is further exemplified by such radicals as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl and the like. Lower-alkyl
15 groups can optionally be substituted, e.g. by OH, CN, halogen, lower-alkoxy, or aminocarbonyl. Unsubstituted lower-alkyl groups are preferred.

 The term "fluoro-lower-alkyl" refers to lower-alkyl groups which are mono- or multiply substituted with fluorine. Examples of fluoro-lower-alkyl groups are e.g. CFH₂, CF₂H, CF₃, CF₃CH₂, CF₃(CH₂)₂, (CF₃)₂CH and CF₂H-CF₂.

20 The term "cycloalkyl" refers to a monovalent carbocyclic radical of 3 to 10 carbon atoms, preferably 3 to 6 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

 The term "alkoxy" refers to the group R'-O-, wherein R' is an alkyl. The term "lower-alkoxy" refers to the group R'-O-, wherein R' is a lower-alkyl.

25 The term "fluoro-lower-alkoxy" refers to the group R"-O-, wherein R" is fluoro-lower-alkyl. Examples of fluoro-lower-alkoxy groups are e.g. CFH₂-O, CF₂H-O, CF₃-O, CF₃CH₂-O, CF₃(CH₂)₂-O, (CF₃)₂CH-O, and CF₂H-CF₂-O.

 The term "amino", alone or in combination, signifies a primary, secondary or tertiary amino group bonded via the nitrogen atom, with the secondary amino group
30 carrying an alkyl or cycloalkyl substituent and the tertiary amino group carrying two similar or different alkyl or cycloalkyl substituents or the two nitrogen substituents together forming a ring, such as, for example, -NH₂, methylamino, ethylamino, dimethylamino, diethylamino, methyl-ethylamino, pyrrolidin-1-yl or piperidino etc.,

preferably primary amino, dimethylamino and diethylamino and particularly dimethylamino.

The term "alkylene" refers to a straight chain or branched divalent saturated aliphatic hydrocarbon group of 1 to 20 carbon atoms, preferably 1 to 16 carbon atoms, more preferably up to 10 carbon atoms. Lower-alkylene groups as described below also are preferred alkylene groups. The term "lower-alkylene" refers to a straight chain or branched divalent saturated aliphatic hydrocarbon group of 1 to 7, preferably 1 to 6 or 3 to 6 carbon atoms. Straight chain alkylene or lower-alkylene groups are preferred.

The term "aryl", alone or in combination, relates to the phenyl or naphthyl group, preferably the phenyl group, which can optionally be substituted by 1 to 5, preferably 1 to 3, substituents independently selected from the group consisting of lower-alkyl, lower-alkoxy, halogen, hydroxy, CN, CF₃, amino, aminocarbonyl, carboxy, NO₂, dioxo-lower-alkylene (forming e.g. a benzodioxyl group), lower-alkylsulfonyl, aminosulfonyl, lower-alkylcarbonyl, lower-alkylcarbonyloxy, lower-alkylcarbonyl-NH, lower-alkoxycarbonyl, fluoro-lower-alkyl, fluoro-lower-alkoxy, cycloalkyl, phenyloxy and methyl-oxadiazolyl. Preferred substituents are halogen, lower-alkyl, fluoro-lower-alkyl and CN.


The term "heteroaryl" refers to an aromatic 5 to 6 membered monocyclic ring or 9 to 10 membered bicyclic ring which can comprise 1, 2 or 3 atoms selected from nitrogen, oxygen and/or sulphur, such as furyl, pyridinyl, pyridazinyl, oxo-pyridazinyl, pyrimidinyl, 2-oxo-pyridinyl, 2-oxo-pyrimidinyl, pyrazinyl, thienyl, isoxazolyl, oxazolyl, oxadiazolyl, imidazolyl, pyrrolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, isothiazolyl, 1,2,3-thiadiazolyl, benzoimidazolyl, indolyl, indazolyl. A preferred heteroaryl group is pyridinyl. A heteroaryl group may have a substitution pattern as described earlier in connection with the term "aryl".

Compounds of formula (I) can form pharmaceutically acceptable acid addition salts. Examples of such pharmaceutically acceptable salts are salts of compounds of formula (I) with physiologically compatible mineral acids, such as hydrochloric acid, sulphuric acid, sulphurous acid or phosphoric acid; or with organic acids, such as methanesulphonic acid, p-toluenesulphonic acid, acetic acid, lactic acid, trifluoroacetic acid, citric acid, fumaric acid, maleic acid, tartaric acid, succinic acid or salicylic acid. The term "pharmaceutically acceptable salts" refers to such salts. Compounds of formula (I) in which a COOH group is present can further form salts with bases. Examples of such salts are alkaline, earth-alkaline and ammonium salts such as e.g. Na-, K-, Ca- and Trimethylammoniumsalt. The term "pharmaceutically acceptable salts" also refers to such salts. Salts obtained by the addition of an acid are preferred.

The compounds of formula I can also be solvated, e.g. hydrated. The solvation can be effected in the course of the manufacturing process or can take place e.g. as a consequence of hygroscopic properties of an initially anhydrous compound of formula I (hydration). The term pharmaceutically acceptable salts also includes physiologically acceptable solvates.

5

In detail, the present invention relates to compounds of formula (I)



wherein

- R^1 is hydrogen, lower-alkyl, aryl, or aryl-lower-alkyl;
- 5 R^2 is aryl, aryl-lower-alkyl, heteroaryl, heteroaryl-lower-alkyl, cycloalkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$;
- 10 R^3 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^4 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^5 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- 15 R^6 is hydrogen or lower-alkyl;
- R^7 is aryl, heteroaryl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and cycloalkyl;
- 20 R^8 and R^9 , independently from each other, are selected from the group consisting of hydrogen and lower-alkyl;

and pharmaceutically acceptable salts thereof.

The compounds of formula I can have one or more asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, 25 diastereoisomeric racemates or mixtures of diastereoisomeric racemates. The invention embraces all of these forms. Preferred are the compounds of formula I.

Preferred compounds of formula (I) as described above are those wherein

- R^1 is hydrogen, lower-alkyl, aryl, or aryl-lower-alkyl;
- R^2 is aryl, aryl-lower-alkyl, heteroaryl, heteroaryl-lower-alkyl, cycloalkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$;
- R^3 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^4 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^5 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^6 is hydrogen or lower-alkyl;
- R^7 is aryl, heteroaryl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen and lower-alkoxy;
- R^8 and R^9 , independently from each other, are selected from the group consisting of hydrogen and lower-alkyl;

and pharmaceutically acceptable salts thereof.

- A preferred embodiment of the present invention relates to compounds of formula (I) as described above, wherein R^1 is hydrogen or lower-alkyl, preferably wherein R^1 is hydrogen.

- Another preferred object of the present invention are the compounds of formula (I), wherein R^2 is aryl-lower-alkyl, heteroaryl-lower-alkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$, wherein R^8 and R^9 are as defined above. Preferably, R^2 is aryl-lower-alkyl or heteroaryl-lower-alkyl, more preferably R^2 is benzyl or pyridinylmethyl, wherein benzyl can optionally be substituted with 1 or 2 halogen. Especially preferred are those compounds wherein R^2 is benzyl, 4-fluoro-benzyl, 4-chloro-benzyl, 3,4-difluoro-benzyl, or pyridin-2-ylmethyl. Other preferred compounds are those, wherein R^2 is aryl, particularly phenyl.

Also preferred are the compounds of formula (I), wherein R³ is hydrogen. Further preferred are those compounds according to formula (I), wherein R⁴ is hydrogen or halogen, particularly wherein R⁴ is hydrogen, fluorine, or chlorine.

Another preferred aspect of the present invention are compounds of formula (I),
5 wherein R⁵ is hydrogen or halogen, particularly wherein R⁵ is hydrogen. Other preferred compounds of formula (I) as defined above are those wherein R⁶ is hydrogen.

In a further preferred embodiment of the present invention, R⁷ is lower-alkyl or aryl. Particularly preferred are those compounds, wherein R⁷ is lower-alkyl, phenyl or naphthyl, wherein phenyl can optionally be substituted with 1 to 3 substituents independently
10 selected from the group consisting of lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, fluoro-lower-alkoxy, dioxo-lower-alkylene, halogen, cyano, phenoxy and 5-methyl-[1,3,4]-oxadiazol-2-yl. More preferably, R⁷ is phenyl substituted with 1 to 2 substituents selected from the group consisting of lower-alkyl, fluoro-lower-alkyl, halogen and cyano,
15 particularly 3-chloro-2-methyl-phenyl, 2,3-dichloro-phenyl, 3-chloro-4-fluoro-phenyl, 3-trifluoromethyl-4-fluoro-phenyl, 3-cyano-phenyl, 2,5-difluoro-phenyl, 3-chloro-2-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, or 2-chloro-phenyl. Other preferred compounds of formula (I) as described above are those, wherein R⁷ is fluoro-lower-alkyl or lower-alkyl substituted with cyclopropyl, especially those wherein R⁷ is cyclopropyl-methyl.

Examples of preferred compounds of formula (I) are those selected from the group
20 consisting of:
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-fluoro-benzenesulfonamide,
25 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-benzenesulfonamide,
Naphthalene-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-methyl-benzenesulfonamide,
30 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-6-methyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluoro-benzenesulfonamide,
35 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-propyl-benzenesulfonamide,

- 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-2,5-dimethyl-benzenesulfonamide,
- 5 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-3-trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-5-trifluoromethyl-
- 10 benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-phenoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-(5-methyl-[1,3,4]oxadiazol-2-yl)-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-benzenesulfonamide,
- 15 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-methoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3,5-bis-trifluoromethyl-benzenesulfonamide,
- 20 Propane-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-N-dimethyl-benzenesulfonamide,
N-(3-Oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide,
- 25 2,3-Dichloro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
3-Chloro-4-fluoro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
3-Chloro-2-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
- 30 3-Chloro-4-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
4-Chloro-2,5-dimethyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
N-(3-Oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
- 35 benzenesulfonamide,
N-(1-Isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide,

- 3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methyl-
benzenesulfonamide,
3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-
benzenesulfonamide,
- 5 3-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
3-Difluoromethoxy-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
4-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
4-Fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
- 10 3-Chloro-4-fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
2,4-Dichloro-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methyl-benzenesulfonamide,
3-Chloro-N-(1-ethyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-benzenesulfonamide,
3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-2-methyl-
- 15 benzenesulfonamide,
3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-
benzenesulfonamide,
(1-Cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
- 20 3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
2,4-Dichloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methyl-
benzenesulfonamide,
3-Chloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-
benzenesulfonamide,
- 25 4-Fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethyl-
benzenesulfonamide,
3-Chloro-4-fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
benzenesulfonamide,
2,4-Dichloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-
- 30 benzenesulfonamide,
3-Chloro-2-methyl-N-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide,
4-Fluoro-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-indazol-5-yl]-3-trifluoromethyl-
benzenesulfonamide,
- 35 4-Fluoro-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-3-
trifluoromethyl-benzenesulfonamide,
2,4-Dichloro-6-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-

- benzenesulfonamide,
3-Chloro-2-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
4-Fluoro-N-[1-(2-hydroxy-2-methyl-propyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-
5 trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-fluoro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
10 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichloro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-benzenesulfonamide,
15 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-3-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxy-
20 benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxy-
benzenesulfonamide,
25 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,5-difluoro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluoro-
30 benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-
benzenesulfonamide,
35 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-
benzenesulfonamide,
(1-Benzyl-4-chloro-3-oxo-2,3-dihydro-indazol-5-yl)-3-chloro-2-methyl-

- benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichloro-
5 benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-
benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluoro-
benzenesulfonamide,
10 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-
benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxy-
benzenesulfonamide,
3-Chloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
15 benzenesulfonamide,
3-Difluoromethoxy-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide,
4-Fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethyl-
benzenesulfonamide,
20 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-
1H-indazol-5-yl]-amide,
2,4-Dichloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methyl-
benzenesulfonamide,
5-Chloro-2,4-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
25 benzenesulfonamide,
4-Chloro-2,5-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide,
3-Chloro--[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-
benzenesulfonamide,
30 3-Cyano-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
Naphthalene-1-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
amide,
3-Chloro-4-fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide,
35 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-
benzenesulfonamide,
3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-

- benzenesulfonamide,
[1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
5 [1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethoxy-
benzenesulfonamide,
2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-5-methyl-
benzenesulfonamide,
3,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
10 benzenesulfonamide,
4,5-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-fluoro-
benzenesulfonamide,
2,4,5-Trichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
benzenesulfonamide,
15 2,3-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
benzenesulfonamide,
2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-
benzenesulfonamide,
3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-
20 benzenesulfonamide,
N-[1-(2,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,
3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-
benzenesulfonamide,
25 N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
3-Chloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-
benzenesulfonamide,
2,3-Dichloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
30 benzenesulfonamide,
Naphthalene-1-sulfonic acid [1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
amide,
3-Chloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-
methyl-benzenesulfonamide,
35 Naphthalene-1-sulfonic acid [1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-
indazol-5-yl]-amide,
2,3-Dichloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-

- benzenesulfonamide,
 N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide,
 N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-
 5 difluoromethoxy-benzenesulfonamide,
 N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethoxy-benzenesulfonamide,
 3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-benzenesulfonamide,
 10 N-[1-(2-Cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide,
 2,4-Dichloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methyl-benzenesulfonamide,
 3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-
 15 benzenesulfonamide,
 N-[6-Chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide,
 5-Chloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-fluoro-benzenesulfonamide,
 20 2,3-Dichloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
 3-Chloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-2-methyl-benzenesulfonamide,
 N-(6-Chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-
 25 trifluoromethyl-benzenesulfonamide,
 2-[5-(3-Chloro-2-methyl-benzenesulfonylamino)-3-oxo-2,3-dihydro-indazol-1-yl]-methyl-acetamide,
 2,4-Dichloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-benzenesulfonamide, and
 30 3-Chloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-benzenesulfonamide,
 and pharmaceutically acceptable salts thereof.

Examples of particularly preferred compounds of formula (I) are those selected from the group consisting of:

- 35 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-benzenesulfonamide,

- 17 -

- N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,5-difluorobenzenesulfonamide,
 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluorobenzenesulfonamide,
 5 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chlorobenzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluorobenzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichlorobenzenesulfonamide,
 10 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluorobenzenesulfonamide,
 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide,
 15 3-Cyano-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
 2,3-Dichloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
 3-Chloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methylbenzenesulfonamide,
 20 4-Fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethylbenzenesulfonamide,
 3-Chloro-4-fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methylbenzenesulfonamide,
 25 N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide, and
 3-Chloro-2-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
 30 and pharmaceutically acceptable salts thereof.

Other examples of preferred compounds of formula (I) are those selected from the group consisting of:

- 2,3-Dichloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
 35 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluorobenzenesulfonamide,

- 2,4-Difluoro-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
 3-Chloro-2-fluoro-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-
 benzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-benzenesulfonamide,
 5 Propane-2-sulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
 3-Cyano-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
 2,2,2-Trifluoro-ethanesulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-
 yl)-amide,
 Propane-2-sulfonic acid (1-benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
 10 Propane-2-sulfonic acid [6-chloro-1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-
 yl]-amide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-trifluoro-
 methanesulfonamide,
 Propane-2-sulfonic acid (1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
 15 Propane-2-sulfonic acid [1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
 amide,
 Propane-2-sulfonic acid [1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
 amide, and
 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-C-cyclopropyl-methanesulfonamide
 20 and pharmaceutically acceptable salts thereof.


Another example of a particularly preferred compound of formula (I) is:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-C-cyclopropyl-methanesulfonamide
 and pharmaceutically acceptable salts thereof.


- It will be appreciated that the compounds of the present invention may be
 25 derivatised at functional groups to provide derivatives which are capable of conversion
 back to the parent compound in vivo. Physiologically acceptable and metabolically labile
 compounds, which are capable of producing the parent compounds of general formula (I)
 in vivo are also within the scope of the present invention.

- A further aspect of the present invention is the process for the preparation of
 30 compounds as defined above comprising the reaction of a compound of formula (II)

- 19 -



with a compound of formula (III)



wherein R¹ to R⁷ are as defined above.

5 Such a process can be carried out under conditions known to the person skilled in the art, e.g. in the presence of a base such as triethylamine or (4-dimethylamino)-pyridine (DMAP) in a solvent such THF, ethanol, methylene chloride DMF or DMSO, or in pyridine as a solvent, with or without the addition of a base such as triethylamine or DMAP, at room temperature or at elevated temperatures.


10 The invention further relates to compounds of formula (I) as described above, when prepared by a process as defined above.

General description of the processes:

The preparation of compounds of formula I of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following schemes. The skills required for carrying out the reaction and purification of the resulting products are known to those persons skilled in the art. The substituents and indices used in the following description of the processes have the significance given above unless indicated to the contrary.

In general, compounds of type I are readily accessible by sulfonylation of appropriately substituted 5-amino-1,2-dihydro-indazol-3-one of formula IIa (R6=H) with sulfonyl chlorides, under various conditions that are known to persons skilled in the art. Examples of such conditions are – as indicated in Scheme 1 below – e.g. pyridine at elevated temperatures or THF under reflux conditions in the presence of a base such as potassium carbonate, sodium carbonate, sodium hydride, triethyl amine or the like. The sulfonyl chlorides are either commercially available or known in the literature or available in analogy to known procedures to those persons skilled in the art

Scheme 1:




Optionally, compounds of formula Ia with R6 = H obtained in this way can be further substituted at the sulfonamide nitrogen by treatment with a base such as sodium hydride, cesium carbonate, potassium carbonate or the like in a solvent such as DMF or THF or similar followed by alkylation of the resulting anion with an alkyl halide such as methyl iodide, ethyl bromide, benzyl bromide or the like in order to introduce the desired R6 substituent. Optionally, R6 can be introduced via alkylation of compound of formula IIa by procedures known to those persons skilled in the art.

Appropriately substituted 5-amino-1,2-dihydro-indazol-3-ones of formula IIa are either known in the literature or can be made in analogy to literature procedures from known starting materials according to scheme 2– e.g. by condensation of the appropriately functionalized 2-fluorobenzoic acid of formula III with hydrazine hydrate in ethanol and cyclisation under acidic conditions, followed by alkylation with R2-X to afford compounds of formula IVa (R6 = H), an optional alkylation with R1-X (under thermal or basic conditions in analogy to procedures described in *Heterocycles* 1997, 45, 129-36 or *J. Med.*

Chem. 1991, 34, 1492-1503) to afford compounds of formula IV, and hydrogenation of the nitro group. (Schindler et al., *Arch. Pharm. Pharm. Med. Chem.*, 1998, 331,13-21).


Alternatively, appropriately substituted 5-amino-1,2-dihydro-indazol-3-ones of formula II are prepared according to the general Scheme 3 via a one-operation condensation of the appropriately functionalized 2-fluorobenzoic acid of formula III with an appropriately substituted hydrazine R₁NHNH₂ of formula V in the presence of a coupling agent in DMF, followed by an optional alkylation with R₁-X (under thermal or basic conditions in analogy to procedures described in *Heterocycles* 1997, 45, 129-36 or *J. Med. Chem.* 1991, 34, 1492-1503) and a reduction of the nitro group. 5-Amino-1,2-dihydro-indazol-3-ones of formula II with R₂ = aryl, heteroaryl can also be prepared in analogy to Menon et al., *Combin. Chem. and High-Throughput Screen.* 2003, 6, 471-480. The appropriately substituted starting materials of formula III are either commercially available or are known in the literature or were prepared in analogy to literature procedures from known starting materials. The corresponding substituted hydrazines of formula V are either commercially available or are known in the literature or synthesised in analogy to literature procedures (such as *J. Org. Chem.* 1984, 49, 336-42; *J. Am. Chem. Soc.* 1986, 108, 7981-4 or *Bioorg. Med. Chem.* 2004, 12, 1357-1366).

Scheme 2:



20

Scheme 3:



As described above, the compounds of formula (I) of the present invention can be used as medicaments for the treatment and/or prevention of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1 (11bHSD1). Examples of such diseases are metabolic disorders, obesity, dyslipidemiae, hypertension
5 and/or diabetes, particularly diabetes type II. The compounds of this invention can further be used in the prophylaxis and/or treatment of high ocular eye pressure, cognition, Alzheimer and/or neurodegeneration.

The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

10 Further, the invention relates to compounds as defined above for use as therapeutic active substances, particularly as therapeutic active substances for the treatment and/or prevention of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1 (11bHSD1), particularly as therapeutic active substances for the treatment and/or prevention of metabolic disorders, obesity, dyslipidemiae,
15 hypertension and/or diabetes, particularly diabetes type II.

The invention further relates to a method for the treatment and/or prophylaxis of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1, particularly for the treatment and/or prophylaxis of metabolic disorders, obesity, dyslipidemiae, hypertension and/or diabetes, particularly diabetes type II, which
20 method comprises administering an effective amount of a compound as defined above.

In another embodiment, the present invention relates to the use of compounds as defined above for the therapeutic and/or prophylactic treatment of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1, particularly for therapeutic and/or prophylactic treatment of metabolic disorders, obesity,
25 dyslipidemiae, hypertension and/or diabetes, particularly type II diabetes.

The invention further relates to the use of compounds as defined above for the preparation of medicaments for the treatment and prophylaxis of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1, particularly for the treatment and prophylaxis of metabolic disorders, obesity,
30 dyslipidemiae, hypertension and/or diabetes, particularly type II diabetes.

Prevention and/or treatment of type II diabetes is the preferred indication.

Assay Procedures

Transient expression and partial Purification:

The cDNA encoding the human 11beta-HSD1 protein was cloned into the expression vector pcDNA3 (Stratagene). This construct (for details see Alex Odermatt et al.; J Biol Chem.,1999, Vol. 274, Issue 40, 28762-28770) was used to transiently express the protein in HEK293 cells (ATCC number: CRL-1573, described in Graham, F.L., Smiley, J., Russell, W.C., Nairn, R.; (1977)) using lipofectamine. 48h after transfection cells were washed twice with ice-cold PBS (Phosphate buffered Saline). To 1 volume of cell suspension in PBS 2 volumes of ice-cold lysis buffer (50mM Tris; pH7.5; 1mM EDTA; 100mM NaCl) were added. The cells were lysed by Potter-homogenization (20 strokes). The resulting homogenate was sonicated with a tip sonicator (10% output; 2 x 30 sec.) and cleared by a low speed centrifugation (10min x 9000g; 4°C). The microsomal fraction was collected by a high speed centrifugation (60 min x 110'000g). The resulting pellet was resuspended in storage buffer (20mM Tris pH 7.5; 1 mM EDTA; 10% Glycerol) and the centrifugation was repeated. The resulting pellet containing the microsomal fraction was again taken up into storage buffer and aliquots were kept frozen in liquid Nitrogen until use.

Generation of stable cell lines expressing 11beta-HSD1:

The same construct used for transient expression of human 11beta-HSD1 was also used to establish cell lines stably expressing the protein. Briefly, (HEK293) cells were transfected with 11beta-HSD1 construct using the lipofectamine reagent (Gibco BRL) according to the manufacturer's instruction. Two days after transfection, geneticin selection (0.8 mg/ml) was initiated and several stable clones were isolated. One clone was further used for pharmacological characterization.

Microsome Assay

Microsomes isolated from HEK293 cells transiently expressing human 11beta-HSD1 (for details see above) were incubated in assay buffer (100 mM NaCl; 1mM EDTA; 1mM EGTA; 1mM MgCl₂; 250 mM Sucrose; 20 mM Tris pH 7.4; Cortisone 50-200nM and NADPH 1mM) together with different concentrations of test substances. After 60 min. of incubation at 37°C the assay was stopped by heating to 80°C (5 min.) and by addition of the inhibitor Carbenoxolone (1 µM). The amount of Cortisol produced in this assay was determined using a commercially available, ELISA-based Cortisol-detection kit (Distributed by Assay Design, Inc.). Inhibitors were characterized by their IC₅₀ values, e.g. the concentration at which the production of cortisol was 50% reduced.

Cellular Assay

To measure the effect of inhibitors in intact cells HEK293 cells stably expressing human 11beta-HSD1 (see above) were cultivated in 96 well plates in DMEM. First inhibitors and 60 min later Cortisone was added to the cells. After 60 min of incubation at 37°C in a 5% CO₂ atmosphere part of the medium was removed and the conversion from Cortisone to Cortisol was measured using a commercially available ELISA kit (Distributed by Assay Design, Inc.).

Results obtained in the microsome assay using representative compounds of the invention as the test compounds are shown in the following table:

<u>Compound</u>	<i>h</i> 11-beta-HSD 1 IC ₅₀ (nM)
Example 2	0.007
Example 35	0.101

10

Compounds as described above have IC₅₀ values below 1000 nM; preferred compounds have IC₅₀ values below 100 nM. More preferred compounds have IC₅₀ values below 10 nM. These results have been obtained by using the foregoing test.

The compounds of formula I and their pharmaceutically acceptable salts and esters can be used as medicaments (e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also be effected parentally, such as intramuscularly or intravenously (e.g. in the form of injection solutions).

The compounds of formula I and their pharmaceutically acceptable salts and esters can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatin capsules. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts etc. can be used, for example, as such adjuvants for tablets, dragées and hard gelatin capsules.

Suitable adjuvants for soft gelatin capsules, are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

Suitable adjuvants for the production of solutions and syrups are, for example, water, polyols, saccharose, invert sugar, glucose, etc.

Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

In accordance with the invention the compounds of formula I and their pharmaceutically acceptable salts can be used for the prophylaxis and treatment of arthritis, cardiovascular diseases, diabetes, renal failure and particularly eating disorders and obesity. The dosage can vary in wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to 4 mg per kg body weight (e.g. about 300 mg per person), divided into preferably 1-3 individual doses, which can consist, for example, of the same amounts,

- 26 -

should be appropriate. It will, however, be clear that the upper limit given above can be exceeded when this is shown to be indicated.

The invention is illustrated hereinafter by examples, which have no limiting character.

- 27 -

Examples

Example 1:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide

5 Step A] 1-Benzyl-5-nitro-1,2-dihydro-indazol-3-one

To a solution of 2-fluoro-5-nitrobenzoic acid (0.5 g) in DMF (9 mL) was added TBTU (1.04 g) followed by N-ethyl-diisopropylamine (2.3 mL). After 10 minutes benzylhydrazine.2HCl (0.63 g) was added. The reaction mixture was stirred at ambient temperature for 22 hours and the reaction was poured onto aqueous 1N HCl solution and
10 extracted with EtOAc. The organic layer was washed with brine, dried over sodium sulfate, filtered and evaporated *in vacuo*. Purification via ISCO combiflash chromatography afforded pure desired 1-benzyl-5-nitro-1,2-dihydro-indazol-3-one (0.26 g) as a yellow solid. MS (ESI⁻): 268.3 ([M-H]⁻).

Step B] 5-Amino-1-benzyl-1,2-dihydro-indazol-3-one

15 To a solution of 1-benzyl-5-nitro-1,2-dihydro-indazol-3-one (0.44 g) in MeOH (50 mL) was added Pd/C(10%) catalyst (0.1 g) and the reaction mixture was stirred for four hours at ambient temperature under a hydrogen atmosphere using a balloon. After this time the reaction mixture was filtered through Celite® which was washed with more EtOAc. The combined organic solution was evaporated *in vacuo* for afford the desired 5-amino-1-
20 benzyl-1,2-dihydro-indazol-3-one (0.39 g) which was taken into the next reaction without further purification. MS (ESI⁺): 240.1 ([M+H]⁺).

Step C] N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide

To a solution of 5-amino-1-benzyl-1,2-dihydro-indazol-3-one (0.04 g) in pyridine (1 mL)
25 was added 3-chloro-2-methylbenzenesulfonylchloride (0.038 g) in one go. The solution was stirred at 60°C for 24 hours. The pyridine was then removed in vacuo and the residue was dissolved in EtOAc/water and separated. The aqueous phase was extracted a further two times with EtOAc and the combined organic phases were washed with brine and dried over sodium sulfate. The drying agent was removed by filtration and the volatiles were
30 removed in vacuo to afford a crude residue. The crude material was purified via flash column chromatography eluting with EtOAc/nHeptane/1%AcOH) to afford the desired N-(1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide (0.015 g) as a pale yellow powder. MS (ESI⁺): 428.4 ([M+H]⁺).

Example 2:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-fluoro-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-chloro-4-fluoro-benzenesulfonyl chloride as an orange solid. MS

5 (ESI⁻): 430.3 ([M-H]⁻).

Example 3:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-(trifluoromethyl)benzenesulfonyl chloride as a light yellow solid. MS

10 (ESI⁺): 448.1 ([M+H]⁺).

Example 4:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2,4-dichlorobenzenesulfonyl chloride as a light yellow solid. MS (ESI⁻):

15 446.1 ([M-H]⁻).

Example 5:

Naphthalene-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 1-naphthalenebenzenesulfonyl chloride as a light yellow solid.

20 MS (ESI⁻): 428.4 ([M-H]⁻).

Example 6:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-methyl-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-chloro-4-methylbenzenesulfonyl chloride as a light yellow solid. MS

25 (ESI⁺): 428.4 ([M+H]⁺).

Example 7:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methoxy-benzenesulfonamide

- 29 -

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 4-methoxybenzenesulfonyl chloride as a light yellow solid.

Example 8:

5 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-6-methylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2,4-dichloro-6-methylbenzenesulfonyl chloride as a brown solid. MS (ESI⁻): 460.1 ([M-H]⁻).

Example 9:

10 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2-trifluoromethylbenzenesulfonyl chloride as a white solid. MS (ESI⁻): 446.1 ([M-H]⁻).

Example 10:

15 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluorobenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2,4-difluorobenzenesulfonyl chloride as a brown solid. MS (ESI⁻): 414.1 ([M-H]⁻).

Example 11:

20 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-propylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 4-propylbenzenesulfonyl chloride as a light yellow solid.

Example 12:

25 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2,3-dihydro-1,4-benzodioxine-6-sulfonyl chloride as an off-white solid. MS (ESI⁺): 438.1 ([M+H]⁺).

Example 13:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-2,5-dimethylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one (0.040 g) and 4-chloro-2,5-dimethylsulfonylchloride as an off-white solid.
5 MS (ESI⁺): 442.1 ([M+H]⁺).

Example 14:

N-(1-Benzyl-3-oxo-2,3-dihydro-H-indazol-5-yl)-4-chloro-3-trifluoromethylbenzenesulfonamide

10 This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-trifluoromethyl-4-chlorobenzenesulfonylchloride as a white solid.
MS (ESI⁺): 482.3 ([M+H]⁺).

Example 15:

15 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 4-fluoro-3-trifluoromethylbenzenesulfonylchloride as a light brown solid. MS (ESI⁻): 464.4 ([M-H]⁻).

Example 16:

20 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-5-trifluoromethylbenzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 2-chloro-5-trifluoromethylbenzenesulfonyl chloride as a light brown solid. MS (ESI⁻): 480.3[M-H]⁻).

25 Example 17:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-phenoxy-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-phenoxy-benzenesulfonyl chloride as a light brown solid. MS (ESI⁻): 470.3[M-H]⁻).

Example 18:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-(5-methyl-[1,3,4]oxadiazol-2-yl)-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-(5-methyl-1,3,4-oxadiazol-2-yl)benzenesulfonyl chloride as a light brown solid. MS (ESI⁻): 460.4[M-H]⁻.

Example 19:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-chloro-benzenesulfonyl chloride as a light brown solid. MS (ESI⁻): 412.1[M-H]⁻.

Example 20:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-methoxy-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-methoxy-benzenesulfonyl chloride as a light brown solid. MS (ESI⁻): 408.0[M-H]⁻.

Example 21:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxy-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3-(difluoromethoxy)-benzenesulfonyl chloride as a colorless solid. MS (ESI⁻): 444.1[M-H]⁻.

Example 22:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3,5-bis-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3,5-bis(trifluoromethyl)benzenesulfonyl chloride (step C) as a white solid. MS (ESI⁻): 514.3 [M-H]⁻.

Example 23:

Propane-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

This compound was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and 3 propane-2-sulfonyl chloride (step C) as a white solid. MS (ESI⁻): 344.3 [M-H]⁻).

5 Example 24:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-N-dimethyl-benzenesulfonamide

To a solution of NaH (0.005 g of a 60% dispersion in mineral oil) in DMF (2 mL) at 0°C was added N-(1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-
10 benzenesulfonamide in DMF (1 mL) dropwise. After 15 minutes MeI (0.017 g) was added and the reaction mixture was allowed to warm to ambient temperature. The reaction was quenched with water and diluted with EtOAc. The phases were separated and the aqueous phase was extracted with more EtOAc. The combined organic phases were washed with
15 brine, dried over sodium sulfate, filtered and reduced *in vacuo*. Flash column chromatography over silica (EtOAc/nheptane) afforded the desired N-(1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-N-dimethyl-benzenesulfonamide as an off-white solid (13 mg). MS (ESI⁺): 442.4[M+H]⁺).

Example 25:

20 N-(3-Oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

Step A] 1-Allyl-5-nitro-1,2-dihydro-indazol-3-one

5-Nitro-1,2-dihydro-indazol-3-one (prepared according to *Org. Synth.* 1949, 29, 54 or *Chem Ber.* 1942, 75, 1104) (9.36 g) was suspended in 40 ml water and 57.5 ml 1N KOH. Allyl bromide (6.32 g) was added in one portion. The mixture was stirred at 75-80°C for
25 1.5 hours. Then NaOH (15%, 5 mL) and allyl bromide (1 g) was added. The reaction mixture was stirred for a further 30 min. The mixture was neutralized with 3N HCl at <10°C and filtered. The solid was washed with water and dried. The solid was then suspended in 10 ml ethyl acetate and stirred at ambient temperature for 3 hours, then filtered and dried to afford the desired 1-allyl-5-nitro-1,2-dihydro-indazol-3-one (7.63 g)
30 as a yellow solid.

Step B] 5-Amino-1-propyl-1,2-dihydro-indazol-3-one

- 33 -

1-Allyl-5-nitroindazolone (1.42 g) and Pd/C (10%, 250 mg) were suspended in MeOH (50 mL) and hydrogenated (hydrogen balloon) at RT for 4-5 h (or overnight.) After filtration and removal of MeOH, the solid was dried in vacuo to afford the desired 5-amino-1-propyl-1,2-dihydro-indazol-3-one (1.2 g) as a crude oil. This material was used in the next
5 reaction without any further purification.

Step C] N-(3-Oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

This material was obtained in analogy to example 1 using 5-amino-1-propyl-1,2-dihydro-indazol-3-one and 2,3-dichlorobenzenesulfonyl chloride.

10 Example 26:

2,3-Dichloro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This material was obtained in analogy to example 25 using 5-amino-1-propyl-1,2-dihydro-indazol-3-one and 2,3-dichlorobenzenesulfonyl chloride.

Example 27:

15 3-Chloro-4-fluoro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This material was obtained in analogy to example 25 using 5-amino-1-propyl-1,2-dihydro-indazol-3-one and 3-chloro-4-fluoro-benzenesulfonyl chloride.

Example 28:

20 3-Chloro-2-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 1 using phenethylhydrazine (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a light yellow solid. MS (ESI⁺): 442.3 [M+H]⁺.

25 Example 29:

3-Chloro-4-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

- 34 -

This compound was obtained in analogy to example 1 using phenethylhydrazine (step A) and 3-chloro-4-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 442.3[M+H]⁺.

Example 30:

- 5 4-Chloro-2,5-dimethyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 1 using phenethylhydrazine (step A) and 4-chloro-2,5-dimethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 456.4[M+H]⁺.

- 10 Example 31:

N-(3-Oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 1 using phenethylhydrazine (step A) and 3-trifluoromethylsulfonyl chloride (step C) as a red solid. MS (ESI⁺): 462.1[M+H]⁺.

- 15 Example 32:

N-(1-Isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

Step A] 1-Isobutyl-5-nitro-1,2-dihydro-indazol-3-one

- 20 To a solution of 2-fluoro-5-nitrobenzoic acid (1.5 g) in DMF (50 mL) was added EDCI.HCl (1.71 g) followed by N-ethyl-diisopropylamine (5.51 mL). After 10 minutes isobutylhydrazine.p-toluenesulfonic acid salt (2.32 g) was added. The reaction mixture was stirred at ambient temperature for 22 hours and the reaction was poured onto aqueous 1N HCl solution and extracted with EtOAc. The organic layer was washed with brine, dried over sodium sulfate, filtered and evaporated *in vacuo*. Purification via ISCO combiflash
25 chromatography afforded pure desired 1-isobutyl-5-nitro-1,2-dihydro-indazol-3-one (0.9 g) as a yellow solid. MS (ESI⁻): 234.1 ([M-H]⁻).

Step B] 5-Amino-1-isobutyl-1,2-dihydro-indazol-3-one

- 30 To a solution of 1-isobutyl-5-nitro-1,2-dihydro-indazol-3-one (0.9 g) in MeOH (50 mL) was added Pd/C(10%) catalyst (0.3 g) and the reaction mixture was stirred for 1 hour at 40°C under a hydrogen atmosphere using a balloon. After this time the reaction mixture was filtered through Celite® which was washed with more EtOAc. The combined organic

- 35 -

solution was evaporated *in vacuo* for afford the desired 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one (0.6 g) which was taken into the next reaction without further purification. MS (ESI⁺): 206.1 ([M+H]⁺).

Step C] N-(1-Isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
5 benzenesulfonamide

To a solution of 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one (0.04 g) in pyridine (1 mL) was added 3-trifluoromethylbenzenesulfonylchloride (0.041 g) in one go. The solution was stirred at 60°C for 24 hours. The pyridine was then removed in vacuo and the residue was dissolved in EtOAc/water and separated. The aqueous phase was extracted a further
10 two times with EtOAc and the combined organic phases were washed with brine and dried over sodium sulfate. The drying agent was removed by filtration and the volatiles were removed in vacuo to afford a crude residue. The crude material was purified via flash column chromatography eluting with EtOAc/nHeptane/1%AcOH) to afford the desired
N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
15 benzenesulfonamide (0.015 g) as a pale yellow powder. MS (ESI⁺): 414.4[M+H]⁺.

Example 33:

3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methyl-
benzenesulfonamide

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-
20 dihydro-indazol-3-one and 3-chloro-4-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 394.1[M+H]⁺.

Example 34:

3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-
benzenesulfonamide

25 This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 394.1[M+H]⁺.

Example 35:

3-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 3-cyano-benzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 371.1 [M+H]⁺.

Example 36:

5 **3-Difluoromethoxy-(1-isobutyl-3-oxo-2,3-dihydro--indazol-5-yl)-benzenesulfonamide**

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 3-difluoromethoxy-benzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 412.4 [M+H]⁺.

Example 37:

10 **4-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide**

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 4-cyano-benzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 371.1 [M+H]⁺.

Example 38:

15 **4-Fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide**

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 432.4 [M+H]⁺.

20 **Example 39:**

3-Chloro-4-fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as an off-
25 white solid. MS (ESI⁺): 398.1 [M+H]⁺.

Example 40:

2,4-Dichloro-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methyl-benzenesulfonamide

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 2,4-dimethyl-5-methylbenzenesulfonyl chloride (step C) as an orange solid. MS (ESI⁺): 428.3 [M+H]⁺.

Example 41:

5 3-Chloro-N-(1-ethyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-benzenesulfonamide

This compound was obtained in analogy to example 1 using ethylhydrazine.oxalate salt (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as light yellow solid. MS (ESI⁺): 366.0 [M+H]⁺.

Example 42:

10 3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-2-methyl-benzenesulfonamide

This compound was obtained in analogy to example 32 using cyclopropylmethylhydrazine (step A) (prepared as described in *J. Am. Chem. Soc.* 1986, 108, 7981-4) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁺): 392.0
15 [M+H]⁺.

Example 43:

3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-benzenesulfonamide

This compound was obtained in analogy to example 32 using cyclopropylmethylhydrazine (step A) (prepared as described in *J. Am. Chem. Soc.* 1986, 108, 7981-4) and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a pink solid. MS (ESI⁺): 396.3
20 [M+H]⁺.

Example 44:

(1-Cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-3-trifluoromethyl-benzenesulfonamide
25

This compound was obtained in analogy to example 32 using cyclopropylmethylhydrazine (step A) (prepared as described in *J. Am. Chem. Soc.* 1986, 108, 7981-4) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁺): 430.4 [M+H]⁺.

30 Example 45:

3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 32 using cyclopropylmethylhydrazine (step A) (prepared as described in *J. Am. Chem. Soc.* 1986, 108, 7981-4) and 3-chlorobenzenesulfonyl chloride (step C) as an orange solid. MS (ESI⁺): 378.3 [M+H]⁺.

5 Example 46:**2,4-Dichloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methylbenzenesulfonamide**

This compound was obtained in analogy to example 32 using cyclopropylmethylhydrazine (step A) (prepared as described in *J. Am. Chem. Soc.* 1986, 108, 7981-4) and 2,4-dichloro-5-methylbenzenesulfonyl chloride (step C) as an orange solid. MS (ESI⁺): 426.1 [M+H]⁺.

Example 47:**3-Chloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methylbenzenesulfonamide**

15 This compound was obtained in analogy to example 32 using (2-methoxy-ethyl)hydrazine (step A) (prepared as described in *J. Org. Chem.* 1984, 49, 336-42) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 396.3 [M+H]⁺.

Example 48:**20 4-Fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethylbenzenesulfonamide**

This compound was obtained in analogy to example 32 using (2-methoxy-ethyl)hydrazine (step A) (prepared as described in *J. Org. Chem.* 1984, 49, 336-42) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 434.3 [M+H]⁺.

25 Example 49:**3-Chloro-4-fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide**

This compound was obtained in analogy to example 32 using (2-methoxy-ethyl)hydrazine (step A) (prepared as described in *J. Org. Chem.* 1984, 49, 336-42) and 3-chloro-

4-fluorobenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 400.3 [M+H]⁺.

Example 50:

2,4-Dichloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-
5 benzenesulfonamide

This compound was obtained in analogy to example 32 using (2-methoxy-ethyl)-hydrazine (step A) (prepared as described in *J. Org. Chem.* 1984, 49, 336-42) and 2,4-dichloro-6-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 430.3 [M+H]⁺.

10 Example 51:

3-Chloro-2-methyl-N-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide

This compound was obtained in analogy to example 32 using (2,2,2-trifluoro-ethyl)-hydrazine (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a red solid.
15 MS (ESI⁺): 420.3 [M+H]⁺.

Example 52:

4-Fluoro-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-indazol-5-yl]-3-trifluoromethyl-
benzenesulfonamide

This compound was obtained in analogy to example 32 using (2,2,2-trifluoro-ethyl)-
20 hydrazine (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a brown solid. MS (ESI⁺): 458.4 [M+H]⁺.

Example 53:

4-Fluoro-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-3-
trifluoromethyl-benzenesulfonamide

25 This compound was obtained in analogy to example 32 using pyridin-2-ylmethyl-hydrazine (step A) (prepared in analogy to examples described in *J. Org. Chem.* 1984, 49, 336-42) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as off-white solid. MS (ESI⁺): 467.4 [M+H]⁺.

Example 54:

2,4-Dichloro-6-methyl-*N*-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1*H*-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 32 using pyridin-2-ylmethylhydrazine (step A) (prepared in analogy to examples described in *J. Org. Chem.* 1984, 49, 336-42) and 2,4-dichloro-6-methylbenzenesulfonyl chloride (step C) as off-white solid.
5 MS (ESI⁺): 463.3 [M+H]⁺.

Example 55:

3-Chloro-2-methyl-*N*-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1*H*-indazol-5-yl)-benzenesulfonamide

10 This compound was obtained in analogy to example 32 using pyridin-2-ylmethylhydrazine (step A) (prepared in analogy to examples described in *J. Org. Chem.* 1984, 49, 336-42) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as off-white solid. MS (ESI⁺): 429.4 [M+H]⁺.

Example 56:

15 4-Fluoro-*N*-[1-(2-hydroxy-2-methyl-propyl)-3-oxo-2,3-dihydro-1*H*-indazol-5-yl]-3-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 32 using pyridin-2-ylmethylhydrazine (step A) (prepared as described in *Bioorg. Med. Chem.* 2004, 12, 1357-1366) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as off-white solid. MS (ESI⁺):
20 448.3 [M+H]⁺.

Example 57:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1*H*-indazol-5-yl)-3-chloro-4-fluorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 464.1 [M-H]⁻.
25

Example 58:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1*H*-indazol-5-yl)-4-fluoro-3-trifluoromethylbenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 498.0 [M-H]⁺).

5 Example 59:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3-chloro-2-methylbenzenesulfonyl
10 chloride (step C) as a light brown solid. MS (ESI⁺): 460.0 [M-H]⁺).

Example 60:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichlorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 2,3-dichloro-benzenesulfonyl chloride
15 (step C) as a light brown solid. MS (ESI⁺): 480.1 [M-H]⁺).

Example 61:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3-cyano-benzenesulfonyl chloride (step
20 C) as a light brown solid. MS (ESI⁺): 437.0 [M-H]⁺).

Example 62:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3-chloro-benzenesulfonyl chloride (step
25 C) as a white solid. MS (ESI⁺): 446.1 [M-H]⁺).

Example 63:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-3-trifluoromethylbenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzyldiazine.2HCl (step A) and 4-chloro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid.

MS (ESI⁺): 514.3 [M-H]⁺.

5 Example 64:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxybenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzyldiazine.2HCl (step A) and 4-chloro-3-

10 trifluoromethylbenzenesulfonyl chloride (step C) as a white solid.

MS (ESI⁺): 478.0 [M-H]⁺.

Example 65:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethylbenzenesulfonamide

15 This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzyldiazine.2HCl (step A) and 2-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 480.0 [M-H]⁺.

Example 66:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxybenzenesulfonamide

20

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzyldiazine.2HCl (step A) and 3-trifluoromethoxybenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 496.4 [M-H]⁺.

Example 67:

25 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,5-difluorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitrobenzoic acid, benzyldiazine.2HCl (step A) and 2,5-difluorobenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁺): 447.9 [M-H]⁺.

30 Example 68:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzyldiazine.2HCl (step A) and 2,4-dichlorobenzenesulfonyl chloride
5 (step C) as a light brown solid. MS (ESI⁺): 482.4 [M+H]⁺).

Example 69:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluoro-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-
10 benzoic acid, benzyldiazine.2HCl (step A) and 3-chloro-2-fluorobenzenesulfonyl chloride (step C) as a yellow solid. MS (ESI⁻): 464.1 [M-H]⁻).

Example 70:

N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-benzenesulfonamide

15 This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzyldiazine.2HCl (step A) and 3-trifluoromethylbenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁻): 480.0 [M-H]⁻).

Example 71:

20 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzyldiazine.2HCl (step A) and 3-chloro-6-fluorobenzenesulfonyl chloride (step C) as a brown solid. MS (ESI⁻): 464.1 [M-H]⁻).

Example 72:

25 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-benzenesulfonamide

This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzyldiazine.2HCl (step A) and 2-chlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 446.1 [M-H]⁻).

Example 73:

(1-Benzyl-4-chloro-3-oxo-2,3-dihydro-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 32 using 2-chloro-6-fluoro-3-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 460.1 [M-H]⁻.

Example 74:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethylbenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 484.5 [M+H]⁺.

Example 75:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichlorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 2,3-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 464.1 [M-H]⁻.

Example 76:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichlorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 2,4-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 464.1 [M-H]⁻.

Example 77:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitrobenzoic acid, benzylhydrazine.2HCl (step A) and 3-chloro-2-fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 447.9 [M-H]⁻.

Example 78:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic
5 acid, benzylhydrazine.2HCl (step A) and 3-chloro-6-fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 447.9 [M-H]⁻).

Example 79:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxybenzenesulfonamide

10 This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3-trifluoromethoxybenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 482.4 [M+H]⁺).

Example 80:

15 3-Chloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

Step A] 1-(4-Fluoro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one

5-Nitro-1,2-dihydro-indazol-3-one (prepared according to *Org. Synth.* 1949, 29, 54 or *Chem Ber.* 1942, 75, 1104) (0.7 g) was suspended in 1 N NaOH (4.2 mL) and stirred for 10 minutes at 75°C. 4-Fluorobenzylbromide (0.48 mL) was then added dropwise over 2
20 hours. The reaction was stirred overnight and was quenched with 2 N HCl (aq.) solution while cooling in an ice bath. The precipitate was collected by filtration and the crude solid was either purified by flash column chromatography or by trituration with ether to give pure 1-(4-fluoro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one. MS (ESI⁻): 286.1 [M-H]⁻).

Step B] 5-Amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one

25 To a solution of 1-(4-fluoro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one (0.81 g) in MeOH (100 mL) was added Pd/C(10%) catalyst (0.1 g) and the reaction mixture was stirred for three hours at 40°C under a hydrogen atmosphere using a balloon. After this time the reaction mixture was filtered through Celite® which was washed with more EtOAc. The combined organic solution was evaporated *in vacuo* for afford the desired 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one (0.617 g) which was taken into the next
30 reaction without further purification. MS (ESI⁺): 258.3 ([M+H]⁺).

- 46 -

Step C] 3-Chloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

To a solution of 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one (0.100 g) in pyridine (1 mL) was added 3-chlorobenzenesulfonylchloride (0.082 g) in one go. The solution was stirred at 60°C for 24 hours. The pyridine was then removed in vacuo and the residue was dissolved in EtOAc/water and separated. The aqueous phase was extracted a further two times with EtOAc and the combined organic phases were washed with brine and dried over sodium sulfate. The drying agent was removed by filtration and the volatiles were removed in vacuo to afford a crude residue. The crude material was purified via flash column chromatography eluting with EtOAc/nHeptane/1%AcOH) to afford the desired 3-chloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide (0.087 g) as a brown solid. MS (ESI⁺): 432.4 [M+H]⁺.

Example 81:

3-Difluoromethoxy-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one and 3-difluoromethoxybenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 464.4 [M+H]⁺.

Example 82:

4-Fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethylbenzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 482.3 [M-H]⁻.

Example 83:

2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one and 3-dihydro-1,4-benzodioxine-6-sulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 456.5 [M-H]⁻.

Example 84:

2,4-Dichloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methylbenzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluorobenzyl)-1,2-dihydro-indazol-3-one and 2,4-dichloro-6-methylbenzenesulfonyl chloride
5 (step C) as a light brown solid. MS (ESI⁺): 480.4 [M+H]⁺.

Example 85:

5-Chloro-2,4-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluorobenzyl)-1,2-dihydro-indazol-3-one and 3-chloro-4,6-difluorobenzenesulfonyl chloride
10 (step C) as an off-white solid. MS (ESI⁺): 468.4 [M+H]⁺.

Example 86:

4-Chloro-2,5-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluorobenzyl)-1,2-dihydro-indazol-3-one and 4-chloro-2,5-difluorobenzenesulfonyl chloride
15 (step C) as an off-white solid. MS (ESI⁺): 468.4 [M+H]⁺.

Example 87:

3-Chloro--[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methylbenzenesulfonamide
20

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluorobenzyl)-1,2-dihydro-indazol-3-one and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 446.0 [M+H]⁺.

Example 88:

3-Cyano-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide
25

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluorobenzyl)-1,2-dihydro-indazol-3-one and 3-cyanobenzenesulfonyl chloride (step C) as light brown solid. MS (ESI⁺): 423.1 [M+H]⁺.

Example 89:

Naphthalene-1-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one and 1-naphthalenesulfonyl chloride (step C) as light
5 brown solid. MS (ESI⁺): 448.3 [M+H]⁺.

Example 90:

3-Chloro-4-fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 5-amino-1-(4-fluoro-benzyl)-1,2-dihydro-indazol-3-one and 3-chloro-4-fluorobenzenesulfonyl chloride (step
10 C) as light brown foam. MS (ESI⁺): 449.9 [M+H]⁺.

Example 91:

3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide
15 (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 464.3 [M+H]⁺.

Example 92:

3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-
20 benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 468.4 [M+H]⁺.

Example 93:

[1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide
25

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 502.1 [M+H]⁺.

Example 94:

3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3-chlorobenzenesulfonyl chloride (step C) as an orange solid. MS (ESI⁺):

5 450.3 [M+H]⁺).

Example 95:

[1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethoxy-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3-trifluoromethoxybenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 500.3 [M+H]⁺).

10

Example 96:

2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-5-methyl-benzenesulfonamide

15 This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 2,4-dichloro-5-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 498.4 [M+H]⁺).

Example 97:

20 3,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3,4-dichloro-benzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 484.4 [M+H]⁺).

Example 98:

25 4,5-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-fluoro-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3,4-dichloro-6-fluorobenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 502.1 [M+H]⁺).

Example 99:

2,4,5-Trichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 3,4,6-trichloro-benzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 520.3 [M+H]⁺.

Example 100:

2,3-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 2,3-dichlorobenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 484.4 [M+H]⁺.

Example 101:

2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-benzenesulfonamide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide (step A) and 2,4-dichloro-6-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 500.3 [M+H]⁺.

Example 102:

3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluorobenzenesulfonamide

This compound was obtained in analogy to example 80 using 2,4-difluorobenzylbromide (step A) and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a pink solid. MS (ESI⁺): 468.4 [M+H]⁺.

Example 103:

N-[1-(2,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 80 using 2,4-difluorobenzylbromide (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 502.1 [M+H]⁺.

Example 104:

- 5 3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 80 using 2,4-difluorobenzylbromide (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 464.3 [M+H]⁺.

- 10 Example 105:

N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide

Step A] 1-(4-Chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one

- 15 This compound was obtained in analogy to step A in example 77 using 5-nitro-1,2-dihydro-indazol-3-one (prepared according to *Org. Synth.* 1949, 29, 54 or *Chem Ber.* 1942, 75, 1104) and 4-chlorobenzylbromide to afford the desired 1-(4-chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one as a brown solid. MS (ESI⁻): 302.3 [M-H]⁻.

Step B] 5-Amino-1-(4-chloro-benzyl)-1,2-dihydro-indazol-3-one

- 20 To 1-(4-chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one (0.45 g) was added HCl (37% aq.) solution (5 mL) followed by SnCl₂.dihydrate (1.9 g). The mixture was heated to 85°C for one hour. The reaction was then neutralized with sodium bicarbonate solution, filtered, and the filtrate extracted with EtOAc. The combined organic phases were washed with brine, dried over sodium sulfate, filtered and reduced in vacuo. Flash column chromatography of the residue over silica gel eluting with methylene chloride/methanol
25 afforded the desired 5-amino-1-(4-chloro-benzyl)-1,2-dihydro-indazol-3-one (0.31 g) as a dark brown solid. MS (ESI⁺): 274.4 [M+H]⁺.

Step C] N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide

- 30 This compound was obtained in analogy to step C in example 77 using 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 500.3 [M+H]⁺.

Example 106:

3-Chloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 105 using 5-amino-1-(4-chloro-benzyl)-1,2-dihydro-indazol-3-one and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 462.1 [M+H]⁺.

Example 107:

2,3-Dichloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 105 using 5-amino-1-(4-chloro-benzyl)-1,2-dihydro-indazol-3-one and 2,3-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 482.4 [M+H]⁺.

Example 108:

Naphthalene-1-sulfonic acid [1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

This compound was obtained in analogy to example 105 using 5-amino-1-(4-chloro-benzyl)-1,2-dihydro-indazol-3-one and 1-naphthalenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 464.1 [M+H]⁺.

Example 109:

3-Chloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methylbenzenesulfonamide

This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 480.4 [M+H]⁺.

Example 110:

Naphthalene-1-sulfonic acid [1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

- 53 -

This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 1-naphthalenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 482.4 [M+H]⁺.

Example 111:

- 5 2,3-Dichloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 2,3-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 500.1 [M+H]⁺.

- 10 Example 112:

N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide

- 15 This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁻): 516.2 [M-H]⁻.

Example 113:

N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-difluoromethoxy-benzenesulfonamide

- 20 This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 3-difluoromethoxybenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁻): 496.3 [M-H]⁻.

Example 114:

N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethoxy-benzenesulfonamide

- 25 This compound was obtained in analogy to example 105 using 2-chloro-4-fluorobenzylbromide (step A) and 3-trifluoromethoxybenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁻): 514.3 [M-H]⁻.

Example 115:

3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-benzenesulfonamide

Step A] 3-(5-Nitro-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile

5-Nitro-1,2-dihydro-indazol-3-one (prepared according to *Org. Synth.* 1949, 29, 54 or
5 *Chem Ber.* 1942, 75, 1104) (0.6 g) was suspended in water (8 mL) and 2N K₂CO₃ (aq.) (1.8 g) was added to the flask and the mixture was heated to 50°C for 10 minutes. Acrylonitrile (0.24 mL) was then added dropwise. The reaction was stirred overnight and was quenched with 2 N HCl (aq.) solution while cooling in an ice bath. The aqueous phase was extracted with EtOAc and the combined phases were washed with brine, dried over sodium sulfate,
10 filtered and reduced in vacuo. Flash column chromatography over silica gel (eluting with EtOAc, nheptane, 3% AcOH) afforded pure desired 3-(5-Nitro-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile as a white solid. MS (ESI⁺: 233.3[M+H]⁺).

Step B] 3-(5-Amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile

To a solution of 3-(5-Nitro-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile (0.521 g) in
15 MeOH (50 mL) was added Pd/C(10%) catalyst (0.13 g) and the reaction mixture was stirred for 20 hours at ambient temperature under a hydrogen atmosphere using a balloon. After this time the reaction mixture was filtered through Celite® which was washed with more EtOAc. The combined organic solution was evaporated *in vacuo* for afford the desired 3-(5-amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile (0.45 g) which was
20 taken into the next reaction without further purification. MS (ESI⁺): 203.4 ([M+H]⁺).

Step C] 3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-benzenesulfonamide

To a solution of 3-(5-amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile (0.08 g) in pyridine (1 mL) was added 3-chloro-2-methylbenzenesulfonylchloride (0.089 g) in one go.
25 The solution was stirred at 60°C for 24 hours. The pyridine was then removed in vacuo and the residue was dissolved in EtOAc/water and separated. The aqueous phase was extracted a further two times with EtOAc and the combined organic phases were washed with brine and dried over sodium sulfate. The drying agent was removed by filtration and the volatiles were removed in vacuo to afford a crude residue. The crude material was
30 purified via flash column chromatography eluting with EtOAc/nHeptane/1%AcOH) to afford the desired 3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-benzenesulfonamide (0.087 g) as a light brown solid. MS (ESI⁺): 391.0 [M+H]⁺).

Example 116:

N-[1-(2-Cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide

This compound was obtained in analogy to example 115 using 3-(5-amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 429.5 [M+H]⁺.

Example 117:

2,4-Dichloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methylbenzenesulfonamide

This compound was obtained in analogy to example 115 using 3-(5-amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile and 2,4-dichloro-6-methylbenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁺): 425.3 [M+H]⁺.

Example 118:

3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluorobenzenesulfonamide

This compound was obtained in analogy to example 115 using 3-(5-amino-3-oxo-2,3-dihydro-indazol-1-yl)-propionitrile and 3-chloro-4-fluorobenzenesulfonyl chloride (step C) as a light red solid. MS (ESI⁺): 395.3 [M+H]⁺.

Example 119:

N-[6-Chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide

Step A] 6-Chloro-1-(2-chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one

6-Chloro-5-nitro-1,2-dihydro-indazol-3-one (prepared in analogy to *Org. Synth.* 1949, 29, 54 or *Chem Ber.* 1942, 75, 1104 from 4-chloro-2-fluoro-5-nitro-benzoic acid over two steps) (0.6 g) was suspended in water (5 mL) and 2N K₂CO₃ (aq.) (1 g) was added to the flask and the mixture was heated to 50°C for 10 minutes. 2-Chlorobenzylbromide (0.40 g) was then added dropwise. The reaction was stirred overnight and was neutralised with 2 N HCl (aq.) while cooling in an ice bath. The precipitate was collected by filtration and triturated with diethylether to afford pure desired 6-chloro-1-(2-chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one as a brown solid. MS (ESI⁻): 336.3[M-H]⁻.

Step B] 5-Amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one

- 56 -

To 6-chloro-1-(2-chloro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one (0.49 g) was added HCl (37% aq.) solution (4 mL) followed by $\text{SnCl}_2 \cdot \text{dihydrate}$ (2.6 g). The mixture was heated to 85°C for one hour. The reaction was then neutralized with sodium bicarbonate solution, filtered, and the filtrate extracted with EtOAc. The combined organic phases were washed with brine, dried over sodium sulfate, filtered and reduced in vacuo. Flash column chromatography of the residue over silica gel eluting with methylene chloride/methanol afforded the desired 5-amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one (0.33 g) as a yellow solid. MS (ESI^+): 308.3 $[\text{M}+\text{H}]^+$.

Step C] N-[6-Chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide

To a solution of 5-amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one (0.070 g) in pyridine (1 mL) was added 4-fluoro-3-trifluoromethylbenzenesulfonylchloride (0.072 g) in one go. The solution was stirred at 60°C for 24 hours. The pyridine was then removed in vacuo and the residue was dissolved in EtOAc/water and separated. The aqueous phase was extracted a further two times with EtOAc and the combined organic phases were washed with brine and dried over sodium sulfate. The drying agent was removed by filtration and the volatiles were removed in vacuo to afford a crude residue. The crude material was purified via flash column chromatography eluting with EtOAc/nHeptane/1%AcOH to afford the desired N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide (0.024 g) as a light brown solid. MS (ESI^+): 534.3 $[\text{M}+\text{H}]^+$.

Example 120:

5-Chloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-fluorobenzenesulfonamide

This compound was obtained in analogy to example 119 using 5-amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one and 3-chloro-6-fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI^+): 500.3 $[\text{M}+\text{H}]^+$.

Example 121:

2,3-Dichloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide

This compound was obtained in analogy to example 119 using 5-amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one and 2,3-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI^+): 518.0 $[\text{M}+\text{H}]^+$.

Example 122:

3-Chloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-2-methyl-benzenesulfonamide

This compound was obtained in analogy to example 119 using 2-chloromethyl-pyridine (step A) and 3-chloro-2-methylbenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 463.1[M+H]⁺.

Example 123:

N-(6-Chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 119 using 2-chloromethyl-pyridine (step A) and 4-fluoro-3-trifluoromethylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 501.4[M+H]⁺.

Example 124:

2-[5-(3-Chloro-2-methyl-benzenesulfonylamino)-3-oxo-2,3-dihydro-indazol-1-yl]-methyl-acetamide

Part A] (5-Nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetic acid ethyl ester

This compound was obtained in analogy to example 1 (step A) using hydrazino-acetic acid ethyl ester.HCl as an orange solid. MS (ESI⁻): 264.1[M-H]⁻.

Part B] (5-Nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetic acid

To a solution of (5-nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetic acid ethyl ester (0.24 g) in THF (9 mL), was added 2 N NaOH (aq.) (1.8 mL). The reaction mixture was stirred at ambient temperature for 30 minutes. The reaction mixture was acidified with 2 N HCl (aq.) and extracated with EtOAc. Evaporation of the organic phases afforded the desired crude (5-nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetic acid (0.146 g) which was taken into the next step without further purification.

Part C] N-Methyl-2-(5-nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetamide

To a solution of crude (5-nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetic acid (0.14 g) in DMF, was added TBTU (0.20 g), followed by N,N-ethyldiisopropylamine (0.40 mL) and after 10 minutes methylamine (0.32 mL of a 2 M solution) was added. The reaction was stirred overnight and was quenched with 2N HCl (aq.) and extracted with EtOAc. The

combined organic layers were washed with brine, dried over sodium sulfate, filtered and evaporated. Flash column chromatography of the residue over silica gel afforded the desired N-methyl-2-(5-nitro-3-oxo-2,3-dihydro-indazol-1-yl)-acetamide (0.040 g) as a yellow solid. MS (ESI⁻): 249.1 ([M-H]⁻).

- 5 Part D and E] 2-[5-(3-Chloro-2-methyl-benzenesulfonylamino)-3-oxo-2,3-dihydro-indazol-1-yl]-methyl-acetamide

This compound was obtained in analogy to example 1 (step B and C) using 3-chloro-2-methylbenzenesulfonyl chloride (step C) as an off-white solid. MS (ESI⁺): 409.1 [M+H]⁺.

Example 125:

- 10 2,4-Dichloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-benzenesulfonamide

Part A] 1-(3,4-Difluoro-benzyl)-2-methyl-5-nitro-1,2-dihydro-indazol-3-one

- To a solution of 1-(3,4-difluoro-benzyl)-5-nitro-1,2-dihydro-indazol-3-one (prepared in analogy to example 80 using 3,4-difluorobenzylbromide (step A)) (0.2 g) in DMF was
15 added NaH (0.019 g) and the solution was stirred for 1 hour. Iodomethane (0.14 g) was then added and the mixture was stirred overnight. The reaction mixture was quenched with water and extracted with EtOAc. The combined organic layers were washed with brine, dried over sodium sulfate, filtered and evaporated in vacuo to afford a crude orange residue. Flash column chromatography of the crude residue over silica gel afforded the
20 desired 1-(3,4-difluoro-benzyl)-2-methyl-5-nitro-1,2-dihydro-indazol-3-one (0.11 g) as a yellow solid. MS (ESI⁻): 318.1 ([M-H]⁻).

Part B] 12,4-Dichloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-benzenesulfonamide

- This compound was obtained in analogy to example 1 (step B and C) using 2,4-dichloro-
25 6-methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁺): 534.2 [M+Na]⁺.

Example 126:

3-Chloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-benzenesulfonamide

- This compound was obtained in analogy to example 125 using 3-chloro-2-
30 methylbenzenesulfonyl chloride (step C) as a white solid. MS (ESI⁻): 478.3 [M-H]⁻.

Example 127:

2,3-Dichloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 119 using 2-chloromethyl-pyridine (step A) and 2,3-dichlorobenzenesulfonyl chloride (step C) as a light brown solid. MS
5 (ESI⁺): 482.9[M+H]⁺.

Example 128:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluorobenzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic
10 acid, benzylhydrazine.2HCl (step A) and 2,4-difluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 434.1[M+H]⁺.

Example 129:

N-(6-Chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-benzenesulfonamide

15 This compound was obtained in analogy to example 32 (step B and C) using 5-nitro-1-phenyl-1,2-dihydro-indazol-3-one (prepared according to *Combinatorial Chemistry and High Throughput Screening* 2003, 6(5), 471-480) in step B and 2,4-difluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 402.3[M+H]⁺.

Example 130:

20 N-(6-Chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-benzenesulfonamide

This compound was obtained in analogy to example 32 (step B and C) using 5-nitro-1-phenyl-1,2-dihydro-indazol-3-one (prepared according to *Combinatorial Chemistry and High Throughput Screening* 2003, 6(5), 471-480) in step B and and 3-chloro-2-
25 fluorobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 418.2[M+H]⁺.

Example 131:

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-benzenesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3-cyanobenzenesulfonyl chloride (step C) as a
30 light brown solid. MS (ESI⁺): 423.0[M+H]⁺.

Example 132:

Propane-2-sulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3 propane-2-sulfonyl chloride (step C) as a light
5 brown solid. MS (ESI⁺): 364.1[M+H]⁺.

Example 133:

3-Cyano-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide

This compound was obtained in analogy to example 32 (step B and C) using 5-nitro-1-phenyl-1,2-dihydro-indazol-3-one (prepared according to *Combinatorial Chemistry and*
10 *High Throughput Screening* 2003, 6(5), 471-480) in step B and and 3-cyanobenzenesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 391.1[M+H]⁺.

Example 134:

2,2,2-Trifluoro-ethanesulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

15 This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) 2,2,2-trifluoro-ethanesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 404.1[M+H]⁺.

Example 135:

Propane-2-sulfonic acid (1-benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

20 This compound was obtained in analogy to example 32 using 4-chloro-2-fluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and 3 propane-2-sulfonyl chloride (step C) as a light beige solid. MS (ESI⁺): 380.1 [M+H]⁺.

Example 136:

25 Propane-2-sulfonic acid [6-chloro-1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

This compound was obtained in analogy to example 119 using 4-chlorobenzylbromide and 5-amino-6-chloro-1-(2-chloro-benzyl)-1,2-dihydro-indazol-3-one (step A) and 3 propane-2-sulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 414.3 [M+H]⁺.

Example 137:

- 61 -

N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-trifluoromethanesulfonamide

This compound was obtained in analogy to example 32 using 2,4-difluoro-5-nitro-benzoic acid, benzylhydrazine.2HCl (step A) and trifluoromethanesulfonyl chloride (step C) as a
5 light brown solid. MS (ESI⁺): 390.1 [M+H]⁺.

Example 138:

Propane-2-sulfonic acid (1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide

This compound was obtained in analogy to example 32 using 5-amino-1-isobutyl-1,2-dihydro-indazol-3-one and 3 propane-2-sulfonyl chloride (step C) as a beige solid. MS
10 (ESI⁺): 312.1 [M+H]⁺.

Example 139:

Propane-2-sulfonic acid [1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

This compound was obtained in analogy to example 80 using 3,4-difluorobenzylbromide
15 (step A) and 3 propane-2-sulfonyl chloride (step C) as a light yellow solid. MS (ESI⁺): 382.3 [M+H]⁺.

Example 140:

Propane-2-sulfonic acid [1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide

20 This compound was obtained in analogy to example 32 using (2-methoxy-ethyl)-hydrazine (step A) (prepared as described in *J. Org. Chem.* 1984, 49, 336-42) and 3 propane-2-sulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 314.0 [M+H]⁺.

Example 141:

N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-C-cyclopropyl-methanesulfonamide

25 This compound was obtained in analogy to example 1 using 5-amino-1-benzyl-1,2-dihydro-indazol-3-one and cyclopropyl-methanesulfonyl chloride (step C) as a light brown solid. MS (ESI⁺): 358.4 ([M+H]⁺).

- 62 -

Example A

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

Ingredients*Per tablet*

Kernel:

Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg

Film Coat:

Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxyde (yellow)	0.8 mg	1.6 mg
Titan dioxide	0.8 mg	1.6 mg

- 5 The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesium stearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aqueous solution / suspension of the above mentioned film coat.

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg

- 5 The components are sieved and mixed and filled into capsules of size 2.

Example C

Injection solutions can have the following composition:

Compound of formula (I)	3.0 mg
Polyethylene Glycol 400	150.0 mg
Acetic Acid	q.s. ad pH 5.0
Water for injection solutions	ad 1.0 ml

- 10 The active ingredient is dissolved in a mixture of Polyethylene Glycol 400 and water for injection (part). The pH is adjusted to 5.0 by Acetic Acid. The volume is adjusted to 1.0 ml by addition of the residual amount of water. The solution is filtered, filled into vials using an appropriate overage and sterilized.

- 64 -

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

Capsule contents

Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg

Gelatin capsule

Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titan dioxide	0.4 mg
Iron oxide yellow	1.1 mg

- 5 The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E

Sachets containing the following ingredients can be manufactured in a conventional manner:


Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidon K 30	10.0 mg
Magnesiumstearate	10.0 mg
Flavoring additives	1.0 mg

- 5 The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidon in water. The granulate is mixed with magnesiumstearate and the flavouring additives and filled into sachets.

- 66 -

CLAIMS

1. Compounds of formula (I)



wherein

- 5 R^1 is hydrogen, lower-alkyl, aryl, or aryl-lower-alkyl;
- R^2 is aryl, aryl-lower-alkyl, heteroaryl, heteroaryl-lower-alkyl, cycloalkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$;
- 10 R^3 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^4 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- 15 R^5 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^6 is hydrogen or lower-alkyl;
- R^7 is aryl, heteroaryl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and cycloalkyl;
- 20 R^8 and R^9 , independently from each other, are selected from the group consisting of hydrogen and lower-alkyl;

and pharmaceutically acceptable salts thereof.

2. Compounds according to claim 1, wherein

R^1 is hydrogen, lower-alkyl, aryl, or aryl-lower-alkyl;

- 67 -

- R^2 is aryl, aryl-lower-alkyl, heteroaryl, heteroaryl-lower-alkyl, cycloalkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$;
- 5 R^3 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^4 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- 10 R^5 is hydrogen, halogen, lower-alkyl, fluoro-lower-alkyl, lower-alkoxy, or fluoro-lower-alkoxy;
- R^6 is hydrogen or lower-alkyl;
- R^7 is aryl, heteroaryl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen and lower-alkoxy;
- 15 R^8 and R^9 , independently from each other, are selected from the group consisting of hydrogen and lower-alkyl;
- and pharmaceutically acceptable salts thereof.
3. Compounds according to any of claims 1 to 2, wherein R^1 is hydrogen or lower-alkyl.
- 20 4. Compounds according to any of claims 1 to 3, wherein R^1 is hydrogen.
5. Compounds according to any of claims 1 to 4, wherein R^2 is aryl-lower-alkyl, heteroaryl-lower-alkyl, cycloalkyl-lower-alkyl, fluoro-lower-alkyl, or lower-alkyl, which lower-alkyl is optionally substituted with 1 to 3 substituents selected from the group consisting of OH, CN, halogen, lower-alkoxy and $C(O)NR^8R^9$, wherein R^8 and
- 25 R^9 are as defined in claim 1.
6. Compounds according to any of claims 1 to 5, wherein R^2 is aryl-lower-alkyl or heteroaryl-lower-alkyl.
7. Compounds according to any of claims 1 to 6, wherein R^2 is benzyl or pyridinylmethyl, wherein benzyl can optionally be substituted with 1 or 2 halogen.

- 68 -

8. Compounds according to any of claims 1 to 7, wherein R^2 is benzyl, 4-fluoro-benzyl, 4-chloro-benzyl, 3,4-difluoro-benzyl, or pyridin-2-ylmethyl.
9. Compounds according to any of claims 1 to 4, wherein R^2 is aryl.
10. Compounds according to any of claims 1 to 9, wherein R^3 is hydrogen.
- 5 11. Compounds according to any of claims 1 to 10, wherein R^4 is hydrogen or halogen.
12. Compounds according to any of claims 1 to 11, wherein R^4 is hydrogen, fluorine, or chlorine.
13. Compounds according to any of claims 1 to 12, wherein R^5 is hydrogen or halogen.
14. Compounds according to any of claims 1 to 13, wherein R^5 is hydrogen.
- 10 15. Compounds according to any of claims 1 to 14, wherein R^6 is hydrogen.
16. Compounds according to any of claims 1 to 15, wherein R^7 is lower-alkyl or aryl.
17. Compounds according to any of claims 1 to 16, wherein R^7 is lower-alkyl, phenyl or naphthyl, wherein phenyl can optionally be substituted with 1 to 3 substituents independently selected from the group consisting of lower-alkyl, fluoro-lower-alkyl, 15 lower-alkoxy, fluoro-lower-alkoxy, dioxo-lower-alkylene, halogen, cyano, phenoxy and 5-methyl-[1,3,4]-oxadiazol-2-yl.
18. Compounds according to any one of claims 1 to 17, wherein R^7 is phenyl substituted with 1 to 2 substituents selected from the group consisting of lower-alkyl, fluoro-lower-alkyl, halogen and cyano.
- 20 19. Compounds according to claim 18, wherein R^7 is 3-chloro-2-methyl-phenyl, 2,3-dichloro-phenyl, 3-chloro-4-fluoro-phenyl, 3-trifluoromethyl-4-fluoro-phenyl, 3-cyano-phenyl, 2,5-difluoro-phenyl, 3-chloro-2-fluoro-phenyl, 5-chloro-2-fluoro-phenyl, or 2-chloro-phenyl.
20. Compounds according to any of claims 1 to 15, wherein R^7 is fluoro-lower-alkyl or 25 lower-alkyl substituted with cyclopropyl.
21. Compounds according to claim 20, wherein R^7 is cyclopropyl-methyl.
22. Compounds according to any of claims 1 to 21 selected from the group consisting of:
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-

- benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-fluoro-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
5 benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-
benzenesulfonamide,
Naphthalene-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-methyl-
10 benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-6-methyl-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethyl-
15 benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluoro-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-propyl-benzenesulfonamide,
2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-
20 indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-2,5-dimethyl-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-3-trifluoromethyl-
benzenesulfonamide,
25 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-5-trifluoromethyl-
benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-phenoxy-benzenesulfonamide,
30 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-(5-methyl-[1,3,4]oxadiazol-2-
yl)-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-methoxy-benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxy-
35 benzenesulfonamide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3,5-bis-trifluoromethyl-
benzenesulfonamide,

- Propane-2-sulfonic acid (1-benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-N-dimethyl-
benzenesulfonamide,
N-(3-Oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
5 benzenesulfonamide,
2,3-Dichloro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
3-Chloro-4-fluoro-N-(3-oxo-1-propyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
10 3-Chloro-2-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
3-Chloro-4-methyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
4-Chloro-2,5-dimethyl-N-(3-oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-
15 benzenesulfonamide,
N-(3-Oxo-1-phenethyl-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
N-(1-Isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
20 3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-methyl-
benzenesulfonamide,
3-Chloro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-
benzenesulfonamide,
3-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
25 3-Difluoromethoxy-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-
benzenesulfonamide,
4-Cyano-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-benzenesulfonamide,
4-Fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
30 3-Chloro-4-fluoro-N-(1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
2,4-Dichloro-(1-isobutyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methyl-
benzenesulfonamide,
3-Chloro-N-(1-ethyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-methyl-
35 benzenesulfonamide,
3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-2-methyl-
benzenesulfonamide,

- 3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-
benzenesulfonamide,
(1-Cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-4-fluoro-3-trifluoromethyl-
benzenesulfonamide,
5 3-Chloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-
benzenesulfonamide,
2,4-Dichloro-(1-cyclopropylmethyl-3-oxo-2,3-dihydro-indazol-5-yl)-5-methyl-
benzenesulfonamide,
3-Chloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-
10 benzenesulfonamide,
4-Fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethyl-
benzenesulfonamide,
3-Chloro-4-fluoro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
benzenesulfonamide,
15 2,4-Dichloro-[1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-
benzenesulfonamide,
3-Chloro-2-methyl-N-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-1H-indazol-5-
yl]-benzenesulfonamide,
4-Fluoro-[3-oxo-1-(2,2,2-trifluoro-ethyl)-2,3-dihydro-indazol-5-yl]-3-
20 trifluoromethyl-benzenesulfonamide,
4-Fluoro-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-3-
trifluoromethyl-benzenesulfonamide,
2,4-Dichloro-6-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-
yl)-benzenesulfonamide,
25 3-Chloro-2-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-
benzenesulfonamide,
4-Fluoro-N-[1-(2-hydroxy-2-methyl-propyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
3-trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-4-fluoro-
30 benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-
benzenesulfonamide,
35 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichloro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-

- benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-chloro-3-
5 trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-difluoromethoxy-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-trifluoromethyl-
benzenesulfonamide,
10 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxy-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,5-difluoro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-
15 benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluoro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethyl-
benzenesulfonamide,
20 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-
benzenesulfonamide,
N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chloro-
benzenesulfonamide,
(1-Benzyl-4-chloro-3-oxo-2,3-dihydro-indazol-5-yl)-3-chloro-2-methyl-
25 benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichloro-
benzenesulfonamide,
30 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-dichloro-
benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluoro-
benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-
35 benzenesulfonamide,
N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-trifluoromethoxy-
benzenesulfonamide,

- 3-Chloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
- 3-Difluoromethoxy-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
- 5 4-Fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-trifluoromethyl-benzenesulfonamide,
- 2,3-Dihydro-benzo[1,4]dioxine-6-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide,
- 2,4-Dichloro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methyl-benzenesulfonamide,
- 10 5-Chloro-2,4-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
- 4-Chloro-2,5-difluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
- 15 3-Chloro-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-benzenesulfonamide,
- 3-Cyano-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
- Naphthalene-1-sulfonic acid [1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-amide,
- 20 3-Chloro-4-fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
- 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-benzenesulfonamide,
- 25 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-benzenesulfonamide,
- [1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-4-fluoro-3-trifluoromethyl-benzenesulfonamide,
- 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
- 30 [1-(3,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-3-trifluoromethoxy-benzenesulfonamide,
- 2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-5-methyl-benzenesulfonamide,
- 35 3,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-benzenesulfonamide,
- 4,5-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-fluoro-

- benzenesulfonamide,
2,4,5-Trichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
benzenesulfonamide,
2,3-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
5 benzenesulfonamide,
2,4-Dichloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-6-methyl-
benzenesulfonamide,
3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-
fluoro-benzenesulfonamide,
10 N-[1-(2,4-Difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,
3-Chloro-N-[1-(2,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-
methyl-benzenesulfonamide,
N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
15 trifluoromethyl-benzenesulfonamide,
3-Chloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-
benzenesulfonamide,
2,3-Dichloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
benzenesulfonamide,
20 Naphthalene-1-sulfonic acid [1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-
5-yl]-amide,
3-Chloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-
methyl-benzenesulfonamide,
Naphthalene-1-sulfonic acid [1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-
25 indazol-5-yl]-amide,
2,3-Dichloro-N-[1-(2-chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-
yl]-benzenesulfonamide,
N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,
30 N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-
difluoromethoxy-benzenesulfonamide,
N-[1-(2-Chloro-4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-
trifluoromethoxy-benzenesulfonamide,
3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-
35 benzenesulfonamide,
N-[1-(2-Cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
trifluoromethyl-benzenesulfonamide,


- 2,4-Dichloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-6-methylbenzenesulfonamide,
 3-Chloro-N-[1-(2-cyano-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluorobenzenesulfonamide,
 5 N-[6-Chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-trifluoromethylbenzenesulfonamide,
 5-Chloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-fluorobenzenesulfonamide,
 2,3-Dichloro-N-[6-chloro-1-(2-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-benzenesulfonamide,
 10 3-Chloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-2-methylbenzenesulfonamide,
 N-(6-Chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-yl)-4-fluoro-3-trifluoromethylbenzenesulfonamide,
 15 2-[5-(3-Chloro-2-methylbenzenesulfonylamino)-3-oxo-2,3-dihydro-indazol-1-yl]-methylacetamide,
 2,4-Dichloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-6-methylbenzenesulfonamide, and
 3-Chloro-[1-(3,4-difluoro-benzyl)-2-methyl-3-oxo-2,3-dihydro-indazol-5-yl]-2-methylbenzenesulfonamide,
 20 and pharmaceutically acceptable salts thereof.
23. Compounds according to any one of claims 1 to 22 selected from the group consisting of:
- N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methylbenzenesulfonamide,
 25 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,5-difluorobenzenesulfonamide,
 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluorobenzenesulfonamide,
 30 N-(1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2-chlorobenzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-fluorobenzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,3-dichlorobenzenesulfonamide,
 35 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-5-chloro-2-fluoro-

- 76 -


- benzenesulfonamide,
 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-chloro-2-methyl-
 benzenesulfonamide,
 3-Cyano-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-
 5 benzenesulfonamide,
 2,3-Dichloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
 benzenesulfonamide,
 3-Chloro-N-[1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-2-methyl-
 benzenesulfonamide,
 10 4-Fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-3-
 trifluoromethyl-benzenesulfonamide,
 3-Chloro-4-fluoro-N-[1-(4-fluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
 benzenesulfonamide,
 3-Chloro-[1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-indazol-5-yl]-2-methyl-
 15 benzenesulfonamide,
 N-[1-(4-Chloro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-4-fluoro-3-
 trifluoromethyl-benzenesulfonamide, and
 3-Chloro-2-methyl-N-(3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-
 yl)-benzenesulfonamide,
 20 and pharmaceutically acceptable salts thereof.
24. Compounds according to any one of claims 1 to 21 selected from the group
 consisting of:
 2,3-Dichloro-N-(6-chloro-3-oxo-1-pyridin-2-ylmethyl-2,3-dihydro-1H-indazol-5-
 yl)-benzenesulfonamide,
 25 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-2,4-difluoro-
 benzenesulfonamide,
 2,4-Difluoro-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-
 benzenesulfonamide,
 3-Chloro-2-fluoro-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-
 30 benzenesulfonamide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-3-cyano-
 benzenesulfonamide,
 Propane-2-sulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-
 amide,
 35 3-Cyano-N-(3-oxo-1-phenyl-2,3-dihydro-1H-indazol-5-yl)-benzenesulfonamide,
 2,2,2-Trifluoro-ethanesulfonic acid (1-benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-

- 77 -

- indazol-5-yl)-amide,
 Propane-2-sulfonic acid (1-Benzyl-6-chloro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-
 amide,
 Propane-2-sulfonic acid [6-chloro-1-(4-chloro-benzyl)-3-oxo-2,3-dihydro-1H-
 5 indazol-5-yl]-amide,
 N-(1-Benzyl-6-fluoro-3-oxo-2,3-dihydro-1H-indazol-5-yl)-trifluoro-
 methanesulfonamide,
 Propane-2-sulfonic acid (1-isobutyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-amide,
 Propane-2-sulfonic acid [1-(3,4-difluoro-benzyl)-3-oxo-2,3-dihydro-1H-indazol-5-
 10 yl]-amide,
 Propane-2-sulfonic acid [1-(2-methoxy-ethyl)-3-oxo-2,3-dihydro-1H-indazol-5-yl]-
 amide, and
 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-C-cyclopropyl-
 methanesulfonamide
 15 and pharmaceutically acceptable salts thereof.
25. The compound according to any one of claims 1 to 21 which is:
 N-(1-Benzyl-3-oxo-2,3-dihydro-1H-indazol-5-yl)-C-cyclopropyl-
 methanesulfonamide
 and pharmaceutically acceptable salts thereof.
- 20 26. A process for the preparation of compounds according to any of claims 1 to 25
 comprising the reaction of a compound of formula (II)



with a compound of formula (III)



- 25 wherein R¹ to R⁷ are as defined in any of claims 1 to 25.

- 78 -

27. Compounds according to any of claims 1 to 25, when prepared by a process as defined in claim 26.
28. Pharmaceutical compositions comprising a compound according to any of claims 1 to 19 and a pharmaceutically acceptable carrier and/or adjuvant.
- 5 29. Compounds according to any of claims 1 to 25 for use as therapeutic active substances.
30. Compounds according to any of claims 1 to 25 for use as therapeutic active substances for the treatment and/or prophylaxis of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1.
- 10 31. A method for the treatment and/or prophylaxis of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1, particularly for the treatment and/or prophylaxis of metabolic disorders, obesity, dyslipidemiae, hypertension and/or diabetes, particularly diabetes type II, which method comprises administering an effective amount of a compound as defined in
15 any of claims 1 to 25.
32. The use of compounds according to any of claims 1 to 25 for the therapeutic and/or prophylactic treatment of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1.
33. The use of compounds according to any of claims 1 to 25 for the therapeutic and/or
20 prophylactic treatment of metabolic disorders, obesity, dyslipidemiae, hypertension and/or diabetes.
34. The use of compounds according to any of claims 1 to 25 for the preparation of medicaments for the treatment and prophylaxis of diseases which are caused by disorders associated with the enzyme 11beta-hydroxysteroid dehydrogenase 1.
- 25 35. The use of compounds according to any of claims 1 to 25 for the preparation of medicaments for the treatment and prophylaxis of metabolic disorders, obesity, dyslipidemiae, hypertension and/or diabetes.
36. The invention as hereinbefore described.
