VEGF TRAPS AND THERAPEUTIC USES THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] The invention encompasses fusion polypeptides capable of binding vascular endothelial cell growth factor (VEGF), VEGF family members, and splice variants with specifically desirable characteristics, as well as therapeutic methods of use.

BRIEF SUMMARY OF THE INVENTION

[0002] In a first aspect, the invention features an isolated nucleic acid molecule encoding a fusion polypeptide comprising receptor components $(R1R2)_X$ and/or $(R1R3)_Y$, wherein R1 is vascular endothelial cell growth factor (VEGF) receptor component Ig domain 2 of Flt-1 (Flt1D2), R2 is VEGF receptor component Ig domain 3 of Flk-1 (Flk1D3), and R3 is VEGF receptor component Ig domain 3 of Flt-4 (Flt1D3 or R3), and wherein $X \geq 1$ and $Y \geq 1$.

[0003] In a related second aspect, the invention features a monomeric VEGF trap or fusion polypeptide comprising VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$ wherein $X \geq 1$, $Y \geq 1$, and R1, R2, and R3 are as defined above. The VEGF receptor components R1, R2, and R3, may be connected directly to each other or connected via one or more spacer sequences. In one specific embodiment, the monomeric VEGF trap is $(R1R2)_X$, were X=2. In a more specific embodiment, the monomeric VEGF trap is SEQ ID NO:24, or a functionally equivalent amino acid variant thereof. The invention encompasses a monomeric VEGF trap consisting essentially of VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$, and functionally equivalent amino acid variants thereof.

[0004] In a third aspect, the invention features an isolated nucleic acid molecule encoding a fusion polypeptide comprising VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$, and a fusion partner (FP) component selected from the group consisting of a multimerizing component (MC), a serum protein, or a molecule capable of binding a serum protein. In a preferred embodiment, FP is a multimerizing component (MC) capable of interacting with a multimerizing component on another fusion polypeptide to form a multimeric structure, e.g., a dimer or trimer. Most preferably, the MC is selected from the group consisting of (i) a multimerizing component comprising a cleavable region (C-region), (ii) a truncated multimerizing component, (iii) an amino acid sequence between 1 to about 200 amino acids in length having at least one cysteine residue, (iv) a leucine zipper, (v) a helix loop motif, (vi) a coil-coil motif, and (vii) an immunoglobulin domain. Further encompassed are fusion polypeptides consisting essentially of $(R1R2)_X$ and/or $(R1R3)_Y$, and FP. In a preferred embodiment, the fusion polypeptide consists essentially of $(R1R2)_X$ and MC.

[0005] In a fourth aspect, the invention features a fusion polypeptide comprising VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$, and FP, as described above. The receptor components may be arranged in different orders, for example, $(R1R2)_X$-FP; $(R1R2)_X$-FP-$(R1R2)_X$; FP-$(R2R1)_X$, etc. The components of the fusion polypeptide may be connected directly to each other, or connected via a spacer sequence.

[0006] In a fifth aspect, the invention features a VEGF trap, comprising a multimer of two or more fusion polypeptides consisting of VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$, and FP, wherein the FP component is a multimerizing component (MC) comprising a C-region. The C-region may be naturally occurring or artificial, and may occur at any point within the multimerizing component, and functions to allow cleavage of a parent MC to a truncated MC. A VEGF trap composed of two or more fusion polypeptides having at least one truncated MC is termed a "truncated mini-trap."

[0007] The C-region may be created in MC by insertion, deletion, or mutation, such that an enzymatically or chemically cleavable site is created. The C-region may be created in any MC and at any position within the MC; preferably, the C-region is created in a full length Fc domain, or a fragment thereof, or a C_H3 domain. The C-region may be a site cleavable by an enzyme, such as, thrombin, ficin, pepsin, matrilysin, or prolidase or cleavable chemically by, for example, formic acid or $CuCl_2$.

[0008] In a sixth related aspect, the invention features a truncated VEGF mini-trap which is a multimeric protein comprising two or more fusion polypeptides consisting of $(R1R2)_X$ and/or $(R1R3)_Y$ and a multimerizing component which is a truncated by cleavage from a parent MC comprising a C-region (tMC).

[0009] In a seventh aspect, the invention features a fusion polypeptide consisting of VEGF receptor components $(R1R2)_X$ and/or $(R1R3)_Y$ and a MC, wherein the MC is an amino acid sequence between 1 to about 200 amino acids in length comprising at least one cysteine residue, wherein the at least one cysteine residue is capable of forming a disulfide bond with a cysteine residue present in the MC of another fusion polypeptide (cMC). In a preferred embodiment, cMC is an amino acid sequence between 1-50 amino acids in length comprising at least one cysteine residue. In a more preferred embodiment, cMC is an amino acid sequence between 1-15 amino acids in length comprising at least one amino acid. In an even more preferred embodiment, cMC is an amino acid sequence between 1-10 amino acids in length comprising 1-2 cysteine residues. One exemplification of this embodiment of the invention is shown in SEQ ID NO:27 having a signal sequence (1-26) followed by R1 (27-129) and R2 (130-231) components, followed by a nine amino acid sequence ending in a cysteine residue. In another embodiment, shown in SEQ ID NO:28, a signal sequence (1-26) is followed by R1 (27-129) and R2 (130-231) components, followed by a six amino acid sequence ending in a cysteine residue.

[0010] In an eighth aspect, the invention features a VEGF mini-trap, comprising a multimer of two or more fusion polypeptides consisting of $(R1R2)_X$ and/or $(R1R3)_Y$ and a cMC. In a more specific embodiment, the mini-trap is a dimer. One exemplification of this embodiment of the mini-trap of the invention is a dimer of the fusion polypeptide shown in SEQ ID NO:2, wherein each fusion polypeptide (R1R2-cMC) has a molecular weight of 23.0 kD and a pI of 9.22.

[0011] In another embodiment, cMC is 4 amino acids in length consisting of two cysteine residues, for example, XCXC (SEQ ID NO:3). In one exemplification of this embodiment of the invention, the mini-trap consists of the VEGF receptor components of the invention, and a cMC consisting of ACGC (SEQ ID NO:4). One exemplification of this embodiment of the mini-trap of the invention is

a dimer of the fusion polypeptide shown in SEQ ID NO:5, wherein each monomer has a molecular weight of 23.2 kD and a pI of 9.22. Another exemplification of this embodiment of the invention is shown in SEQ ID NO:26 having a signal sequence (1-26) followed by R1 (27-129) and R2 (130-231) components, followed by a nine amino acid sequence ending in CPPC.

[0012] In all embodiments of the VEGF trap of the invention (including truncated VEGF mini-trap, VEGF mini-traps, and monomeric VEGF mini-traps), a signal sequence (S) may be included at the beginning (or N-terminus) of the fusion polypeptide of the invention. The signal sequence may be native to the cell, recombinant, or synthetic. When a signal sequence is attached to the N-terminus of a first receptor component, thus a fusion polypeptide may be designated as, for example, S-$(R1R2)_X$.

[0013] The components of the fusion polypeptide may be connected directly to each other or be connected via spacers. In specific embodiments, one or more receptor and/or fusion partner components of the fusion polypeptide are connected directly to each other without spacers. In other embodiments, one or more receptor and/or fusion partner components are connected with spacers.

[0014] The invention encompasses vectors comprising the nucleic acid molecules of the invention, including expression vectors comprising the nucleic acid molecule operatively linked to an expression control sequence. The invention further encompasses host-vector systems for the production of a fusion polypeptide which comprise the expression vector, in a suitable host cell; host-vector systems wherein the suitable host cell is a bacterial, yeast, insect, mammalian cell; an *E. coli* cell, or a COS or CHO cell. Additional encompassed are VEGF traps of the invention modified by acetylation or pegylation. Methods for acetylating or pegylating a protein are well known in the art.

[0015] In a related ninth aspect, the invention features a method of producing a VEGF trap of the invention, comprising culturing a host cell transfected with a vector comprising a nucleic acid sequence of the invention, under conditions suitable for expression of the protein from the host cell, and recovering the fusion polypeptides so produced.

[0016] The VEGF traps of the invention are therapeutically useful for treating any disease or condition which is improved, ameliorated, or inhibited by removal, inhibition, or reduction of VEGF. A non-exhaustive list of specific conditions improved by inhibition or reduction of VEGF include, for example, undesirable plasma leakage or vascular permeability, undesirable blood vessel growth, e.g., such as in a tumor, edema associated with inflammatory disorders such as psoriasis or arthritis, including rheumatoid arthritis; asthma; generalized edema associated with burns; ascites and pleural effusion associated with tumors, inflammation or trauma; chronic airway inflammation; asthma; capillary leak syndrome; sepsis; kidney disease associated with increased leakage of protein; pancreatic ductal adenocarcinoma (PDAC) and eye disorders such as age related macular degeneration and diabetic retinopathy. The VEGF mini-trap is particularly useful in treatment of eye disorders, and as an adjuvant to eye surgeries, including glaucoma surgery; and the treatment of intra-ocular tumors, such as for example, uveal melanoma, retinoblastoma, via intravitreal delivery.

[0017] Accordingly, in a tenth aspect, the invention features a therapeutic method for the treatment of a VEGF-related disease or condition, comprising administering a VEGF trap of the invention to a subject suffering from a VEGF-related disease or condition. Although any mammal

can be treated by the therapeutic methods of the invention, the subject is preferably a human patient suffering from or at risk of suffering from a condition or disease which can be improved, ameliorated, inhibited or treated with a VEGF trap.

[0018] In a eleventh aspect, the invention further features diagnostic and prognostic methods, as well as kits for detecting, quantitating, and/or monitoring VEGF with the mini-traps of the invention.

[0019] In a twelfth aspect, the invention features pharmaceutical compositions comprising a VEGF trap of the invention with a pharmaceutically acceptable carrier. Such pharmaceutical compositions may comprise a dimeric fusion polypeptide trap, or nucleic acids encoding the fusion polypeptide. The mini-traps of the invention find specific uses in conditions in which a VEGF trap with reduced serum half life (e.g., faster clearance), and/or increased tissue penetration due to smaller size is desirable. Specific applications for the VEGF mini-trap include, for example, diseases where local administration to a specific tissue or cell is desirable. Examples of such a condition or disease are ocular diseases of the eye.

[0020] Other objects and advantages will become apparent from a review of the ensuing detailed description.

DETAILED DESCRIPTION OF THE INVENTION

[0021] Before the present methods are described, it is to be understood that this invention is not limited to particular methods, and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only the appended claims.

[0022] As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, a reference to "a method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure and so forth.

[0023] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to describe the methods and/or materials in connection with which the publications are cited.

General Description

[0024] The invention encompasses a VEGF trap capable of binding and inhibiting VEGF activity which is a monomer or multimer of one or more fusion polypeptides. The molecules of the invention bind and inhibit the biological action of VEGF and/or the physiological reaction or response. For a description of VEGF-receptor-based antagonist VEGF traps Flt1D2.Flk1D3.FcΔC1(a) (SEQ ID NOs:7-8) and VEGFR1R2-FcΔC1(a) (SEQ ID NOs:9-10), see PCT WO/0075319, the contents of which is incorporated in its entirety herein by reference.

[0025] The mini-trap of the invention is smaller than the full sized trap, e.g., about 50 - 60 kD

versus 120 kD of the parent trap, and include monomeric traps consisting essentially of VEGF receptor domains $(R1R2)_X$, $(R1R3)_Y$, or combinations thereof, traps generated by cleavage of a portion of a parent multimerized trap having a fusion partner component which is a multimerizing component (MC) containing a cleavage region (C-region); or by attaching a cysteine residue or amino acid sequence containing one or more cysteine residues to or between receptor component domains. In specific embodiments, the mini-trap of the invention is less than about 60 kD as measured by SDS-PAGE analysis; more preferably, about 50 kD; even more preferably about 20-30 kD; or is about 25 kD and capable of binding VEGF with an affinity comparable to a full-sized parent trap described in PCT/US00/14142.

Nucleic Acid Constructs and Expression

[0026] The present invention provides for the construction of nucleic acid molecules encoding fusion polypeptides capable of binding VEGF alone or multimerized VEGF traps. The nucleic acid molecules of the invention may encode wild-type R1, R2, and/or R3 receptor components, or functionally equivalent variants thereof. Amino acid sequence variants of the R1, R2 and/or R3 receptor components of the traps of the invention may also be prepared by creating mutations in the encoding nucleic acid molecules. Such variants include, for example, deletions from, or insertions or substitutions of, amino acid residues within the amino acid sequence of R1, R2 and/or R3. Any combination of deletion, insertion, and substitution may be made to arrive at a final construct, provided that the final construct possesses the ability to bind and inhibit VEGF.

[0027] These nucleic acid molecules are inserted into a vector that is able to express the fusion polypeptides when introduced into an appropriate host cell. Appropriate host cells include, but are not limited to, bacterial, yeast, insect, and mammalian cells. Any of the methods known to one skilled in the art for the insertion of DNA fragments into a vector may be used to construct expression vectors encoding the fusion polypeptides of the invention under control of transcriptional/translational control signals.

[0028] Expression of the nucleic acid molecules of the invention may be regulated by a second nucleic acid sequence so that the molecule is expressed in a host transformed with the recombinant DNA molecule. For example, expression may be controlled by any promoter/enhancer element known in the art. Promoters which may be used to control expression of the chimeric polypeptide molecules include, but are not limited to, a long terminal repeat (Squinto et al. (1991) Cell 65:1-20); SV40 early promoter region, CMV, M-MuLV, thymidine kinase promoter, the regulatory sequences of the metallothionine gene; prokaryotic expression vectors such as the b-lactamase promoter, or the tac promoter (see also Scientific American (1980) 242:74-94); promoter elements from yeast or other fungi such as Gal 4 promoter, ADH, PGK, alkaline phosphatase, and tissue-specific transcriptional control regions derived from genes such as elastase I.

[0029] Expression vectors capable of being replicated in a bacterial or eukaryotic host comprising the nucleic acid molecules of the invention are used to transfect the host and thereby direct expression of such nucleic acids to produce the fusion polypeptides of the invention, which form traps capable of binding to VEGF. Transfected cells may transiently or, preferably, constitutively

and permanently express the VEGF traps of the invention.

[0030] The traps of the invention may be purified by any technique which allows for the subsequent formation of a stable, biologically active trap. For example, and not by way of limitation, the factors may be recovered from cells either as soluble proteins or as inclusion bodies, from which they may be extracted quantitatively by 8M guanidinium hydrochloride and dialysis (see, for example, US Patent No. 5,663,304). In order to further purify the factors, conventional ion exchange chromatography, hydrophobic interaction chromatography, reverse phase chromatography or gel filtration may be used.

VEGF Receptor Components

[0031] The VEGF receptor components of the VEGF mini trap consist of the Ig domain 2 of Flt-1 (Flt1D2) (R1), the Ig domain 3 of Flk-1 (Flk1D3) (R2) (together, R1R2), and/or R1 and Ig domain 3 of Flt-4 (Flt1D3) (R3) (together, R1R3). The term "Ig domain" of Flt-1, Flt-4, or Flk-1 is intended to encompass not only the complete wild-type domain, but also insertional, deletional, and/or substitutional variants thereof which substantially retain the functional characteristics of the intact domain. It will be readily apparent to one of skill in the art that numerous variants of the above Ig domains can be obtained which will retains substantially the same functional characteristics as the wild-type domain.

[0032] The term "functional equivalents" when used in reference to R1, R2, or R3, is intended to encompass an R1, R2, or R3 domain with at least one alteration, e.g., a deletion, addition, and/or substitution, which retains substantially the same functional characteristics as does the wild type R1, R2, or R3 domain, that is, a substantially equivalent binding to VEGF. It will be appreciated that various amino acid substitutions can be made in R1, R2, or R3 without departing from the spirit of the invention with respect to the ability of these receptor components to bind and inactivate VEGF. The functional characteristics of the traps of the invention may be determined by any suitable screening assay known to the art for measuring the desired characteristic. Examples of such assays are described in the experimental section below which allow determination of binding characteristics of the traps for VEGF (Kd), as well as their half-life of dissociation of the trap-ligand complex ($T_{1/2}$). Other assays, for example, a change in the ability to specifically bind to VEGF can be measured by a competition-type VEGF binding assay. Modifications of protein properties such as thermal stability, hydrophobicity, susceptibility to proteolytic degradation, or tendency to aggregate may be measured by methods known to those of skill in the art.

[0033] The components of the fusion polypeptide may be connected directly to each other or be connected via spacers. Generally, the term "spacer" (or linker) means one or more molecules, e.g., nucleic acids or amino acids, or non-peptide moieties, such as polyethylene glycol, which may be inserted between one or more component domains. For example, spacer sequences may be used to provide a desirable site of interest between components for ease of manipulation. A spacer may also be provided to enhance expression of the fusion polypeptide from a host cell, to decrease steric hindrance such that the component may assume its optimal tertiary structure and/or interact appropriately with its target molecule. For spacers and methods of identifying desirable spacers, see,

for example, George et al. (2003) Protein Engineering 15:871-879, herein specifically incorporated by reference. A spacer sequence may include one or more amino acids naturally connected to a receptor component, or may be an added sequence used to enhance expression of the fusion polypeptides, provide specifically desired sites of interest, allow component domains to form optimal tertiary structures and/or to enhance the interaction of a component with its target molecule. In one embodiment, the spacer comprises one or more peptide sequences between one or more components which is (are) between 1-100 amino acids, preferably 1-25.

[0034] In the most specific embodiments, R1 is amino acids 27-126 of SEQ ID NO:8, or 1-126 of SEQ ID NO:8 (including the signal sequence 1-26); or amino acids 27-129 of SEQ ID NO:10, or 1-129 of SEQ ID NO:10 (including the signal sequence at 1-26). In the most specific embodiments, R2 is amino acids 127-228 of SEQ ID NO:8, or amino acids 130-231 of SEQ ID NO:10. In the most specific embodiments, R3 is amino acids 127-225 of SEQ ID NO: 13 (without a signal sequence). When, for example, R2 is placed at the N-terminus of the fusion polypeptide, a signal sequence may desirably precede the receptor component. The receptor component(s) attached to the multimerizing component may further comprise a spacer component, for example, the GPG sequence of amino acids 229-231 of SEQ ID NO:7.

Fusion Partner and Multimerizing Components

[0035] The fusion partner is any component that enhances the functionality of the fusion polypeptide. Thus, for example, an fusion partner may enhance the biological activity of the fusion polypeptide, aid in its production and/or recovery, or enhance a pharmacological property or the pharmacokinetic profile of the fusion polypeptide by, for example, enhancing its serum half-life, tissue penetrability, lack of immungenicity, or stability. In preferred embodiments, the fusion partner is selected from the group consisting of a multimerizing component, a serum protein, or a molecule capable of binding a serum protein.

[0036] When the fusion partner is a serum protein or fragment thereof, it is selected from the group consisting of α-1-microglobulin, AGP-1, orosomuciod, α-1-acid glycoprotein, vitamin D binding protein (DBP), hemopexin, human serum albumin (hSA), transferrin, ferritin, afamin, haptoglobin, α-fetoprotein thyroglobulin, α-2-HS-glycoprotein, β-2-glycoprotein, hyaluronan-binding protein, syntaxin, C1R, C1q a chain, galectin3-Mac2 binding protein, fibrinogen, polymeric Ig receptor (PIGR), α-2-macroglobulin, urea transport protein, haptoglobin, IGFBPs, macrophage scavenger receptors, fibronectin, giantin, Fc, α-1-antichyromotrypsin, α-1-antitrypsin, antithrombin III, apolipoprotein A-I, apolipoprotein B, β-2-microglobulin, ceruloplasmin, complement component C3 or C4, CI esterase inhibitor, C-reactive protein, cystatin C, and protein C. In a more specific embodiment, fusion partner is selected from the group consisting of α-1-microglobulin, AGP-1, orosomuciod, α-1-acid glycoprotein, vitamin D binding protein (DBP), hemopexin, human serum albumin (hSA), afamin, and haptoglobin. The inclusion of a fusion partner component may extend the serum half-life of the fusion polypeptide of the invention when desired. See, for example, US Patent Nos. 6,423,512, 5,876,969, 6,593,295, and 6,548,653, herein specifically incorporated by

reference in their entirety, for examples of serum albumin fusion polypeptides. hSA is widely distributed throughout the body, particularly in the intestinal and blood components, and has an important role in the maintenance of osmolarity and plasma volume. It is slowly cleared in the liver, and typically has an *in vivo* half-life of 14-20 days in humans (Waldmann et al. (1977) Albumin, Structure Function and Uses; Pergamon Press; pp. 255-275).

[0037] When a fusion partner is a molecule capable of binding a serum protein, the molecule may be a synthetic small molecule, a lipid or liposome, a nucleic acid, including a synthetic nucleic acid such as an aptomer, a peptide, or an oligosaccharide. The molecule may further be a protein, such as, for example, FcγR1, FcγR2, FcγR3, polymeric Ig receptor (PIGR), ScFv, and other antibody fragments specific for a serum protein.

[0038] When the fusion partner is a multimerizing component (MC), it is any natural or synthetic sequence capable of interacting with another MC to form a higher order structure, e.g., a dimer, a trimer, etc. Suitable MCs may include a leucine zipper, including leucine zipper domains derived from c-jun or c-fos; sequences derived from the constant regions of kappa or lambda light chains; synthetic sequences such as helix-loop-helix motifs (Müller et al. (1998) FEBS Lett. 432:45-49), coil-coil motifs, etc., or other generally accepted multimerizing domains known to the art. In some embodiments, the fusion component comprises an immunoglobulin-derived domain from, for example, human IgG, IgM or IgA. In specific embodiments, the immunoglobulin-derived domain may be selected from the group consisting of the Fc domain of IgG, the heavy chain of IgG, and the light chain of IgG. The Fc domain of IgG may be selected from the isotypes IgG1, IgG2, IgG3, and IgG4, as well as any allotype within each isotype group. In one example of the VEGF trap of the invention, the multimerizing component is an IgG4 Fc domain (SEQ ID NO:29).

Generation of Truncated VEGF Mini-Traps

[0039] In one embodiment of the trap of the invention, a truncated VEGF mini-trap comprising two or more fusion polypeptides of the invention, is generated by subjecting a parent trap having C-region-containing MCs to conditions under which one or more of the C-region-containing MCs is (are) cleaved. The resulting truncated mini-trap may be a full and partial cleavage product of a parent trap.

[0040] The C-region-containing MC may be any MC capable of interacting with another MC to form a higher order structure, e.g., a dimer or a trimer. The C-region may be created within an MC at any desired location. In light of the guidance provided in the examples below, one of skill in the art would be able to select a desired site for creation of a C-region based on the desired properties of the resulting truncated traps, e.g., molecular weight, monomeric or dimeric, etc.

[0041] In a specific embodiment, the C-region is a thrombin cleavage site (LVPRGS) (SEQID NO:6) inserted into an FcΔC1 domain following the N-terminal CPPC sequence (SEQ ID NO:1). In this embodiment, a full-sized parent VEGF trap construct is expressed in a cell as an Fc-tagged protein, thus allowing capture and purification by, for example, a Protein A column. Following formation of a dimer and covalent bonding between one or both of the cysteine residues of the CPPC sequence

(SEQ ID NO:1), the dimer is exposed to thrombin under conditions which cleave one or both of the FcΔC1 domains such that truncated dimeric mini-traps are generated, having a molecular weight of approximately 50 kD – 90 kD, and has an affinity for VEGF comparable to that of the parent trap. The conditions of cleavage may be controlled by one of skill in the art to favor formation of the partial cleavage product or the fully cleaved product, the choice of cleavage conditions selected by desire for a particular product having specific properties such as molecular weight.

[0042] In a specific embodiment, the C-region is a thrombin cleavage site (LVPRGS) (SEQID NO:6) inserted into an FcΔC1 domain N-terminal to the CPPC sequence (SEQ ID NO:1). Following formation of a dimer and covalent bonding between one or both of the cysteine residues of the CPPC sequence (SEQ ID NO:1), the dimer is exposed to thrombin under conditions in which one or both of the FcΔC1 domain occur and truncated monomeric mini-traps are generated. The monomeric truncated mini-trap thus generated comprises a receptor component, and a small fragment of the Fc, and is approximately 25 kD in size and exhibits a reduced affinity for VEGF relative to the truncated dimeric trap and the full length parent trap. A similar monomeric trap produced as a recombinant protein has been shown to have a K_D of about 1 nM.

Generation of VEGF Mini-Traps

[0043] In one embodiment, the invention features VEGF mini-traps having one or more receptor component domains $(R1R2)_X$ and/or $R1R3)_Y$, wherein $X \geq 1$, $Y \geq 1$, and R1, R2, and R3 are as defined above, and optionally, a fusion partner which is preferably a MC domain which is an amino acid sequence between 1 to about 200 amino acids in length comprising at least one cysteine residue, wherein the at least one cysteine residue is capable of forming a disulfide bond with a cysteine residue present in the MC of another fusion polypeptide (cMC). The cMC may occur at the N-terminus or C-terminus of a fusion polypeptide, or between two receptor component domains. In one specific embodiment, cysteine is added to the C-terminus of a VEGF receptor component, e.g., $R1R2_C$, which allows the fusion polypeptide to form covalent dimers through formation of a covalent disulfide bond between the cysteine residue at the C-terminus of one fusion polypeptide and the cysteine residue at the C-terminus of another fusion polypeptide. In this exemplification, the mini-trap is a dimer of the fusion polypeptide shown in SEQ ID NO:2, wherein each fusion polypeptide (R1R2-cMC or $R1R2_C$) has a molecular weight of about 23.0 kD.

[0044] In another embodiment, the cMC is a sequence of 4 amino acids (XXXX) (SEQ ID NO:11) wherein X is any amino acid and the sequence comprises at least one cysteine residue. In a specific embodiment, the cMC is added to the C-terminus of a receptor component domain. In a more specific embodiment, the 4 amino acid sequence is ACGC (SEQ ID NO:4) and the cMC forms two disulfide bonds with the cysteine residues present in a second fusion polypeptide. As shown below (Table 2), both the exemplified mini-traps exhibit an affinity for VEGF comparable to the parent trap.

Therapetic Uses

[0045] The VEGF mini-traps of the invention are therapeutically useful for treating any disease or

condition which is improved, ameliorated, inhibited or prevented by removal, inhibition, or reduction of VEGF. A non-exhaustive list of specific conditions improved by inhibition or reduction of VEGF include, clinical conditions that are characterized by excessive vascular endothelial cell proliferation, vascular permeability, edema or inflammation such as brain edema associated with injury, stroke or tumor; edema associated with inflammatory disorders such as psoriasis or arthritis, including rheumatoid arthritis; asthma; generalized edema associated with burns; ascites and pleural effusion associated with tumors, inflammation or trauma; chronic airway inflammation; capillary leak syndrome; sepsis; kidney disease associated with increased leakage of protein; and eye disorders such as age related macular degeneration and diabetic retinopathy.

[0046] The compositions of the invention are therapeutically useful for treating a wide variety of diseases associated with increased VEGF levels. For example, exaggerated Th2 inflammation and airway remodeling are characteristic in the pathogenesis of asthma (see, for example, Elias et al. (1999) J. Clin. Invest. 104:1001-6). Elevated VEGF levels have been detected in tissues and biologic samples from patients with asthma, which correlate directly with disease activity (Lee et al. (2001) J. Allergy Clin. Immunol. 107:1106-1108) and inversely with airway caliber and airway responsiveness. Further, VEGF has been postulated to contribute to asthmatic tissue edema.

[0047] Another disease associated with increased VEGF is pancreatic ductal adenocarcinoma (PDAC). This malignancy often exhibits enhanced foci of endothelial cell proliferation and frequently overexpresses VEGF (Ferrara (1999) J. Mol. Med. 77:527-543). PDAC is responsible for over 20% of deaths due to gastrointestinal malignancies, making it the fourth most common cause of cancer-related mortality in the U.S. and other industrialized countries. Experimental evidence supports an important role for VEGF in pancreatic cancer, thus a VEGF inhibitor has promise as a therapeutic to attenuate intrapancreatic tumor growth and regional and distal metastasis.

[0048] A smaller, non-glycosylated mini-trap expressed in *E. coli* (Example 4), a glycosylated mini-trap expressed in CHO cells (Example 5), or a receptor-based monomeric trap (Example 6) has optimized characteristics for local/intra-vitreal delivery, ie. a shorter serum half life for faster clearance and minimizing unwanted systemic exposure. In addition due to its smaller size, the mini-trap has the ability to penetrate through the inner-limiting membrane (ILM) in the eye, and diffuse through the vitreous to the retina/retinal pigment epithelial (RPE) layer which will help to treat retinal disease. Additionally, the mini-trap can be used for local administration for the treatment of ocular disease such as choroidal neovascularization, diabetic macular edema, proliferative diabetic retinopathy, corneal neovascularization/transplant rejection. Still further, the mini-trap can be used in any situation where transient (short-term) blocking of VEGF is required, e.g., to avoid chronic exposure to VEGF blockade, such as, for example, in the treatment of psoriasis.

[0049] A serious problem leading to failure following glaucoma surgery is early inflammation and angiogenesis, as well as too aggressive wound healing. Accordingly, the VEGF traps of the invention may be usefully employed is as an adjuvant to glaucoma surgery to prevent early hem- and lymphangiogenesis and macrophage recruitement to the filterig bleb after glaucoma surgery, and improve surgical outcome.

Combination Therapies

[0050] In numerous embodiments, a VEGF trap may be administered in combination with one or more additional compounds or therapies, including a second VEGF trap molecule, a chemotherapeutic agent, surgery, catheter devices, and radiation. Combination therapy includes administration of a single pharmaceutical dosage formulation which contains a VEGF trap and one or more additional agents; as well as administration of a VEGF trap and one or more additional agent(s) in its own separate pharmaceutical dosage formulation. For example, a VEGF trap and a cytotoxic agent, a chemotherapeutic agent or a growth inhibitory agent can be administered to the patient together in a single dosage composition such as a combined formulation, or each agent can be administered in a separate dosage formulation. Where separate dosage formulations are used, the VEGF-specific fusion polypeptide of the invention and one or more additional agents can be administered concurrently, or at separately staggered times, i.e., sequentially.

[0051] The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. I^{131}, I^{125}, Y^{90} and Re^{186}), chemotherapeutic agents, and toxins such as enzymatically active toxins of bacterial, fungal, plant or animal origin, or fragments thereof.

[0052] A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclosphosphamide (Cytoxan®); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphaoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabicin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®;

razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g. paclitaxel (Taxol®, Bristol-Myers Squibb Oncology, Princeton, N.J.) and docetaxel (Taxotere®; Aventis Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY 117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0053] A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, especially a cancer cell either *in vitro* or *in vivo*. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest. Classical M-phase blockers include the vincas (vincristine and vinblastine), Taxol ®, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C.

Methods of Administration

[0054] The invention provides methods of treatment comprising administering to a subject an effective amount of a VEGF trap of the invention. In a preferred aspect, the trap is substantially purified (*e.g.*, substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably a mammal, and most preferably a human.

[0055] Various delivery systems are known and can be used to administer an agent of the invention, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, *e.g.*, Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction can be enteral or parenteral and include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, intraocular, and oral routes. The compounds may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. Administration can be acute or chronic (e.g. daily, weekly, monthly, etc.) or in combination with other agents. Pulmonary administration can also be employed,

e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[0056] In another embodiment, the active agent can be delivered in a vesicle, in particular a liposome, in a controlled release system, or in a pump. In another embodiment where the active agent of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by use of a retroviral vector (see, for example, U.S. Patent No. 4,980,286), by direct injection, or by use of microparticle bombardment, or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see *e.g.*, Joliot et al., 1991, Proc. Natl. Acad. Sci. USA 88:1864-1868), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

[0057] In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved, for example, and not by way of limitation, by local infusion during surgery, topical application, *e.g.*, by injection, by means of a catheter, or by means of an implant, the implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, fibers, or commercial skin substitutes.

[0058] A composition useful in practicing the methods of the invention may be a liquid comprising an agent of the invention in solution, in suspension, or both. The term "solution/suspension" refers to a liquid composition where a first portion of the active agent is present in solution and a second portion of the active agent is present in particulate form, in suspension in a liquid matrix. A liquid composition also includes a gel. The liquid composition may be aqueous or in the form of an ointment. Further, the composition can take the form of a solid article that can be inserted in the eye, such as for example between the eye and eyelid or in the conjunctival sac, where the VEGF trap is released. Release from such an article is usually to the cornea, either via the lacrimal fluid, or directly to the cornea itself, with which the solid article is generally in direct contact. Solid articles suitable for implantation in the eye are generally composed primarily of bioerodible or nonbioerodible polymers. An aqueous solution and/or suspension can be in the form of eye drops. A desired dosage of the active agent can be measured by administration of a known number of drops into the eye. For example, for a drop volume of 25 µl, administration of 1-6 drops will deliver 25-150 µl of the composition.

[0059] An aqueous suspension or solution/suspension useful for practicing the methods of the invention may contain one or more polymers as suspending agents. Useful polymers include water-soluble polymers such as cellulosic polymers and water-insoluble polymers such as cross-linked carboxyl-containing polymers. An aqueous suspension or solution/suspension of the present invention is preferably viscous or muco-adhesive, or even more preferably, both viscous or mucoadhesive.

[0060] In another embodiment, the composition useful in practicing the methods of the invention is an *in situ* gellable aqueous composition. Such a composition comprises a gelling agent in a concentration effective to promote gelling upon contact with the eye or with lacrimal fluid. Suitable

gelling agents include but are not limited to thermosetting polymers. The term "*in situ* gellable" as used herein is includes not only liquids of low viscosity that form gels upon contact with the eye or with lacrimal fluid, but also includes more viscous liquids such as semi-fluid and thixotropic gels that exhibit substantially increased viscosity or gel stiffness upon administration to the eye.

Diagnostic and Screening Methods

[0061] The VEGF traps of the invention may be used diagnostically and/or in screening methods. For example, the trap may be used to monitor levels of VEGF during a clinical study to evaluate treatment efficacy. In another embodiment, the methods and compositions of the present invention are used to screen individuals for entry into a clinical study to identify individuals having, for example, too high or too low a level of VEGF. The traps can be used in methods known in the art relating to the localization and activity of VEGF, *e.g.*, imaging, measuring levels thereof in appropriate physiological samples, in diagnostic methods, etc.

[0062] The traps of the invention may be used in *in vivo* and *in vitro* screening assay to quantify the amount of non-bound VEGF present, e.g., for example, in a screening method to identify test agents able to decrease the expression of VEGF. More genenerally, the traps of the invention may be used in any assay or process in which quantification and/or isolation of VEGF is desired.

Pharmaceutical Compositions

[0063] The present invention also provides pharmaceutical compositions comprising a VEGF mini-trap of the invention. Such compositions comprise a therapeutically effective amount of one or more mini-traps, and a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin.

[0064] The VEGF mini-trap of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[0065] Further more, aqueous compositions useful for practicing the methods of the invention have ophthalmically compatible pH and osmolality. One or more ophthalmically acceptable pH adjusting agents and/or buffering agents can be included in a composition of the invention, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, and sodium lactate; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases, and buffers are included in an amount required to maintain pH of the composition in an ophthalmically acceptable range. One or more ophthalmically acceptable salts can be included in the composition in an amount sufficient to bring osmolality of the composition into an ophthalmically acceptable range. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions.

[0066] The amount of the trap that will be effective for its intended therapeutic use can be determined by standard clinical techniques based on the present description. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. Generally, suitable dosage ranges for intravenous administration are generally about 50-5000 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0067] For systemic administration, a therapeutically effective dose can be estimated initially from *in vitro* assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Initial dosages can also be estimated from *in vivo* data, e.g., animal models, using techniques that are well known in the art. One having ordinary skill in the art could readily optimize administration to humans based on animal data.

[0068] Dosage amount and interval may be adjusted individually to provide plasma levels of the compounds that are sufficient to maintain therapeutic effect. In cases of local administration or selective uptake, the effective local concentration of the compounds may not be related to plasma concentration. One having skill in the art will be able to optimize therapeutically effective local dosages without undue experimentation.

[0069] The amount of compound administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician. The therapy may be repeated intermittently while symptoms are detectable or even when they are not detectable. The therapy may be provided alone or in combination with other drugs.

Cellular Transfection and Gene Therapy

[0070] The present invention encompasses the use of nucleic acids encoding the fusion polypeptides of the invention for transfection of cells *in vitro* and *in vivo*. These nucleic acids can be inserted into any of a number of well-known vectors for transfection of target cells and organisms. The nucleic acids are transfected into cells *ex vivo* and *in vivo*, through the interaction of the vector and the

target cell. The compositions are administered (e.g., by injection into a muscle) to a subject in an amount sufficient to elicit a therapeutic response. An amount adequate to accomplish this is defined as "a therapeutically effective dose or amount."

[0071] In another aspect, the invention provides a method of reducing VEGF levels in a human or other animal comprising transfecting a cell with a nucleic acid encoding a fusion polypeptide of the invention, wherein the nucleic acid comprises an inducible promoter operably linked to the nucleic acid encoding the fusion polypeptide or mini-trap. For gene therapy procedures in the treatment or prevention of human disease, see for example, Van Brunt (1998) Biotechnology 6:1149-1154.

Kits

[0072] The invention also provides an article of manufacturing comprising packaging material and a pharmaceutical agent contained within the packaging material, wherein the pharmaceutical agent comprises at least one VEGF trap composed of two or more fusion polypeptides of the invention, and wherein the packaging material comprises a label or package insert which indicates that the VEGF-specific fusion polypeptide can be used for treating a VEGF-mediated disease or condition.

Transgenic Animals

[0073] The invention includes transgenic non-human animals expressing a trap of the invention. A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the transgene to particular cells. A transgenic non-human animal expressing a fusion polypeptide or mini-trap of the invention is useful in a variety of applications, including as a means of producing such a fusion polypeptide.. Further, the transgene may be placed under the control of an inducible promoter such that expression of the fusion polypeptide or mini-trap may be controlled by, for example, administration of a small molecule.

Specific Embodiments

[0074] In the experiments described below, smaller VEGF traps were generated and their ability to bind VEGF was investigated. Such mini-traps are preferably uses in specific applications. For example, certain conditions or diseases may be preferably treated with local administration of a VEGF trap to a specific organ, tissue, or cell, rather than by systemic administration. In one exemplification of the mini-traps of the invention, a smaller VEGF trap was generated by directed cleavage of a dimerized VEGF trap having a cleavage region (C-region) generated in a Fc domain (Example 2). The truncated trap exhibited comparable affinity for VEGF and half-life as the full-sized parent trap. Examples 3-5 describe construction of fusion polypeptides having a VEGF receptor component and a multimerizing component consisting of one or two cysteine residues. Affinity measurements showed that the non-glycosylated fusion polypeptides expressed in *E. coli* or

the glycosylated polypeptides expressed in CHO cells had comparable binding affinity for VEGF as the full-sized parent trap. Example 6 further illustrates a monomeric VEGF trap consisting of $(R1R2)_2$ which is capable of binding and inhibiting VEGF. Example 7 describes the construction of a VEGF mini-trap (SEQ ID NO:26) exhibiting high affinity binding for VEGF comparable to the full length trap (SEQ ID NO:10).

[0075] Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

[0076] The following example is put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1. Construction of Flt1D2.Flk1D3.FcΔC1(a)

[0077] The construction of a parent VEGF trap, Flt1D2.Flk1D3.FcΔC1(a) (SEQ ID NOs:7-8), VEGFR1R2.FcΔC1(a) (SEQ ID NOs:9-10), and Flt1D2.VEGFR3D3.FcΔC1(a) (SEQ ID NOs:12-13) is described in detail in PCT publication WO/0075319, herein specifically incorporated by reference in its entirety. Also described in WO/0075319 are methods of constructing and expressing nucleic acid constructs encoding VEGF traps, methods of detecting and measuring VEGF trap binding to VEGF, methods of determining the stoichiometry of VEGF binding by BIAcore analysis, and pharmacokinetic analyses.

Example 2: Thrombin-cleaved dimeric VEGF mini-trap

[0078] The VEGFR1R2.FcΔC1(a) (SEQ ID NOs:9-10) construct was modified by insertion of a thrombin cleavage following the CPPC (SEQ ID NO:1) of the Fc domain. Purified VEGF trap (5 μg) was incubated with thrombin (Novagen) in 20 mM Tris-HCl, pH 8.4, 50 mM NaCl, 2.5 mM $CaCl_2$ for 16 hrs at 37° C. Controls included cleavage control protein (CCP) and parent VEGF trap protein incubated without thrombin. SDS-PAGE analysis (Tris-Glycine 4-20% gel; 5 μg protein per lane) verified correct cleavage (results not shown).

[0079] Affinity determination. The Kd of binding of each VEGF trap to hVEGF165 was determined as described in WO/0075319, for the parent VEGF trap, uncleaved VEGF trap containing a thrombin cleavage site ("uncleaved VEGF trap"), cleaved VEGF mini-trap and recombinant monomeric R1R2-myc myc his. More specifically, the ability of the traps to block $VEGF_{165}$-dependent receptor phosphorylation was determined using primary human endothelial cells (HUVECs). $VEGF_{165}$ was incubated in the presence of varying concentrations of the test traps, and the mixture was added to

HUVECs to stimulate tyrosine phosphorylation of VEGFR2. At sub-stoichiometric concentrations of VEGF trap, unbound VEGF induced receptor phosphorylation. However, at a 1:1 molar ratio of greater of a VEGF trap to ligand, complete blocking of receptor signaling was observed, establishing that a single molecule of a trap dimer is capable of blocking a single molecule of human $VEGF_{165}$. Thus, the high binding affinity of the VEGF trap for VEGF results in formation of a complex that prevents VEGF from interaction with cell surface receptors. Equivalent results were obtained for identical phosphorylation inhibition experiments for the parent VEGF trap, uncleaved VEGF trap, and cleaved VEGF mini-trap The results are shown in Table 1.

TABLE 1

Trap	Kinetic Dissociation Rate (1/s)	$T_{1/2}$ (hr)
parent VEGF trap	$5.51 \times 10^{-5} \pm 0.94\%$	3.5
uncleaved VEGF trap	$4.93 \times 10^{-5} \pm 0.70\%$	3.9
cleaved VEGF mini-trap	$5.46 \times 10^{-5} \pm 0.62\%$	3.53
R1R2-myc myc his monomer	$6.74 \times 10^{-3} \pm 0.38\%$	0.028

Example 3. Construction of Plasmids Encoding VEGF Mini-Traps

[0080] VEGF mini-traps were constructed from a precursor of the parent VEGF trap, VEGFR1R2.FcΔC1(a) (SEQ ID NOs:9-10), in which the three amino acids glycine-alanine-proline served as a linker between the Flk1 D3 and FcΔC1(a). This plasmid, pTE115 was used in the construction of the VEGF mini-traps because the linker DNA sequence included a Srf I restriction endonuclease recognition sequence that facilitated engineering the VEGF trap. In all other respects, the VEGF trap encoded by pTE115 is identical to that of the VEGF trap, VEGFR1R2.FcΔC1(a) (SEQ ID NOs:9-10) described in detail in PCT publication WO/0075319.

[0081] Two VEGF mini-traps were constructed with multimerization domains consisting of either a single cysteine residue ($R1R2_C$) (SEQ ID NO:2) or the amino acids ACGC (SEQ ID NO:4) ($R1R2_{ACGC}$) (SEQ ID NO:5) added to the C-terminus of receptor components Flt1D2.Flk1D3. Both of these constructs are capable of forming homo-dimeric molecules stabilized by one ($R1R2_C$) or two ($R1R2_{ACGC}$) intermolecular disulfides.

[0082] The plasmid pTE517 was made by removing the 690 bp fragment generated by digestion of pTE115 DNA with Srf I and Not I and inserting the synthetic DNA fragment formed by annealing the oligos R1R2NC (SEQ ID NO:14) and R1R2CC (SEQ ID NO:15). The resulting plasmid encodes $R1R2_C$, which consists of the Flt1D2.Flk1D3 domains followed by a cysteine residue (SEQ ID NO:23). Similarly, the plasmid pTE518 was made by removing the 690 bp fragment generated by digestion of pTE115 DNA with Srf I and Not I, followed by ligation with the synthetic DNA fragment formed by annealing the oligos R1R2NACGC (SEQ ID NO:16) and R1R2CACGC (SEQ ID NO:17). The resulting plasmid encodes $R1R2_{ACGC}$, which consists of the Flt1D2.Flk1D3 domains followed by the amino acids ACGC (SEQ ID NO:25).

[0083] Plasmids were also constructed to direct the expression of these mini-traps in *E. coli*. The primers R1R2N-Nco1 (SEQ ID NO:18) and R1R2CNot1 (SEQ ID NO:19) were used to amplify a DNA fragment from pTE115 that encodes amino acids G30 to K231, relative to the parental VEGF trap (SEQ ID NO:10). Amplification of this sequence resulted in fusion of an initiating methionine

codon at the 5' end and fusion of the codon for cysteine, followed by a stop codon, at the 3' end (SEQ ID NO:2). This DNA fragment was then cloned into the Nco I and Not I sites of the *E. coli* expression plasmid pRG663 to yield pRG1102 such that expression of $R1R2_C$ was dependent on transcription from the phage T7 Φ1.1 promoter. Induction of gene expression from pRG1102 results in accumulation of R1R2cys in the cytoplasm of the *E. coli* host strain RFJ238. Similarly, the primers R1R2N-Nco1 (SEQ ID NO:18) and R1R2ACGC-N ot1 (SEQ ID NO:20) were used to amplify a DNA fragment from pTE115 that encodes amino acids G30 to K231 (SEQ ID NO:10) resulting in fusion of an initiating methionine codon at the 5' end and fusion of codons for ACGC (SEQ ID NO:4), followed by a stop codon, at the 3' end (SEQ ID NO:5). This fragment was then cloned into the Nco I and Not I sites of the *E. coli* expression plasmid pRG663 to yield pRG1103 such that expression of $R1R2_{ACGC}$ was dependent on transcription from the phage T7 Φ1.1 promoter. Induction of gene expression from both pRG1102 and pRG1103 resulted in accumulation of $R1R2_C$ or $R1R2_{ACGC}$, respectively, in the cytoplasm of the *E. coli* host strain RFJ238.

Example 4. Purification and characterization of VEGF mini-traps from *E. coli*

[0084] Both $R1R2_C$ and $R1R2_{ACGC}$ were expressed as cytoplasmic proteins in *E. coli* and were purified by the same method. Induction of the phage T7 Φ1.1 promoter on either pRG1102 or pRG1103 in the *E. coli* K12 strain RFJ238 resulted in accumulation of the protein in the cytoplasm. After induction, cells were collected by centrifugation, resuspended in 50 mM Tris-HCl, pH 7.5, 20 mM EDTA, and lysed by passage through a Niro-Soavi cell homogenizer. Inclusion bodies were collected from lysed cells by centrifugation, washed once in distilled H_2O, then solubilized in 8 M guanidinium-HCl, 50 mM Tris-HCl, pH 8.5, 100 mM sodium sulfite,10 mM sodium tetrathionate and incubated at room temperature for 16 hours. Clarified supernatant was fractionated on an S300 column equilibrated with 6 M guanidinium-HCl, 50 mM Tris-HCl, pH 7.5. Fractions containing $R1R2_C$ were pooled and dialyzed against 6M Urea, 50 mM Tris-HCl, pH 7.5. Dialyzed protein was diluted to 2M Urea, 50 mM Tris-HCl, pH 8.5, 2 mM cysteine then stirred slowly for 7 days at 4°C. Refolded protein was dialyzed against 50 mM Tris-HCl, pH 7.5 then loaded onto an SP-sepharose column equilibrated with 50 mM Tris-HCl, pH 7.5 and eluted with a NaCl gradient from 0 to 1 M in 50 mM Tris-HCl, pH 7.5. Fractions containing $R1R2_C$ were pooled, concentrated, and loaded onto a Superdex 200 column equilibrated with 50 mM Tris-HCl, pH 7.5, 150 mM NaCl. Fractions containing mini-trap dimer were collected and pooled. The molecular weight of purified mini-trap was estimated to be about 46 kD by SDS-PAGE.

[0085] BIAcore assays were conducted (as described in WO/0075319) to determine trap affinity for VEGF, and the results showed that the $R1R2_C$ and $R1R2_{ACGC}$ mini-traps had VEGF affinity comparable to the full length VEGF trap (Table 2).

TABLE 2

Trap	Kinetic Dissociation Rate (1/s)	$T_{1/2}$ (hr)
VEGF trap	4.23×10^{-5}	4.53
$R1R2_C$	3.39×10^{-5}	5.68
$R1R2_{ACGC}$	3.41×10^{-5}	5.65

Example 5. Expression of VEGF mini-traps in CHO K1

[0086] Expression of the VEGF mini-traps encoded by pTE517 and pTE518 is dependent on transcription from the human CMV-MIE promoter and results in secretion of the mini-traps into the culture medium when expressed in CHO cells. When expressed as secreted proteins in CHO K1, both mini-traps were found in the conditioned media and estimation of their molecular weight by SDS-PAGE suggested, as expected, that the proteins were glycosylated. Analysis by SDS-PAGE also indicated that the mini-traps were capable of forming homo-dimeric molecules stabilized by intermolecular disulfide(s) between the C-terminal cysteine(s). Specifically, the $R1R2_C$ mini-trap efficiently formed covalent dimers when expressed as a secreted protein in CHO cells.

Example 6. Construction and expression of a single chain VEGF mini-trap

[0087] A VEGF mini-trap was also constructed that did not require a multimerization domain (SEQ ID NO:24). This mini-trap was constructed by direct fusion of one Flt1D2.Flk1D3 domain (R1R2) (amino acids 30-231 of SEQ ID NO:24) to a second Flt1D2.Flk1D3 domain (R1R2) (amino acids 234-435 of SEQ ID NO:24) with a Gly-Pro linker between the tandem receptor domains (amino acids 232-233 of SEQ ID NO:24).

[0088] To construct a gene encoding tandem Flt1D2.Flk1D3 domains, a DNA fragment was synthesized (Blue Heron Biotechnology) that encoded one Flt1D2.Flk1D3 domain that minimized DNA homology with the Flt1D2.Flk1D3 domain-encoding DNA found in pTE115. This synthetic DNA fragment was cloned as a Srf I-Not I fragment into the Srf I-Not I sites of pTE115 to yield pTE570, which expresses the R1R2-R1R2 VEGF mini-trap from the CMV-MIE promoter. When this plasmid is transfected into CHO K1 cells the R1R2-R1R2 VEGF mini-trap accumulates in the culture medium.

Example 7. Construction and expression of a VEGF mini-trap

[0089] A VEGF mini-trap was constructed as described above, by direct fusion of one Flt1D2.Flk1D3 domain (R1R2) (amino acids 30-231 of SEQ ID NO:26) with a C-terminal nine amino acid sequence terminating in CPPC. When this plasmid is transfected into CHO K1 cells the VEGF mini-trap of SEQ ID NO:26 is secreted into the culture medium. Subsequent purification by non-reducing SDS-PAGE electrophoresis as well as native light-scattering analysis identified a trap molecule with molecular weight approximately 64 kDa. This molecular weight indicates that a covalent dimer was formed between two fusion polypeptides of SEQ ID NO:26. Similar experiments were conducted with plasmids encoding the fusion polypeptides of SEQ ID NOS:27 and 28, and similarly showed these molecules formed homodimeric traps. Affinity determinations for human VEGF-165 binding to EGF traps composed of dimers of SEQ ID NO:10 and SEQ ID NO:26 are shown in Table 3.

TABLE 3

VEGF Trap	ka (1/Ms)	kd (1/s)	KD (M)
SEQ ID NO:10	$2.73 \times 10^{+7}$	1.79×10^{-5}	6.55×10^{-13}
SEQ ID NO:26	$2.00 \times 10^{+7}$	6.56×10^{-6}	3.28×10^{-13}
SEQ ID NO:26	$2.61 \times 10^{+7}$	5.77×10^{-6}	2.21×10^{-13}

Claims

1. An isolated nucleic acid molecule encoding a fusion polypeptide consisting of components $(R1R2)_X$ and a multimerizing component (MC), wherein $X \geq 1$, R1 is vascular endothelial cell growth factor (VEGF) receptor component Ig domain 2 of Flt-1 and R2 is Ig domain 3 of Flk-1.

2. The isolated nucleic acid of claim 1, wherein the MC is selected from the group consisting of (i) a multimerizing component comprising a cleavable region (C-region), (ii) a truncated multimerizing component, (iii) an amino acid sequence between 1 to about 200 amino acids in length having at least one cysteine residue, (iv) a leucine zipper, (v) a helix loop motif, (vi) a coil-coil motif, and (vii) an immunoglobulin domain.

3. A fusion polypeptide encoded by the nucleic acid molecule of claims 1 to 2.

4. The fusion polypeptide of claim 3, having the amino acid sequence of SEQ ID NO:26, 27, or 28.

5. A replicable expression vector capable in a transformed host cell comprising the nucleic acid molecule of claims 1 to 2.

6. A method of producing a VEGF fusion polypeptide, comprising the steps of introducing into a suitable expression system the expression vector of claim 5, and effecting expression of the VEGF fusion polypeptide.

7. A vascular endothelial cell growth factor (VEGF) trap, comprising a multimer of two or more fusion polypeptides of claim 3.

8. The VEGF trap of claim 7, which is a dimer.

9. A dimeric VEGF trap comprising two fusion polypeptides comprising the amino acid sequence of SEQ ID NO:26, 27, or 28.

10. A pharmaceutical composition comprising the fusion polypeptide of claims 7 or 8, and a pharmaceutically acceptable carrier.

11. A method of treating a disease or condition which is improved, ameliorated, or inhibited by removal or inhibition of vascular endothelial growth factor (VEGF), comprising administering the pharmaceutical composition of claim 10 to a subject in need thereof.

12. The method of claim 11, wherein the disease or condition is an ocular disease or condition.

13. The method of claim 12, wherein the ocular disease or condition is age related macular degeneration.

14. An isolated nucleic acid molecule encoding a fusion polypeptide consisting of receptor components $(R1R2)_X$, wherein $X \geq 1$, R1 is vascular endothelial cell growth factor (VEGF) receptor component Ig domain 2 of Flt-1 and R2 is Ig domain 3 of Flk-1.

15. A fusion polypeptide capable of binding vascular endothelial growth factor (VEGF) encoded by the nucleic acid molecule of claims 14.

16. The fusion polypeptide of claim 15, comprising the amino acid sequence of SEQ ID NO:24.

17. A fusion polypeptide consisting of receptor components $(R1R2)_X$, wherein $X \geq 1$, R1 is vascular endothelial cell growth factor (VEGF) receptor component Ig domain 2 of Flt-1 and R2 is Ig domain 3 of Flk-1.

18. The fusion polypeptide of claim 17, having the amino acid sequence of SEQ ID NO:24.

19. An article of manufacturing comprising:

(a) packaging material; and

(b) a pharmaceutical agent contained within said packaging material;

wherein the pharmaceutical agent comprises at least one VEGF trap according to any one claims 4, 7, 8, 9, 15, 16, 17, or 18 and wherein the packaging material comprises a label or package insert which indicates that said VEGF-specific fusion polypeptide can be used for treating a VEGF-mediated disease or condition.