



## CLAIMS

1. A compound corresponding to formula (I):




in which:


- n represents 0 or 1;
- Y represents a bridged morpholine chosen from



(a)



(b)



(c)

➤ L represents a linker  $-\text{CH}_2-\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or a  $(\text{C}_1-\text{C}_2)$  alkyl, said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1-\text{C}_3)$  alkyl group and a hydroxyl group;

➤  $\text{R}_1$  represents:

- a linear, branched, cyclic or partially cyclic  $(\text{C}_1-\text{C}_5)$  alkyl group, optionally substituted with one or more substituents chosen from a hydroxyl group, an aryl group and a trifluoromethyl group,
- a  $(\text{C}_3-\text{C}_6)$  cycloalkyl group, optionally substituted with a hydroxyl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, a cyano group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1-\text{C}_4)$  alkyl, a morpholine group, a group of formula  $-\text{SO}_2-(\text{C}_1-\text{C}_5)$  alkyl, a  $(\text{C}_1-\text{C}_5)$  alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:

- a halogen atom,

- a hydroxyl group or a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
  - a group -COR<sub>3</sub>, in which R<sub>3</sub> represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
  - a group -CONR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
  - a group -NR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
  - a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
  - a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group;
- 5
- 10
- a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - a halogen atom,
    - a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
    - a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
    - a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,
  - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an oxygen atom,
  - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group,
  - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
- 15
- 20
- 25
- 30
- 35
- R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when


n represents 0;

➤  $R_4$  and  $R_{4'}$ , independently, which may be identical or different, represent a hydrogen atom or a  $(C_1-C_3)$ alkyl group, in the form of the base or of an addition salt with an acid or with a base.

5

2. The compound of formula (I) as claimed in claim 1, characterized in that:

- n represents 0 or 1;
- Y represents a bridged morpholine chosen from



10

➤ L represents a linker  $-CH_2-CO-$  such that the carbonyl function is attached to the substituent  $R_1$ , or a  $(C_1-C_2)$ alkyl, said alkyl being optionally substituted with one or more substituents chosen from a  $(C_1-C_3)$ alkyl group and a hydroxyl group;

15

➤  $R_1$  represents:

- a linear or branched  $(C_1-C_5)$  alkyl group, optionally substituted with one or more substituents chosen from a hydroxyl group and an aryl group,
- a  $(C_3-C_6)$ cycloalkyl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, a cyano group, an  $-NH_2$  group, a urea group of formula  $-NH-CO-NH-(C_1-C_4)$ alkyl, a morpholine group, a group of formula  $-SO_2-(C_1-C_5)$ alkyl, a  $(C_1-C_5)$ alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:

20

25

- a halogen atom,
- a hydroxyl group or a  $(C_1-C_5)$ alkoxy group,
- a group  $-COR_3$ , in which  $R_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
- a group  $-CONR_4R_{4'}$  in which  $R_4$  and  $R_{4'}$  are as defined below,
- a group  $-NR_4R_{4'}$  in which  $R_4$  and  $R_{4'}$  are as defined below,
- a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,


30

- a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group;
  - 5 - a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - a halogen atom,
    - a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
    - 10 ○ a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
    - 15 ○ a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,
    - 20 - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an oxygen atom,
    - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group and an acetyl group,
    - 25 - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
      - R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
      - 30 ➤ R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,
- in the form of the base or of an addition salt with an acid or with a base.

3. The compound of formula (I) as claimed in claim 1 or 2, characterized in that:

- 35 ➤ Y represents a bridged morpholine (a)






(a)

in the form of the base or of an addition salt with an acid or with a base.

4. The compound of formula (I) as claimed in claim 1 or 2, characterized in that:

- 5
- n represents 0 or 1;
  - Y represents a bridged morpholine (a)



(a)

- 10
- L represents a linker  $-\text{CH}_2-\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or  $(\text{C}_1-\text{C}_2)\text{alkyl}$ , said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1-\text{C}_3)\text{alkyl}$  group;

- 15
- $\text{R}_1$  represents:
    - a linear, branched, cyclic or partially cyclic  $(\text{C}_1-\text{C}_5)\text{alkyl}$  group, optionally substituted with one or more substituents chosen from a hydroxyl group, an aryl group, a trifluoromethyl group and a  $(\text{C}_3-\text{C}_6)\text{cycloalkyl}$  group,
    - a  $(\text{C}_3-\text{C}_6)\text{cycloalkyl}$  group, optionally substituted with a hydroxyl group,
    - an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1-\text{C}_4)\text{alkyl}$ , a morpholinyl group, a group of formula  $-\text{SO}_2-(\text{C}_1-\text{C}_5)\text{alkyl}$ , a  $(\text{C}_1-\text{C}_5)\text{alkoxy}$  group, said alkoxy being optionally substituted with one or more substituents chosen from:


- 20
- a halogen atom,
    - a hydroxyl group or a  $(\text{C}_1-\text{C}_5)\text{alkoxy}$ ,
    - a group  $-\text{COR}_3$ , in which  $\text{R}_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
    - a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
    - a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_1-\text{C}_3)\text{alkyl}$  group, a hydroxyl group and
- 25

an -NH<sub>2</sub> group;

- a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - 5      ○ a halogen atom,
    - a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
    - a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a
    - 10      heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
    - a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen
    - 15      atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,
  - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an
  - 20      oxygen atom,
  - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group,
  - 25      - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
    - R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
    - R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a
    - 30      hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,
- in the form of the base or of an addition salt with an acid or with a base.

5. The compound of formula (I) as claimed in any one of claims 1 to 4, characterized in that:

- 35      ➤ n represents 0 or 1;
- Y represents a bridged morpholine (a)



(a)

➤ L represents a linker  $-\text{CH}_2\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or  $(\text{C}_1\text{-C}_2)$ alkyl, said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1\text{-C}_3)$ alkyl group;

➤  $\text{R}_1$  represents:

- a linear, branched, cyclic or partially cyclic  $(\text{C}_1\text{-C}_5)$ alkyl group, optionally substituted with one or more substituents chosen from a hydroxyl group and an aryl group,
- a  $(\text{C}_3\text{-C}_6)$ cycloalkyl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1\text{-C}_4)$ alkyl, a morpholinyl group, a group of formula  $-\text{SO}_2-(\text{C}_1\text{-C}_5)$ alkyl, a  $(\text{C}_1\text{-C}_5)$ alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:
  - a halogen atom,
  - a hydroxyl group or a  $(\text{C}_1\text{-C}_5)$ alkoxy,
  - a group  $-\text{COR}_3$ , in which  $\text{R}_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
  - a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
  - a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_1\text{-C}_3)$ alkyl group, a hydroxyl group and an  $-\text{NH}_2$  group,
- a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
  - a halogen atom,
  - a  $(\text{C}_1\text{-C}_3)$ alkyl group optionally substituted with one or more halogen atoms,
  - a  $(\text{C}_1\text{-C}_5)$ alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_3\text{-C}_5)$ cycloalkyl group, a heteroaryl group optionally substituted with one or more substituents

chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,

5      ○ a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,

10      - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an oxygen atom,

10      - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group and an acetyl group,

15      - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,

➤ R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;

➤ R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,


20      in the form of the base or of an addition salt with an acid or with a base.

6. The compound of formula (I) as claimed in claim 1 or 2, characterized in that:


➤ n represents 0 or 1;

➤ Y represents a bridged morpholine chosen from (b) and (c)

25



(b)



(c)

30      ➤ L represents a linker -CH<sub>2</sub>-CO- such that the carbonyl function is attached to the substituent R<sub>1</sub>, or (C<sub>1</sub>-C<sub>2</sub>)alkyl, said alkyl being optionally substituted with a hydroxyl group;

➤ R<sub>1</sub> represents:

- a linear or branched (C<sub>1</sub>-C<sub>5</sub>)alkyl group, optionally substituted with an aryl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:
  - a group -CONR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
  - a group -NR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
- a heteroaryl group comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more (C<sub>1</sub>-C<sub>3</sub>)alkyl groups, optionally substituted with one or more halogen atoms,
  - R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
  - R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, in the form of the base or of an addition salt with an acid or with a base.

7. The compound as claimed in any one of claims 1 to 6, characterized in that the linker L represents -CH<sub>2</sub>-CO, in the form of the base or of an addition salt with an acid or with a base.

8. The compound as claimed in any one of claims 1 to 7, characterized in that n represents 1, in the form of the base or of an addition salt with an acid or with a base.

9. The compound as claimed in any one of claims 1 to 7, characterized in that n represents 0, in the form of the base or of an addition salt with an acid or with a base.

10. The compound as claimed in any one of claims 1 to 9, characterized in that R<sub>1</sub> represents a heteroaryl group, in the form of the base or of an addition salt with an acid or with a base.

11. The compound as claimed in any one of claims 1 to 9, characterized in that R<sub>1</sub> represents a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group, in the form of the base or of an addition salt with an acid or with a base.

12. The compound as claimed in any one of the preceding claims 1 to 11, characterized in that it is chosen from:

- 1 (8S)-9-(2-Methyl-2-pyrid-4-ylpropyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 2 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-4-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 3 (8S)-9-[2-(6-Aminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 4 (8S)-9-[2-(6-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 (8S)-9-[2-(6-Methylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 6 (8S)-9-[2-(6-Dimethylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 7 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 8 1-[2-(6-Dimethylaminopyrid-3-yl)-2-oxoethyl]-2-(S)-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 9 2-(S)-Methyl-1-[2-(6-methylaminopyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 10 (8S)-1-[2-(4-Methoxyphenyl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-

azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

11 (S)-1-[2-(6-Aminopyrid-3-yl)-2-oxoethyl]-2-methyl-7-(1S,4S)-2-oxa-5-  
5 azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

12 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

13 2-Methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-1-(2-pyrid-3-ylethyl)-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

10 14 (8S)-9-{2-[6-(2-Hydroxyethylamino)pyrid-3-yl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

15 15 (8S)-9-[2-(5-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

16 2-Methyl-1-[2-(5-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

20 17 2-Methyl-1-[2-(6-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

18 2-Methyl-1-[2-(2-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

25 19 (8S)-9-[2-(2-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

20 (8S)-9-[2-(4-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-

4-one

- 21 2-Methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-1-(2-oxo-2-pyrid-3-yl)ethyl)-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 22 (8S)-9-[2-(6-Cyclopropylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 23 1-Ethyl-3-{4-[2-((S)-8-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)ethyl]phenyl}urea
- 24 1-Ethyl-3-{4-[2-((S)-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-5-oxo-2-trifluoromethyl-2,3-dihydro-5H-imidazo[1,2-a]pyrimidin-1-yl)ethyl]phenyl}urea
- 25 (8S)-9-[2-(4-Methylthiazol-5-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 26 2-Methyl-1-[2-(4-methylthiazol-5-yl)ethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 27 (8S)-9-[2-(3,5-Dimethyl-1H-pyrazol-4-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 28 1-[2-(3,5-Dimethyl-1H-pyrazol-4-yl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 29 (8S)-9-(3,3-Dimethyl-2-oxobutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 30 1-(3,3-Dimethyl-2-oxobutyl)-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 31 (8S)-9-[2-(6-Amino-5-methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one



- 32 1-[2-(4-Aminophenyl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 33 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(6-trifluoromethylpyrid-3-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5
- 34 (8S)-9-[2-{6-[(2-Hydroxyethyl)methylamino]pyrid-3-yl}-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 35 (8S)-9-[2-(6-Ethoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10
- 36 (8S)-9-[2-(6-Amino-4,5-dimethylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 37 (S)-9-[2-(4-Difluoromethoxyphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15
- 38 (8S)-9-[2-(3,4-Dihydro-2H-pyrido[3,2-b][1,4]oxazin-7-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20
- 39 (8S)-9-[2-(4-Methyloxazol-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 40 (S)-9-[2-(3,4-Difluorophenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25
- 41 (8S)-9-[2-(4-Morpholin-4-ylphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 42 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]benzonitrile
- 43 (8S)-9-[2-(4-Methylthiazol-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 44 (8S)-9-[2-(5-Chloropyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 45 (8S)-9-[2-(6-Methoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 46 (8S)-9-[2-(3-Methylisoxazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 47 (8S)-9-[2-(2-Benzo[1,2,3]thiadiazol-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 48 (8S)-9-[2-(2,4-Difluorophenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 49 (8S)-9-(3-Ethyl-3-hydroxypentyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 50 (8S)-9-(3-Hydroxy-3-methylbutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 51 (8S)-9-(1-Methyl-1H-indazol-3-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 52 (8S)-9-[2-(2-Cyclopropylmethoxypyrimidin-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 53 (8S)-9-[2-(3,5-Dimethylisoxazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 54 (8S)-9-(2-Ethyl-2-hydroxybutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 55 3-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]benzonitrile
- 56 (8S)-9-(3-Methyl-2-oxobutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 57 {5-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]pyrid-2-yl}carbamic acid ethyl ester
- 58 {5-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]pyrid-2-yl}carbamic acid methyl ester
- 15
- 59 (8S)-9-(5-Methyl-[1,2,4]oxadiazol-3-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 60 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(2-trifluoromethylpyrid-3-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20
- 61 (8S)-9-(2-Benzo[1,2,5]thiadiazol-5-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 62 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(tetrahydropyran-4-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 63 (8S)-9-{2-[6-(2-Fluoroethoxy)pyrid-3-yl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 64 (8S)-9-{2-[3-Fluoro-4-(2-fluoroethoxy)phenyl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 65 (8S)-9-[2-(2-Methoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 66 (8S)-9-[2-(3-Methyl-3H-imidazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 67 (8S)-9-(2-Cyclopropyl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 68 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-2-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 69 (8S)-9-[2-(2-Methyl-2H-pyrazol-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 70 N,N-Dimethyl-2-(4-{2-[(S)-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-2-trifluoromethyl-2,3-dihydro-5H-imidazo[1,2-a]pyrimidin-1-yl]ethyl}phenoxy)acetamide
- 20 71 (8S)-9-[(S)-2-(4-Fluoro-2-methoxyphenyl)-2-hydroxyethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 72 (2S)-1-[2-(4-Hydroxyphenyl)ethyl]-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 25 73 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-phenylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 74 (2S)-1-{2-[4-(2-Dimethylaminoethoxy)phenyl]ethyl}-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

- 75 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-4-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 76 (S)-1-[2-(4-Methoxyphenyl)ethyl]-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 5 77 (S)-2-Methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-1-(3-phenylpropyl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 78 (S)-1-[2-[4-(3-Dimethylaminopropoxy)phenyl]ethyl]-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 10 79 (2S)-1-((S)-2-Hydroxy-2-phenylethyl)-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 80 (8S)-9-((S)-2-Hydroxy-2-phenylethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 81 (8S)-9-[2-(4-Methoxyphenyl)ethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 82 (8S)-9-((R)-2-Benzo[b]thiophen-2-yl-2-hydroxyethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 83 (8S)-9-[2-(4-Hydroxyphenyl)ethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 84 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(3-phenylpropyl)-8-trifluoromethylmethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 85 (8S)-2-(3-Oxa-8-azabicyclo[3.2.1]oct-8-yl)-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 86 (8S)-9-(1-Difluoromethyl-1H-pyrazol-3-ylmethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 87 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 88 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-2-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 89 (S)-9-[2-(1-Acetylpiperid-4-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 90 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)ethyl]piperidine-1-carbaldehyde
- 91 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]piperidine-1-carboxylic acid
- 10 ethyl ester
- 92 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(tetrahydropyran-4-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 93 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(tetrahydropyran-4-ylmethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15
- 94 (8S)-9-(1-Acetylpiperid-4-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 95 4-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-ylmethyl)piperidine-1-carbaldehyde
- 20 96 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(3,3,3-trifluoro-2-hydroxy-2-trifluoromethylpropyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 97 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(4,4,4-trifluoro-3-hydroxy-3-trifluoromethylbutyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 one
- 98 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-oxoethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 99 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-2-oxoethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 100 (8S)-9-[2-(1-Hydroxycyclopentyl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 101 (8S)-9-(1-Hydroxycyclopentylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 102 (8S)-9-(3,3-Dicyclopropyl-3-hydroxypropyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 103 (8S)-9-(2,2-Dicyclopropyl-2-hydroxyethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 104 (8S)-9-(1-Hydroxycyclopropylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 105 (8S)-9-[2-(1-Hydroxycyclopropyl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 106 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-quinolin-5-ylmethyl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 107 (8S)-9-[2-(3-Methylisothiazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 108 (8S)-9-[2-(4-Methanesulfonylphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 109 (8S)-9-Isoquinolin-5-ylmethyl-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 110 (8S)-9-(2-Morpholin-4-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-

yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one


- 111 (8S)-9-{2-[4-(2-Morpholin-4-ylethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 112 N-Methoxy-N-methyl-2-((S)-8-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetamide
- 113 (8S)-9-(2-Imidazo[1,2-a]pyrid-6-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 114 (8S)-9-[2-(6-Difluoromethoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 115 (S)-9-{2-[4-(2-Morpholin-4-yl-2-oxoethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 116 (8S)-9-(1-Methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 117 (8S)-9-{2-[4-(2-Dimethylaminoethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 118 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]piperidine-1-carbaldehyde
- 25 119 (8S)-9-[2-(1-Acetylpiperid-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

in the form of the base or of an addition salt with an acid or with a base.

13. A process for preparing a compound of formula (I) as claimed in any one of claims




1 to 12, comprising the reaction of a compound of formula E



in which n represents 0 or 1 and R<sub>2</sub> represents a hydrogen atom when n = 1 or a methyl group when n = 0, with a bridged morpholine Y, chosen from (a), (b) and (c) as


5 defined in claim 1, to obtain a compound of formula I



and the alkylation reaction by addition to I of a compound of formula J = R<sub>1</sub>-L-Lg in which R<sub>1</sub> and L are as defined in any one of the preceding claims and Lg is a leaving group.


10

14. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, comprising the alkylation reaction of a compound of formula E



in which n represents 0 or 1 and R<sub>2</sub> represents a hydrogen atom when n = 1 or a methyl group when n = 0, by addition of a compound of formula J = R<sub>1</sub>-L-Lg in which R<sub>1</sub> and L are as defined in any one of claims 1 to 8 and Lg is a leaving group, to obtain a compound of formula K


15



in which R<sub>1</sub>, R<sub>2</sub>, L and n are as defined in one of claims 1 to 8, and a reaction on a


compound K with a compound of formula Y being a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1.

15. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker L is an ethyl group,  $R_1$  is a linear or branched ( $C_1$ - $C_5$ )alkyl group substituted with a hydroxyl group, Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1, n represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising a Michael addition reaction of a compound of formula E



E

in which n represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , on a compound of formula  $M = CH_2=CH_2-CO_2Alkyl$ , to obtain a compound of formula N,




N

- 15 in which n represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , and Y is a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,  
and a reaction of alkyl on a compound of formula N with a compound of formula  $O = Z-Mg-X$  in which Z represents a linear or branched alkyl radical and X is a halogen atom.

20


16. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker L is a methyl group,  $R_1$  is a linear or branched ( $C_1$ - $C_5$ )alkyl group substituted with a hydroxyl group, Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1, n represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising an addition reaction of a compound of formula E

25



in which  $n$  represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , with a compound of formula  $P = X-CH_2-CO_2Alkyl$  with  $X$  being a halogen atom, to obtain a compound of formula Q

5




in which  $Y$  is a bridged morpholine chosen from (a), (b) and (c),  $n$  represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ ,

and an alkylation reaction on a compound of formula Q with a compound of formula  $O = Z-Mg-X$  in which  $Z$  represents a linear or branched alkyl radical and  $X$  is a halogen atom.


17. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker  $L$  is a methyl group,  $R_1$  is a group  $-NR_6R_{6'}$  with  $R_6$  et  $R_{6'}$  being either different and representing an alkyl group and an alkoxy group, or  $R_6$  and  $R_{6'}$  together forming a monocyclic or bicyclic heterocycloalkyl,  $Y$  represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,  $n$  represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising a hydrolysis reaction of a compound of formula Q

20



in which  $Y$  is a bridged morpholine chosen from (a), (b) and (c),  $n$  represents 0 or 1 and


$R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , to obtain a compound of formula S



in which Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,  $n$  represents 1 or 0 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ ,

and a coupling reaction between a compound of formula S and a compound of formula  $HNR_6R_6'$  with  $R_6$  and  $R_6'$  being either different and representing an alkyl group and an alkoxy group, or  $R_6$  and  $R_6'$  together forming a monocyclic or bicyclic heterocycloalkyl.

18. Compounds of formulae I, N, Q and S:



in which  $n$ ,  $R_2$  and Y are as defined in claim 1.

19. A medicament, characterized in that it comprises a compound as claimed in any one of claims 1 to 12, or an addition salt of this compound with a pharmaceutically acceptable acid or base.

20. The compound as claimed in any one of claims 1 to 12, as a medicament.

21. A pharmaceutical composition, characterized in that it comprises a compound as claimed in any one of claims 1 to 12, or a pharmaceutically acceptable salt of this compound, and also at least one pharmaceutically acceptable excipient.

5

22. The use of a compound as claimed in any one of claims 1 to 12, for the preparation of a medicament for treating parasite-induced malaria.


10

23. The compound as claimed in any one of claims 1 to 12, for its use in the treatment of malaria induced by all species of *Plasmodium*, such as *P. falciparum*, *P. vivax*, *P. malariae*, *P. ovale* and *P. knowlesi*, by all species of *Trypanosoma* and by all species of *Leishmania*, in the treatment of sleeping sickness, the treatment of Chagas disease, the various forms of leishmaniasis and the treatment of other parasitic infections, such as schistosomiasis (bilharzia), toxoplasmosis and coccidiosis.

15

## CLAIMS


1. A compound corresponding to formula (I):




5

in which:


- n represents 0 or 1;
- Y represents a bridged morpholine chosen from



(a)



(b)



(c)

10

➤ L represents a linker  $-\text{CH}_2-\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or a  $(\text{C}_1-\text{C}_2)$ alkyl, said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1-\text{C}_3)$ alkyl group and a hydroxyl group;

15

➤  $\text{R}_1$  represents:

- a linear, branched, cyclic or partially cyclic  $(\text{C}_1-\text{C}_5)$ alkyl group, optionally substituted with one or more substituents chosen from a hydroxyl group, an aryl group and a trifluoromethyl group,
- a  $(\text{C}_3-\text{C}_6)$ cycloalkyl group, optionally substituted with a hydroxyl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, a cyano group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1-\text{C}_4)$ alkyl, a morpholine group, a group of formula  $-\text{SO}_2-(\text{C}_1-\text{C}_5)$ alkyl, a  $(\text{C}_1-\text{C}_5)$ alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:

25

- a halogen atom,

- a hydroxyl group or a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
- a group -COR<sub>3</sub>, in which R<sub>3</sub> represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
- a group -CONR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
- 5      ○ a group -NR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
- a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
- a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and
- 10      an -NH<sub>2</sub> group;
- a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
  - a halogen atom,
  - 15      ○ a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
  - a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a heteroaryl group optionally substituted with one or more substituents
  - 20      chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
  - a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally
  - 25      substituted with one or more hydroxyl groups,
- a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an oxygen atom,
- 30      - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group,
- a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
- 35      ➤ R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when


n represents 0;

➤  $R_4$  and  $R_{4'}$ , independently, which may be identical or different, represent a hydrogen atom or a  $(C_1-C_3)$ alkyl group, in the form of the base or of an addition salt with an acid or with a base.

5

2. The compound of formula (I) as claimed in claim 1, characterized in that:

- n represents 0 or 1;
- Y represents a bridged morpholine chosen from



10

➤ L represents a linker  $-CH_2-CO-$  such that the carbonyl function is attached to the substituent  $R_1$ , or a  $(C_1-C_2)$ alkyl, said alkyl being optionally substituted with one or more substituents chosen from a  $(C_1-C_3)$ alkyl group and a hydroxyl group;

15

➤  $R_1$  represents:

- a linear or branched  $(C_1-C_5)$  alkyl group, optionally substituted with one or more substituents chosen from a hydroxyl group and an aryl group,
- a  $(C_3-C_6)$ cycloalkyl group,
- an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, a cyano group, an  $-NH_2$  group, a urea group of formula  $-NH-CO-NH-(C_1-C_4)$ alkyl, a morpholine group, a group of formula  $-SO_2-(C_1-C_5)$ alkyl, a  $(C_1-C_5)$ alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:

20

25

- a halogen atom,
- a hydroxyl group or a  $(C_1-C_5)$ alkoxy group,
- a group  $-COR_3$ , in which  $R_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
- a group  $-CONR_4R_{4'}$  in which  $R_4$  and  $R_{4'}$  are as defined below,
- a group  $-NR_4R_{4'}$  in which  $R_4$  and  $R_{4'}$  are as defined below,
- a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,


30



- a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group;
  - 5     - a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - a halogen atom,
    - a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
    - 10     ○ a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
    - 15     ○ a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,
    - 20     - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an oxygen atom,
    - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group and an acetyl group,
    - 25     - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
      - R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
    - 30     ➤ R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,
- in the form of the base or of an addition salt with an acid or with a base.

3. The compound of formula (I) as claimed in claim 1 or 2, characterized in that:

- 35     ➤ Y represents a bridged morpholine (a)




(a)

in the form of the base or of an addition salt with an acid or with a base.

4. The compound of formula (I) as claimed in claim 1 or 2, characterized in that:

- 5
- n represents 0 or 1;
  - Y represents a bridged morpholine (a)



(a)

- 10
- L represents a linker  $-\text{CH}_2-\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or  $(\text{C}_1-\text{C}_2)\text{alkyl}$ , said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1-\text{C}_3)\text{alkyl}$  group;


- 15
- $\text{R}_1$  represents:
    - a linear, branched, cyclic or partially cyclic  $(\text{C}_1-\text{C}_5)\text{alkyl}$  group, optionally substituted with one or more substituents chosen from a hydroxyl group, an aryl group, a trifluoromethyl group and a  $(\text{C}_3-\text{C}_6)\text{cycloalkyl}$  group,
    - a  $(\text{C}_3-\text{C}_6)\text{cycloalkyl}$  group, optionally substituted with a hydroxyl group,
    - an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1-\text{C}_4)\text{alkyl}$ , a morpholinyl group, a group of formula  $-\text{SO}_2-(\text{C}_1-\text{C}_5)\text{alkyl}$ , a  $(\text{C}_1-\text{C}_5)\text{alkoxy}$  group, said alkoxy being optionally substituted with one or more substituents chosen from:
      - 20
        - a halogen atom,
        - a hydroxyl group or a  $(\text{C}_1-\text{C}_5)\text{alkoxy}$ ,
        - a group  $-\text{COR}_3$ , in which  $\text{R}_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
        - 25
          - a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
          - a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_1-\text{C}_3)\text{alkyl}$  group, a hydroxyl group and

an -NH<sub>2</sub> group;

- a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - 5           ○ a halogen atom,
    - a (C<sub>1</sub>-C<sub>3</sub>)alkyl group optionally substituted with one or more halogen atoms,
    - a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group, a
    - 10           heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a (C<sub>1</sub>-C<sub>3</sub>)alkyl group, a hydroxyl group and an -NH<sub>2</sub> group,
    - a group -NR<sub>5</sub>R<sub>5'</sub> in which R<sub>5</sub> and R<sub>5'</sub>, independently, which may be identical or different, represent a substituent chosen from a hydrogen
    - 15           atom, a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>3</sub>)alkyl group, a (C<sub>3</sub>-C<sub>5</sub>)cycloalkyl group and a linear or branched (C<sub>1</sub>-C<sub>3</sub>)alkyl group, said alkyl group being optionally substituted with one or more hydroxyl groups,
    - a pyridine group bearing two linked adjacent groups forming, together with the two carbons that bear them, a heterocycle comprising a nitrogen atom and an
    - 20           oxygen atom,
    - a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group,
    - 25           - a group -NR<sub>6</sub>R<sub>6'</sub> in which R<sub>6</sub> and R<sub>6'</sub>, which are different, represent a (C<sub>1</sub>-C<sub>5</sub>)alkyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group,
      - R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
      - R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a
      - 30           hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,
- in the form of the base or of an addition salt with an acid or with a base.

5. The compound of formula (I) as claimed in any one of claims 1 to 4, characterized in that:

- 35           ➤ n represents 0 or 1;
- Y represents a bridged morpholine (a)



(a)

- L represents a linker  $-\text{CH}_2\text{CO}-$  such that the carbonyl function is attached to the substituent  $\text{R}_1$ , or  $(\text{C}_1\text{-C}_2)\text{alkyl}$ , said alkyl being optionally substituted with one or more substituents chosen from a  $(\text{C}_1\text{-C}_3)\text{alkyl}$  group;
- $\text{R}_1$  represents:
- a linear, branched, cyclic or partially cyclic  $(\text{C}_1\text{-C}_5)\text{alkyl}$  group, optionally substituted with one or more substituents chosen from a hydroxyl group and an aryl group,
  - a  $(\text{C}_3\text{-C}_6)\text{cycloalkyl}$  group,
  - an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group, an  $-\text{NH}_2$  group, a urea group of formula  $-\text{NH}-\text{CO}-\text{NH}-(\text{C}_1\text{-C}_4)\text{alkyl}$ , a morpholinyl group, a group of formula  $-\text{SO}_2-(\text{C}_1\text{-C}_5)\text{alkyl}$ , a  $(\text{C}_1\text{-C}_5)\text{alkoxy}$  group, said alkoxy being optionally substituted with one or more substituents chosen from:
    - a halogen atom,
    - a hydroxyl group or a  $(\text{C}_1\text{-C}_5)\text{alkoxy}$ ,
    - a group  $-\text{COR}_3$ , in which  $\text{R}_3$  represents a substituent chosen from a heterocycloalkyl group and a hydroxyl group,
    - a heterocycloalkyl group comprising one or two heteroelements chosen from a nitrogen atom and an oxygen atom,
    - a heteroaryl group optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_1\text{-C}_3)\text{alkyl}$  group, a hydroxyl group and an  $-\text{NH}_2$  group,
  - a heteroaryl group, comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more substituents chosen from:
    - a halogen atom,
    - a  $(\text{C}_1\text{-C}_3)\text{alkyl}$  group optionally substituted with one or more halogen atoms,
    - a  $(\text{C}_1\text{-C}_5)\text{alkoxy}$  group, optionally substituted with one or more substituents chosen from a halogen atom, a  $(\text{C}_3\text{-C}_5)\text{cycloalkyl}$  group, a heteroaryl group optionally substituted with one or more substituents



- a linear or branched (C<sub>1</sub>-C<sub>5</sub>)alkyl group, optionally substituted with an aryl group,
  - an aryl group, optionally substituted with one or more substituents chosen from a halogen atom, a hydroxyl group and a (C<sub>1</sub>-C<sub>5</sub>)alkoxy group, said alkoxy being optionally substituted with one or more substituents chosen from:
    - a group -CONR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
    - a group -NR<sub>4</sub>R<sub>4'</sub> in which R<sub>4</sub> and R<sub>4'</sub> are as defined below,
  - a heteroaryl group comprising one or more heteroatoms chosen from a nitrogen atom, a sulfur atom and an oxygen atom, optionally substituted with one or more (C<sub>1</sub>-C<sub>3</sub>)alkyl groups, optionally substituted with one or more halogen atoms,
- R<sub>2</sub> represents a hydrogen atom when n represents 1 and a methyl group when n represents 0;
- R<sub>4</sub> and R<sub>4'</sub>, independently, which may be identical or different, represent a hydrogen atom or a (C<sub>1</sub>-C<sub>3</sub>)alkyl group,
- in the form of the base or of an addition salt with an acid or with a base.

7. The compound as claimed in any one of claims 1 to 6, characterized in that the linker L represents -CH<sub>2</sub>-CO, in the form of the base or of an addition salt with an acid or with a base.

8. The compound as claimed in any one of claims 1 to 7, characterized in that n represents 1, in the form of the base or of an addition salt with an acid or with a base.

9. The compound as claimed in any one of claims 1 to 7, characterized in that n represents 0, in the form of the base or of an addition salt with an acid or with a base.

10. The compound as claimed in any one of claims 1 to 9, characterized in that R<sub>1</sub> represents a heteroaryl group, in the form of the base or of an addition salt with an acid or with a base.

11. The compound as claimed in any one of claims 1 to 9, characterized in that R<sub>1</sub> represents a heterocycloalkyl group comprising one or more heteroatoms chosen from oxygen and nitrogen atoms, said nitrogen atom being optionally substituted with a substituent chosen from a formyl group, an acetyl group and a -CO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl group, in the form of the base or of an addition salt with an acid or with a base.

12. The compound as claimed in any one of the preceding claims 1 to 11, characterized in that it is chosen from:

- 1 (8S)-9-(2-Methyl-2-pyrid-4-ylpropyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 2 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-4-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 3 (8S)-9-[2-(6-Aminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 4 (8S)-9-[2-(6-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 (8S)-9-[2-(6-Methylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 6 (8S)-9-[2-(6-Dimethylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 7 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 8 1-[2-(6-Dimethylaminopyrid-3-yl)-2-oxoethyl]-2-(S)-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 9 2-(S)-Methyl-1-[2-(6-methylaminopyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 10 (8S)-1-[2-(4-Methoxyphenyl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-

- azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 11 (S)-1-[2-(6-Aminopyrid-3-yl)-2-oxoethyl]-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 12 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 13 2-Methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-1-(2-pyrid-3-ylethyl)-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 10 14 (8S)-9-{2-[6-(2-Hydroxyethylamino)pyrid-3-yl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 15 (8S)-9-[2-(5-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 16 2-Methyl-1-[2-(5-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 17 2-Methyl-1-[2-(6-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 20 18 2-Methyl-1-[2-(2-methylpyrid-3-yl)-2-oxoethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 25 19 (8S)-9-[2-(2-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 (8S)-9-[2-(4-Methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-



4-one

- 21 2-Methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-1-(2-oxo-2-pyrid-3-ylethyl)-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 22 (8S)-9-[2-(6-Cyclopropylaminopyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 23 1-Ethyl-3-{4-[2-((S)-8-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)ethyl]phenyl}urea
- 24 1-Ethyl-3-{4-[2-((S)-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-5-oxo-2-trifluoromethyl-2,3-dihydro-5H-imidazo[1,2-a]pyrimidin-1-yl)ethyl]phenyl}urea
- 25 (8S)-9-[2-(4-Methylthiazol-5-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 26 2-Methyl-1-[2-(4-methylthiazol-5-yl)ethyl]-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 27 (8S)-9-[2-(3,5-Dimethyl-1H-pyrazol-4-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 28 1-[2-(3,5-Dimethyl-1H-pyrazol-4-yl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 29 (8S)-9-(3,3-Dimethyl-2-oxobutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 30 1-(3,3-Dimethyl-2-oxobutyl)-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 31 (8S)-9-[2-(6-Amino-5-methylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 32 1-[2-(4-Aminophenyl)ethyl]-2-methyl-7-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-2-((S)-trifluoromethyl)-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 33 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(6-trifluoromethylpyrid-3-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5
- 34 (8S)-9-[2-{6-[(2-Hydroxyethyl)methylamino]pyrid-3-yl}-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 35 (8S)-9-[2-(6-Ethoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10
- 36 (8S)-9-[2-(6-Amino-4,5-dimethylpyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 37 (S)-9-[2-(4-Difluoromethoxyphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15
- 38 (8S)-9-[2-(3,4-Dihydro-2H-pyrido[3,2-b][1,4]oxazin-7-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20
- 39 (8S)-9-[2-(4-Methyloxazol-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 40 (S)-9-[2-(3,4-Difluorophenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25
- 41 (8S)-9-[2-(4-Morpholin-4-ylphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 42 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]benzonitrile
- 43 (8S)-9-[2-(4-Methylthiazol-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 4-one
- 44 (8S)-9-[2-(5-Chloropyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 45 (8S)-9-[2-(6-Methoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 4-one
- 46 (8S)-9-[2-(3-Methylisoxazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 47 (8S)-9-(2-Benzo[1,2,3]thiadiazol-5-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 4-one
- 48 (8S)-9-[2-(2,4-Difluorophenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 4-one
- 49 (8S)-9-(3-Ethyl-3-hydroxypentyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 50 (8S)-9-(3-Hydroxy-3-methylbutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 51 (8S)-9-(1-Methyl-1H-indazol-3-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 4-one
- 52 (8S)-9-[2-(2-Cyclopropylmethoxypyrimidin-5-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 53 (8S)-9-[2-(3,5-Dimethylisoxazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 54 (8S)-9-(2-Ethyl-2-hydroxybutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 55 3-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]benzonitrile
- 56 (8S)-9-(3-Methyl-2-oxobutyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 57 {5-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]pyrid-2-yl}carbamic acid ethyl ester
- 58 {5-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]pyrid-2-yl}carbamic acid methyl ester
- 15
- 59 (8S)-9-(5-Methyl-[1,2,4]oxadiazol-3-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 60 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(2-trifluoromethylpyrid-3-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20
- 61 (8S)-9-(2-Benzo[1,2,5]thiadiazol-5-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 62 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-oxo-2-(tetrahydropyran-4-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 63 (8S)-9-{2-[6-(2-Fluoroethoxy)pyrid-3-yl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 64 (8S)-9-{2-[3-Fluoro-4-(2-fluoroethoxy)phenyl]-2-oxoethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 65 (8S)-9-[2-(2-Methoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 66 (8S)-9-[2-(3-Methyl-3H-imidazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 67 (8S)-9-(2-Cyclopropyl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 68 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(2-oxo-2-pyrid-2-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 69 (8S)-9-[2-(2-Methyl-2H-pyrazol-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 70 N,N-Dimethyl-2-(4-{2-[(S)-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-5-oxo-2-trifluoromethyl-2,3-dihydro-5H-imidazo[1,2-a]pyrimidin-1-yl]ethyl}phenoxy)acetamide
- 71 (8S)-9-[(S)-2-(4-Fluoro-2-methoxyphenyl)-2-hydroxyethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 72 (2S)-1-[2-(4-Hydroxyphenyl)ethyl]-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 73 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-phenylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 74 (2S)-1-{2-[4-(2-Dimethylaminoethoxy)phenyl]ethyl}-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one

- 75 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-4-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 76 (S)-1-[2-(4-Methoxyphenyl)ethyl]-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 5 77 (S)-2-Methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-1-(3-phenylpropyl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 78 (S)-1-{2-[4-(3-Dimethylaminopropoxy)phenyl]ethyl}-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 10 79 (2S)-1-((S)-2-Hydroxy-2-phenylethyl)-2-methyl-7-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-trifluoromethyl-2,3-dihydro-1H-imidazo[1,2-a]pyrimidin-5-one
- 80 (8S)-9-((S)-2-Hydroxy-2-phenylethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 81 (8S)-9-[2-(4-Methoxyphenyl)ethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 82 (8S)-9-((R)-2-Benzo[b]thiophen-2-yl-2-hydroxyethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 83 (8S)-9-[2-(4-Hydroxyphenyl)ethyl]-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 84 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(3-phenylpropyl)-8-trifluoromethylmethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 85 (8S)-2-(3-Oxa-8-azabicyclo[3.2.1]oct-8-yl)-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 25 86 (8S)-9-(1-Difluoromethyl-1H-pyrazol-3-ylmethyl)-2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 87 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-3-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 88 (8S)-2-(8-Oxa-3-azabicyclo[3.2.1]oct-3-yl)-9-(2-oxo-2-pyrid-2-ylethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 89 (S)-9-[2-(1-Acetylpiperid-4-yl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 90 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)ethyl]piperidine-1-carbaldehyde
- 91 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]piperidine-1-carboxylic acid  
10 ethyl ester
- 92 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(tetrahydropyran-4-yl)ethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 93 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(tetrahydropyran-4-ylmethyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one  
15
- 94 (8S)-9-(1-Acetylpiperid-4-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 95 4-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-ylmethyl)piperidine-1-carbaldehyde
- 20 96 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(3,3,3-trifluoro-2-hydroxy-2-trifluoromethylpropyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 97 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-(4,4,4-trifluoro-3-hydroxy-3-trifluoromethylbutyl)-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one  
25
- 98 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(8-oxa-3-azabicyclo[3.2.1]oct-3-yl)-2-oxoethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 99 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-[2-(3-oxa-8-azabicyclo[3.2.1]oct-8-yl)-2-oxoethyl]-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 100 (8S)-9-[2-(1-Hydroxycyclopentyl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 101 (8S)-9-(1-Hydroxycyclopentylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 102 (8S)-9-(3,3-Dicyclopropyl-3-hydroxypropyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 103 (8S)-9-(2,2-Dicyclopropyl-2-hydroxyethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 104 (8S)-9-(1-Hydroxycyclopropylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 105 (8S)-9-[2-(1-Hydroxycyclopropyl)ethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 106 (8S)-2-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-9-quinolin-5-ylmethyl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 107 (8S)-9-[2-(3-Methylisothiazol-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 108 (8S)-9-[2-(4-Methanesulfonylphenyl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 109 (8S)-9-Isoquinolin-5-ylmethyl-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 110 (8S)-9-(2-Morpholin-4-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-




yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

- 111 (8S)-9-{2-[4-(2-Morpholin-4-ylethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 5 112 N-Methoxy-N-methyl-2-((S)-8-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetamide
- 113 (8S)-9-(2-Imidazo[1,2-a]pyrid-6-yl-2-oxoethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 10 114 (8S)-9-[2-(6-Difluoromethoxypyrid-3-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 15 115 (S)-9-{2-[4-(2-Morpholin-4-yl-2-oxoethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 116 (8S)-9-(1-Methyl-3-trifluoromethyl-1H-pyrazol-4-ylmethyl)-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 20 117 (8S)-9-{2-[4-(2-Dimethylaminoethoxy)phenyl]ethyl}-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one
- 118 4-[2-((S)-8-(1S,4S)-2-Oxa-5-azabicyclo[2.2.1]hept-5-yl-6-oxo-2-trifluoromethyl-3,4-dihydro-2H,6H-pyrimido[1,2-a]pyrimidin-1-yl)acetyl]piperidine-1-carbaldehyde
- 25 119 (8S)-9-[2-(1-Acetylpiperid-4-yl)-2-oxoethyl]-2-(1S,4S)-2-oxa-5-azabicyclo[2.2.1]hept-5-yl-8-trifluoromethyl-6,7,8,9-tetrahydropyrimido[1,2-a]pyrimidin-4-one

in the form of the base or of an addition salt with an acid or with a base.


13. A process for preparing a compound of formula (I) as claimed in any one of claims

1 to 12, comprising the reaction of a compound of formula E



in which n represents 0 or 1 and R<sub>2</sub> represents a hydrogen atom when n = 1 or a methyl group when n = 0, with a bridged morpholine Y, chosen from (a), (b) and (c) as


5 defined in claim 1, to obtain a compound of formula I



and the alkylation reaction by addition to I of a compound of formula J = R<sub>1</sub>-L-Lg in which R<sub>1</sub> and L are as defined in any one of the preceding claims and Lg is a leaving group.


10

14. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, comprising the alkylation reaction of a compound of formula E



in which n represents 0 or 1 and R<sub>2</sub> represents a hydrogen atom when n = 1 or a methyl group when n = 0, by addition of a compound of formula J = R<sub>1</sub>-L-Lg in which R<sub>1</sub> and L are as defined in any one of claims 1 to 8 and Lg is a leaving group, to obtain a compound of formula K


15



in which R<sub>1</sub>, R<sub>2</sub>, L and n are as defined in one of claims 1 to 8, and a reaction on a


compound K with a compound of formula Y being a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1.

15. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker L is an ethyl group,  $R_1$  is a linear or branched ( $C_1$ - $C_5$ )alkyl group substituted with a hydroxyl group, Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1, n represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising a Michael addition reaction of a compound of formula E



10

in which n represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , on a compound of formula  $M = CH_2=CH_2-CO_2Alkyl$ , to obtain a compound of formula N,




- 15 in which n represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , and Y is a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,

and a reaction of alkyl on a compound of formula N with a compound of formula  $O = Z-Mg-X$  in which Z represents a linear or branched alkyl radical and X is a halogen atom.

20


16. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker L is a methyl group,  $R_1$  is a linear or branched ( $C_1$ - $C_5$ )alkyl group substituted with a hydroxyl group, Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1, n represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising an addition reaction of a compound of formula E

25



in which  $n$  represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , with a compound of formula  $P = X-CH_2-CO_2Alkyl$  with  $X$  being a halogen atom, to obtain a compound of formula Q

5




in which  $Y$  is a bridged morpholine chosen from (a), (b) and (c),  $n$  represents 0 or 1 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ ,

and an alkylation reaction on a compound of formula Q with a compound of formula  $O = Z-Mg-X$  in which  $Z$  represents a linear or branched alkyl radical and  $X$  is a halogen atom.


17. A process for preparing a compound of formula (I) as claimed in any one of claims 1 to 12, in which the linker  $L$  is a methyl group,  $R_1$  is a group  $-NR_6R_{6'}$  with  $R_6$  and  $R_{6'}$  being either different and representing an alkyl group and an alkoxy group, or  $R_6$  and  $R_{6'}$  together forming a monocyclic or bicyclic heterocycloalkyl,  $Y$  represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,  $n$  represents 1 or 0, and  $R_2$  represents a hydrogen atom when  $n = 1$  and a methyl group when  $n = 0$ , comprising a hydrolysis reaction of a compound of formula Q

20



in which  $Y$  is a bridged morpholine chosen from (a), (b) and (c),  $n$  represents 0 or 1 and


$R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ , to obtain a compound of formula S



in which Y represents a bridged morpholine chosen from (a), (b) and (c) as defined in claim 1,  $n$  represents 1 or 0 and  $R_2$  represents a hydrogen atom when  $n = 1$  or a methyl group when  $n = 0$ ,

and a coupling reaction between a compound of formula S and a compound of formula  $HNR_6R_{6'}$  with  $R_6$  and  $R_{6'}$  being either different and representing an alkyl group and an alkoxy group, or  $R_6$  and  $R_{6'}$  together forming a monocyclic or bicyclic heterocycloalkyl.

18. Compounds of formulae I, N, Q and S:



in which  $n$ ,  $R_2$  and Y are as defined in claim 1.

19. A medicament, characterized in that it comprises a compound as claimed in any one of claims 1 to 12, or an addition salt of this compound with a pharmaceutically acceptable acid or base.

20. The compound as claimed in any one of claims 1 to 12, as a medicament.

21. A pharmaceutical composition, characterized in that it comprises a compound as claimed in any one of claims 1 to 12, or a pharmaceutically acceptable salt of this compound, and also at least one pharmaceutically acceptable excipient.

5

22. The use of a compound as claimed in any one of claims 1 to 12, for the preparation of a medicament for treating parasite-induced malaria.

10

23. The compound as claimed in any one of claims 1 to 12, for use in the treatment of malaria induced by all species of *Plasmodium*, such as *P. falciparum*, *P. vivax*, *P. malariae*, *P. ovale* and *P. knowlesi*, by all species of *Trypanosoma* and by all species of *Leishmania*, in the treatment of sleeping sickness, the treatment of Chagas disease, the various forms of leishmaniasis and the treatment of other parasitic infections, such as schistosomiasis (bilharzia), toxoplasmosis and coccidiosis.

15