[CLAIMS]

[Claim 1]

A form-II crystal of 2-{4-[N-(5,6-diphenylpyrazin-2-yl)-N-isopropylamino]butyloxy}acetic acid, which shows diffraction peaks at diffraction angles (2θ) of 9.6°, 11.4°, 11.7°, 16.3°, 17.5°, 18.5°, 18.7°, 19.9°, 20.1°, 21.0°, and 24.6° in a powder X-ray diffraction spectrum obtained using a Cu-Kα radiation (λ=1.54 Å).

[Claim 2]

The form-II crystal of 2-{4-[N-(5,6-diphenylpyrazin-2-yl)-N-isopropylamino]butyloxy}acetic acid according to claim 1, which shows absorption peaks at wavenumbers of 2867 cm^{-1}, 1749 cm^{-1}, 1568 cm^{-1}, 1382 cm^{-1}, 1131 cm^{-1}, and 701 cm^{-1} in an infrared absorption spectrum.

[Claim 3]

The form-II crystal of 2-{4-[N-(5,6-diphenylpyrazin-2-yl)-N-isopropylamino]butyloxy}acetic acid according to claim 1, which has an endothermic peak at 147°C in differential scanning calorimetry.

[Claim 4]

A pharmaceutical composition, comprising the crystal according to any one of claims 1 to 3 as an active ingredient.

[Claim 5]

A PGI2 receptor agonist, comprising the crystal according to any one of claims 1 to 3 as an active ingredient.

[Claim 6]

A therapeutic agent comprising the crystal according to any one of claims 1 to 3 as an active ingredient, for use in treating symptoms associated with diabetic neuropathy, diabetic gangrene, a peripheral circulatory disturbance, chronic arterial occlusion, intermittent claudication, scleroderma, thrombosis, pulmonary hypertension, myocardial infarction, angina pectoris, glomerulonephritis, diabetic nephropathy, chronic renal failure, bronchial asthma, interstitial pneumonia (pulmonary fibrosis), a chronic obstructive pulmonary disease, tubulointerstitial nephritis, an inflammatory bowel disease, or spinal canal stenosis.

[Claim 7]

Use of the crystal according to any one of claims 1 to 3 in the manufacture of a therapeutic agent for treating symptoms associated with diabetic neuropathy, diabetic gangrene, a peripheral circulatory disturbance, chronic arterial occlusion, intermittent claudication, scleroderma, thrombosis, pulmonary hypertension, myocardial infarction, angina pectoris, glomerulonephritis, diabetic nephropathy, chronic renal failure, bronchial asthma, interstitial pneumonia (pulmonary fibrosis), a chronic obstructive pulmonary disease, tubulointerstitial nephritis, an inflammatory bowel disease, or spinal canal stenosis.