

ENGINEERED DUAL BINDING ANTIBODIES AND USES THEREOF**SEQUENCE LISTING STATEMENT**

[1] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on May 25, 2022, is named P-605548-PC.txt and is 14.8 Kilo bytes in size.

FIELD OF INTEREST

[2] The disclosure relates in general to dual binding antibodies that bind to IL-13 and TSLP. In one embodiment, the antibodies can be used for treating allergic or respiratory conditions.

BACKGROUND

[3] IL-13 is a 12.3kDa monomeric protein of the class I cytokines. IL-13 has a 4 alpha helical bundle core topology typical of class I short helical cytokines; it is similar in structure to its closely related cytokine IL-4 sharing low sequence identity but high structural identity. Along with IL-4, IL-13 has been shown to control immunoglobulin class switching to IgE in B cells and is involved in mast cell recruitment. IL-13 is secreted by CD4⁺ Th2 cells, as well as type 2 innate lymphoid cells ILC2. It has been demonstrated that IL-13 can trigger the production of (TGF- β) and in bronchial epithelial cells induces gene expression of *MUC5AC* and production of mucin. IL-13 can also enhance contraction in smooth bronchial muscle cells. IL-13 binds the IL-4Ra/IL-13Ra1 heterodimeric complex, and upon binding it triggers a JAK-signal transducer and STAT6 dependent signaling cascade, which in turn triggers Th2 helper T-cell differentiation, polarization of macrophages to the M2 “alternatively activated” phenotype, epithelial mucus production, smooth muscle contractility and chemokine release.

[4] IL-13 has been shown to be involved in protection against parasites, wherein it was demonstrated that in knockout IL-13 mice model, clearance of *N. brasiliensis* is severely delayed. Also, the expulsion of *Trichuris muris* is abolished completely, in spite of an otherwise intact Th2-type response. Further research demonstrated that IL-13 is a double-edged sword, on the one hand it has a major role in protection against parasites, however IL-13 function in situations of dysregulated immune system is also known.

[5] IL-13 has implicated in the pathogenesis of human asthma as elevated levels of IL-13 mRNA and protein have been detected in lungs of asthmatic patients, which correlate with severity of the disease. In addition, human IL-13 genetic polymorphisms, which lead to elevated IL-13 levels, have been identified and are associated with asthma and atopy, and elevated IL-13 levels

have been detected in the lung of asthma patients.

[6] Although IL-13 and IL-4 share similar receptors and signaling pathway, IL-13 has distinguished role in asthma, which is independent of IL-4. It has been shown in mice models that administration of IL-13 alone is sufficient to induce eosinophil derived inflammation, and mucus cell hyperplasia. Moreover, a specific blockade of IL-13 but not IL-4 is sufficient to reverse airway hyperreactivity and mucus production in mice models. Additionally, polymorphism in the human IL-13 locus is known to be associated with high susceptibility for asthma.

[7] Specific inhibition of IL-13 signaling could therefore have positive therapeutic effect on asthma patients, or patients with other known allergic or respiratory conditions.

[8] Thymic stromal lymphopoietin (TSLP) is a cytokine that signals through a heterodimeric receptor consisting of the IL-7R α subunit and TSLP-R, a unique component with homology to the common γ -receptor-like chain. TSLP is expressed by epithelial cells in the thymus, lung, skin, intestine, and tonsils, as well as airway smooth muscle cells, lung fibroblasts, and stromal cells. These cells produce TSLP in response to proinflammatory stimuli, and TSLP drives allergic inflammatory responses through its activity on a number of innate immune cells, including dendritic cells. TSLP can also promote proliferation of naive T cells and drive their differentiation into Th2 cells expressing high levels of IL-4, IL-5, and IL-13. High level of TSLP expression has been found in asthmatic lung epithelial cells and chronic atopic dermatitis lesions, suggesting a role for TSLP in allergic inflammation. Recent evidence also implicates TSLP in the differentiation of Th17 cells and Th17-driven inflammatory processes. Chronic allergic (atopic) asthma is often characterized by Th2-type inflammation, while non-allergic asthmatic inflammation is predominately neutrophilic with a mixed Th1 and Th17 cytokine milieu. Antagonists to TSLP would be expected to be useful for treating inflammatory conditions.

[9] Thus, there remains an unmet need for compositions and methods of treatment of diseases and conditions triggered by IL-13 and TSLP activation, for example but not limited to allergic and respiratory conditions, including but not limited to asthma.

SUMMARY

[10] In one embodiment, the present disclosure provides an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences of SEQ ID NOs:149-154. In another embodiment, the dual binding antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL) having the amino acid sequences of SEQ ID Nos:155 and 156, or SEQ ID Nos:157 and 158.

[11] Disclosed herein, in one aspect is an isolated dual binding antibody, wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:359, 360 and 361 respectively.

[12] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 371 respectively.

[13] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:362, 360 and 384 respectively.

[14] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 384 respectively.

[15] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.

[16] Disclosed herein, in one aspect is an isolated dual binding antibody, said dual binding antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain, wherein said VH domain comprises a set of complementarity-determining regions (CDRs), HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid; or wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid; or a combination of (a) and (b).

[17] In a related aspect, wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is selected from the group consisting of W and S; wherein the amino acid sequence of HCDR3 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group

consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M; wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G.

[18] In a further related aspect of the isolated dual binding antibody, HX1 is W, HX2 is selected from the group consisting of A and S, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX12 is selected from the group I and L, LX1 is L, LX2 is I, LX3 is L, LX4 is selected from the group consisting of S and G, LX5 is S, LX6 is S, and LX7 is selected from the group consisting of H and G.

[19] In yet another related aspect of isolated dual binding antibody, an isolated dual binding antibody comprises CDRs wherein HX1 is W, HX2 is A, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX12 is I, LX4 is S, and LX7 is G; or HX1 is W, HX2 is A, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX12 is L, LX4 is S, and LX7 is H; or HX1 is W, HX2 is S, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX12 is L, LX4 is G, and LX7 is G.

[20] In another related aspect of the isolated dual binding antibody, said VH domain comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); said VL domain comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of the VH domain set forth in (a) and the VL domain set forth in (b); wherein the total number of variant positions in said VH domain, said VL domain, or said combination thereof of said dual binding antibody, is at least 2.

[21] In a further related aspect, said at least one variant amino acid in said VH domain

comprises a variant at position 106 of SEQ ID NO: 1 (IMTG position 112). In another further related aspect, the amino acid sequence of said VH domain is selected from the sequences set forth in SEQ ID Nos: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54. In yet another further related aspect, said at least one amino acid variant in said VL domain comprises a variant amino acid in a CDR region. In still another further related aspect, said variant amino acid in said VL domain comprises a variant at any of positions 26, 27, 31, or 96 of SEQ ID NO: 2, or a combination thereof (IMGT positions: 27, 28, 38, or 115, or a combination thereof). In another further related aspect, there are at least two variants in said VL domain and the second variant comprises a variant amino acid in a framework region. In yet another further related aspect, said variant amino acid in a framework region comprises a variant at position 56 or 77 of SEQ ID NO: 2, or a combination thereof (IMGT positions: 70 or 94, or a combination thereof).

[22] In another related aspect, the amino acid sequence of said VL domain is selected from the sequences set forth in SEQ ID Nos: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53. In still another related aspect, the amino acid sequences of a VH domain – VL domain pair are selected from the pair sequences set forth in SEQ ID Nos: 4 and 3, SEQ ID Nos: 6 and 5, SEQ ID Nos: 8 and 7, SEQ ID Nos: 10 and 9, SEQ ID Nos: 12 and 11, SEQ ID Nos: 14 and 13, SEQ ID Nos: 16 and 15, SEQ ID Nos: 18 and 17, SEQ ID Nos: 20 and 19, SEQ ID Nos: 22 and 21, SEQ ID Nos: 24 and 23, SEQ ID Nos: 26 and 25, SEQ ID Nos: 28 and 27, SEQ ID Nos: 30 and 29, SEQ ID Nos: 32 and 31, SEQ ID Nos: 34 and 33, SEQ ID Nos: 36 and 35, SEQ ID Nos: 38 and 37, SEQ ID Nos: 40 and 39, SEQ ID Nos: 42 and 41, SEQ ID Nos: 44 and 43, SEQ ID Nos: 46 and 45, SEQ ID Nos: 48 and 47, SEQ ID Nos: 50 and 49, SEQ ID Nos: 52 and 51, and SEQ ID Nos: 54 and 53.

[23] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:209 and 210.

[24] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:219 and 220.

[25] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:249 and 250.

[26] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and

VL comprise the amino acid sequences of SEQ ID Nos:337 and 338.

[27] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences as shown in Table 1 or Table 10.

[28] In another related aspect, the dual binding antibody comprises an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody. In a further related aspect, the IgG comprises IgG1, IgG2, IgG3, or an IgG4. In still another related aspect, said IgG comprises a mutated IgG which is unable to bind to antibody-dependent cellular cytotoxicity components.

[29] Disclosed herein, in one aspect is a composition comprising the isolated dual binding antibody and a pharmaceutically acceptable carrier.

[30] Disclosed herein, in one aspect is a nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding antibody, said antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain, wherein said VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid; or wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid; or a combination of (a) and (b).

[31] In a related aspect of the nucleic acid, the encoded amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is selected from the group consisting of W and S; wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M; wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is

selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G.

[32] In a related aspect of the nucleic acid, the encoded amino acid for HX1 is W, HX2 is selected from the group consisting of A and S, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX12 is selected from the group I and L, LX1 is L, LX2 is I, LX3 is L, LX4 is selected from the group consisting of S and G, LX5 is S, LX6 is S, and LX7 is selected from the group consisting of H and G. A further related aspect, wherein the encoded amino acid for HX1 is W, HX2 is A, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX12 is I, LX4 is S, and LX7 is G; or HX1 is W, HX2 is A, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX12 is L, LX4 is S, and LX7 is H; or HX1 is W, HX2 is S, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX12 is L, LX4 is G, and LX7 is G.

[33] In a related aspect of the nucleic acid construct, said VH domain comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); said VL domain comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of VH domain set forth in (a) and the VL domain set forth in (b); wherein the total number of variant positions in the encoded VH domain, the encoded VL domain, or a combination thereof, is at least 2. In a further related aspect, said sequence comprises two nucleic acid sequences, one encoding the variant dual binding antibody VH domain, and one encoding the variant dual binding antibody VL domain. In a further related aspect, the nucleic acid sequence encoding said VH domain is selected from the sequences set forth in SEQ ID Nos: 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 105, and 107. In another further related aspect, the nucleic acid sequence encoding said VL domain is selected from the sequences set forth in SEQ ID Nos: 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, and 108. In yet another further related aspect,

the nucleic acid sequences encoding the dual antibody VH domain – VL domain pair are selected from the paired sequences set forth in SEQ ID Nos: 57 and 58, SEQ ID Nos: 59 and 60, SEQ ID Nos: 61 and 62, SEQ ID Nos: 63 and 64, SEQ ID Nos: 65 and 66, SEQ ID Nos: 67 and 68, SEQ ID Nos: 69 and 70, SEQ ID Nos: 71 and 72, SEQ ID Nos: 73 and 74, SEQ ID Nos: 75 and 76, SEQ ID Nos: 77 and 78, SEQ ID Nos: 79 and 80, SEQ ID Nos: 81 and 82, SEQ ID Nos: 83 and 84, SEQ ID Nos: 85 and 86, SEQ ID Nos: 87 and 88, SEQ ID Nos: 89 and 90, SEQ ID Nos: 91 and 92, SEQ ID Nos: 93 and 94, SEQ ID Nos: 95 and 96, SEQ ID Nos: 97 and 98, SEQ ID Nos: 99 and 100, SEQ ID Nos: 101 and 102, SEQ ID Nos: 103 and 104, SEQ ID Nos: 105 and 106, and SEQ ID Nos: 107 and 108.

[34] In a related aspect of the nucleic acid construct, said antibody comprises an IgG, a Fv, a scFv, a Fab, a F(ab')₂, a minibody, a diabody, or a triabody. In a further related aspect, said IgG comprises a mutated IgG which is unable to bind to antibody-dependent cellular cytotoxicity components.

[35] In another related aspect, the nucleic acid construct further comprises a regulatory sequence operably linked to said nucleic acid sequence.

[36] Disclosed herein, in one aspect is an expression vector comprising the nucleic acid construct encoding a dual binding antibody, said antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain.

[37] Disclosed herein, in one aspect is a host cell comprising the expression vector comprising a nucleic acid construct encoding a dual binding antibody, said antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain.

[38] Disclosed herein, in one aspect is a composition comprising a nucleic acid construct encoding a dual binding antibody, said antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain and a pharmaceutically acceptable carrier.

[39] Disclosed herein, in one aspect is a method of producing a dual binding antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain, said method comprising culturing the host cell comprising the expression vector comprising a nucleic acid construct encoding a dual binding antibody, said antibody comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain; expressing said nucleic acid construct from said vector; isolating said dual binding antibody.

[40] Disclosed herein, in one aspect is a library of immunoglobulins or fragments thereof comprising an antibody antigen-binding domain site comprising a heavy chain variable region (VH) domain and a light chain variable region (VL) domain, wherein said VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid; or wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid.

[41] In a related aspect of the library, the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is selected from the group consisting of W and S; wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M; wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G.

[42] In a further related aspect of the library, HX1 is W, HX2 is selected from the group consisting of A and S, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX12 is selected from the group I and L, LX1 is L, LX2 is I, LX3 is L, LX4 is selected from the group consisting of S and G, LX5 is S,

LX6 is S, and LX7 is selected from the group consisting of H and G.

[43] In yet another further related aspect of the library, HX1 is W, HX2 is A, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX12 is I, LX4 is S, and LX7 is G; HX1 is W, HX2 is A, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX12 is L, LX4 is S, and LX7 is H; or HX1 is W, HX2 is S, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX12 is L, LX4 is G, and LX7 is G. In still another further related aspect of the library, said VH domain comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); said VL domain comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of the VH domain set forth in (a) and the VL domain set forth in (b); wherein the total number of variant positions in the VH domain, the VL domain, or a combination thereof is at least 2.

[44] In another related aspect of the library, the immunoglobulin comprises an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody.

[45] In another related aspect of the library, the IgG comprises a mutated IgG which is unable to bind to antibody-dependent cellular cytotoxicity components.

[46] Disclosed herein in one aspect, is a method of treating a subject suffering from a disease or condition comprising an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, said method comprising administering to said subject an isolated dual binding antibody disclosed herein.

[47] In one related aspect of the method of treating a subject, said allergic or respiratory condition is asthma; allergic asthma; nonallergic asthma; severe asthma; mild asthma; chronic obstructive pulmonary disease (COPD); a condition involving airway inflammation including eosinophilia, fibrosis and excess mucus production, cystic fibrosis, allergic lung disease, airway hyperresponsiveness, goblet cell metaplasia, mucus hypersecretion, airway remodeling, pulmonary fibrosis; atopic disorders including atopic dermatitis, urticaria, eczema, allergic enterogastritis, and allergic rhinitis; or a combination thereof; or said inflammatory and/or autoimmune conditions including inflammatory bowel diseases (IBD) and liver conditions including cirrhosis or fibrosis; or a combination thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[48] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[49] The subject matter of engineered dual binding antibodies is particularly pointed out and distinctly claimed in the concluding portion of the specification. These dual binding antibodies, however, both as to their generation and method of use, together with objects, features, and advantages thereof, may best be understood by reference to the following detailed description when read with the accompanying drawings in which:

[50] **Figures 1A and 1B** present the template antibody heavy chain (SEQ ID NO: 1) (**Figure 1A**) and light chain (SEQ ID NO: 2) amino acid sequences (**Figure 1B**), respectively, indicating the framework (FR) and complementarity-determining regions (CDR) regions. For the Heavy (H) chain, the different regions are labeled FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4, and in some embodiments are referred to as HFR1, HCDR1, HFR2, HCDR2, HFR3, HCDR3, and HFR4. For the Light (L) chain, the different regions are labeled FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4, and in some embodiments are referred to as LFR1, LCDR1, LFR2, LCDR2, LFR3, LCDR3, and LFR4. Below the template amino acid sequences, the variant amino acids of the engineered dual binding clones are displayed and aligned within the CDR and FR regions.

[51] **Figures 2A and 2B** present bar graphs showing binding of re-epitoped antibodies displayed on yeast to recombinant human IL-13 (rh-IL-13) (**Figure 2A**) or recombinant human TSLP (rhTSLP) (**Figure 2B**). **Figure 2A** shows binding of isolated yeast-surface display anti-IL13 clones to 10nM rh-IL-13. **Figure 2B** shows binding of isolated yeast-surface display anti-TSLP clones to 10nM rhTSLP. Data was normalized to the yeast surface expression levels of each clone, and to an anti-hIL-13 and anti-hTSLP mean fluorescence intensity (MFI) binding signal of positive control yeast clones.

[52] **Figures 3A-3F** presents size exclusion chromatography (SEC) scans of a human standard IgG1 (**Figure 3A**), BDG33.003 (**Figure 3B**), BDG33.004 (**Figure 3C**), BDG 33.005 (**Figure 3D**), BDG33.023 (**Figure 3E**), and BDG33.025 (**Figure 3F**). The purified IgGs were run on a GE Superdex®200 10/300 increase (column volume (CV)=25ml) in PBS buffer at 0.5ml/min. In the antibody scans shown in **Figures 3B-3D**, the leading peak corresponds to (0.36CV) that typical of a large aggregate, and a second peak with retention of approximately 13.2ml (0.528CV) that is typical of an ordinary human IgG. Area Under the Curve (AUC) peak ratio is approximately 23% misfolded/77% folded IgG fraction, respectively. For the antibody scans shown in **Figures 3E-3F**, the leading peak corresponds to (0.36CV) that typical of a large

diameter aggregate, and a second peak with retention of approximately 13.8ml (0.55CV) that is typical of an ordinary human IgG. Area Under the Curve (AUC) peak ratio is 97.3% folded/2.8% misfolded and 98.5% folded/1.5% misfolded for BDG33.023 (**Figure 3E**) and BDG33.025 (**Figure 3F**) respectively.

[53] **Figures 4A and 4B** present Differential Scanning Fluorimetry (DSF) analysis of the melting point of indicated IgGs BDG33.023 (**Figure 4A**) and BDG33.025 (**Figure 4B**). Light gray dashed line in the upper graph represents the T-onset and bold gray dashed lines represents the Tm1 and Tm2. The lower graph is the 1st derivative of the measurement. **Figure 4A:** DSF of BDG33.023 T-onset of 64.2°C and first transition point at 67.7°C. **Figure 4B:** DSF of BDG33.025 T-onset of 56.4°C, first transition point at 60.9°C and second transition point at 67.4°C.

[54] **Figures 5A-5F** presents Surface Plasmon Resonance (SPR) analysis of antibodies binding to human IL-13, Cyno IL-13, and human TSLP. Representative SPR sonograms of BDG33.003 and BDG33.004 binding to IL-13 are presented in **Figures 5A-5D**. Recombinant human IL-13 (rh-IL-13) was tested at 800nM with a 2-fold dilution (**Figures 5A – 5B**). Recombinant cyno IL-13 (rc-IL-13) was tested at 200nM with a 2-fold dilution (**Figures 5C – 5D**). Representative SPR sensorgrams of BDG33.003 and BDG33.004 binding to human TSLP (h-TSLP) are presented in **Figures 5E and 5F**. hTSLP served as analyte at concentrations of 3.2nM to 0.2nM with two-fold dilutions (**Figures 5E-F**). Representative SPR sensorgrams of BDG33.023 and BDG33.025 binding to human IL-13 (h-IL-13) are presented in **Figures 5G and 5H**. hIL-13 served as analyte at concentrations of 20nM to 0.6nM with tow fold dilutions (**Figures 5G-5H**).

[55] **Figures 6A-6E** present ELISA EC50 binding of BDG33.023 and BDG33.025 to human TSLP, cytomegaly monkey (cyno) TSLP or cytomegaly monkey (cyno) IL-13. Binding of BDG33.023 (filled circles) and BDG33.025 (filled squares) to human TSLP (**Figure 6A-human TSLP**). Binding of BDG33.023 to cyno-TSLP (**Figures 6B-33.023 cyno TSLP**). Binding of BDG33.025 to cyno-TSLP (**Figures 6C-33.025 cyno TSLP**). Binding of BDG33.023 to cyno-IL-13 (**Figures 6D-33.023 cyno IL-13**). Binding of BDG33.025 to cyno-IL-13 (**Figures 6E-33.025 cyno IL-13**).

[56] **Figures 7A-7D** present competitive binding assay of antibodies to hTSLP or hIL-13. **Figure 7A:** Indicated antibodies (anti-TSLP-control; anti-IL-13 control; BDG33.023; BDG33.025) were pre-incubated with increasing levels of hIL-13 and added to a plate that was pre-coated with hIL-13. BDG33.023 and BDG33.025 binding to plate-bound hIL-13 was inhibited as soluble hIL-13 concentration increased. **Figure 7B:** Indicated antibodies (anti-TSLP-

control; anti-IL-13 control; BDG33.023; BDG33.025) were pre-incubated with increasing levels of hTSLP and added to a plate that was pre-coated with hTSLP. BDG33.023 and BDG33.025 binding to plate-bound hTSLP was inhibited as soluble hTSLP concentration increased. **Figure 7C:** Antibodies (anti-TSLP-control; anti-IL-13 control; BDG33.023; BDG33.025) were pre-incubated with increasing levels of hTSLP and added to a plate pre-coated with hIL-13, BDG33.023 binding to IL-13 was inhibited as soluble hTSLP concentration increased. **Figure 7D:** Indicated antibodies (anti-TSLP-control; anti-IL-13 control; BDG33.023; BDG33.025) were pre-incubated with increasing levels of hIL-13 and added to a plate pre-coated with hTSLP. BDG33.023 binding to plate-bound hTSLP was inhibited as soluble hIL-13 concentration increased. Anti-TSLP and anti-IL-13 control antibodies only showed binding with their respective ligands, and only competed with their respective ligands.

[57] **Figure 8** presents the results of an ELISA specificity test that compared non-specific binding to specific binding of BDG330.23 and BDG33.025. The ELISA plate was coated with hIL-13, hTSLP, and the non-related cytokines IL-2, IL-17, and IL-4. BSA binding signal corresponds to assay background level.

[58] **Figure 9** presents the results of an IC₅₀ inhibition assay that measured IgG specific blocking of hTSLP from binding to an ELISA plate coated with TSLP receptor (TSLP-R). The X-axis represents concentration of competitor. Competitors: TSLP-R (black circles) with a resultant IC₅₀= 3nM; and BGD33.023 (black triangles) with a resultant IC₅₀=0.41nM.

[59] **Figure 10** presents a schematic representation of the HEK-Blue IL-13 system downstream signaling.

[60] **Figures 11A-11D** present hIL-13 pSTAT6 signaling inhibition data. The results are based on stimulation of HEK-Blue cell's IL-13 activation pathway by recombinant rh-IL-13 and inhibition of this stimulation by indicated IgGs. HEK-Blue IL-13 cells (50,000cells/well) were incubated with rh-IL-13 at a range of concentrations (0nM-8nM). IL-13 downstream signaling was quantified with QUANTI-Blue 24h post incubation (**Figure 11A**). hIL-13 downstream inhibition on HEK-BLUE IL-13 cells by engineered dual binding antibodies was analyzed as follows. rh-IL-13 (0.4 nM) was incubated with indicated antibodies at an antibody concentration range of 0nM-750nM . Antibodies assayed were BDG33.002 (positive control), BDG33.003 (Clone C2), and BDG33.006 (negative control), respectively (**Figure 11B**). Clones BDG33.023 and BDG33.025 were assayed at an antibody concentration range of 0nM-100nM (**Figures 11C** and **11D** show 33.023 IL-13 pSTAT6 inhibition, and 33.025 IL-13 pSTAT6 inhibition, respectively. After the incubation the hIL-13/IgG mixture was added to the cells, secreted embryonic alkaline phosphatase (SEAP) activity was quantified with QUANTI-Blue 24h post

incubation. Data shown is the mean of triplicate experiments, and error bars represent standard deviation.

[61] **Figures 12A-12C** present TSLP signaling pathway inhibition data. TSLP dependent pSTAT5 signaling activation pathway, and inhibition of the activation by BDG 33.023 in human leukemia MUTZ5 cells. Figure 12A shows flow-cytometry analysis of MUTZ 5 CD127 (IL-7a) receptor and TSLP-R receptor expression, as follows. Unstained cells (panel a), cells stained for CD127+ wherein approximately 36% of the total cell population was labeled (panel b), cells stained for TSLP-R+, wherein approximately 96% of the total cell population was labeled (panel c), and cells stained for both TSLP-R+ and CD127+ wherein approximately 41% of total cell population was labeled (panel d) (**Figure 12A**). **Figure 12B** shows MUTZ5 pSTAT5 activation. EC₅₀ of hTSLP phosphor-STAT5 (pSTAT5) activation in MUTZ5 cells. Percent (%) positive cells represents pSTAT5 positive cells as a percentage of the parent population. **Figure 12C** shows inhibition of MUTZ5 pSTAT5 activation. IC₅₀ of BDG33.023 inhibition of TSLP dependent pSTAT5 activation in MUTZ5 cells. TSLP was pre-incubated for 30min with 0.48pM to 500pM of BDG33.023 and added to MUTZ5 cells. Positive cells are representing pSTAT5 positive population as a percentage of the parent population (**Figure 12C**).

[62] **Figure 13** shows retention time and calculated pI for some of the dual binding antibodies disclosed herein. IgG marker retention time was 4.77 min.

[63] **Figures 14A-14C** shows competitive ELISA of some of the dual binding antibodies and 33.001 (Tezepelumab) over TSLP. ELISA plates were coated over night at 4°C with 50 ng/well of 33.001. Dual binding antibodies were double diluted and pre-incubated with 7nM constant concentration of TSLP-HIS for 1 hour at room temperature. After blocking and washing steps, the dual binding antibodies-TSLP mix were subjected over the plates, incubated for 10 minutes and washed again before 30 minutes incubation with anti-HIS. The results show all tested dual binding antibodies presented a similar IC₅₀.

[64] **Figure 15** shows results of nanoscale differential scanning fluorimetry (nanoDSF) analysis of some of the dual binding antibodies disclosed herein. T_m threshold for lambda chain was >65°C and T-onset >60°C.

[65] **Figures 16A-16F** show the results of SPR (Surface Plasmon Resonance) analysis for some of the dual binding antibodies disclosed herein for human IL-13 and TSLP.

[66] **Figures 17A and B** show size exclusion chromatography (SEC) scans (**Figure 17A**), and nano-differential scanning fluorimetry (DSF) analysis of the melting point (**Figure 17B**) for antibody BDG38.074. Representative analysis of the melting point of indicated IgGs were analyzed in duplicate. Light gray dashed line represents the T-onset and bold gray dashed lines

represents the Tm1 and Tm2. **Figure 17B** shows the 1st derivative of the measurement. DSF values are summarized in **Figure 15**.

[67] **Figures 18A – 18D** show binding affinities of representative clone BDG38.74. **Figure 18A** shows binding affinities of antibody BDG38.074 to IL-13. The results show antibody BDG38.074 binds human IL-13 with double digit picomolar affinities. **Figure 18B** shows binding affinities of antibody BDG38.074 to human TSLP. The results show antibody BDG38.074 binds human TSLP with double digit picomolar affinities. **Figure 18C** shows binding affinities of antibody BDG38.074 to cyno IL-13. **Figure 18D** shows binding affinities of antibody BDG38.074 to cyno TSLP.

[68] **Figures 19A and 19B** shows size exclusion chromatography (SEC) scans (**Figure 19A**), and nano-differential scanning fluorimetry (DSF) analysis of the melting point (**Figure 19B**) for antibody BDG38.079. Shown representative DSF analysis of the melting point of indicated IgGs (analyzed in duplicates). Light gray dashed line represents the T-onset and bold gray dashed lines represents the Tm1 and Tm2. **Figure 19B** is a graph of the 1st derivative of the measurement. DSF values are summarized in **Figure 15**.

[69] **Figures 20A-20D** show IL-13 and TSLP binding of representative clone BDG38.079. **Figure 20A** shows binding affinities of antibody BDG38.079 to human IL-13. The results show antibody BDG38.079 binds human IL-13 with double digit picomolar affinities. **Figure 20B** shows binding affinities of antibody BDG38.079 to human TSLP. The results show antibody BDG38.079 binds human TSLP with single digit picomolar affinities. **Figure 20C** shows binding affinities of antibody BDG38.079 to cyno IL-13. **Figure 20D** shows binding affinities of antibody BDG38.079 to cyno TSLP.

[70] **Figures 21A and 21B** show SPR (Surface Plasmon Resonance) analysis of antibodies BDG38.074 and BDG38.079 for human or cyno IL-13 or TSLP.

[71] **Figure 22** shows antibodies BDG38.074 and BDG38.079 inhibit IL-13 function in HEK reporter cell line with double digit picomolar affinity. hIL-13 pSTAT6 signaling inhibition data. The results are based on stimulation of HEK-Blue cell's IL-13 activation pathway by recombinant rh-IL-13 and inhibition of this stimulation by indicated IgGs. rh-IL-13 (0.4 nM) was incubated with indicated antibodies at an antibody concentration range of 0nM-100nM After the incubation the hIL-13/IgG mixture was added to the cells, secreted embryonic alkaline phosphatase (SEAP) activity was quantified with QUANTI-Blue 24h post incubation. Data shown is the mean of triplicate experiments, and error bars represent standard deviation. Antibodies assayed were Tralokinumab, BDG38.074 and BDG38.079 respectively.

[72] **Figure 23** shows antibodies BDG38.074 and BDG38.079 exhibit similar functional

inhibition to anti-TSLP benchmarks in MUTZ-5 cell line. MUTZ5 cells were stimulated with human TSLP (hTSLP) and phosphorylated STAT5 (pSTAT5) staining was evaluated by phospho-flow cytometry.

[73] **Figures 24A and 24B** show inhibition results for representative clones **Figure 24A** shows antibodies BDG38.074 and BDG38.079 exhibit similar inhibition of CD23 expression to the anti-IL-13 benchmark (Tralokinumab). IC₅₀ of antibody inhibition of IL-13 was determined by measuring CD23 expression level in monocytes. At the end of 48 hours incubation of the cells with different concentrations of antibodies, monocytes were detached from the bottom of the wells and stained with CD3 (Bio Legend, CAT: 300450), CD14 (Bio Legend, CAT: 301814), CD19 (Bio Legend, CAT: 302212) and CD23 (Bio Legend, CAT: 338506) antibodies. CD23 percentage of CD14+ population was measured using CytoFLEX flow cytometer (Beckman Coulter). **Figure 24B** shows antibodies BDG38.074 and BDG38.079 inhibit TARC expression similarity to anti-TSLP benchmark (Tezepelumab). IC₅₀ of antibody inhibition of hTSLP was determined by TARC inhibition. TARC levels were determined using TARC DUOSET ELISA kit DY364 (R&D systems) according to kit instructions. ELISA plates were read at 450nm. Values were analyzed using standard sample curve.

[74] **Figure 25A and 25B** shows size exclusion chromatography (SEC) scans (**Figure 25A**), and nano-differential scanning fluorimetry (DSF) analysis of the melting point (**Figure 25B**) for antibody BDG38.094. Shown representative DSF analysis of the melting point of indicated IgGs (analyzed in duplicates). Light gray dashed line in the upper graph represents the T-onset and bold gray dashed lines represents the T_{m1} and T_{m2}. The **Figure 25B** is the 1st derivative of the measurement. DSF values are summarized in **Table 11**

[75] **Figures 26A-26D** show binding affinities of a representative clone. **Figure 26A** shows binding affinities of antibody BDG38.094 to human IL-13. **Figure 26B** shows binding affinities of antibody BDG38.094 to human TSLP. **Figure 26C** shows binding affinities of antibody BDG38.094 to cyno IL-13. **Figure 26B** shows binding affinities of antibody BDG38.094 to cyno TSLP.

[76] **Figures 27A and 27B** shows size exclusion chromatography (SEC) scans (**Figure 27A**), and nano-differential scanning fluorimetry (DSF) analysis of the melting point (**Figure 27B**) for antibody BDG38.138. Shown representative DSF analysis of the melting point of indicated IgGs (analyzed in duplicates). Light gray dashed line in the upper graph represents the T-onset and bold gray dashed lines represents the T_{m1} and T_{m2}. **Figure 27B** is the 1st derivative of the measurement. DSF values are summarized in **Table 11**.

[77] **Figures 28A-28D** show binding affinities of a representative clone. **Figure 28A** shows

binding affinities of antibody BDG38.138 to human IL-13. **Figure 28B** shows binding affinities of antibody BDG38.138 to human TSLP. **Figure 28C** shows binding affinities of antibody BDG38.138 to cyno IL-13. **Figure 28D** shows binding affinities of antibody BDG38.138 to cyno TSLP.

DETAILED DESCRIPTION

[78] In the following detailed description, numerous specific details are set forth in order to provide a thorough understanding of the engineered dual binding antibodies disclosed herein, including a description of their heavy chain and light chain variable regions. However, it will be understood by those skilled in the art that preparation and use dual binding antibodies may in certain cases, be practiced without these specific details. In other instances, well-known methods, procedures, and components have not been described in detail so as not to obscure the disclosure presented herein.

[79] Antigen binding sequences are conventionally located within the heavy chain and light chain variable region sequences of an antibody. These heavy and light chain variable regions may, in certain instances, be manipulated to create new binding sites, for example to create antibodies or fragments thereof, that bind to a different antigen or an epitope of a different antigen thereof. In some embodiments, as described herein, manipulating the sequence of a heavy chain variable region or the sequence of a light chain variable region, or both, creates a new binding site for an epitope while maintaining antibody functionality. In one embodiment, twenty-one specific sites within the heavy and light chain variable regions are identified, wherein the presence of variant amino acids at these sites, in certain embodiments, creates an engineered dual binding antibody or fragment thereof. In some embodiments, the 21 potential variant sites provide a unique platform from which to engineer dual binding antibodies or fragments thereof.

[80] Disclosed herein are engineered dual binding antibodies or fragments thereof, wherein either a heavy chain variable region, or a light chain variable region, or both, have been mutated to include variant amino acids. In some embodiments, these engineered dual binding antibodies may be identified and selected from a library created to include variant amino acid residues at particular sites within a variable heavy chain or variable light region, or both. In some embodiments, these engineered dual binding antibodies may be produced by specifically mutating target amino acid sites within a variable heavy chain or variable light region, or both. In some embodiments, these engineered dual binding antibodies may be used in a therapeutic method for treating a subject suffering from an allergic or respiratory condition.

Engineered Dual Binding Antibodies

[81] As used herein, the term “dual binding antibodies” refers to antibodies that have two binding specificities. In certain embodiments, the dual binding antibodies disclosed herein bind to IL-13 and TSLP.

[82] In some embodiments, the present disclosure provides an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3) (see e.g. Tables 8 and 9). In some embodiments, the CDRs have the sequences of SEQ ID NOs:149-154. In some embodiments, the dual binding antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL) having the amino acid sequences of SEQ ID Nos:155 and 156, or SEQ ID Nos:157 and 158.

[83] In one embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:359, 360 and 361 respectively.

[84] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 371 respectively.

[85] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:362, 360 and 384 respectively.

[86] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 384 respectively.

[87] In another embodiment, the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.

[88] In some embodiments, disclosed herein is an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein

- (i) the HCDR1 comprises the amino acid sequence of SEQ ID NO:349 or 355, or the amino acid sequence of SEQ ID NO:149 or SEQ ID NO:136;
- (ii) the HCDR2 comprises the amino acid sequence of one of SEQ ID NOs:350, 352, 354 and 356, or the amino acid sequence of SEQ ID NO:150 or the sequence set forth as: I HX1 Y D G S N K (SEQ ID NO:142), wherein HX1 is any amino acid;

(iii) the HCDR3 comprises the amino acid sequence of one of SEQ ID NOs:351, 353, 357, and 358, or the amino acid sequence of SEQ ID NO:151 or the sequence set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO:143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid;

(iv) the LCDR1 comprises the amino acid sequence of one of SEQ ID NOs:359, 362, 364, 366, 369, and 375, or the amino acid sequence of SEQ ID NO:152 or the sequence set forth as LX1, LX2, G S K LX3 V (SEQ ID NO:144), wherein LX1, LX2, and LX3 are any amino acid;

(v) the LCDR2 comprises the amino acid sequence of SEQ ID NO:360 or 367, or the amino acid sequence of SEQ ID NO:153 or the sequence set forth as D D LX4 (SEQ ID NO:145), wherein LX4 is any amino acid; and

(vi) the LCDR3 comprises the amino acid sequence of one of SEQ ID NOs:361, 363, 365, 368, 370-374, 376-407, or the amino acid sequence of SEQ ID NO:154 or the sequence set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO:146), wherein LX5, LX6, and LX7 are any amino acid.

[89] In some embodiments, disclosed herein is an isolated dual binding antibody wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences of SEQ ID NOs:359, 360 and 361 respectively; or

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences of SEQ ID NOs:364, 360 and 371 respectively; or

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences of SEQ ID NOs:362, 360 and 384 respectively; or

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences of SEQ ID NOs:364, 360 and 384 respectively.

[90] In some embodiments, disclosed herein is an isolated dual binding antibody, wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.

[91] In some embodiments, disclosed herein is an isolated dual binding antibody, wherein HX1 is W or S; HX2 is A or S; HX3 is P; HX4 is Q; HX5 is W; HX6 is E, Q, M, L, or V; HX7

is L, W, or Y; HX8 is V or T; HX9 is H, A, or S; HX10 is E; HX11 is A; HX12 is I, L, or M; wherein LX1 is N, L, or I; LX2 is L or I; LX3 is S or L; LX4 is S or G; LX5 is S or T; LX6 is S or G; LX7 is H or G.

[92] In some embodiments, disclosed herein is an isolated dual binding antibody, wherein HX1 is W, HX2 is A or S, HX6 is E or M, HX7 is L or W, HX8 is V or T, HX9 is H or A, HX12 is I or L, LX1 is L, LX2 is I, LX3 is L, LX4 is S or G, LX5 is S, LX6 is S, LX7 is H or G.

[93] In some embodiments, disclosed herein is an isolated dual binding antibody, wherein

(a) HX1 is W, HX2 is A, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX12 is I, LX4 is S, and LX7 is G; or

(b) HX1 is W, HX2 is A, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX12 is L, LX4 is S, and LX7 is H; or

(c) HX1 is W, HX2 is S, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX12 is L, LX4 is G, and LX7 is G.

[94] In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any position or a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any position, or a combination thereof, wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or said combination thereof, is at least 2. In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2, wherein there are at least two amino acid variants within the heavy chain variable region or the light chain variable region or the combination thereof. In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least two amino acid variants at any position and any light chain variable region. In some embodiments, disclosed herein is an isolated dual binding antibody comprising a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least two amino acid variants at any position and any heavy chain variable region.

[95] As used herein, the term “heavy chain variable region” may be used interchangeable with the term “VH domain” or the term “VH”, having all the same meanings and qualities. As used herein, the term “light chain variable region” may be used interchangeable with the term “VL domain” or the term “VL”, having all the same meanings and qualities.

[96] In certain embodiments, a specific variant VH and/or VL domain, described herein, may be used to screen a library of the complementary variable region to identify VH/VL, respectively, with desirable properties, such as increased affinity for an antigen. Such methods are described, for example, in Portolano et al., J. Immunol. (1993) 150:880-887; Clarkson et al., Nature (1991) 352:624-628. Fischer et al., (2015) Exploiting light chains for the scalable generation and platform purification of native human bispecific IgG. *Nature Communications* volume 6, Article number: 6113.

[97] Other methods may also be used to mix and match VH and VL domains to identify a Fab or F'(ab)₂ having desired dual binding activity. For example: Klimka et al., British Journal of Cancer (2000) 83: 252-260, describe a screening process using a mouse VL and a human VH library with CDR3 and FR4 retained from the mouse VH. After obtaining antibodies, the VH was screened against a human VL library to obtain antibodies that bound antigen. Beiboer et al., J. Mol. Biol. (2000) 296:833-849 describe a screening process using an entire mouse heavy chain and a human light chain library. After obtaining antibodies, one VL was combined with a human VH library with the CDR3 of the mouse retained. Antibodies capable of binding antigen were obtained. Rader et al., PNAS (1998) 95:8910-8915 describe a process similar to Beiboer et al above.

[98] These just-described techniques are, in and of themselves, known as such in the art. The skilled person will, however, be able to use such techniques to obtain antigen-binding fragments of antibodies according to several embodiments of the disclosure described herein, using routine methodology in the art.

[99] A skilled artisan would appreciate that dual binding antibody encompasses in its broadest sense an antibody that specifically binds an antigenic determinant of IL-13 and TSLP. The skilled artisan would appreciate that specificity for binding to IL-13 or TSLP reflects that the binding is selective for the antigen and can be discriminated from unwanted or nonspecific interactions. In certain embodiments, the dual binding antibody comprises an antibody fragment or fragments.

[100] In some embodiments, an antigenic determinant comprises an IL-13 or TSLP epitope. The term "epitope" includes any determinant, in certain embodiments, a polypeptide determinant, capable of specific binding to an anti-IL-13 or anti-TSLP binding domain. An epitope is a region of an antigen that is bound by an antibody or an antigen-binding fragment thereof. In some embodiments, the antigen-binding fragment of an antibody comprises a heavy chain variable region, a light chain variable region, or a combination thereof as described herein.

[101] In certain embodiments, epitope determinants include chemically active surface

groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl, and may in certain embodiments have specific three-dimensional structural characteristics, and/or specific charge characteristics. In certain embodiments, the dual binding antibody is said to specifically bind an IL-13 or TSLP epitope when it preferentially recognizes IL-13 or TSLP in a complex mixture of proteins and/or macromolecules. The dual binding antibody is said to specifically bind an epitope when the equilibrium dissociation constant is $\leq 10^{-5}$, 10^{-6} , or 10^{-7} M. In some embodiments, the equilibrium dissociation constant may be $\leq 10^{-8}$ M or 10^{-9} M. In some further embodiments, the equilibrium dissociation constant may be $\leq 10^{-10}$ M, 10^{-11} M, or 10^{-12} M. In some embodiments, the equilibrium dissociation constant may be in the range of $\leq 10^{-5}$ M to 10^{-12} M.

[102] An antibody binding domain can be a fragment of an antibody or a genetically engineered product of one or more fragments of the antibody, which fragment is involved in specifically binding with the antigen. By "specifically binding" is meant that the binding is selective for the antigen of interest, for example for IL-13 or TSLP in embodiments described herein and can be discriminated from unwanted or nonspecific interactions. As used herein, the term "dual binding antibody" may in certain embodiments, encompass complete immunoglobulin structures, fragments thereof, or domains thereof.

[103] Examples of antibody binding domains include, without limitation, a complementarity determining region (CDR), a variable region (Fv), a VH domain, a light chain variable region (VL), a heavy chain, a light chain, a single chain variable region (scFv), and a Fab fragment. A skilled artisan would appreciate that an scFv is not actually a fragment of an antibody, but instead is a fusion polypeptide comprising the variable heavy chain (VH) and variable light chain (VL) regions of an immunoglobulin, connected by a short linker peptide of for example but not limited to ten to about 25 amino acids. The skilled artisan would also appreciate that the term "Fab" with regard to an antibody, generally encompasses that portion of the antibody consisting of a single light chain (both variable and constant regions) bound to the variable region and first constant region of a single heavy chain by a disulfide bond.

[104] In some embodiments, an antibody encompasses whole antibody molecules, including monoclonal, polyclonal and multispecific (e.g., bispecific) antibodies. In some embodiments, an antibody encompasses an antibody fragment or fragments that retain binding specificity including, but not limited to, variable heavy chain (VH) fragments, variable light chain (VL) fragments, Fab fragments, F(ab')₂ fragments, scFv fragments, Fv fragments, minibodies, diabodies, triabodies, and tetrabodies (see, e.g., Hudson and Souriau, *Nature Med.* 9: 129-134 (2003) (hereby incorporated by reference in their entirety)). Also encompassed are humanized, primatized, and chimeric antibodies.

[105] As used herein, in some embodiments, the term “Antibody” may be used interchangeably with the term “Immunoglobulin” having all the same qualities and meanings. Similarly, as used herein, in some embodiments, the term “Antibody or fragments thereof” may be used interchangeably with the term “Immunoglobulin or fragments thereof” having all the same qualities and meanings. Thus, a skilled artisan would appreciate that in some embodiments, “an antibody or fragments thereof”, or “immunoglobulins or fragments thereof” may encompass IgG immunoglobulins or fragments thereof or structures comprising a fragment or fragments thereof, including but not limited to an IgG, an scFv fragment, an Fab fragment, an F(ab')₂ fragment, Fv fragments, minibodies, diabodies, triabodies, and tetrabodies.

[106] A skilled artisan would recognize that a “Heavy chain variable region” or “VH” with regard to an antibody encompasses the fragment of the heavy chain that contains three CDRs interposed between flanking stretches known as framework (FR) regions, which are more highly conserved than the CDRs, and form a scaffold to support the CDRs. In certain embodiments, the terms a “Heavy chain variable region” or a “VH” may be used interchangeably with “VH domain”.

[107] A skilled artisan would recognize that a “Light chain variable region” or “VL” with regard to an antibody encompasses the fragment of the light chain that contains three CDRs interposed between framework (FR) regions. In certain embodiments, the terms a “Light chain variable region” or a “VL” may be used interchangeably with “VL domain”.

[108] Disclosed herein are a number of amino acid sequences for HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, LCDR3, as well as VH and VL regions for dual binding antibodies that bind to IL-13 and TSLP. Discussion on some embodiments of representative sequences disclosed herein is presented below. **Figure 1A** presents the template VH domain amino acid sequence, set forth in SEQ ID NO: 1, and the location of the three heavy-chain (H) CDR regions (HCDR1, HCDR2, HCDR3) and four FR regions (HFR1, HFR2, HFR3, HFR4), while **Figure 1B** presents the template VL domain amino acid sequence, set forth in SEQ ID NO: 2, and the location of the three light-chain (L) CDR regions (LCDR1, LCDR2, LCDR3) and four FR regions (LFR1, LFR2, LFR3, LFR4). The amino acid residues including variant residues, present in each of the CDR regions and each of the FR regions of the re-epitoped clones, are clearly identified by comparing the linear schematic representation of the template VH or template VL sequence with the numbering and amino acids provided below (**Figures 1A and 1B**).

[109] In some embodiments, an isolated dual binding antibody comprises an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of

HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid. A skilled artisan would recognize the 12 unique sites within the VH domain presented in **Figure 1A**, wherein a variant amino acid may be found, which are herein identified as HX and may in certain embodiments, encompass the presence of a variant amino acid within the template sequence of the heavy chain.

[110] In some embodiments, a VH domain of a dual binding antibody comprises HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 142), and HCDR3 (SEQ ID NO: 143), wherein the VH domain comprises a variant amino acid at, at least one of HX1, HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12.

[111] In some embodiments, a VH domain of a dual binding antibody comprises HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is selected from the group consisting of W and S; and HCDR3 (SEQ ID NO: 138) wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M. In certain embodiments, the isolated dual binding antibody comprises variant amino acids comprising CDR1 (SEQ ID NO: 136), CDR2 (SEQ ID NO: 137) wherein HX1 is W, CDR3 (SEQ ID NO: 138) wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX10 is E, HX11 is A, and HX12 is selected from the group I and L.

[112] In some embodiments, the dual binding antibody may have a VH domain comprising an HCDR1 (SEQ ID NO: 136), an HCDR2 (SEQ ID NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX10 is E, HX11 is A, and HX12 is I; or an HCDR1 (SEQ ID NO: 136), an HCDR2 (SEQ ID NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX10 is E, HX11 is A, and HX12 is L; or an HCDR1 (SEQ ID NO: 136), an HCDR2 (SEQ ID

NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is S, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX10 is E, HX11 is A, and HX12 is L.

[113] Engineered antibody clones having variants in the VH domain as described above, are presented in **Figure 1A**.

[114] In some embodiments, an isolated dual binding antibody comprises an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said in some embodiments the VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid. A skilled artisan would recognize the 7 unique sites within the VL domain presented in **Figure 1B**, wherein a variant amino acid may be found within a CDR, which are herein identified as LX and may in certain embodiments, encompass the presence of a variant amino acid within the template sequence of the light chain.

[115] In some embodiments, a variant amino acid within the light chain may reside in one of the framework regions. In some embodiments, a variant amino acid within the VL domain is in the LFR3 region.

[116] In some embodiments, a VL domain of the dual binding antibody comprises LCDRs wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; and wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G. In certain embodiments, the isolated dual binding antibody comprises variant amino acids wherein LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is selected from the group consisting of S and G, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is selected from the group consisting of H and G.

[117] In some embodiments, the dual binding antibody may have a VL domain comprising an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID

NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G; or an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is H, or an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID NO: 140) wherein LX4 is G, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G.

[118] Engineered antibody clones having variants in the VL domain as described above, are presented in **Figure 1B**.

[119] In some embodiments, disclosed herein is an isolated antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), said VH and VL comprise the amino acid sequences of SEQ ID Nos:209 and 210, SEQ ID Nos:219 and 220, SEQ ID Nos:249 and 250, SEQ ID Nos:337 and 338, SEQ ID Nos:155 and 156, SEQ ID Nos:157 and 158, SEQ ID Nos:4 and 3, SEQ ID Nos:6 and 5, SEQ ID Nos:8 and 7, SEQ ID Nos:10 and 9, SEQ ID Nos:12 and 11, SEQ ID Nos:14 and 13, SEQ ID Nos:16 and 15, SEQ ID Nos:18 and 17, SEQ ID Nos:20 and 19, SEQ ID Nos:22 and 21, SEQ ID Nos:24 and 23, SEQ ID Nos:26 and 25, SEQ ID Nos:28 and 27, SEQ ID Nos:30 and 29, SEQ ID Nos:32 and 31, SEQ ID Nos:34 and 33, SEQ ID Nos:36 and 35, SEQ ID Nos:38 and 37, SEQ ID Nos:40 and 39, SEQ ID Nos:42 and 41, SEQ ID Nos:44 and 43, SEQ ID Nos:46 and 45, SEQ ID Nos:48 and 47, SEQ ID Nos:50 and 49, SEQ ID Nos:52 and 51, or SEQ ID Nos:54 and 53.

[120] In some embodiments, disclosed herein is an isolated antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), said antibody comprising the sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99% identical) to the sequences set forth in any of SEQ ID Nos:209 and 210, SEQ ID Nos:219 and 220, SEQ ID Nos:249 and 250, SEQ ID Nos:337 and 338, SEQ ID Nos:155 and 156, SEQ ID Nos:157 and 158, SEQ ID Nos:4 and 3, SEQ ID Nos:6 and 5, SEQ ID Nos:8 and 7, SEQ ID Nos:10 and 9, SEQ ID Nos:12 and 11, SEQ ID Nos:14 and 13, SEQ ID Nos:16 and 15, SEQ ID Nos:18 and 17, SEQ ID Nos:20 and 19, SEQ ID Nos:22 and 21, SEQ ID Nos:24 and 23, SEQ ID Nos:26 and 25, SEQ ID Nos:28 and 27, SEQ ID Nos:30 and 29, SEQ ID Nos:32 and 31, SEQ ID Nos:34 and 33, SEQ ID Nos:36 and 35, SEQ ID Nos:38 and 37, SEQ ID Nos:40 and 39, SEQ ID Nos:42 and 41, SEQ ID Nos:44 and 43, SEQ ID Nos:46 and 45, SEQ ID Nos:48 and 47, SEQ ID Nos:50 and 49, SEQ ID Nos:52 and 51, or SEQ ID Nos:54 and 53.

[121] In some embodiments, disclosed herein is an isolated, wherein the antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), said VH and VL comprise the amino acid sequences as shown in Table 10 or Table 1. In some embodiments,

disclosed herein is an isolated antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), said antibody comprising the sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99% identical) to the sequences set forth in Table 10 or Table 1.

[122] In some embodiments, disclosed herein is an isolated antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein

(a) said VH domain comprises the amino acid sequence set forth in SEQ ID NO: 1 with amino acid variants at two or more of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and

(b) said VL domain comprises the amino acid sequence set forth in SEQ ID NO: 2 with amino acid variants at two or more of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof).

[123] In one embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:209 and 210.

[124] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:219 and 220.

[125] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:249 and 250.

[126] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:337 and 338.

[127] In another embodiment, the isolated dual binding antibody disclosed herein comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences as shown in Table 1 or Table 10.

[128] In another embodiment, the isolated dual binding antibody disclosed herein comprises VH and VL sequences that are at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH and VL sequences disclosed herein.

[129] In some embodiments, an isolated dual binding antibody comprising an antibody

antigen-binding domain site comprising a VH domain and a VL domain comprising a combination of VH domain HCDRs and VL domain LCDRs described above. For example, but not limited to, in certain embodiments, a VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid; and wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid.

[130] In some embodiments, a VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136, wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is selected from the group consisting of W and S; wherein the amino acid sequence of HCDR3 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M; and a VL domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G.

[131] In certain embodiments, the isolated dual binding antibody comprises a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of

HCDR1 is set forth in SEQ ID NO: 136, wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is W, wherein the amino acid sequence of HCDR3 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, HX3 is P, HX4 is Q, HX5 is W, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX10 is E, HX11 is A, HX12 is selected from the group I and L, and comprising a VL domain comprising a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139 wherein LX1 is L, LX2 is I, LX3 is L, wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140 wherein LX4 is selected from the group consisting of S and G, wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141 wherein LX5 is S, LX6 is S, and LX7 is selected from the group consisting of H and G.

[132] In some embodiments, the dual binding antibody comprises a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, and a VL domain comprising a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequences of each CDR are as set forth in Figures 1A and 1B for the clones set forth there, for example but not limited to:

Clone C2: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX10 is E, HX11 is A, HX12 is I, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G;

Clone C6: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX10 is E, HX11 is A, HX12 is L, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is H; or

Clone C9: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is S, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX10 is E, HX11 is A, HX12 is L, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is G, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G.

[133] In some embodiments, the dual binding antibody comprises a set of HCDRs as disclosed herein, and any VL domain. In some embodiments, the dual binding antibody comprises a set of LCDRs as disclosed herein, and any VH domain. In some embodiments, the dual binding

antibody comprises a paired set of HCDRs -LCDRs, as disclosed herein.

[134] In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs as described herein may be encoded by a nucleic acid construct. In certain embodiments, the dual binding antibody comprising a VL domain comprising LCDRS as described herein may be encoded by a nucleic acid construct. In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs and a VL domain comprising HCDRs as described herein, may be encoded by a nucleic acid construct.

[135] In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs as described herein, may be encoded by a nucleic acid construct. In certain embodiments, the dual binding antibody comprising a VL domain comprising LCDRS as described herein may be encoded by a nucleic acid construct. In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs and a VL domain comprising HCDRs as described herein, may be encoded by a nucleic acid construct.

[136] In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs as described herein, may be comprised within a library of immunoglobulins. In certain embodiments, the dual binding antibody comprising a VL domain comprising LCDRS as described herein may be comprised within a library of immunoglobulins. In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs and a VL domain comprising HCDRs as described herein, may be comprised within a library of immunoglobulins.

[137] In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs as described herein, may be produced by expressing a nucleic acid construct comprising a nucleic acid sequence encoding the HCDRs from a host cell and isolating the antibody. In certain embodiments, the dual binding antibody comprising a VL domain comprising LCDRS as described herein may be produced by expressing a nucleic acid construct comprising a nucleic acid sequence encoding the LCDRS from a host cell and isolating the antibody. In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs and a VL domain comprising HCDRs as described herein, may be produced by expressing a nucleic acid construct comprising a nucleic acid sequence encoding the HCDRs and LCDRS from a host cell and isolating the antibody.

[138] In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs as described herein may be administered in a method of treating a subject in need, wherein said subject suffers from a disease or condition comprising an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs;

scleroderma; or tumors or cancers including Hodgkin's lymphoma. In certain embodiments, the dual binding antibody comprising a VL domain comprising LCDRs as described herein may be administered in a method of treating a subject in need, wherein said subject suffers from a disease or condition comprising an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma. In certain embodiments, the dual binding antibody comprising a VH domain comprising HCDRs and a VL domain comprising LCDRs as described herein, may be administered in a method of treating a subject in need, wherein said subject suffers from a disease or condition comprising an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma.

[139] In some embodiments, an antibody comprising a heavy chain variable region amino acid sequence set forth in SEQ ID NO: 1 or a light chain variable region amino acid sequence set forth in SEQ ID NO: 2, or the combination thereof, does not bind an IL-13 epitope. Accordingly, the dual binding antibodies described herein are engineered to comprise a binding region not previously present in the antibody. In other words, the dual binding antibodies comprise “re-epitoped” antibodies. As used throughout, the term “engineered” and “re-epitoped” may in certain embodiments, be used interchangeably having all the same qualities and meanings. In some embodiments, a “re-epitoped” antibody comprises improved binding compared to available antibodies. In some embodiments, a “re-epitoped” antibody comprises improved association and dissociation constants (K_{on} and K_{off}), compared to the parent antibodies. In some embodiments, a “re-epitoped” antibody comprises improved stability compared with the parent antibodies. In certain embodiments, incorporating variant amino acid residues in at least two of the unique set of 21 variant sites within the CDRs and FR of the VH domain and VL domain, as described herein, results in a “re-epitoped” dual binding antibody comprising improved characteristics compared with the parent antibodies. These re-epitoped antibodies may provide advantageous characteristics.

[140] A skilled artisan would recognize that a “Fv” with regard to an antibody encompasses the smallest fragment of the antibody to bear the complete antigen binding site. An Fv fragment consists of the variable region of a single light chain (VL) bound to the variable region of a single heavy chain (VH).

[141] A skilled artisan would recognize that a “single-chain Fv antibody” or “scFv” with regard to an antibody encompasses an engineered antibody consisting of a VL domain and a VH domain connected to one another directly or via a peptide linker sequence. The skilled artisan

would appreciate that a linker, in some embodiments, may comprise a linear amino acid sequence. In some embodiments, the linear amino acid sequence (“linker”) comprises an enzyme cleavage site and may, in certain embodiments, be termed a “cleavable linker” or a “linker” or a “cleavable peptide”. In some embodiments, a linker may be a cleavable linker. In some embodiments, a linker may be a non-cleavable. In some embodiments, a linker sequence is set forth in SEQ ID NO: 147 (GGGGSGGGSGGGGS; SEQ ID NO: 147).

[142] In some embodiments, peptide linker sequences contain, for example, Gly, Asn, and/or Ser residues, in various combinations. Other near neutral amino acids, such as Thr and Ala, may also be included in the linker sequence.

[143] Other amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46 (1985); Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262 (1986); U.S. Pat. No. 4,935,233 and U.S. Pat. No. 4,751,180; Chaudhary et al., 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:1066-1070; Bird et al., 1988, *Science* 242:423-426, incorporated herein in their entirety.

[144] In some embodiments, coding sequences of VH and VL domains of the dual binding antibody or fragment thereof can be fused directly without any junctional amino acids or by using a flexible polylinker composed.

[145] A peptide linker, in certain embodiments, is designed to enable the correct interaction between two beta-sheets forming the variable region of the single chain antibody. Any suitable linkers can be used to make an indirect link, such as without limitation, peptide linker, polymer linker, and chemical linker. In certain embodiments, the covalent link is an indirect link through a peptide linker.

[146] In some embodiments, an antibody comprises a mutated immunoglobulin. Examples of mutated immunoglobulins include but are not limited to an IgG that does not bind antibody-dependent cellular cytotoxicity (ADCC) components. IgG comprising L₂₃₄A/L₂₃₅A (LALA) mutations cannot bind the Fc receptor (*See*, Xu D, Alegre ML, Varga SS, Rothermel AL, Collins AM, Pulito VL, et al. . *In vitro* characterization of five humanized OKT3 effector function variant antibodies. *Cell Immunol.* (2000) 200:16–26. 10.1006/cimm.2000.1617). In some embodiments, the dual antibody comprises an IgG comprising the L₂₃₄A/L₂₃₅A (LALA) mutations. The mutations as numbered here are based on an EU numbering convention used for the constant region, (*See*, Xu D, Alegre ML, Varga SS, Rothermel AL, Collins AM, Pulito VL, et al. . *In vitro* characterization of five humanized OKT3 effector function variant antibodies. *Cell Immunol.* (2000) 200:16–26. 10.1006/cimm.2000.1617).

[147] In some embodiments, a mutated IgG comprises an IgG1, wherein the Fc region is

engineered. In some embodiments, a mutated IgG comprises an IgG2, wherein the Fc region is engineered. In some embodiments, a mutated IgG comprises an IgG3, wherein the Fc region is engineered. In some embodiments, a mutated IgG comprises an IgG4, wherein the Fc region is engineered. In certain embodiments, mutations within an Fc region of an antibody abolishes immune effector functions of the antibody.

[148] In some embodiments, an isolated dual binding antibody comprises an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody. In some embodiments, an isolated dual binding antibody comprises an IgG, wherein said IgG is IgG1, IgG2, IgG3, or IgG4.

[149] In some embodiments, an isolated dual binding antibody comprises a mutated IgG, wherein said mutant IgG is unable to bind to antibody-dependent cellular cytotoxicity components.

[150] In some embodiments, the dual binding antibody described herein comprises an IgG immunoglobulin. In some embodiments, the dual binding antibody described herein comprises an IgG1 immunoglobulin, an IgG2 immunoglobulin, an IgG3 immunoglobulin, or an IgG4 immunoglobulin. In some embodiments, the dual binding antibody comprises an IgG1 immunoglobulin. In some embodiments, the dual binding antibody comprises an IgG2 immunoglobulin. In some embodiments, the dual binding antibody comprises an IgG3 immunoglobulin. In some embodiments, the dual binding antibody comprises an IgG4 immunoglobulin. In some embodiments, the dual binding antibody comprises an IgG1 immunoglobulin or an IgG4 immunoglobulin.

[151] In some embodiments, the dual binding antibody described herein comprises a Fab immunoglobulin fragment. In some embodiments, the dual binding antibody described herein comprises an F(ab')₂ immunoglobulin fragment. In some embodiments, the dual binding antibody described herein comprises an Fv immunoglobulin construct. In some embodiments, the dual binding antibody described herein comprises an scFv immunoglobulin construct. In some embodiments, the dual binding antibody described herein comprises a minibody immunoglobulin construct comprising a pair of single-chain Fv fragments which are linked via CH3 domains.

[152] In some embodiments, the dual binding antibody described herein comprises a diabody immunoglobulin construct. In some embodiments, a diabody immunoglobulin construct comprises a heavy chain variable (VH) and light chain variable (VL) regions connected by a small peptide linker. In some embodiments, a diabody immunoglobulin construct comprises single-chain (Fv)₂ in which two scFv fragments are covalently linked to each other. In some embodiments, the dual binding antibody described herein comprises a diabody immunoglobulin construct comprising three scFv fragments covalently linked to each other. Diabodies have been

shown in the art to have dissociation constants up to 40-fold lower than corresponding scFvs, meaning that they have a much higher affinity to their target. Consequently, use of a diabody in a method of use as described below, could results in much lower dosing of a diabody or triabody, than an IgG comprising the same VH and VL domains.

[153] In some embodiments, the dual binding antibody comprising a linker or linkers between binding components, for example but not limited to between a VH and a VL in an scFv, a minibody, a diabody, a triabody, or a tetrabody. In some embodiments, the dual binding antibody does not comprise a linker or linkers between binding components, for example but not limited to between a VH and a VL in an scFv, a minibody, a diabody, a triabody, or a tetrabody. In some embodiments, a linker may comprise a single amino acid. In some embodiments, a linker comprises any known linker in the art. In some embodiments, a linker comprises the amino acid sequence set forth in SEQ ID NO: 147.

[154] A skilled artisan would appreciate that the term "variant" encompasses a polypeptide differing from a specifically recited polypeptide sequences, for example the amino acid sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2, by single or multiple amino acid insertions, deletions, and/or substitutions, created using, e.g., recombinant DNA techniques. Variants of the antigen binding molecules disclosed herein include antigen binding molecules wherein one or several of the amino acid residues are modified by at least one substitution, addition and/or deletion in such manner that the antigen binding affinity is newly created in the antigen binding molecules.

[155] The dual binding portion of the antibodies described herein comprises an immunoglobulin heavy chain variable region and an immunoglobulin light chain variable region (VH and VL, respectively), wherein the amino acid sequence set forth in SEQ ID NO: 1 comprises a VH template and the amino acid sequence set forth in SEQ ID NO: 2 comprises a VL template, and the dual binding region comprises at least two variants within the VH template sequence, or within the VL template sequence, or a combination thereof.

[156] A skilled artisan would appreciate that an "isolated dual binding antibody", in certain embodiments, encompasses an antibody that (1) is free of at least some other proteins with which it would typically be found in nature or with which it would typically be found during synthesis thereof, (2) is essentially free of other non-identical binding antibodies from the same source, (3) may be expressed recombinantly by a cell, (4) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is associated in during synthesis, or (5) does not occur in nature, or a combination thereof. Such an isolated antibody may be encoded by genomic DNA, cDNA, mRNA or other RNA, of may be of synthetic origin, or any combination thereof. In certain embodiments, the isolated antibody is substantially free from

proteins or polypeptides or other contaminants that would interfere with its use (therapeutic, diagnostic, prophylactic, research or otherwise). As used throughout, the terms “dual antibody” and “dual binding antibody” may be used interchangeably having all the same meanings and qualities.

[157] In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT numbering of heavy chain variable region variant positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT numbering of light chain variable region variant positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of the variant heavy chain variable region and the variant light chain variable region; wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or said combination thereof, is at least 2.

[158] IMGT[®] is the international ImMunoGeneTics information system[®], (See, Nucleic Acids Res. 2015 Jan;43(Database issue):D413-22. doi: 10.1093/nar/gku1056. Epub 2014 Nov 5 Free article. PMID: 25378316 LIGM:441 and Dev Comp Immunol. 2003 Jan;27(1):55-77). IMGT is a unique numbering system for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains, (Lefranc MP1, Pommié C, Ruiz M, Giudicelli V, Foulquier E, Truong L, Thouvenin-Contet V, Lefranc G. Dev Comp Immunol 27: 55-77. (2003)). IMGT[®] presents a uniform numbering system for these IG and TcR variable domain sequences, based on aligning 5 or more 000 IG and TcR variable region sequences, taking into account and combining the Kabat definition of FRs and CDRs, structural data, and Chothia's characterization of the hypervariable loops. IMGT is considered a universal numbering scheme for antibodies well known in the art.

[159] In describing variant amino acid positions present in the VH and VL domains, in some embodiments the IMGT numbering is used. In some embodiments, variant amino acid positions are presented as the specific positions within a given sequence, for example but not limited to SEQ ID NO: 1 and SEQ ID NO: 2. In some embodiments, variant amino acid positions are identified by both the specific positions within a given SEQ ID NO: sequence and the IMGT numbering system. A skilled artisan would recognize that the actual amino acid position number of an amino acid identified by position number relative to a SEQ ID NO: may differ from the IMGT numbering

system, yet the residue identified is identical. For example, but not limited to the amino acid residue at position 106 of SEQ ID NO: 1 is the identical residue identified at position 112 by the IMGT numbering system. The skilled artisan would recognize that while the same amino acid residue may be identified as having different positions depending what system is used, the amino acid residues location and identity within a contiguous amino acid sequence is clear.

[160] In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variants at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT numbering of heavy chain variable region variant positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof), and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2, wherein optionally said amino acid sequence SEQ ID NO: 2 comprises at least one variant amino acid, wherein the total number of variant positions in said heavy chain variable region, light chain variable region, or a combination thereof is at least 2. In some embodiments, disclosed herein is an isolated dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least two amino acid variants at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT numbering of heavy chain variable region variant positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof), and any light chain variable region.

[161] The amino acid sequences of many light chain variable regions, in and of themselves, are known as such in the art. The skilled person would be able to use such known sequences, and in conjunction with the heavy chain variable regions described herein, analyze for dual binding using routine methodologies and techniques well known in the art (See for example but not limited to, Example 1 below).

[162] In some embodiments, disclosed herein is an isolated dual binding antibody comprising a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variants at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT numbering of light chain variable region variant positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof), and a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, wherein optionally said amino acid sequence SEQ ID NO: 1 comprises at least one variant amino acid, wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or a combination thereof is at least 2. In some embodiments, disclosed herein is an isolated dual

binding antibody comprising a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least two amino acid variants at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT numbering of light chain variable region variant positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof), and any heavy chain variable region.

[163] The amino acid sequences of many heavy chain variable regions, in and of themselves, are known as such in the art. The skilled person would be able to use such known sequences, and in conjunction with the light chain variable regions described herein, analyze for dual binding using routine methodologies and techniques well known in the art (See for example but not limited to, Example 1 below).

[164] In some embodiments, a VH domain described herein comprises the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variants at any position. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises at least two amino acid variants at any position. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises at least between 1- 10 amino acid variants at any position. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises 1-5 amino acid variants at any position. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more amino acid variants at any position.

[165] In some embodiments, a VL domain described herein comprises the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variants at any position. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises at least two amino acid variants at any position. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises at least between 1- 10 amino acid variants at any position. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises 1-5 amino acid variants at any position. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more amino acid variants at any position.

[166] In some embodiments, a VH domain and a VL domain described herein comprising the amino acid sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2, respectfully have a combined number of variant positions of at least 2. In some embodiments, a VH domain and a VL domain described herein comprising the amino acid sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2, respectfully have a combined number of variant positions of between 2-20. In some embodiments, a VH domain and a VL domain described herein comprising the amino acid

sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2, respectfully have a combined number of variant positions of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 17, 18, 19, or 20. In some embodiments, a VH domain and a VL domain described herein comprising the amino acid sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2, respectfully have a combined number of variant positions of more than 20.

[167] In certain embodiments, dual antibody binding regions, as described herein include a heavy chain and a light chain CDR set, respectively, interposed between a heavy chain and a light chain framework region (FR) set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term “CDR set” refers to the three hypervariable regions of a heavy or light chain variable region. Proceeding from the N-terminus of a heavy or light chain polypeptide, these regions are denoted as “CDR1,” “CDR2,” and “CDR3” respectively. An antigen binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain variable region. Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with a bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the CDR regions are primarily responsible for the specificity of an antigen binding site.

[168] As used herein, the term “FR set” refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain variable region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the variable region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, all variable region sequences contain an internal disulfide loop of around 90 amino acid residues. When the variable regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs, which influence the folded shape of the CDR loops into certain “canonical” structures—regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

[169] In some embodiments, an at least one variant in said VH comprises a variant amino acid in a CDR region. In some embodiments, an at least one variant in said VL comprises a variant amino acid in a CDR region. In some embodiments, an at least one variant in said VH comprises a variant amino acid in a FR region. In some embodiments, an at least one variant in said VL comprises a variant amino acid in a FR region. In some embodiments, an at least two variants in

said VH comprises variant amino acids in a CDR region, an FR region, or both. In some embodiments, an at least two variants in said VL comprises variant amino acids in a CDR region, an FR region, or both. In some embodiments, variant positions in the VH include variants in at least one CDR and at least one FR region. In some embodiments, variant positions in the VL include variants in at least one CDR and at least one FR region.

[170] In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT numbering of heavy chain variable region variant positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 52 (IMGT position 57). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 99 (IMGT position 107). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 100 (IMGT position 108). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 101 (IMGT position 109). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 102 (IMGT position 110). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 103 (IMGT position 111). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 104 (IMGT position 111A). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 105 (IMGT position 112A). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 106 (IMGT position 112). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 107 (IMGT position 113). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 108 (IMGT position 114). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 111 (IMGT position 117).

[171] In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 52 (IMGT position 57) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 99 (IMGT position 107)

and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 100 (IMGT position 108) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 101 (IMGT position 109) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 102 (IMGT position 110) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 103 (IMGT position 111) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 104 (IMGT position 111A) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 105 (IMGT position 112A) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 106 (IMGT position 112) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 107 (IMGT position 113) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 108 (IMGT position 114) and between 1-3 additional variant amino acids. In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at position 111 (IMGT position 117) and between 1-3 additional variant amino acids.

[172] In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 105 and 106 (IMGT positions 112A and 112). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 106 and 111 (IMGT positions 112 and 117). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 103 and 106 (IMGT positions 111 and 112). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 104 and 106 (IMGT positions 111A and 112). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 104, 106, and 111 (IMGT positions 111A, 112, and 117). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid

variant at positions 105, 106, and 111 (IMGT positions 112A, 112, and 117). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 103, 106, and 111 (IMGT positions 111, 112, and 117). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 99, 104, and 111 (IMGT positions 107, 111A, and 117). In some embodiments, a VH comprising the amino acid sequence set forth in SEQ ID NO: 1 comprises an amino acid variant at positions 52, 99, 104, and 111 (IMGT positions 57, 107, 111A, and 117).

[173] In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT numbering of light chain variable region variant positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 26 (IMGT position 27). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 27 (IMGT position 28). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 31 (IMGT position 38). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 51 (IMGT position 65). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 56 (IMGT position 70). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 77 (IMGT position 94). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 92 (IMGT position 109). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 93 (IMGT position 110). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 96 (IMGT position 115).

[174] In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 26 (IMGT position 27) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 27 (IMGT position 28) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 31 (IMGT position 38) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant

at position 51 (IMGT position 65) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 56 (IMGT position 70) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 77 (IMGT position 94) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 92 (IMGT position 109) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 93 (IMGT position 110) and between 1-7 additional variant amino acids. In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at position 96 (IMGT position 115) and between 1-7 additional variant amino acids.

[175] In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, and 96 (IMGT positions 27, 28, 38, 70, 94, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, 92, and 96 (IMGT positions 27, 28, 38, 70, 94, 109, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 31, 56, and 77 (IMGT positions 27, 38, 70, and 94). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, and 77 (IMGT positions 27, 28, 38, 70, and 94). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, and 92 (IMGT positions 27, 28, 38, 70, 94, and 109). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 51, 56, 77, and 92 (IMGT positions 27, 28, 38, 65, 70, 94, and 109). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 31, 56, 77, 92, and 96 (IMGT positions 27, 38, 70, 94, 109, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 31, 77, and 92 (IMGT positions 27, 38, 94, and 109). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, and 93 (IMGT positions 27, 28, 38, 70, 94, and 110). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, 93, and 96 (IMGT positions 27, 28, 38, 70, 94, 110, and 115). In some embodiments, a VL

comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 51, 77, 93, and 96 (IMGT positions 27, 28, 38, 65, 94, 110, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 56, 77, 93, and 96 (IMGT positions 27, 28, 38, 70, 94, 110, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 27, 31, 56, 77, and 96 (IMGT positions 28, 38, 70, 94, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 27, 31, 56, 77, 92, and 96 (IMGT positions 28, 38, 70, 94, 109, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, 51, 77, and 96 (IMGT positions 27, 28, 38, 65, 94, and 115). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises an amino acid variant at positions 26, 27, 31, or 96 (IMGT positions 27, 28, 38, and 115), or any combination thereof.

[176] In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises amino acid variants at positions in a framework region at any of positions 56 or 77 of SEQ ID NO: 2, or a combination thereof (IMGT positions: 70 or 94, or a combination thereof). In some embodiments, a VL comprising the amino acid sequence set forth in SEQ ID NO: 2 comprises at least one amino acid variant at a position within a framework region at any of positions 56 or 77 of SEQ ID NO: 2, or a combination thereof (IMGT positions: 70 or 94, or a combination thereof), wherein the variant amino acid at position 56 comprises a leucine, an alanine, an arginine, a lysine, an aspartic acid, a glycine, or a glutamic acid, and or the variant amino acid at position 77 comprises a valine.

[177] In some embodiments, disclosed herein is an isolated dual binding antibody comprising: a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions (IMGT) 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof; and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions (IMGT) 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof; wherein the total number of variant positions in said dual binding antibody is at least 2.

[178] In some embodiments, an isolated dual binding antibody comprises a variant VH or a variant VL, or a combination thereof, with variant amino acids at positions other than (IMGT) 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117 in the heavy chain variable region and or (IMGT) 27, 28, 38, 65, 70, 94, 109, 110, or 115 in the light chain variable region.

[179] In some embodiments, an at least one amino acid variant in a VH comprises a variant amino acid in a CDR region. In some embodiments, an at least one amino acid variant in a VH comprises a variant amino acid in a CDR1 region. In some embodiments, an at least one amino acid variant in a VH comprises a variant amino acid in a CDR2 region. In some embodiments, an at least one amino acid variant in a VH comprises a variant amino acid in a CDR3. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in two different CDR regions. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a same CDR region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a same CDR1 region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a same CDR2 region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a same CDR3 region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a CDR1 and a CDR2 region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a CDR1 and a CDR3 region. In some embodiments, an at least two amino acid variants in a VH comprises variant amino acids in a CDR2 and a CDR3 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a single CDR region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR1 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR2 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR3 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, an at least three amino acid variants in a VH comprises variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a single CDR region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR1 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR2 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR1 region and a

CDR3 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VH comprises variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a single CDR region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise amino acids in a CDR1 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR2 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR3 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, when there are 5 or more amino acid variants in a VH, variant positions comprise variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region.

[180] In some embodiments, an at least one amino acid variant in a VL comprises a variant amino acid in a CDR region. In some embodiments, an at least one amino acid variant in a VL comprises a variant amino acid in a CDR1 region. In some embodiments, an at least one amino acid variant in a VL comprises a variant amino acid in a CDR2 region. In some embodiments, an at least one amino acid variant in a VL comprises a variant amino acid in a CDR3. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in two different CDR regions. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a same CDR region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a same CDR1 region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a same CDR2 region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a same CDR3 region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a CDR1 and a CDR2 region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a CDR1 and a CDR3 region. In some embodiments, an at least two amino acid variants in a VL comprises variant amino acids in a CDR2 and a CDR3 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a single CDR region. In some embodiments, an at least

three amino acid variants in a VL comprises variant amino acids in a CDR1 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR2 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR3 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, an at least three amino acid variants in a VL comprises variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a single CDR region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR1 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR2 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, an at least 4 amino acid variants in a VL comprises variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a single CDR region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise amino acids in a CDR1 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR3 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, when there are 5 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a single CDR region. In some embodiments,

when there are 6 amino acid variants in a VL, variant positions comprise amino acids in a CDR1 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR3 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, when there are 6 amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a single CDR region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise amino acids in a CDR1 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR3 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR2 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region and a CDR3 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR2 region and a CDR3 region. In some embodiments, when there are 7 or more amino acid variants in a VL, variant positions comprise variant amino acids in a CDR1 region, a CDR2 region, and a CDR3 region.

[181] In some embodiments, an amino acid variant comprises a substitution of one amino acid residue for another. In some embodiments, an amino acid variant comprises a substitution of a hydrophobic residue for a non-hydrophobic residue. In some embodiments, an amino acid variant comprises a substitution of a charged residue for a non-charged residue. In some embodiments, an amino acid variant comprises a neutral substitution, wherein the amino acid being substituted has similar qualities. In some embodiments, an amino acid variant comprises a substitution of an aromatic residue for a non-aromatic residue. In some embodiments, natural aromatic amino acids such as Trp, Tyr and Phe, are substituted for synthetic non-natural acid such as Phenylglycine, TIC, naphthylelanine (Nol), ring-methylated derivatives of Phe, halogenated derivatives of Phe or o-methyl-Tyr. In some embodiments, a variant substitution comprises

substituting a modified amino acid or a non-amino acid monomer (e.g. fatty acid, complex carbohydrates etc). A skilled artisan would appreciate that while the choice of amino acid residues at each variant position may in certain embodiments, affect the 3D structure of the VH, VL, and/or combination thereof, the choice of amino acid residues at each variant position is considered independently.

[182] In some embodiments, "amino acid" or "amino acid residue" or "residue" is understood to include the 20 naturally occurring, encoded amino acid residues, and those amino acids often modified post-translationally *in vivo*, including, for example, hydroxyproline, phosphoserine and phosphothreonine; and other unusual amino acid including, but not limited to, 2-aminoadipic acid, hydroxylysine, isodesmosine, nor-valine, nor-leucine and ornithine. In some embodiments, "amino acid" includes both D- and L-amino acid. In some embodiments, an amino acid variant substitution is a D-amino acid. In some embodiments, an amino acid variant substitution is an L-amino acid. In some embodiments, a variant residue comprises a naturally occurring amino acid. In some embodiments, a variant residue comprises a naturally occurring, encoded amino acid residue. In some embodiments, a variant residue comprises a naturally occurring, non-encoded amino acid residue. In some embodiments, a variant residue comprises a non-naturally occurring amino acid.

[183] In some embodiments, a variant residue comprises a non-naturally occurring, non-proteinogenic amino acid.

[184] In some embodiments, the amino acid sequence of a VH domain of the dual binding antibody is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54; and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 4 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 6 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 8 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 10 and any variable light chain region. In some embodiments, an isolated dual binding

[illegible]

variable region comprising the amino acid sequences set forth in SEQ ID NOs: 44 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 46 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 48 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 50 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 52 and any variable light chain region. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 54 and any variable light chain region.

[185] In some embodiments, the amino acid sequence of a VH domain of the dual binding antibody is one of those set forth in Table 1 or Table 10. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region comprising one of the amino acid sequences set forth in Table 1 or Table 10; and any variable light chain region.

[186] In some embodiments, the amino acid sequence of a VL domain of the dual binding antibody is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53; and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 3 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 5 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 7 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 9 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 11 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set

[illegible]

NOs: 45 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 47 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 49 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 51 and any variable heavy chain region. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising the amino acid sequences set forth in SEQ ID NOs: 53 and any variable heavy chain region.

[187] In some embodiments, the amino acid sequence of a VL domain of the dual binding antibody is one of those set forth Table 1 or Table 10. In some embodiments, an isolated dual binding antibody comprises a light chain variable region comprising one of the amino acid sequences set forth in Table 1 or Table 10; and any variable heavy chain region.

[188] A skilled artisan would recognize that when pairing a VH or VL domain comprising a known amino acid sequence, with a VL or VH domain, respectively, to comprise an antigen binding region, such a pairing may be analyzed for binding properties using methods well known in the art (See for example, the disclosure herein and Examples below).

[189] In some embodiments, the amino acid sequence of a heavy chain variable region – light chain variable region pair are selected from, but not limited to, the pair sequences set forth in SEQ ID NOs: 4 and 3, SEQ ID Nos: 6 and 5, SEQ ID Nos: 8 and 7, SEQ ID Nos: 10 and 9, SEQ ID Nos: 12 and 11, SEQ ID Nos: 14 and 13, SEQ ID Nos: 16 and 15, SEQ ID Nos: 18 and 17, SEQ ID Nos: 20 and 19, SEQ ID Nos: 22 and 21, SEQ ID Nos: 24 and 23, SEQ ID Nos: 26 and 25, SEQ ID Nos: 28 and 27, SEQ ID Nos: 30 and 29, SEQ ID Nos: 32 and 31, SEQ ID Nos: 34 and 33, SEQ ID Nos: 36 and 35, SEQ ID Nos: 38 and 37, SEQ ID Nos: 40 and 39, SEQ ID Nos: 42 and 41, SEQ ID Nos: 44 and 43, SEQ ID Nos: 46 and 45, SEQ ID Nos: 48 and 47, SEQ ID Nos: 50 and 49, SEQ ID Nos: 52 and 51, and SEQ ID Nos: 54 and 53. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: 4 and 3. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 6 and 5. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 8 and 7. In some embodiments, an isolated dual binding antibody comprises a heavy

[illegible]

region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 40 and 39. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 42 and 41. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 44 and 43. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: SEQ ID Nos: 46 and 45. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: 48 and 47. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: 50 and 49. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID NOs: 52 and 51. In some embodiments, an isolated dual binding antibody comprises a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in SEQ ID Nos: 54 and 53.

[190] In some embodiments, the amino acid sequences of a heavy chain variable region – light chain variable region pair are selected from the pair sequences set forth in any one of the following: SEQ ID NOs:209 and 210, SEQ ID NOs:211 and 212, SEQ ID NOs:213 and 214, SEQ ID NOs:215 and 216, SEQ ID NOs:217 and 218, SEQ ID NOs:219 and 220, SEQ ID NOs:221 and 222, SEQ ID NOs:223 and 224, SEQ ID NOs:225 and 226, SEQ ID NOs:227 and 228, SEQ ID NOs:229 and 230, SEQ ID NOs:231 and 232, SEQ ID NOs:233 and 234, SEQ ID NOs:235 and 236, SEQ ID NOs:237 and 238, SEQ ID NOs:239 and 240, SEQ ID NOs:241 and 242, SEQ ID NOs:243 and 244, SEQ ID NOs:245 and 246, SEQ ID NOs:247 and 248, SEQ ID NOs:249 and 250, SEQ ID NOs:251 and 252, SEQ ID NOs:253 and 254, SEQ ID NOs:255 and 256, SEQ ID NOs:257 and 258, SEQ ID NOs:259 and 260, SEQ ID NOs:261 and 262, SEQ ID NOs:263 and 264, SEQ ID NOs:265 and 266, SEQ ID NOs:267 and 268, SEQ ID NOs:269 and 270, SEQ ID NOs:271 and 272, SEQ ID NOs:273 and 274, SEQ ID NOs:275 and 276, SEQ ID NOs:277 and 278, SEQ ID NOs:279 and 280, SEQ ID NOs:281 and 282, SEQ ID NOs:283 and 284, SEQ ID NOs:285 and 286, SEQ ID NOs:287 and 288, SEQ ID NOs:289 and 290, SEQ ID NOs:291 and 292, SEQ ID NOs:293 and 294, SEQ ID NOs:295 and 296, SEQ ID NOs:297 and 298, SEQ ID NOs:299 and 300, SEQ ID NOs:301 and 302, SEQ ID NOs:303 and 304, SEQ ID NOs:305 and 306, SEQ ID NOs:307 and 308, SEQ ID NOs:309 and 310, SEQ ID NOs:311 and 312, SEQ

ID NOs:313 and 314, SEQ ID NOs:315 and 316, SEQ ID NOs:317 and 318, SEQ ID NOs:319 and 320, SEQ ID NOs:321 and 322, SEQ ID NOs:323 and 324, SEQ ID NOs:325 and 326, SEQ ID NOs:327 and 328, SEQ ID NOs:329 and 330, SEQ ID NOs:331 and 332, SEQ ID NOs:333 and 334, SEQ ID NOs:335 and 336, SEQ ID NOs:337 and 338, SEQ ID NOs:339 and 340, SEQ ID NOs:341 and 342, SEQ ID NOs:343 and 344, SEQ ID NOs:345 and 346, SEQ ID NOs:347 and 348.

[191] In some embodiments, the amino acid sequence of an scFv fragment comprises the pair of sequences set forth in, but not limited to, any of the following pairs: SEQ ID NOs: 4 and 3, SEQ ID Nos: 6 and 5, SEQ ID Nos: 8 and 7, SEQ ID Nos: 10 and 9, SEQ ID Nos: 12 and 11, SEQ ID Nos: 14 and 13, SEQ ID Nos: 16 and 15, SEQ ID Nos: 18 and 17, SEQ ID Nos: 20 and 19, SEQ ID Nos: 22 and 21, SEQ ID Nos: 24 and 23, SEQ ID Nos: 26 and 25, SEQ ID Nos: 28 and 27, SEQ ID Nos: 30 and 29, SEQ ID Nos: 32 and 31, SEQ ID Nos: 34 and 33, SEQ ID Nos: 36 and 35, SEQ ID Nos: 38 and 37, SEQ ID Nos: 40 and 39, SEQ ID Nos: 42 and 41, SEQ ID Nos: 44 and 43, SEQ ID Nos: 46 and 45, SEQ ID Nos: 48 and 47, SEQ ID Nos: 50 and 49, SEQ ID Nos: 52 and 51, and SEQ ID Nos: 54 and 53.

Nucleotide Sequences Encoding Engineered “Re-Epitoped” VH domains, VL domains, or both VH and VL domains, and Vectors and Host Cells Comprising these Nucleotide Sequences

[192] The present disclosure provides dual binding antibodies comprising a VH domain, a VL domain, or both a VH and VL domain, comprising variant amino acid sequences compared with template VH and VL sequences SEQ ID NO: 1 and SEQ ID NO: 2, respectively. As described in detail above, in some embodiments the dual binding antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in the encoded heavy chain variable region, the encoded light chain variable region, or a combination thereof, is at least 2.

[193] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence

encodes an isolated dual binding antibody comprises an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3 as disclosed herein. In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes an isolated dual binding antibody comprises an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said VH domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3 as disclosed in Table 8 or Table 4. In some embodiments, the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid.

[194] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a VH domain of a dual binding antibody comprises HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 142), and HCDR3 (SEQ ID NO: 143), wherein the VH domain comprises a variant amino acid at, at least one of HX1, HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12.

[195] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a VH domain of a dual binding antibody comprises HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is selected from the group consisting of W and S; and HCDR3 (SEQ ID NO: 138) wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M. In certain embodiments, a nucleic acid construct comprising a nucleic acid sequence encoding a dual binding antibody comprising variant amino acids comprising HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HX2 is selected from the group consisting of A and S, HX3 is P, HX4 is Q, HX5 is W, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX10 is E, HX11 is A, and HX12 is selected from the group I and L.

[196] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a dual binding antibody comprising a VH domain comprising an HCDR1 (SEQ ID NO:

136), an HCDR2 (SEQ ID NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX10 is E, HX11 is A, and HX12 is I; or an HCDR1 (SEQ ID NO: 136), an HCDR2 (SEQ ID NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX10 is E, HX11 is A, and HX12 is L; or an HCDR1 (SEQ ID NO: 136), an HCDR2 (SEQ ID NO: 137) wherein HX1 is W, and an HCDR3 (SEQ ID NO: 138) wherein HX2 is S, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX10 is E, HX11 is A, and HX12 is L.

[197] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes an isolated dual binding antibody comprising an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said in some embodiments the VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCDR3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid.

[198] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a VL domain of a dual binding antibody comprising LCDRs. In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes an isolated dual binding antibody comprising an antibody antigen-binding domain site comprising a VH domain and a VL domain, wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3 as disclosed in Table 9 or Table 5. In some embodiments, the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; and wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G. In certain embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes the isolated dual binding antibody comprising variant amino acids wherein LCDR1 (SEQ ID NO: 139) LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is selected from the group consisting of S and G, LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is selected from the group

consisting of H and G.

[199] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a dual binding antibody comprising a VL domain comprising an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G; or an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is H, or an LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, and LX3 is L, an LCDR2 (SEQ ID NO: 140) wherein LX4 is G, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G.

[200] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes an isolated dual binding antibody comprising an antibody antigen-binding domain site comprising a VH domain and a VL domain comprises a combination of VH domain HCDRs and VL domain LCDRs described above. For example, but not limited to, in certain embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136; wherein the amino acid sequence of HCDR2 is set forth as: I HX1 Y D G S N K (SEQ ID NO: 142), wherein HX1 is any amino acid; and wherein the amino acid sequence of HCDR3 is set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO: 143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid; and wherein said VL domain comprises a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth as LX1, LX2, G S K LX3 V (SEQ ID NO: 144), wherein LX1, LX2, and LX3 are any amino acid; wherein the amino acid sequence of LCDR2 is set forth as D D LX4 (SEQ ID NO: 145), wherein LX4 is any amino acid; and wherein the amino acid sequence of LCD3 is set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO: 146), wherein LX5, LX6, and LX7 are any amino acid.

[201] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136, wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is selected from the group consisting of W and S; wherein the amino acid sequence of HCDR3 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, wherein HX3 is P, wherein HX4 is Q, wherein HX5 is W, wherein HX6 is selected from the group consisting of E, Q, M, L, and V, wherein HX7 is selected from the group consisting of L, W, and Y, wherein HX8 is selected from

the group consisting of V and T, wherein HX9 is selected from the group consisting of H, A, S, wherein HX10 is E, wherein HX11 is A, wherein HX12 is selected from the group consisting of I, L, and M; and a VL domain comprises a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139, wherein LX1 is selected from the group consisting of N, L, and I, wherein LX2 is selected from the group consisting of L and I, wherein LX3 is selected from the group consisting of S and L; wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140, wherein LX4 is selected from the group consisting of S and G; wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141, wherein LX5 is selected from the group consisting of S and T, wherein LX6 is selected from the group consisting of S and G, and wherein LX7 is selected from the group consisting of H and G.

[202] In certain embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes the isolated dual binding antibody comprising a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, wherein the amino acid sequence of HCDR1 is set forth in SEQ ID NO: 136, wherein the amino acid sequence of HCDR2 is set forth in SEQ ID NO: 137, wherein HX1 is W, wherein the amino acid sequence of HCDR3 is set forth in SEQ ID NO: 138, wherein HX2 is selected from the group consisting of A and S, HX3 is P, HX4 is Q, HX5 is W, HX6 is selected from the group consisting of E and M, HX7 is selected from the group consisting of L and W, HX8 is selected from the group consisting of V and T, HX9 is selected from the group consisting of H and A, HX10 is E, HX11 is A, HX12 is selected from the group I and L, and comprising a VL domain comprising a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequence of LCDR1 is set forth in SEQ ID NO: 139 wherein LX1 is L, LX2 is I, LX3 is L, wherein the amino acid sequence of LCDR2 is set forth in SEQ ID NO: 140 wherein LX4 is selected from the group consisting of S and G, wherein the amino acid sequence of LCDR3 is set forth in SEQ ID NO: 141 wherein LX5 is S, LX6 is S, and LX7 is selected from the group consisting of H and G.

[203] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encodes a re-epitoped dual binding antibody comprising a VH domain comprising a set of CDRs, HCDR1, HCDR2, and HCDR3, and a VL domain comprising a set of CDRs, LCDR1, LCDR2, and LCDR3, wherein the amino acid sequences of each CDR are as set forth in **Figures 1A** and **1B** for the clones set forth there, for example but not limited to:

[204] Clone C2: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX10 is E, HX11 is A, HX12 is I, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3

(SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G;

[205] Clone C6: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is A, HX3 is P, HX4 is Q, HX5 is W, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX10 is E, HX11 is A, HX12 is L, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is S, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is H; or

[206] Clone C9: HCDR1 (SEQ ID NO: 136), HCDR2 (SEQ ID NO: 137) wherein HX1 is W, HCDR3 (SEQ ID NO: 138) wherein HX2 is S, HX3 is P, HX4 is Q, HX5 is W, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX10 is E, HX11 is A, HX12 is L, LCDR1 (SEQ ID NO: 139) wherein LX1 is L, LX2 is I, LX3 is L, LCDR2 (SEQ ID NO: 140) wherein LX4 is G, and LCDR3 (SEQ ID NO: 141) wherein LX5 is S, LX6 is S, and LX7 is G.

[207] In certain embodiments, a nucleic acid construct comprises a single nucleic acid sequence. In certain embodiments, a nucleic acid construct comprises two nucleic acid sequence. In certain embodiments, a nucleic acid construct comprises a single nucleic acid sequence, wherein a VH domain and a VL domain are encoded by the nucleic acid sequence. In certain embodiments, a nucleic acid construct comprises two nucleic acid sequences, wherein a VH domain is encoded by one nucleic acid sequence, and a VL domain is encoded by the other nucleic acid sequence.

[208] As described herein, the present disclosure provides the polynucleotide sequences encoding the variant VH, VL, or both VH and VL domains described herein. In certain embodiments, the template VH domain is encoded by the nucleotide sequence set forth in SEQ ID NO: 55 and the template VL domain is encoded by the nucleotide sequence set forth in SEQ ID NO: 56.

[209] In some embodiments, disclosed herein is nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding antibody, said antibody comprising: a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in the encoded heavy chain variable region, the encoded light chain variable

region, or a combination thereof, is at least 2.

[210] In some embodiments, the nucleotide construct sequence comprises two nucleic acid sequences, one encoding a variant heavy chain variable region, and one a variant light chain variable region. In some embodiments, the nucleotide sequence or sequences encoding the dual binding antibody heavy chain variable region, light chain variable region, or both, is optimized for mammalian transcription and translation.

[211] The present disclosure further provides, in certain embodiments, an isolated nucleic acid construct encoding the nucleic acid sequence as described herein. Illustrative polynucleotide sequence encoding variant VH and VL domains are provided in Table 2 below. Illustrative nucleic acid constructs, comprising a nucleic acid sequence encoding variant VH domains linked to VL domains are provided in Table 3 below.

[212] Nucleic acids include DNA and RNA. These and related embodiments may include polynucleotides encoding the dual binding antibody as described herein. The term "isolated polynucleotide" as used herein shall mean a polynucleotide of genomic, cDNA, or synthetic origin, or some combination thereof, which by virtue of its origin the isolated polynucleotide (1) is not associated with all or a portion of a polynucleotide in which the isolated polynucleotide is found in nature, (2) is linked to a polynucleotide to which it is not linked in nature, or (3) does not occur in nature as part of a larger sequence.

[213] A skilled artisan would appreciate that the terms "polynucleotide" and "nucleic acid sequence" may in some embodiments be used interchangeably having all the same meanings and qualities.

[214] In some embodiments, isolated nucleic acid sequences disclosed herein, encode a VH domain comprising set of HCDRs as disclosed throughout and in **Figure 1A**, a VL domain comprising set of a set of LCDRs as disclosed throughout and in **Figure 1B**, a VH domain comprising a set of HCDRs and a VL domain comprising set of set of LCDRs as disclosed throughout and in **Figures 1A and 1B**, a VL domain or a VL domain, or a VH and a VL domain, of a dual binding antibody as described herein throughout in detail.

[215] The term "polynucleotide" as used herein encompasses single-stranded or double-stranded nucleic acid polymers. In certain embodiments, the nucleotides comprising the polynucleotide can be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. Said modifications include base modifications such as bromouridine, ribose modifications such as arabinoside and 2',3'-dideoxyribose and internucleotide linkage modifications such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate and phosphoroamidate. The term

"polynucleotide" specifically includes single and double stranded forms of DNA.

[216] The term "naturally occurring nucleotides" includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" includes oligonucleotide linkages such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. See, e.g., LaPlanche et al., 1986, Nucl. Acids Res., 14:9081; Stec et al., 1984, J. Am. Chem. Soc., 106:6077; Stein et al., 1988, Nucl. Acids Res., 16:3209; Zon et al., 1991, Anti-Cancer Drug Design, 6:539; Zon et al., 1991, OLIGONUCLEOTIDES AND ANALOGUES: A PRACTICAL APPROACH, pp. 87-108 (F. Eckstein, Ed.), Oxford University Press, Oxford England; Stec et al., U.S. Pat. No. 5,151,510; Uhlmann and Peyman, 1990, Chemical Reviews, 90:543, the disclosures of which are hereby incorporated by reference for any purpose. An oligonucleotide can include a detectable label to enable detection of the oligonucleotide or hybridization thereof.

[217] In other related embodiments, polynucleotide variants may have substantial identity to a polynucleotide template sequence, though the template sequence does not encode a dual binding antibody, or fragment thereof, or domain thereof.

[218] In some embodiments, polynucleotide variants will contain one or more substitutions, additions, deletions and/or insertions, such that the binding affinity of a binding domain encoded by the variant polynucleotide newly binds to an epitope, relative to the unmodified template as specifically set forth herein.

[219] In some embodiments, a nucleic acid sequence encodes the heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof). In some embodiments, a nucleic acid sequence encodes the heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least two amino acid variants at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof). In some embodiments, a nucleic acid sequence encodes the heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acid variants at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof).

[220] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence

encoding a heavy chain variable region comprising a sequence selected from the sequences set forth in SEQ ID Nos: 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 105, and 107. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 57. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 59. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 61. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 63. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 65. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 67. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 69. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 71. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 73. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 75. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 77. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 79. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 81. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 83. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 85. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 87. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 89. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 91. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 93. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 95. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 97. In some embodiments, the

nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 99. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 101. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 105. In some embodiments, the nucleic acid sequence encoding a heavy chain variable region comprises the sequences set forth in SEQ ID Nos: 107.

[221] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody heavy chain variable region sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 4. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 6. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 8. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 10. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 12. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 14. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 16. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 18. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 20. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 22. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 24. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 26. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 28. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 30. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 32. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO:

34. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 36. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 38. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 40. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 42. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 44. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 46. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 48. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 50. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 52. In some embodiments, the nucleic acid sequence encodes a dual binding antibody heavy chain variable region sequence set forth in SEQ ID NO: 54.

[222] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody heavy chain variable region sequence set forth in Table 10 or Table 1; for example, the VH may comprise any one of SEQ ID NOs: 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, 253, 255, 257, 259, 261, 263, 265, 267, 269, 271, 273, 275, 277, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345 and 347. In another embodiment, the nucleic acid construct comprises a nucleic acid sequence encoding a VH that is at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH sequences disclosed herein.

[223] In some embodiments, a nucleic acid sequence encodes the light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof). In some embodiments, a nucleic acid sequence encodes the light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof). In some embodiments, a nucleic acid sequence encodes the light chain variable region comprising the amino acid sequence set

forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof).

[224] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence encoding a light chain variable region comprising a sequence selected from the sequences set forth in SEQ ID NOs: 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, and 108. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 58. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 60. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 62. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 64. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 66. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 68. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 70. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 72. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 74. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 76. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 78. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 80. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 82. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 84. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 86. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 88. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 90. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 92. In some embodiments, the nucleic acid sequence encoding a light chain variable region

comprises the sequences set forth in SEQ ID NO: 94. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 96. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 98. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 100. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 102. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 104. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 106. In some embodiments, the nucleic acid sequence encoding a light chain variable region comprises the sequences set forth in SEQ ID NO: 108.

[225] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody light chain variable region sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 3. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 5. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 7. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 9. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 11. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 13. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 15. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 17. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 19. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 21. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 23. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 25. In some embodiments, the nucleic acid sequence

encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 27. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 29. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 31. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 33. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 35. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 37. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 39. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 41. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 43. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 45. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 47. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 49. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 51. In some embodiments, the nucleic acid sequence encodes a dual binding antibody light chain variable region sequence set forth in SEQ ID NO: 53.

[226] In some embodiments, the nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody light chain variable region sequence set forth in Table 10 or Table 1; for example, the VL may comprise one of SEQ ID NOs: 210, 212, 214, 216, 218, 220, 222, 224, 226, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, 254, 256, 258, 260, 262, 264, 266, 268, 270, 272, 274, 276, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346 and 348. In another embodiment, the nucleic acid construct comprises a nucleic acid sequence encoding a VL that is at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VL sequences disclosed herein.

[227] In some embodiments, a nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody heavy chain variable region – light chain variable region pair, said nucleic acid sequence selected from the paired sequences set forth in SEQ ID NOs: 57 and

58, SEQ ID Nos: 59 and 60, SEQ ID Nos: 61 and 62, SEQ ID Nos: 63 and 64, SEQ ID Nos: 65 and 66, SEQ ID Nos: 67 and 68, SEQ ID Nos: 69 and 70, SEQ ID Nos: 71 and 72, SEQ ID Nos: 73 and 74, SEQ ID Nos: 75 and 76, SEQ ID Nos: 77 and 78, SEQ ID Nos: 79 and 80, SEQ ID Nos: 81 and 82, SEQ ID Nos: 83 and 84, SEQ ID Nos: 85 and 86, SEQ ID Nos: 87 and 88, SEQ ID Nos: 89 and 90, SEQ ID Nos: 91 and 92, SEQ ID Nos: 93 and 94, SEQ ID Nos: 95 and 96, SEQ ID Nos: 97 and 98, SEQ ID Nos: 99 and 100, SEQ ID Nos: 101 and 102, SEQ ID Nos: 103 and 104, SEQ ID Nos: 105 and 106, and SEQ ID Nos: 107 and 108.

[228] In some embodiments, a nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody heavy chain variable region – light chain variable region pair as shown in Table 10 or Table 1; for example, the VH and VL pair can be one of the following: SEQ ID NOs:209 and 210, SEQ ID NOs:211 and 212, SEQ ID NOs:213 and 214, SEQ ID NOs:215 and 216, SEQ ID NOs:217 and 218, SEQ ID NOs:219 and 220, SEQ ID NOs:221 and 222, SEQ ID NOs:223 and 224, SEQ ID NOs:225 and 226, SEQ ID NOs:227 and 228, SEQ ID NOs:229 and 230, SEQ ID NOs:231 and 232, SEQ ID NOs:233 and 234, SEQ ID NOs:235 and 236, SEQ ID NOs:237 and 238, SEQ ID NOs:239 and 240, SEQ ID NOs:241 and 242, SEQ ID NOs:243 and 244, SEQ ID NOs:245 and 246, SEQ ID NOs:247 and 248, SEQ ID NOs:249 and 250, SEQ ID NOs:251 and 252, SEQ ID NOs:253 and 254, SEQ ID NOs:255 and 256, SEQ ID NOs:257 and 258, SEQ ID NOs:259 and 260, SEQ ID NOs:261 and 262, SEQ ID NOs:263 and 264, SEQ ID NOs:265 and 266, SEQ ID NOs:267 and 268, SEQ ID NOs:269 and 270, SEQ ID NOs:271 and 272, SEQ ID NOs:273 and 274, SEQ ID NOs:275 and 276, SEQ ID NOs:277 and 278, SEQ ID NOs:279 and 280, SEQ ID NOs:281 and 282, SEQ ID NOs:283 and 284, SEQ ID NOs:285 and 286, SEQ ID NOs:287 and 288, SEQ ID NOs:289 and 290, SEQ ID NOs:291 and 292, SEQ ID NOs:293 and 294, SEQ ID NOs:295 and 296, SEQ ID NOs:297 and 298, SEQ ID NOs:299 and 300, SEQ ID NOs:301 and 302, SEQ ID NOs:303 and 304, SEQ ID NOs:305 and 306, SEQ ID NOs:307 and 308, SEQ ID NOs:309 and 310, SEQ ID NOs:311 and 312, SEQ ID NOs:313 and 314, SEQ ID NOs:315 and 316, SEQ ID NOs:317 and 318, SEQ ID NOs:319 and 320, SEQ ID NOs:321 and 322, SEQ ID NOs:323 and 324, SEQ ID NOs:325 and 326, SEQ ID NOs:327 and 328, SEQ ID NOs:329 and 330, SEQ ID NOs:331 and 332, SEQ ID NOs:333 and 334, SEQ ID NOs:335 and 336, SEQ ID NOs:337 and 338, SEQ ID NOs:339 and 340, SEQ ID NOs:341 and 342, SEQ ID NOs:343 and 344, SEQ ID NOs:345 and 346, SEQ ID NOs:347 and 348. In another embodiment, the VH and VL pair is at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH and VL sequences disclosed herein.

[229] A skilled artisan would appreciate that in some embodiments, the sequence encoding a VH domain and the sequence encoding the VL domain are linked by a sequence encoding a linker

sequence. In some embodiments, a nucleic acid sequence encodes a polypeptide linker. ggcggtggtgtagcggaggcggaggatcaggtggaggcggcagt (SEQ ID NO: 148).

[230] In some embodiments, a nucleic acid construct comprises a nucleic acid sequence encoding a dual binding antibody heavy chain variable region – light chain variable region scFv, said nucleic acid sequence selected from the sequences set forth in SEQ ID NOs: 109-135.

[231] In some embodiments, a nucleic acid construct comprising a nucleic acid sequence encoding a dual antibody described herein, encodes an IgG immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG1 immunoglobulin, an IgG2 immunoglobulin, an IgG3 immunoglobulin, or an IgG4 immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG1 immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG2 immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG3 immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG4 immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG1 immunoglobulin or an IgG4 immunoglobulin.

[232] In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an Fab immunoglobulin fragment. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an F(ab')₂ immunoglobulin fragment. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an Fv immunoglobulin construct. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an scFv immunoglobulin construct. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a minibody immunoglobulin construct comprising a pair of single-chain Fv fragments, which are linked via CH3 domains.

[233] In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a diabody immunoglobulin construct. In some embodiments, a diabody immunoglobulin construct comprises a heavy chain variable (VH) and light chain variable (VL) regions connected by a small peptide linker. In some embodiments, a diabody immunoglobulin construct comprises single-chain (Fv)₂ in which two scFv fragments are covalently linked to each other. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a diabody immunoglobulin construct comprising three scFv fragments covalently linked to each other.

[234] In some embodiments, an isolated polynucleotide construct encodes an isolated dual binding antibody, as disclosed herein.

[235] In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a

mutated immunoglobulin. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a mutant IgG unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes a mutant IgG1 unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG comprising the L₂₃₄A/L₂₃₅A (LALA) mutations. In some embodiments, a nucleic acid sequence encoding a dual antibody encodes an IgG1 comprising the L₂₃₄A/L₂₃₅A (LALA) mutations.

[236] In some embodiments, as disclosed herein, a mutagenesis approach, such as site-specific mutagenesis, may be employed for the preparation of variants VH, VL, or VH and VL nucleic acid sequences encoding the variant VH, VL, or VH and VL amino acid sequences. Template VH and VL nucleic acid sequences SEQ ID NO: 55 and 56, respectively, encode the template amino acid sequences SEQ ID NO: 1 and SEQ ID NO: 2, respectively. In some embodiments, a dual binding antibody comprises a variant VH domain, a variant VL domain, or both, encoded by a variant VH, VL, or VH and VL nucleotide sequences, wherein said nucleotide sequence comprises site-specific mutagenesis of the nucleotide template sequences SEQ ID NO: 55 and SEQ ID NO: 56, respectively. By this approach, specific modifications in a polypeptide sequence can be made through mutagenesis of the underlying polynucleotides that encode them. These techniques provide a straightforward approach to prepare and test sequence variants, for example but not limited to, introducing one or more nucleotide sequence changes into the polynucleotide in view of the amino acid variant sites desired, as described above in detail.

[237] Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

[238] In certain embodiments, mutagenesis of the polynucleotide sequences that encode component parts of the dual binding antibody (VH domains, VL domain, or a combination thereof, as disclosed herein, is contemplated in order to alter the binding properties of the encoded template VH or VL or both, such that the resulting antibody comprises a dual binding affinity. The techniques of site-specific mutagenesis are well-known in the art and are widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising

typically about 14 to about 25 nucleotides or so in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

[239] As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phages are readily commercially available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

[240] In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

[241] The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. In some embodiments, methods of preparing libraries includes those known in the art, for example but not limited to methods described in United States Patent No. 9,889,423, which are included herein in their entirety. In some embodiments, a method for designing the sequence variants within a library comprises designing the variant sequences on a computer and then have the sequence synthesized, a method that involves both chemical and biochemical processes.

[242] As used herein, the term "oligonucleotide directed mutagenesis procedure" encompasses template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" encompasses a process that involves the template-dependent extension of a primer molecule. The term "template dependent process" encompasses nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the

newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U.S. Pat. No. 4,237,224, specifically incorporated herein by reference in its entirety.

[243] In another approach for the production of polypeptide VH and VL variants, recursive sequence recombination, as described in U.S. Pat. No. 5,837,458, may be employed. In this approach, iterative cycles of recombination and screening or selection are performed to "evolve" individual polynucleotide variants having, for example, increased binding affinity. Certain embodiments also provide constructs in the form of plasmids, vectors, transcription or expression cassettes which comprise at least one polynucleotide as described herein.

[244] In certain embodiments, the polynucleotides described above, e.g., VH, VL, or VH and VL variant polynucleotides, fragments and hybridizing sequences, encoding the amino acid VH, VL, or VH and VL variants, are comprised in a dual binding antibody.

[245] The polynucleotides described herein, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative polynucleotide segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful.

[246] In certain embodiments, the isolated polynucleotide is inserted into a vector. In some embodiments, a vector comprises an expression vector comprising a polynucleotide construct disclosed herein.

[247] The term "vector" as used herein encompasses a vehicle into which a polynucleotide encoding a protein may be covalently inserted so as to bring about the expression of that protein and/or the cloning of the polynucleotide. The isolated polynucleotide may be inserted into a vector using any suitable methods known in the art, for example, without limitation, the vector may be digested using appropriate restriction enzymes and then may be ligated with the isolated polynucleotide having matching restriction ends.

[248] Examples of suitable vectors include, without limitation, plasmids, phagemids, cosmids, artificial chromosomes such as yeast artificial chromosome (YAC), bacterial artificial chromosome (BAC), or P1-derived artificial chromosome (PAC), bacteriophages such as lambda phage or M13 phage, and animal viruses. Examples of categories of animal viruses useful as vectors include, without limitation, retrovirus (including lentivirus), adenovirus, adeno-associated virus, herpesvirus (e.g., herpes simplex virus), poxvirus, baculovirus, papillomavirus, and papovavirus (e.g., SV40).

[249] For expression of the dual antibody or components thereof, the vector may be introduced into a host cell to allow expression of the polypeptide within the host cell. The expression vectors may contain a variety of elements for controlling expression, including without limitation, promoter sequences, transcription initiation sequences, enhancer sequences, selectable markers, and signal sequences. These elements may be selected as appropriate by a person of ordinary skill in the art. In some embodiments, these elements may be considered "control" elements.

[250] A skilled artisan would appreciate that the term "control sequence" may encompass polynucleotide sequences that can affect expression, processing or intracellular localization of coding sequences to which they are ligated or operably linked. The nature of such control sequences may depend upon the host organism. In particular embodiments, transcription control sequences for prokaryotes may include a promoter, ribosomal binding site, and transcription termination sequence. In other particular embodiments, transcription control sequences for eukaryotes may include promoters comprising one or a plurality of recognition sites for transcription factors, transcription enhancer sequences, transcription termination sequences and polyadenylation sequences. In certain embodiments, "control sequences" can include leader sequences and/or fusion partner sequences.

[251] In some embodiments, for example but not limited to, the promoter sequences may be selected to promote the transcription of the polynucleotide in the vector. Suitable promoter sequences include, without limitation, T7 promoter, T3 promoter, SP6 promoter, beta-actin promoter, EF1a promoter, CMV promoter, and SV40 promoter. Enhancer sequences may be selected to enhance the transcription of the polynucleotide. Selectable markers may be selected to allow selection of the host cells inserted with the vector from those not, for example, the selectable markers may be genes that confer antibiotic resistance. Signal sequences may be selected to allow the expressed polypeptide to be transported outside of the host cell.

[252] A vector may also include materials to aid in its entry into the cell, including but not limited to a viral particle, a liposome, or a protein coating. In some embodiments, a host cell

comprises an expression vector disclosed herein.

[253] In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a dual antibody or a component thereof, for example but not limited to a VH domain, a VL domain, , a combined VH-VL domain as may be present in Fab elements, F(ab')₂ elements, an scFv, an Fv, a minibody, a diabody, or a triabody, as described above. Dual binding domains and the components thereof have been described in detail above.

[254] In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VH domain. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VL domain. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VH and a VL domain. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding two VH and VL domains. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding three VH and VL domains.

[255] In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VH domain component of a dual antibody. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VL domain component of a dual antibody. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding VH and VL domain components of a dual antibody.

[256] In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VH domain component of a dual IgG antibody or a fragment thereof. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VL domain component of a dual IgG antibody or a fragment thereof. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding VH and VL domain components of a dual IgG antibody or a fragment thereof.

[257] In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VH domain component of a scFv. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding a VL domain component of a scFv. In some embodiments, an expression vector comprises an isolated nucleic acid sequence encoding VH and VL domain components of a scFv.

[258] Dual binding antibodies have been described in detail above. The skilled artisan, using the knowledge in the art and specific details newly described herein would surely appreciate the range of components that may be encoded by an isolated nucleic acid described herein.

[259] For cloning of the polynucleotide, the vector may be introduced into a host cell (an isolated host cell) to allow replication of the vector itself and thereby amplify the copies of the

polynucleotide contained therein. The cloning vectors may contain sequence components generally include, without limitation, an origin of replication, promoter sequences, transcription initiation sequences, enhancer sequences, and selectable markers. These elements may be selected as appropriate by a person of ordinary skill in the art. For example, the origin of replication may be selected to promote autonomous replication of the vector in the host cell.

[260] In certain embodiments, the present disclosure provides isolated host cells containing the vector provided herein. The host cells containing the vector may be useful in expression or cloning of the polynucleotide(s) contained in the vector.

[261] In some embodiments, a recombinant host cell comprises one or more constructs as described above. A nucleic acid encoding any CDR or set of CDR's or VH domain or VL domain or antibody antigen-binding site or antibody molecule, for example but not limited to an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody. In some embodiments, disclosed herein is a method of production of the encoded product, which method comprises expression from encoding nucleic acid constructs. Expression may in some embodiments, be achieved by culturing under appropriate conditions recombinant host cells containing the nucleic acid construct. Following production by expression a VH or VL domain, or a VH-VL pair, or an antibody, may be isolated and/or purified using any suitable technique, then used as appropriate, for example in methods of treatment as described herein.

[262] In some embodiments, dual binding antibodies, VH and/or VL domains, and encoding nucleic acid molecules and vectors according to the present invention may be prepared and isolated and/or purified, in substantially pure or homogeneous form.

[263] In some embodiments, systems for cloning and expression of a polypeptide in a variety of different host cells are well known. Suitable host cells can include, without limitation, prokaryotic cells, fungal cells, yeast cells, or higher eukaryotic cells such as insect cells or mammalian cells.

[264] Suitable prokaryotic cells for this purpose include, without limitation, eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacterales such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescens*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis*, *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*.

[265] The expression of antibodies and antigen-binding fragments in prokaryotic cells such as *E. coli* is well established in the art. For a review, see for example Pluckthun, A. *Bio/Technology* 9: 545-551 (1991). Expression in eukaryotic cells in culture is also available to those skilled in the art as an option for production of antibodies or antigen-binding fragments

thereof, see recent reviews, for example Ref, M. E. (1993) Curr. Opinion Biotech. 4: 573-576; Trill J. J. et al. (1995) Curr. Opinion Biotech 6: 553-560.

[266] Suitable fungal cells for this purpose include, without limitation, filamentous fungi and yeast. Illustrative examples of fungal cells include, *Saccharomyces cerevisiae*, common baker's yeast, *Schizosaccharomyces pombe*, Kluyveromyces hosts such as, eg., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilarum* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[267] Higher eukaryotic cells, in particular, those derived from multicellular organisms can be used for expression of glycosylated VH and VL domains, as provided herein. Suitable higher eukaryotic cells include, without limitation, invertebrate cells and insect cells, and vertebrate cells. Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the K-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein as described herein, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can also be utilized as hosts. Mammalian cell lines available in the art for expression of a heterologous polypeptide include Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney cells, NS0 mouse melanoma cells, YB2/0 rat myeloma cells, human embryonic kidney cells, human embryonic retina cells and many others. Non-limiting examples of vertebrate cells include mammalian host cell lines such as monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); ExpiCHO-S(TM) cells (ThermoFisher Scientific cat. #A29133); mouse Sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRK-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442);

human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[268] In some embodiments, an expression vector comprises a nucleic acid construct described herein. Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator sequences, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. Regulatory sequences may be operably linked to the nucleic acid sequence(s) comprised within a nucleic acid construct. Vectors may be plasmids, viral e.g. phage, or phagemid, as appropriate. For further details see, for example, Molecular Cloning: a Laboratory Manual: 3rd edition, Sambrook and Russell, 2001, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in Current Protocols in Molecular Biology, Second Edition, Ausubel et al. eds., John Wiley & Sons, 1988, Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, Ausubel et al. eds., John Wiley & Sons, 4^{sup}.th edition 1999. The disclosures of Sambrook et al. and Ausubel et al. (both) are incorporated herein by reference.

[269] The vector can be introduced to the host cell using any suitable methods known in the art, including, without limitation, DEAE-dextran mediated delivery, calcium phosphate precipitate method, cationic lipids mediated delivery, liposome mediated transfection, electroporation, microprojectile bombardment, receptor-mediated gene delivery, delivery mediated by polylysine, histone, chitosan, and peptides. Standard methods for transfection and transformation of cells for expression of a vector of interest are well known in the art.

[270] In some embodiments, provided herein is a host cell containing nucleic acid as disclosed herein. Such a host cell may be *in vitro* and may be in culture. Such a host cell may be *in vivo*. *In vivo* presence of the host cell may allow intracellular expression of the dual binding antibodies described herein, as "intrabodies" or intracellular antibodies. Intrabodies may be used for gene therapy.

[271] In certain embodiments, the host cells comprise a first vector encoding a first polypeptide, e.g., a VH domain, and a second vector encoding a second polypeptide, e.g., a VL domain. In certain embodiments, the host cells comprise a vector encoding a first polypeptide, e.g., a VH domain, and a second polypeptide, e.g., a VL domain.

[272] In certain embodiments, the host cells comprise a first vector encoding a variant VH domain and a second vector encoding a variant VL domain. In certain embodiments, the host cells

comprise a single vector encoding a variant VH domain and a variant VL domain.

[273] In some embodiments, an isolated cell comprises an isolated nucleic acid sequence, as disclosed herein. In some embodiments, an isolated cell comprises two isolated nucleic acid sequences as disclosed herein, wherein one nucleic acid encodes a variant VH domain and the other nucleic acid encodes a variant VL domain. In some embodiments, an isolated cell comprises a single isolated nucleic acid sequence as disclosed herein, that encodes a variant VH domain and a variant VL domain.

[274] In certain embodiments, a first vector and a second vector may or may not be introduced simultaneously. In certain embodiments, the first vector and the second vector may be introduced together into the host cell. In certain embodiments, the first vector may be introduced first into the host cell, and then the second vector may be introduced. In certain embodiments, the first vector may be introduced into the host cell, which is then established into a stable cell line expressing the first polypeptide, and then the second vector may be introduced into the stable cell line.

[275] In certain embodiments, the host cells comprise a vector encoding for at least one variant VH domain and at least one a variant VL comprised within a dual binding antibody.

[276] The introduction may be followed by causing or allowing expression from the nucleic acid, e.g. by culturing host cells under conditions for expression of the gene. In certain embodiments, the present disclosure provides methods of expressing the polypeptide provided herein, comprising culturing the host cell containing the vector under conditions in which the inserted polynucleotide in the vector is expressed.

[277] In some embodiments, the nucleic acid is integrated into the genome (e.g. chromosome) of the host cell. Integration may be promoted by inclusion of sequences which promote recombination with the genome, in accordance with standard techniques. In some embodiments, the nucleic acid construct is not integrated into the genome and the vector is episomal.

[278] In some embodiments, disclosed herein is a method which comprises using a construct as stated above in an expression system in order to express a dual binding antibody or fragment thereof, as described herein above.

[279] Suitable conditions for expression of the polynucleotide may include, without limitation, suitable medium, suitable density of host cells in the culture medium, presence of necessary nutrients, presence of supplemental factors, suitable temperatures and humidity, and absence of microorganism contaminants. A person with ordinary skill in the art can select the suitable conditions as appropriate for the purpose of the expression.

Methods of Synthesizing an Engineered, “Re-Epitoped” Dual Antibody

[280] In some embodiments, described herein is a method of producing a dual binding antibody comprising a VH domain comprising HCDRs as described herein. In some embodiments, described herein is a method of producing a dual binding antibody comprising a VL domain comprising LCDRs as described herein. In some embodiments, described herein is a method of producing a dual binding antibody comprising a VH domain comprising HCDRs as described herein and a VL domain comprising LCDRs as described herein.

[281] In some embodiments, a method of producing a dual binding antibody a heavy chain variable region comprising: (a) the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or (c) a combination of the heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or said combination thereof of said dual binding antibody, is at least 2; comprises steps of culturing a cell or cells comprising a nucleic acid sequence encoding at least a VH and a VL of the dual binding antibody, wherein polypeptides comprising the variant VH and variant VL domains are expressed and isolated, and wherein said isolated variant VH and variant VL domains form a heterodimer. As disclosed herein in detail, the isolated nucleic acid sequences encoding variant VH and variant VL domains may be comprised within vectors, wherein the same vector or different vectors are used. In some embodiments, each variant VH domain and variant VL domain may be expressed from a different host cell, wherein dimerization occurs following isolation or purification of the component variant VH and variant VL domains. In some embodiments variant VH and variant VL domains may be expressed from a same host cell, wherein dimerization occurs in culture or following isolation or purification of the component variant VH and variant VL domains.

[282] A skilled artisan would appreciate that producing a dual binding antibody comprises synthesizing amino acid polypeptide components comprising VH domains, VL domains, or both. In some embodiments, said synthesis starts from a nucleic acid construct as described herein in detail. The terms “producing” and “synthesizing” may in some embodiments, be used herein interchangeably having all the same qualities and meanings.

[283] In some embodiments, synthesizing a dual binding antibody comprises synthesizing an

IgG heavy chain comprising a variant VH domain, synthesizing an IgG light chain comprising a variant VL domain, or both. In some embodiments, synthesizing a dual binding antibody comprises synthesizing an IgG heavy chain comprising a variant VH domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing an IgG light chain comprising a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing both an IgG heavy chain comprising a variant VH domain and an IgG light chain comprising a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing an Fab comprising a fragment of an IgG heavy chain comprising a variant VH domain and a fragment of an IgG light chain comprising a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing an F(ab')₂ comprising a fragment of an IgG heavy chain comprising a variant VH domain and a fragment of an IgG light chain comprising a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing an Fv comprising a variant VH domain and a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a scFv comprising a variant VH domain and a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a minibody comprising a variant VH domain and a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a diabody comprising a variant VH domain and a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a triabody comprising a variant VH domain and a variant VL domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a variant VH domain. In some embodiments, synthesizing a dual binding antibody comprises synthesizing a variant VL domain.

[284] In certain embodiments, the polypeptide expressed in the host cell can form a dimer and thus produce a dual binding antibody or the binding component thereof.

[285] In some embodiments, methods of synthesizing a dual binding antibody comprise a step of mutating a nucleic acid sequence encoding a template heavy chain variable region that does not comprise a dual binding VH domain in order to create a variant VH domain that may comprise a dual binding VH domain. In some embodiments, methods of synthesizing a dual binding antibody comprise a step of mutating a nucleic acid sequence encoding a template light chain variable region that does not comprise a dual binding VL domain in order to create a variant VL domain that may comprise a dual binding VL domain. In some embodiments, methods of synthesizing a dual binding antibody comprise a step of mutating a nucleic acid sequence encoding a template heavy chain variable region that does not comprise a dual binding VH domain in order to create a variant VH domain that may comprise a dual binding VH domain, and mutating a

nucleic acid sequence encoding a template light chain variable region that does not comprise a dual binding VL domain in order to create a variant VL domain that may comprise a dual binding VL domain, wherein the variant VH and VL domains comprise a dual variable region of an antibody. Methods of mutating nucleic acid sequences have been described in detail above and are exemplified below in the Examples.

[286] In some embodiments, a template nucleic acid sequence encoding the template heavy chain variable region is set forth in SEQ ID NO: 55. In some embodiments, a template nucleic acid sequence encoding the template light chain variable region is set forth in SEQ ID NO: 56. As stated throughout, template VH and VL sequences do not comprise dual binding regions.

[287] In some embodiments, methods of synthesizing a dual binding antibody comprise introducing at least 2 variant sites within VH and VL domains. In some embodiments, methods of synthesizing a dual binding antibody comprise introducing at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 variant sites within VH and VL domains. Variant sites may be distributed between the VH domain and the VL domain. In some embodiments, variant sites are within CDR regions of a VH domain. In some embodiments, variant sites are within CDR regions of a VL domain. In some embodiments, variant sites are within FR regions of a VH domain. In some embodiments, variant sites are within FR regions of a VL domain. In some embodiments, variant sites are within CDR and or FR regions of a VH domain. In some embodiments, variant sites are within CDR and or FR regions of a VL domain. In some embodiments, variant sites are within CDR and or FR regions of a VH domain, and within CDR and or FR regions of a VL domain.

[288] In certain embodiments, the variant VH and variant VL domains complex may be formed inside the host cell. For example, the variant VH and variant VL domains heterodimer may be formed inside the host cell with the aid of relevant enzymes and/or cofactors. In certain embodiments, the variant VH and variant VL domains polypeptide complex may be secreted out of the cell. In certain embodiments, the variant VH and variant VL domains may be secreted out of the host cell and form a heterodimer outside of the host cell.

[289] In certain embodiments, the variant VH and variant VL domains may be separately expressed and allowed to dimerize under suitable conditions. For example, the variant VH and variant VL domains may be combined in a suitable buffer and allow the variant VH and variant VL domains to dimerize through appropriate interactions such as hydrophobic interactions. For another example, variant VH and variant VL domains may be combined in a suitable buffer containing an enzyme and/or a cofactor which can promote the dimerization of the variant VH and variant VL domains. For another example, the variant VH and variant VL domains may be combined in a suitable vehicle and allow them to react with each other in the presence of a suitable

reagent and/or catalyst.

[290] In certain embodiments, the variant VH and variant VL domains may be comprised within longer polypeptide sequences, which may include for example but not limited to constant regions, hinge regions, linker regions, Fc regions, or disulfide binding regions, or any combination thereof. A constant domain is an immunoglobulin fold unit of the constant part of an immunoglobulin molecule, also referred to as a domain of the constant region (e.g. CH1, CH2, CH3, CH4, C κ , C λ). In some embodiments, the longer polypeptide may comprise multiple copies of a variant VH domain, a variant VL domain, or both, for example but not limited to when the dual binding antibody comprises a diabody or a triabody.

[291] In certain embodiments, the variant VH and variant VL domains are generated by DNA synthesis and PCR, and translation of nucleotide sequences generated thereof. In certain embodiments, the generated sequences may be subcloned into an expression vector. In certain embodiments, the generated sequences may be subcloned into two expression vectors. In certain embodiments, said expression vector is a plasmid. In certain embodiments, said the variant VH and variant VL domains are constructed on an IgG template, wherein said IgG template does not have dual binding capabilities.

[292] In certain embodiments, transient expression is performed by co-transfecting the expression vector encoding the variant VH and variant VL domains or by transfecting an expression vector encoding both into a suitable cell. A skilled artisan would appreciate that there are a number of transfection methods and protocols that can be used for this purpose. In certain embodiments, transfection or co-transfection is executed using the PEI method.

[293] The expressed polypeptides comprising the variant VH and variant VL domains and/or the polypeptide complex can be collected using any suitable methods. The variant VH and variant VL domains and/or the polypeptide complex can be expressed intracellularly, in the periplasmic space or be secreted outside of the cell into the medium. If the polypeptides comprising variant VH and variant VL domains and/or the polypeptide complex is expressed intracellularly, the host cells containing the polypeptides comprising variant VH and variant VL domains and/or the polypeptide complex may be lysed and polypeptide and/or the polypeptide complex may be isolated from the lysate by removing the unwanted debris by centrifugation or ultrafiltration. If the polypeptides comprising variant VH and variant VL domains and/or the polypeptide complex is secreted into periplasmic space of *E. coli*, the cell paste may be thawed in the presence of agents such as sodium acetate (pH 3.5), EDTA, and phenylmethylsulfonylfluoride (PMSF) over about 30 min, and cell debris can be removed by centrifugation (Carter et al., *BioTechnology* 10:163-167 (1992)). If the polypeptides comprising variant VH and variant VL domains and/or the

polypeptide complex is secreted into the medium, the supernatant of the cell culture may be collected and concentrated using a commercially available protein concentration filter, for example, an Amincon or Millipore Pellicon ultrafiltration unit. A protease inhibitor and/or an antibiotic may be included in the collection and concentration steps to inhibit protein degradation and/or growth of contaminated microorganisms.

[294] The expressed polypeptides comprising variant VH and variant VL domains and/or the polypeptide complex can be further purified by a suitable method, such as without limitation, affinity chromatography, hydroxylapatite chromatography, size exclusion chromatography, gel electrophoresis, dialysis, ion exchange fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin sepharose, chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation (see, for review, Bonner, P. L., Protein purification, published by Taylor & Francis, 2007; Janson, J. C., et al, Protein purification: principles, high resolution methods and applications, published by Wiley-VCH, 1998).

[295] In certain embodiments, the polypeptides comprising variant VH and variant VL domains and/or polypeptide dimer complexes can be purified by affinity chromatography. In certain embodiments, protein A chromatography or protein A/G (fusion protein of protein A and protein G) chromatography can be useful for purification of polypeptides and/or polypeptide complexes comprising a component derived from antibody CH2 domain and/or CH3 domain (Lindmark et al., J. Immunol. Meth. 62:1-13 (1983)); Zettlitz, K. A., Antibody Engineering, Part V, 531-535, 2010). In certain embodiments, a dual binding antibody disclosed herein does not bind to protein A. In certain embodiments, protein G chromatography can be useful for purification of polypeptides and/or polypeptide complexes comprising IgG γ 3 heavy chain (Guss et al., EMBO J. 5:1567 1575 (1986)). In certain embodiments, protein L chromatography can be useful for purification of polypeptides and/or polypeptide complexes comprising K light chain (Sudhir, P., Antigen engineering protocols, Chapter 26, published by Humana Press, 1995; Nilson, B. H. K. et al, J. Biol. Chem., 267, 2234-2239 (1992)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a CH3 domain, the Bakerbond ABX resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification.

[296] Following any preliminary purification step(s), the mixture comprising the dual binding antibody and contaminants may be subjected to low pH hydrophobic interaction chromatography

using an elution buffer at a pH between about 2.5-4.5, preferably performed at low salt concentrations (e.g., from about 0-0.25M salt).

[297] In certain embodiments, the polypeptides comprising variant VH and variant VL domains and/or polypeptide dimer complexes can be purified by affinity chromatography and size exclusion chromatography (SEC). A skilled artisan would appreciate that there are a number of methods and protocols suitable for this purpose. In certain embodiments, protein purification by affinity chromatography and SEC is performed using an AKTA pure instrument (GE Lifesciences). In certain embodiments, affinity capture of the dual binding antibody is achieved by passing the harvested supernatants over a column of CaptureSelect™ CH1-XL Affinity Matrix (Thermo Scientific). After washing column with PBS, the protein is eluted with 0.1M Glycine, pH 2.5, and immediately neutralized with 1/6 volume of 1M Tris-HCl, pH 8.0. The affinity purified protein is then concentrated to 5-10mg/ml using Amicon 30kD concentrator (Merck Millipore) and subjected to SEC purification on a Superdex®200 column (GE Lifesciences) equilibrated with PBS. Protein fractions are then collected and analyzed using SDS-PAGE and HPLC-SEC.

[298] Binding to an epitope of the synthesized dual binding immunoglobulins may be analyzed using well known methods in the art, as described herein, including ELISA analysis, SPR analysis, DSF analysis, and cell-based binding assays.

[299] In some embodiments, a method of synthesizing a dual binding antibody comprising: a heavy chain variable region comprising a template amino acid sequence set forth in SEQ ID NO: 1, wherein said template comprises at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and a light chain variable region comprising a template amino acid sequence set forth in SEQ ID NO: 2, wherein said template comprises at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or the combination thereof, is at least 2; comprises the following steps:

- (a) mutating the template heavy chain variable region, the template light chain variable region, or both,
- (i) wherein said mutating said template heavy variable chain region comprises mutating the template heavy chain variable region set forth in SEQ ID NO: 1, wherein said selected template variable chain does not comprise a dual binding region,

- (ii) wherein said mutating said template light chain variable region comprises mutating the template light chain variable region set forth in SEQ ID NO: 2, wherein said selected template variable chain does not comprise a dual binding region;
 - (iii) wherein said mutating both said template heavy chain variable region and said template light chain variable region comprises mutating the template heavy chain variable region set forth in SEQ ID NO: 1 and mutating the template light chain variable region set forth in SEQ ID NO: 2, wherein said selected template variable chains together do not comprise a dual binding region, and wherein said mutating comprises mutating at least two residue position in said template heavy chain variable region, said template light chain variable region or a combination thereof;
 - (b) synthesizing the mutated template variant heavy chain variable chain and the mutated template variant light chain variable chain;
 - (c) formatting said mutated template variant heavy chain variable chain and said mutated template variant light chain variable chain into a human antibody format; and
 - (d) screening the human antibody of (c) for binding to dual antigens;
- thereby producing a dual binding antibody.

[300] As described herein and exemplified below, in some embodiments, that antibody synthesized comprises an IgG immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG1 immunoglobulin, an IgG2 immunoglobulin, an IgG3 immunoglobulin, or an IgG4 immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG1 immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG2 immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG3 immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG4 immunoglobulin. In some embodiments, the antibody synthesized comprises an IgG1 immunoglobulin or an IgG4 immunoglobulin.

[301] In some embodiments, the antibody synthesized comprises an Fab immunoglobulin fragment. In some embodiments, the antibody synthesized comprises an F(ab')₂ immunoglobulin fragment. In some embodiments, the antibody synthesized comprises an Fv immunoglobulin construct. In some embodiments, the antibody synthesized comprises an scFv immunoglobulin construct. In some embodiments, the antibody synthesized comprises a minibody immunoglobulin construct comprising a pair of single-chain Fv fragments which are linked via CH3 domains.

[302] In some embodiments, the antibody synthesized comprises a diabody immunoglobulin construct. In some embodiments, the antibody synthesized comprises a diabody immunoglobulin construct comprising three scFv fragments covalently linked to each other. In some embodiments,

the antibody synthesized comprises a triabody.

[303] In some embodiments, the antibody synthesized comprises a mutated IgG that is unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, the antibody synthesized comprises a mutated IgG1 that is unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, the antibody synthesized comprises a mutated IgG4 that is unable to bind antibody-dependent cellular cytotoxicity components.

Immunoglobulin Libraries

[304] In certain embodiments, disclosed herein is a library of immunoglobulins or fragments thereof, comprising variant VH domains, variant VL domains, or variant VH and VL domains, as described herein in detail (*See*, Examples below). A library of immunoglobulins or fragments thereof comprising a variant VH domains, a variant VL domains, or variant VH and VL domains, may in some embodiments, be screened for dual binding antibodies, fragments thereof, or components thereof.

[305] In some embodiments, a library of immunoglobulins or fragments thereof, comprises a library of variable heavy chain domains. In some embodiments, a library of immunoglobulins or fragments thereof, comprises a library of variable light chain domains. In some embodiments, a library of immunoglobulins or fragments thereof, comprises a library of variable heavy chain domains and variable light chain domains.

[306] In some embodiments, a method for generating a library of dual antigen binding immunoglobulin variable heavy chain regions, for screening for binding to an epitope comprises: (a) selecting the VH template antigen-binding molecule set forth in SEQ ID NO: 1, wherein said selected template does not specifically bind an epitope; (b) selecting at least one residue position from positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof) in said template SEQ ID NO: 1, for mutation; and (c) selecting at least one variant residue to substitute at the at least one residue position selected in (b); such that a library containing a plurality of variants of said template VH is generated. In some embodiments, a method for generating a library of dual antigen binding immunoglobulin variable light chain regions, for screening for binding to an epitope comprises: (a) selecting the VL template antigen-binding molecule set forth in SEQ ID NO: 2, wherein said selected template does not specifically bind an epitope; (b) selecting at least one residue position from positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof) in said template SEQ ID NO: 2, for mutation; and (c) selecting at

least one variant residue to substitute at the at least one residue position selected in (b); such that a library containing a plurality of variants of said template VL is generated.

[307] In some embodiments, a method for generating a library of dual antigen binding immunoglobulin comprising variable heavy chain regions and variable light chain regions, for screening for binding to an epitope comprises: (a) selecting the VH template antigen-binding molecule set forth in SEQ ID NO: 1, wherein said selected template does not specifically bind an epitope; (b) selecting the VL template antigen-binding molecule set forth in SEQ ID NO: 2, wherein said selected template does not specifically bind an epitope; (c) selecting at least one residue position from positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof) in said template SEQ ID NO: 1, for mutation; (d) selecting at least one residue position from positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof) in said template SEQ ID NO: 2, for mutation; and (e) selecting at least one variant residue to substitute at the at least one residue position selected in (c) or selecting at least one variant residue to substitute at the at least one residue position selected in (d), such that the total number of variant residues in each potentially dual binding immunoglobulins is a least 2, and such that a library containing a plurality of variants of said template VH and variants of said template VL is generated.

[308] In some embodiments, methods for constructing a library can be found in the examples. In some embodiments, a library generated as described herein, may be used to identify immunoglobulins binding to dual targets. In some embodiments, a library generated as described herein, may be used to identify immunoglobulins binding to specific epitopes.

[309] In some embodiments, use of a protein library comprising an immunoglobulin comprising a variant VH, a variant VL, or a variant VH and variant VL as described herein in detail, provides a method to identify immunoglobulins binding to dual targets. In some embodiments, use of a protein library comprising an immunoglobulin comprising a variant VH, a variant VL, or a variant VH and variant VL as described herein in detail, provides a method to identify immunoglobulins binding to specific epitopes.

[310] In some embodiments, a protein library comprising an immunoglobulin comprising a variant VH and variant VL comprises a library of antibody molecules. In some embodiments, a protein library comprising an immunoglobulin comprising a variant VH and variant VL comprises a library of IgG molecules. In some embodiments, a protein library comprising an immunoglobulin comprising a variant VH and variant VL comprises a library of IgG1, IgG2,

IgG3, or IgG4 molecules. In some embodiments, the IgG molecule is a mutant IgG molecule, unable to bind antibody-dependent cellular cytotoxicity components.

[311] In some embodiments, a protein library comprising an immunoglobulin comprising a variant VH and variant VL comprises a library of Fab or F(ab')₂ molecules. In some embodiments, a protein library comprising an immunoglobulin comprising a variant VH and variant VL comprises a library of Fv molecules, scFv molecules, minibody molecules, diabody molecules, or triabody molecules.

[312] In some embodiments, existing immunoglobulin VH and VL templates can be changed to introduce variant amino acids at specific positions with the goal of generating dual antigen binding sites in said variant VH and VL domains, wherein a protein library of the variant VH and VL domains comprises at least 10; 100; 1,000; 10,000; 100,000; or 1,000,000 variant VH, variant VL, or variant VH and variant VL domains with at least two variant positions. In some embodiments, a protein library of the variant VH and VL domains comprises between 1,000 to 1,000,000 variant VH, variant VL, or variant VH and variant VL domains with at least two variant positions. In some embodiments, a protein library of the variant VH and VL domains comprises between 10,000 to 1,000,000 variant VH, variant VL, or variant VH and variant VL domains with at least two variant positions.

[313] In some embodiments, a protein library of the variant VH and VL domains comprises between 1,000 to 1,000,000 variant VH, variant VL, or variant VH and variant VL domains with at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant positions. In some embodiments, a protein library of the variant VH and VL domains comprises between 10,000 to 1,000,000 variant VH, variant VL, or variant VH and variant VL domains with at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant positions.

[314] In some embodiments, a protein library of the variant VH and VL domains comprises between 10⁶ to 10¹⁴ variant VH, variant VL, or variant VH and variant VL domains with at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant positions. In some embodiments, a protein library of the variant VH and VL domains comprises between 10⁶ to 10¹⁴ variant VH, variant VL, or variant VH and variant VL domains with at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant positions.

[315] The library is then screened for binding to one or more antigens. After molecular characterization for the desired properties a selected antibody domain or region, for example but not limited to a VH or VL domain or both, is cloned into an immunoglobulin molecule by genetic engineering techniques, so that it replaces the corresponding region. Alternatively, only the DNA coding for the VH, or VL, or both regions, or coding for the mutated amino acids may be

exchanged to obtain an immunoglobulin with the additional binding site for a molecule. In some embodiments, selection of the immunoglobulin molecule into which the variant regions are cloned, may be selected from an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody. In some embodiments, an IgG is an IgG1, an IgG2, an IgG3, or an IgG4. In some embodiments, an IgG comprises a mutant IgG unable to bind antibody-dependent cellular cytotoxicity components.

[316] In some embodiments, the CDRs expressed are as described above for HCDR1, HCDR2, HCD3, LCDR1, LCDR2, and LCDR3, wherein certain positions comprise variant amino acids, as described in detail above and is shown in **Figure 1A** and **1B**.

[317] The sites for mutation are describe above, and in certain embodiments include from positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof) in said template SEQ ID NO: 1, and positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof) in said template SEQ ID NO: 2. In some embodiments, additional sites within said VH template or said VL template may be mutated.

[318] In certain embodiments, the method of generating a library further comprises synthesizing the template variants (VH, VL, or both VH and VL) from said nucleic acid constructs, described above in detail, to form the library.

[319] The result of generating a library as described above comprises a library of immunoglobulins comprising: (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or (c) a combination of the heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in the heavy chain variable region, the light chain variable region, or a combination thereof is at least 2.

[320] Mammalian cell expression systems have been described above. These expression systems offer a number of potential advantages for therapeutic antibody generation to create a library of potential dual binding immunoglobulins, including the ability to co-select for key manufacturing- related properties such as high-level expression and stability, while displaying

functional glycosylated IgGs on the cell surface.

[321] In some embodiments, a library of immunoglobulins comprises IgG molecules, Fab molecules, F(ab')₂ molecules, FV molecules, VH molecules, VL molecules, scFv molecules, diabodies, minibodies, or triabodies. In some embodiments, an IgG molecule comprises an IgG1, an IgG2, an IgG3, or an IgG4. In some embodiments, an IgG comprises a mutated IgG that is unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, an IgG1 comprises a mutated IgG1 that is unable to bind antibody-dependent cellular cytotoxicity components.

[322] In some embodiments, disclosed herein are methods directed to screening the library with antigen molecules or a portion thereof, in order to select for dual-binding molecules that have desired properties (e.g., binding affinity, stability, etc.). In some embodiments, a portion of an antigen comprises an least one IL-13 antigenic epitope. In some embodiments, disclosed herein is a dual-binding molecule isolated from the library after said screening.

[323] In some embodiments, disclosed herein is a method for screening a library of immunoglobulins as described, for dual-binding molecules, comprising: (a) screening the library with an antigen molecules or fragment thereof to identify dual-binding molecules that bind said epitopes of interest; (b) sequencing the binders identified in step (a) to determine which residues are variants and which variant residues are enriched in the binding immunoglobulins; (c) using the information from step (b) to synthesize an optimized library of variants of the dual binders; and (d) repeating steps (a)-(c) using the optimized library. In some embodiments, disclosed herein is a method for screening a library of immunoglobulins as described, for dual-binding molecules, comprising: (a) screening the library with epitopes of interest to identify dual-binding molecules that bind said epitope of interest; (b) sequencing the binders identified in step (a) to determine which residues are variants and which variant residues are enriched in the binding immunoglobulins; (c) using the information from step (b) to synthesize an optimized library of variants of the dual binders; and (d) repeating steps (a)-(c) using the optimized library.

[324] According to some embodiments, the specific binding of the variant immunoglobulins to the antigen molecule is determined by a binding assay selected from the group consisting of immunological assays, including but not limited to enzyme linked immunosorbent assays (ELISA), surface plasmon resonance assays, saturation transfer difference nuclear magnetic resonance spectroscopy, transfer NOE (trNOE) nuclear magnetic resonance spectroscopy, competitive assays, tissue binding assays, live cell binding assays and cellular extract assays.

[325] Binding assays can be carried out using a variety of methods known in the art, including but not limited to FRET (Fluorescence Resonance Energy Transfer) and BRET (Bioluminescence

Resonance Energy Transfer)-based assays, AlphaScreen.TM. (Amplified Luminescent Proximity Homogeneous Assay), Scintillation Proximity Assay, ELISA (Enzyme-Linked Immunosorbent Assay), SPR (Surface Plasmon Resonance, also known as BIACORE®), isothermal titration calorimetry, differential scanning calorimetry, gel electrophoresis, and chromatography including gel filtration. These and other methods may take advantage of some fusion partner or label.

[326] The variant immunoglobulin is, in some embodiments, conjugated to a label selected from the group consisting of organic molecules, enzyme labels, radioactive labels, colored labels, fluorescent labels, chromogenic labels, luminescent labels, haptens, digoxigenin, biotin, metal complexes, metals, colloidal gold and mixtures thereof. Conjugation to a label may in certain embodiments, allow the simple detection of said conjugate in, for instance, binding assays (e.g. ELISA) and binding studies.

Compositions of Use

[327] In some embodiments, described herein are pharmaceutical compositions comprising the dual binding antibody, as described herein in detail, which provides a therapeutic agent. In some embodiments, described herein are pharmaceutical compositions comprising the dual binding antibody comprising a therapeutic agent comprising a mutant IgG unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments, described herein are pharmaceutical compositions comprising a dual binding antibody having therapeutic properties against allergic or respiratory conditions.

[328] In some embodiments, a pharmaceutical composition comprises a dual binding antibody comprising a variant VH, a variant VL, or a variant VH and a variant VL, and a pharmaceutically acceptable carrier. The amino acid sequences of variant VH and variant VL domains, and pair thereof, have been described in detail above (See for example, but not limited to Table 1).

[329] In certain embodiments, a composition comprises any of the isolated dual binding antibodies disclosed herein, and a pharmaceutically acceptable carrier.

[330] In one embodiment, the pharmaceutical composition comprises a dual binding antibody having HCDR1, HCDR2 and HCDR3 comprising the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and LCDR1, LCDR2 and LCDR3 comprising the amino acid sequence of SEQ ID NOs:359, 360 and 361 respectively.

[331] In another embodiment, the pharmaceutical composition comprises a dual binding antibody having HCDR1, HCDR2 and HCDR3 comprising the amino acid sequence of SEQ ID NOs:349, 356 and 351 respectively, and LCDR1, LCDR2 and LCDR3 comprising the amino acid

sequence of SEQ ID NOs:364, 360 and 371 respectively.

[332] In another embodiment, the pharmaceutical composition comprises a dual binding antibody having HCDR1, HCDR2 and HCDR3 comprising the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and LCDR1, LCDR2 and LCDR3 comprising the amino acid sequence of SEQ ID NOs:362, 360 and 384 respectively.

[333] In another embodiment, the pharmaceutical composition comprises a dual binding antibody having HCDR1, HCDR2 and HCDR3 comprising the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and LCDR1, LCDR2 and LCDR3 comprising the amino acid sequence of SEQ ID NOs:364, 360 and 384 respectively.

[334] In another embodiment, the pharmaceutical composition comprises a dual binding antibody having HCDR1, HCDR2 and HCDR3 comprising the amino acid sequence as shown in Table 8 or Table 4, and LCDR1, LCDR2 and LCDR3 comprising the amino acid sequence as shown in Table 9 or Table 5.

[335] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:209 and 210.

[336] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:219 and 220.

[337] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:249 and 250.

[338] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:337 and 338.

[339] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences as shown in Table 10 or Table 1.

[340] In another embodiment, the pharmaceutical composition comprises a dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL are at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH and VL sequences disclosed herein.

[341] In some embodiments, a pharmaceutical composition comprising a dual binding antibody comprises any dual antibody described herein comprising a variant VH, a variant VL, or

a variant VH and a variant VL. In some embodiments, a pharmaceutical composition comprising a dual binding antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of the heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in the heavy chain variable region, the light chain variable region, or a combination thereof is at least 2.

[342] A skilled artisan would recognize that in some embodiments, the term “dual binding antibody” may be used interchangeably with the term “drug” or “agent” having all the same meanings and qualities. In some embodiments, a drug comprising a dual binding antibody comprises a pharmaceutical composition.

[343] In some embodiments, described herein are compositions comprising the dual binding antibody as described herein and administration of such composition in a variety of therapeutic settings.

[344] Administration of the dual binding antibodies described herein, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration of agents for serving similar utilities. The pharmaceutical compositions can be prepared by combining a dual binding antibody or a dual binding antibody-containing composition with an appropriate physiologically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. In addition, other pharmaceutically active ingredients and/or suitable excipients such as salts, buffers and stabilizers may, but need not, be present within the composition. Administration may be achieved by a variety of different routes, including oral, parenteral, nasal, intravenous, intradermal, subcutaneous or topical. In some embodiments, modes of administration depend upon the nature of the condition to be treated or prevented. An amount that, following administration, reduces, inhibits, prevents or delays the progression and/or metastasis of a cancer is considered effective. A skilled artisan would appreciate that the term “physiologically acceptable carrier, diluent or excipient”, may in some embodiments be used interchangeably with the term “pharmaceutically acceptable carrier” having all the same means

and qualities.

[345] In some embodiments, a pharmaceutical composition described herein comprises a nucleotide sequence encoding a dual binding antibody. In some embodiments, a nucleotide sequence encoding a dual binding antibody disclosed herein, comprises a single linear nucleotide sequence. In some embodiments, a nucleotide sequence encoding a dual binding antibody disclosed herein, comprises two nucleotide sequences. In some embodiments, a nucleotide sequence encoding a dual binding antibody disclosed herein, comprises two nucleotide sequences present on the same vector. In some embodiments, a nucleotide sequence encoding a dual binding antibody disclosed herein, comprises two nucleotide sequences present on different vectors.

[346] In some embodiments, the nucleotide sequence encodes a variant VH or a variant VL domain or a combination thereof. In some embodiments, the same nucleotide sequence encodes a variant VH or a variant VL domain or a combination thereof. In some embodiments, different nucleotide sequences encode a variant VH or a variant VL domain or a combination thereof. In some embodiments, one nucleotide sequence encodes a variant VH domain and another nucleotide sequence encodes a variant VL domain. In some embodiments, one nucleotide sequence encodes variant VH domain and another nucleotide sequence encodes a variant VL domain having a linker sequence between them, thus allowing a variant VH and a variant VL domain to hetero-dimerize, as described in Duperret EK et al., Cancer Res, Oct. 4 (doi: 10.1158/0008-5472.CAN-18-1429).

[347] In some embodiments, a method of treating an allergic or respiratory condition in a subject, or a combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody comprising (a) a variant VH domain comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and (b) a variant VL domain comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); to a subject in need, wherein the method treats the allergic or respiratory condition, or a combination thereof, compared with a subject not administered said pharmaceutical composition.

[348] In some embodiments, a method of treating an allergic or respiratory condition in a subject, or a combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody comprising (a variant VH domain comprising the amino acid

sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); to a subject in need, wherein the method treats the allergic or respiratory condition, or a combination thereof, compared with a subject not administered said pharmaceutical composition.

[349] In some embodiments, a method of treating an allergic or respiratory condition in a subject, or a combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody comprising a variant VL domain comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); to a subject in need, wherein the method treats the allergic or respiratory condition, or a combination thereof, compared with a subject not administered said pharmaceutical composition.

Methods of Use

[350] In some embodiments, disclosed herein is a method of treating a subject suffering from a disease or condition, said method comprises administering to said subject a composition comprising an isolated dual binding antibody as disclosed herein. In some embodiments, the disease or condition is an allergic or respiratory condition, an inflammatory or autoimmune condition, or tumors or cancers. In some embodiments, said disease or condition is asthma, allergic asthma, nonallergic asthma, severe asthma, mild asthma, chronic obstructive pulmonary disease (COPD), a condition involving airway inflammation, cystic fibrosis, allergic lung disease, airway hyperresponsiveness, goblet cell metaplasia, mucus hypersecretion, airway remodeling, pulmonary fibrosis, atopic dermatitis, urticaria, eczema, allergic enterogastritis, allergic rhinitis, inflammatory bowel diseases, liver cirrhosis or fibrosis, or a combination thereof.

[351] In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising a heavy chain variable region comprising HCDRs (HCDR1, HCDR2, HCDR3 as described herein in detail; for example, see Table 8 or Table 4). In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a

subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising a light chain variable region comprising LCDRs (LCDR1, LCDR2, LCDR3 as described herein in detail; for example, see Table 9 or Table 5). In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising a heavy chain variable region comprising HCDRs (HCDR1, HCDR2, HCDR3) and LCDRs (LCDR1, LCDR2, LCDR3 as described herein in detail).

[352] In certain embodiments, a method of treating a subject suffering from a disease or condition comprises administering a dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:359, 360 and 361 respectively; or the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 371 respectively; or the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:362, 360 and 384 respectively; or the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 384 respectively; or the CDRs having the sequences of SEQ ID NOs:149-154.

[353] In some embodiments a method of treating a subject suffering from a disease or condition comprises administering a dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.

[354] In some embodiments a method of treating a subject suffering from a disease or condition comprises administering a dual binding antibody comprising VH and VL having the

sequences of SEQ ID Nos:209 and 210, SEQ ID Nos:219 and 220, SEQ ID Nos:249 and 250, SEQ ID Nos:337 and 338, SEQ ID NOs:155 and 156, SEQ ID NOs:157 and 158. In some embodiments a method of treating a subject suffering from a disease or condition comprises administering a dual binding antibody comprising VH and VL domains having the sequences as shown in Table 10 or Table 1.

[355] In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or a combination of the heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or said combination thereof, is at least 2, to a subject in need, wherein the method treats the allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in said subject, compared with a subject not administered said dual binding antibody, or pharmaceutical composition thereof.

[356] In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof), wherein the total number of variant positions in said heavy chain variable region is at least 2, to a subject in need, wherein the method treats the allergic

or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in said subject, compared with a subject not administered said dual binding antibody, or pharmaceutical composition thereof.

[357] In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof), wherein the total number of variant positions in said light chain variable region is at least 2, to a subject in need, wherein the method treats the allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in said subject, compared with a subject not administered said dual binding antibody, or pharmaceutical composition thereof.

[358] In some embodiments, a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in a subject, or any combination thereof, comprises a step of administering a pharmaceutical composition comprising a dual binding antibody or a pharmaceutical composition thereof, said dual binding antibody comprising (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); wherein the total number of variant positions in said heavy chain variable region, said light chain variable region, or said combination thereof is at least 2, to a subject in need, wherein the method treats the allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma in said subject, compared with a subject not administered said dual binding antibody, or pharmaceutical

composition thereof.

[359] In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, the amino acid sequence of the variant VH domain is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54. In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, a dual binding antibody comprises a heavy chain variable region comprising the amino acid sequences set forth in, but not limited to, any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54; and any variable light chain region. In some embodiments of a method disclosed herein, the amino acid sequence of the variant VH domain comprises the sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99% identical) to the sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54.

[360] In some embodiments of method disclosed herein, the VH domain of the dual binding antibody is selected from the sequences set forth in any one of SEQ ID NOs: 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, 253, 255, 257, 259, 261, 263, 265, 267, 269, 271, 273, 275, 277, 279, 281, 283, 285, 287, 289, 291, 293, 295, 297, 299, 301, 303, 305, 307, 309, 311, 313, 315, 317, 319, 321, 323, 325, 327, 329, 331, 333, 335, 337, 339, 341, 343, 345 and 347. In another embodiment, the VH domain is at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH sequences disclosed herein.

[361] One skilled in the art would appreciate that percent sequence identity may be determined using any of a number of publicly available software application, for example but not limited to BlastP software of the National Center of Biotechnology Information (NCBI) using default parameters.

[362] In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, the amino acid sequence of the variant light chain variable region (VL) is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53. In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, a dual binding antibody comprises a light chain

variable region comprising the amino acid sequences set forth in, but not limited to, any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53, and any variable heavy chain region. In some embodiments of a method disclosed herein, the amino acid sequence of the variant VH domain comprises the sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99% identical) to the sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53, and any variable heavy chain region.

[363] In some embodiments of methods disclosed herein, the VL domain of the dual binding antibody is selected from the sequences set forth in any one of SEQ ID NOs: 210, 212, 214, 216, 218, 220, 222, 224, 226, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, 254, 256, 258, 260, 262, 264, 266, 268, 270, 272, 274, 276, 278, 280, 282, 284, 286, 288, 290, 292, 294, 296, 298, 300, 302, 304, 306, 308, 310, 312, 314, 316, 318, 320, 322, 324, 326, 328, 330, 332, 334, 336, 338, 340, 342, 344, 346 and 348. In another embodiment, the VL domain is at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VL sequences disclosed herein.

[364] In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, the amino acid sequence of the variant VH domain is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, and 54; and the amino acid sequence of the variant light chain variable region (VH) is selected from, but not limited to, the sequences set forth in any of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, 17, 19 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, and 53. In some embodiments of a method of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, the amino acid sequences of a heavy chain variable region – light chain variable region pair are selected from, but not limited to, the pair sequences set forth in SEQ ID Nos: 4 and 3, SEQ ID Nos: 6 and 5, SEQ ID Nos: 8 and 7, SEQ ID Nos: 10 and 9, SEQ ID Nos: 12 and 11, SEQ ID Nos: 14 and 13, SEQ ID Nos: 16 and 15, SEQ ID Nos: 18 and 17, SEQ ID Nos: 20 and 19, SEQ ID Nos: 22 and 21, SEQ ID Nos: 24 and 23, SEQ ID Nos: 26 and 25, SEQ ID Nos: 28 and 27, SEQ ID Nos: 30 and 29, SEQ ID Nos: 32 and 31, SEQ ID Nos: 34 and 33, SEQ ID Nos: 36 and 35, SEQ ID Nos: 38 and 37, SEQ ID Nos: 40 and 39, SEQ ID Nos: 42 and 41, SEQ ID Nos: 44 and 43, SEQ ID Nos: 46 and 45, SEQ ID Nos: 48 and 47, SEQ ID Nos: 50 and 49, SEQ ID Nos: 52 and 51, and SEQ ID Nos: 54 and 53. In some embodiments, the amino acid sequences of the VH-VL pair are selected from the pair sequences set forth in any one of the following: SEQ ID

NOs:209 and 210, SEQ ID NOs:211 and 212, SEQ ID NOs:213 and 214, SEQ ID NOs:215 and 216, SEQ ID NOs:217 and 218, SEQ ID NOs:219 and 220, SEQ ID NOs:221 and 222, SEQ ID NOs:223 and 224, SEQ ID NOs:225 and 226, SEQ ID NOs:227 and 228, SEQ ID NOs:229 and 230, SEQ ID NOs:231 and 232, SEQ ID NOs:233 and 234, SEQ ID NOs:235 and 236, SEQ ID NOs:237 and 238, SEQ ID NOs:239 and 240, SEQ ID NOs:241 and 242, SEQ ID NOs:243 and 244, SEQ ID NOs:245 and 246, SEQ ID NOs:247 and 248, SEQ ID NOs:249 and 250, SEQ ID NOs:251 and 252, SEQ ID NOs:253 and 254, SEQ ID NOs:255 and 256, SEQ ID NOs:257 and 258, SEQ ID NOs:259 and 260, SEQ ID NOs:261 and 262, SEQ ID NOs:263 and 264, SEQ ID NOs:265 and 266, SEQ ID NOs:267 and 268, SEQ ID NOs:269 and 270, SEQ ID NOs:271 and 272, SEQ ID NOs:273 and 274, SEQ ID NOs:275 and 276, SEQ ID NOs:277 and 278, SEQ ID NOs:279 and 280, SEQ ID NOs:281 and 282, SEQ ID NOs:283 and 284, SEQ ID NOs:285 and 286, SEQ ID NOs:287 and 288, SEQ ID NOs:289 and 290, SEQ ID NOs:291 and 292, SEQ ID NOs:293 and 294, SEQ ID NOs:295 and 296, SEQ ID NOs:297 and 298, SEQ ID NOs:299 and 300, SEQ ID NOs:301 and 302, SEQ ID NOs:303 and 304, SEQ ID NOs:305 and 306, SEQ ID NOs:307 and 308, SEQ ID NOs:309 and 310, SEQ ID NOs:311 and 312, SEQ ID NOs:313 and 314, SEQ ID NOs:315 and 316, SEQ ID NOs:317 and 318, SEQ ID NOs:319 and 320, SEQ ID NOs:321 and 322, SEQ ID NOs:323 and 324, SEQ ID NOs:325 and 326, SEQ ID NOs:327 and 328, SEQ ID NOs:329 and 330, SEQ ID NOs:331 and 332, SEQ ID NOs:333 and 334, SEQ ID NOs:335 and 336, SEQ ID NOs:337 and 338, SEQ ID NOs:339 and 340, SEQ ID NOs:341 and 342, SEQ ID NOs:343 and 344, SEQ ID NOs:345 and 346, SEQ ID NOs:347 and 348.

[365] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences of SEQ ID NOs:149-154. In another embodiment, the dual binding antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL) having the amino acid sequences of SEQ ID Nos:155 and 156, or SEQ ID Nos:157 and 158.

[366] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences set forth in SEQ ID NOs:349, 350 and 351, respectively, and SEQ ID NOs:359, 360, and 361, respectively.

[367] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences set forth in SEQ ID NOs:349, 356, and 351, respectively, and SEQ ID NOs:364, 360 and 371 respectively.

[368] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences set forth in SEQ ID NOs:349, 350, and 351, respectively, and SEQ ID NOs:362, 360 and 384 respectively.

[369] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences set forth in SEQ ID NOs:349, 350, and 351, respectively, and SEQ ID NOs:364, 360 and 384 respectively.

[370] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the CDRs have the sequences set forth in Table 8 or Table 4, and Table 9 or Table 5.

[371] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein the VH and VL comprise the amino acid sequences of SEQ ID Nos:209 and 210.

[372] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising the amino acid sequences of SEQ ID Nos:219 and 220.

[373] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an

isolated dual binding antibody comprising amino acid sequences of SEQ ID Nos:249 and 250.

[374] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising the amino acid sequences of SEQ ID Nos:337 and 338.

[375] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody comprising the amino acid sequences as shown in Table 10 or Table 1.

[376] In some embodiments, a method of treating one or more conditions in a subject as described herein comprises a step of administering a pharmaceutical composition comprising an isolated dual binding antibody, wherein the VH and VL are at least 80%, 85%, 90%, 95%, 98%, or 99% identical to the VH and VL sequences disclosed herein.

[377] The variant VH and VL domains have been described in detail above, including methods of creating these variant VH and VL domains by re-epitoping template sequence. That disclosure is incorporated herein in full, wherein methods of treating an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma comprise use of any variant VH or VL domain described herein.

[378] A skilled artisan would appreciate that the term “treating” and grammatical forms thereof, may in some embodiments encompass both therapeutic treatment and prophylactic or preventative measures with respect to an allergic or respiratory condition, as described herein, wherein the object is to treat, prevent, reduce, or alleviate, the allergic or respiratory condition, or symptoms thereof, or a combination thereof. Thus, in some embodiments of methods disclosed herein, treating may include directly affecting or curing, suppressing, inhibiting, preventing, reducing the severity of, delaying the onset of, reducing symptoms associated with the disease, disorder or condition, or a combination thereof; for example, when said disease or disorder comprises an allergic or respiratory condition. In some embodiments, “treating” encompasses enhancing the ability of host immune cells to destroy intracellular pathogens. In some embodiments, “treating” encompasses interference with the IL-13receptor/IL-4receptor signaling cascade. In some embodiments, “treating” encompasses inhibition of IL-13 activities. In some embodiments, “treating” encompasses reduction of IL-13 activities.

[379] In some embodiments, “preventing” encompasses delaying the onset of symptoms or an allergic or respiratory condition. In some embodiments, “suppressing” or “inhibiting”, encompass reducing the severity of symptoms, reducing the severity of an acute episode, reducing

the number of symptoms, reducing the incidence of disease-related symptoms, reducing the latency of symptoms, ameliorating symptoms, reducing secondary symptoms, reducing secondary infections, prolonging patient survival, or a combination thereof.

[380] In one embodiment, the subject is a mammal, e.g., a human suffering from one or more IL-13-associated disorders, including but not limited to respiratory disorders or conditions (e.g., asthma (e.g., allergic and nonallergic asthma for example but not limited to asthma due to infection with, e.g., respiratory syncytial virus (RSV), e.g., in younger children), severe asthma, mild asthma), chronic obstructive pulmonary disease (COPD), and other conditions involving airway inflammation, eosinophilia, fibrosis and excess mucus production (e.g., cystic fibrosis and pulmonary fibrosis); atopic disorders (e.g., atopic dermatitis, urticaria, eczema, allergic enterogastritis, and allergic rhinitis); inflammatory and/or autoimmune conditions of, the skin, gastrointestinal organs (e.g., inflammatory bowel diseases (IBD), such as ulcerative colitis and/or Crohn's disease), and liver (e.g., cirrhosis, fibrosis); scleroderma; or tumors or cancers, e.g., Hodgkin's lymphoma.

[381] In some embodiments, methods of treating comprising treating, reducing, preventing, or ameliorating, symptoms of an allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma, or any combination thereof. For example, symptoms of asthma may include, but are not limited to, wheezing, shortness of breath, bronchoconstriction, airway hyperreactivity, decreased lung capacity, fibrosis, airway inflammation, and mucus production. The method comprises administering to the subject a dual binding antibody or a pharmaceutical composition thereof, as described herein, in an amount sufficient to treat (e.g., reduce, ameliorate) or prevent one or more symptoms. The dual binding antibody can be administered therapeutically or prophylactically, or both. The dual binding antibody can be administered to the subject, alone or in combination with other therapeutic modalities.

[382] The precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by testing the compositions in model systems known in the art and extrapolating therefrom. Controlled clinical trials may also be performed. Dosages may also vary with the severity of the condition to be alleviated. A pharmaceutical composition is generally formulated and administered to exert a therapeutically useful effect while minimizing undesirable side effects. The composition may be administered one time or may be divided into a number of smaller doses to be administered at intervals of time. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need.

[383] In some embodiments, compositions comprising a nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding antibody comprising: (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); or (c) a combination of heavy chain variable region set forth in (a) and the light chain variable region set forth in (b); wherein the total number of variant positions in the encoded heavy chain variable region, the encoded light chain variable region, or a combination thereof, is at least 2, may be administered alone or in combination with other known allergic or respiratory condition, an inflammatory and/or autoimmune condition of the skin or gastrointestinal organs; scleroderma; or tumors or cancers including Hodgkin's lymphoma treatments.

[384] In some embodiments, methods of treatment comprise administration of a composition comprising a nucleic acid construct comprising a dual binding antibody comprising an VH domain comprising HCDRs (HCDR1, HCDR2, HCDR3 as described herein; for example, see Table 8 or Table 4). In some embodiments, methods of treatment comprise administration of a composition comprising a nucleic acid construct comprising a dual binding antibody comprising an VL domain comprising LCDRs (LCDR1, LCDR2, LCDR3 as described herein; for example, see Table 9 or Table 5). In some embodiments, methods of treatment comprise administration of a composition comprising a nucleic acid construct comprising a dual binding antibody comprising an VH domain comprising HCDRs (HCDR1, HCDR2, HCDR3 as described herein) and a VL domain comprising LCDRs (LCDR1, LCDR2, LCDR3 as described herein).

[385] In some embodiments, compositions comprising a nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding antibody comprising: (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); wherein the total number of variant positions in the encoded heavy chain variable region is at least 2, may be administered alone or in combination with other known allergic or respiratory condition treatments. In some embodiments, compositions comprising a nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding

antibody comprising: a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); wherein the total number of variant positions in the encoded light chain variable region is at least 2, may be administered alone or in combination with other known allergic or respiratory condition treatments. In some embodiments, compositions comprising a nucleic acid construct, comprising a nucleic acid sequence encoding a dual binding antibody comprising: (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1 with at least one amino acid variant at any of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2 with at least one amino acid variant at any of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof); wherein the total number of variant positions in the encoded heavy chain variable region, the encoded light chain variable region, or a combination thereof, is at least 2, may be administered alone or in combination with other known allergic or respiratory condition treatments.

[386] In some embodiments of a method of treating an allergic or respiratory condition, the nucleotide construct encoding the variant VH domain is selected from the sequences set forth in any of SEQ ID NOs: 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 105, and 107. In some embodiments of a method of treating an allergic or respiratory condition, the nucleotide construct encoding the variant light chain variable region (VH) is selected from the sequences set forth in any of SEQ ID NOs: 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, and 108. In some embodiments of a method of treating an allergic or respiratory condition, the nucleotide construct encoding the variant VH domain is selected from the sequences set forth in any of SEQ ID NOs: 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, 89, 91, 93, 95, 97, 99, 101, 105, and 107; and, the nucleotide construct encoding the variant light chain variable region (VH) is selected from the sequences set forth in any of SEQ ID NOs: 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, and 108. In some embodiments of a method of treating an allergic or respiratory condition, the nucleotide construct encoding a dual binding antibody heavy chain variable region – light chain variable region scFv, is selected from the sequences set forth in SEQ ID NOs: 109-135.

[387] The nucleotide sequences encoding variant VH and VL domains have been described

in detail above, including methods of creating these nucleotide sequences encoding the variant VH and VL domains by mutating template sequence. That disclosure is incorporated herein in full, wherein methods of treating an allergic or respiratory condition comprise use of any nucleic acid construct encoding variant VH or VL domain described herein.

[388] In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody described herein comprises an IgG immunoglobulin. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an IgG1 immunoglobulin, an IgG2 immunoglobulin, an IgG3 immunoglobulin, or an IgG4 immunoglobulin. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an IgG1 immunoglobulin. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an IgG2 immunoglobulin. In some embodiments, a dual binding antibody comprises an IgG3 immunoglobulin. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an IgG4 immunoglobulin. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an IgG1 immunoglobulin or an IgG4 immunoglobulin.

[389] In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a mutated IgG, said mutant IgG unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a mutated IgG1, said mutant IgG1 unable to bind antibody-dependent cellular cytotoxicity components. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a mutated IgG4, said mutant IgG4 unable to bind antibody-dependent cellular cytotoxicity components.

[390] In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an Fab immunoglobulin fragment. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an F(ab')₂ immunoglobulin fragment. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an Fv immunoglobulin construct. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises an scFv immunoglobulin construct. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a minibody immunoglobulin construct comprising a pair of single-chain Fv fragments which are linked via CH3 domains.

[391] In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a diabody immunoglobulin construct. In some embodiments of a method of treating an allergic or respiratory condition, a dual binding antibody comprises a triabody immunoglobulin construct.

[392] Typical routes of administering these and related dual binding antibodies or pharmaceutical compositions thereof, include, without limitation, oral, topical, transdermal, inhalation, parenteral, sublingual, buccal, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Dual binding antibodies or pharmaceutical compositions thereof according to certain embodiments as described herein, are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Dual binding antibodies or pharmaceutical compositions thereof that will be administered to a subject or patient may take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a herein described dual binding antibody in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington: The Science and Practice of Pharmacy, 20th Edition (Philadelphia College of Pharmacy and Science, 2000). The dual binding antibodies or pharmaceutical compositions thereof to be administered will, in any event, contain a therapeutically effective amount of a dual binding antibody of the present disclosure, for treatment of an allergic or respiratory condition.

[393] A pharmaceutical composition may be in the form of a solid or liquid. In some embodiments, the pharmaceutically acceptable carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The pharmaceutically acceptable carrier(s) may be liquid, with the compositions being, for example, an oral oil, injectable liquid or an aerosol, which is useful in, for example, inhalatory administration. When intended for oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

[394] As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like. Such a solid composition will typically contain one or more inert diluents or edible pharmaceutically acceptable carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as

alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent. When the pharmaceutical composition is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid pharmaceutically acceptable carrier such as polyethylene glycol or oil.

[395] The pharmaceutical composition may be in the form of a liquid, for example, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

[396] The liquid pharmaceutical compositions, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

[397] A liquid pharmaceutical composition intended for either parenteral or oral administration should contain an amount of a dual binding antibody as herein disclosed such that a suitable dosage will be obtained.

[398] The pharmaceutical composition may be intended for topical administration, in which case the pharmaceutically acceptable carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device. The pharmaceutical composition may be intended for

rectal administration, in the form, for example, of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

[399] The pharmaceutical composition may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredient (a dual binding antibody) may be encased in a gelatin capsule. The pharmaceutical composition in solid or liquid form may include an agent that binds to the antibody as disclosed herein, and thereby assists in the delivery of the compound. Suitable agents that may act in this capacity include other monoclonal or polyclonal antibodies, one or more proteins or a liposome. The pharmaceutical composition may consist essentially of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, subcontainers, and the like, which together may form a kit. One of ordinary skill in the art, without undue experimentation may determine preferred aerosols.

[400] The pharmaceutical compositions may be prepared by methodology well known in the pharmaceutical art. For example, a pharmaceutical composition intended to be administered by injection can be prepared by combining a composition that comprises a dual binding antibody as described herein and optionally, one or more of salts, buffers and/or stabilizers, with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the dual binding antibody composition so as to facilitate dissolution or homogeneous suspension of the dual binding antibody in the aqueous delivery system.

[401] The compositions may be administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the dual binding antibody employed; the metabolic stability and length of action of the dual binding antibody; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular allergic or respiratory disorder or condition; and the subject undergoing therapy.

[402] Compositions comprising the dual binding antibody of the present disclosure or comprising a nucleotide sequence encoding the dual binding antibody may also be administered simultaneously with, prior to, or after administration of one or more other therapeutic agents. Such combination therapy may include administration of a single pharmaceutical dosage formulation which contains a dual binding antibody as disclosed herein, and one or more additional active agents, as well as administration of compositions comprising dual binding antibody as disclosed herein, and each active agent in its own separate pharmaceutical dosage formulation. For example, a dual binding antibody or comprising a nucleotide sequence encoding the dual binding antibody, as described herein, and the other active agent can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Similarly, a dual binding antibody or comprising a nucleotide sequence encoding the dual binding antibody, as described herein, and the other active agent can be administered to the patient together in a single parenteral dosage composition such as in a saline solution or other physiologically acceptable solution, or each agent administered in separate parenteral dosage formulations. Where separate dosage formulations are used, the compositions comprising dual binding antibody or comprising a nucleotide sequence encoding the dual binding antibody, and one or more additional active agents can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially and in any order; combination therapy is understood to include all these regimens.

[403] Thus, in certain embodiments, also contemplated is the administration of dual binding antibody compositions of this disclosure or comprising a nucleotide sequence encoding the dual binding antibody, in combination with one or more other therapeutic agents. Such therapeutic agents may be accepted in the art as a standard treatment for a particular disease state as described herein, such as an allergic or respiratory condition.

[404] In some embodiments, a pharmaceutically acceptable carrier may be liquid, semi-liquid or solid. Solutions or suspensions used for parenteral, intradermal, subcutaneous or topical application may include, for example, a sterile diluent (such as water), saline solution, fixed oil, polyethylene glycol, glycerin, propylene glycol or other synthetic solvent; antimicrobial agents (such as benzyl alcohol and methyl parabens, phenols or cresols, mercurials, chlorobutanol, methyl and propyl p-hydroxybenzoic acid esters, thimerosal, benzalkonium chloride and benzethonium chloride); antioxidants (such as ascorbic acid and sodium bisulfite; methionine, sodium thiosulfate, platinum, catalase, citric acid, cysteine, thioglycerol, thioglycolic acid, thiosorbitol, butylated hydroxyanisole, butylated hydroxytoluene, and/or propyl gallate) and chelating agents (such as ethylenediaminetetraacetic acid (EDTA)); buffers (such as acetates,

citrates and phosphates). If administered intravenously, suitable pharmaceutically acceptable carriers include physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, polypropylene glycol and mixtures thereof.

[405] The compositions comprising a dual binding antibody as described herein, may be prepared with pharmaceutically acceptable carriers that protect the dual binding antibody against rapid elimination from the body, such as time release formulations or coatings. Such pharmaceutically acceptable carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others known to those of ordinary skill in the art.

[406] As used herein, the singular form "a", "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "an immunoglobulin" or "at least one immunoglobulin" may include a plurality of immunoglobulins, including mixtures thereof.

[407] Throughout this application, various embodiments may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed sub ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

[408] Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals there between.

EXAMPLES

Example 1: Experimental Procedures

[409] *Objective:* To generate unique, dual binding antibodies.

[410] *Library Design Methods:*

[411] To generate a dual binding antibody that binds IL-13 and TSLP, a "re-epitoping" approach was applied to an existing antibody. The re-epitoping process allows for the introduction

of new specificity to an existing antibody, e.g. a known antibody with 3-dimensional structure and well established biochemical and biophysical properties. The re-epitoped antibody is likely therefore to have both a new specificity and desirable developability properties. Briefly, re-epitoping is an engineering approach that allows the redirection of an existing antibody toward a new epitope, possibly on a new antigen unrelated to the cognate antigen of the original antibody. The computational process of re-epitoping requires two steps: (i) using any computational analysis that identifies putative contacts between an existing antibody and a new epitope, and (ii) application of any computational analysis or tool that can suggest the introduction of specific mutations into the antibody that are predicted to enhance its binding to the new, desired epitope. In (Ref : Ofra Y et al, US20180068055A1; Nimrod G, et al, Cell Rep. 2018 Nov 20;25(8):2121-2131) some examples of possible such computational processes are presented. In particular, two libraries were designed using the sequence of the variable domains of a template antibody (SEQ ID NO: 1 -template variable heavy chain sequence; SEQ ID NO: 2 – template variable light chain sequence)) as a starting point.

[412] Template Variable Heavy Chain:

QMQLVESGGGVVQPGRSLRLSCAASGFTFRTYGMHWVRQAPGKGLEWVAVIWDYG
SNKHYADSVKGRFTITRDNSKNLTNLQMNSLRAEDTAVYYCARAPQWELVHEAFDI

WGQGTMVTSS (SEQ ID NO: 1). Template Variable Light Chain:

SYVLTPPPSVSVAPGQTARITCGGNLGSKSVHWYQQKPGQAPVLVVYDDSDRPSWI
PERFSGSNSGTATLTISRGEAGDEADYYCQVWDSSSDHVVFGGGTKLTVL (SEQ ID
NO: 2).

[413] Each library contained 21 positions that were chosen for variation with respect to the template original sequences. These positions are located in both CDRs (H2, H3, L1, L2, and L3) and framework (**Figures 1A-1B**). The following positions were chosen for variation in the libraries (IMGT® numbering scheme [the international ImMunoGeneTics information system® <http://www.imgt.org>):

Variable H chain (**SEQ ID NO: 1**): 57(H2), 107(H3), 108(H3), 109(H3), 110(H3), 111(H3), 111A(H3), 112A(H3), 112(H3), 113(H3), 114(H3), 117(H3).

[414] Variable L chain (**SEQ ID NO: 2**): 27(L1), 28(L1), 38 (FR2), 65(L2), 70(FR3), 94(FR3), 109(L3), 110(L3), 115(L3).

[415] The resulting IL13/TSLP binding antibodies comprising variant heavy chain/variant light chain pairs, included a clone (C2) that contained 8 mutations relative to the template starting sequences (See, **Figures 1A** and **1B**).

[416] *Library Construction Methods:*

[417] Libraries were constructed on the 5J13 template (PDB5J13) by overlapping extension PCR with degenerate oligonucleotides encoding the diversity 2×10^{14} . PCR to introduce diversity was done using Phusion high fidelity DNA polymerase (New England Biolabs USA, Cat: M0530) according to manufacturer instructions in a 3-step reaction (98°C for 30 sec, 65°C for 20 sec, 72°C for 30 sec, 30 cycles). The PCR products were gel purified by gel purification kit and assembled (100ng from each) in equimolar ratios in a 3-step PCR reaction, as above, in the absence of primers. The assembled PCR product was reused as the template for PCR amplifying the full scFv library, as above, using forward and reverse primers adding vector sequences 5' and 3' to the scFv library to efficiently perform homologous recombination in yeast cells.

[418] Library transformation was carried out as published (Benatuil *et al.*, (2010) An improved yeast transformation method for the generation of very large human antibody libraries. *Protein Eng. Des. Sel.* **23**, 155–159. 400µl of a yeast suspension (EBY100, ATCC, USA) per 0.2cm cuvette (cell projects) was electroporated (BioRad, USA, GenePulser) with 4µg linearized vector (pCTcon3) and 12µg DNA insert (scFv Library) in a 1:3 vector to insert ratio (Chao, G. *et al.* Isolating and engineering human antibodies using yeast surface display. *Nat. Protoc.* **1**, 755–768 (2006)). The number of transformants of each library was determined to $\sim 1 \times 10^8$ by serial dilutions of transformed cells (Benatuil *et al.* (2010) *ibid*)

[419] *Methods of Screening and Selection using Yeast Surface display:*

[420] Yeast-displayed scFv libraries were grown in a SDCAA selective medium and induced for expression with 2% w/v galactose at 30 °C overnight according to established protocols (Chao *et al.*, (2006) *ibid*) The library was screened on BioRad S3e Fluorescence Activated Cell Sorter for high affinity binders of rh-IL-13-Fc (Reprokine, Israel) using mouse anti Myc-FITC (Santa Cruze, USA) and goat anti human Fc-APC (Jackson Immuno research, USA). Isolated clones from the final sort were sequenced by extraction of plasmid DNA from the yeast clones using a Zymoprep kit (Zymo Research, USA) and the DNA was sequenced. The chosen clones were incubated with either 10nM recombinant human IL-13 (rh-IL-13)-Fc or 10nM recombinant human TSLP (hTSLP)-Fc for 1 hour at room temperature. Cells were washed and resuspended in ice-cold PBS 0.1% BSA buffer containing a fluorescent labeled secondary antibody as described above for 20 min and analyzed using a flow cytometer. The values obtained were normalized to expression levels and to a positive control (an anti-IL-13 or anti TSLP binding antibody).

[421] *Methods of IgG production - Production of the IgGs including the light chain (LC) and heavy chain (HC) variable regions:*

[422] Sequences of the selected clones were synthesized as GeneBlock (GB) with 5' 25bp region homologous to the cloning regions of pSF-CMV-HuIgG1_HC and pSF-CMV-

HuLambda_LC (Oxford genetics, Oxford UK), the GB codon usage was optimized for mammalian expression (integrated DNA Technologies. Coralville, Iowa USA). The pSF-CMV-HuIgG1_HC and pSF-CMV-HuLambda_LC were digested with using BseRI and NcoI, and the LC and HC variable region DNA fragments were cloned into the expression corresponding vectors using NEBuilder (NEB Ipswich, Massachusetts, USA). The expression vectors were transfected and expressed in ExpiCHO Expression System (ThermoFisher Scientific, USA) according to the manufacturer's instructions. Briefly: 25ml CHO cells were grown at 37°C to a density of 6×10^6 cell/ml, 25µg expression vector 1:2 HC/LC ratio were transfected into CHO cells, 20 hours post transfection the cells growth conditions were changed to 32°C with 120 rpm shaking for 10 days. Subsequently the cells were centrifuged and IgGs were purified from the supernatant using proteinA beads, followed by size exclusion chromatography on a Superdex® 200 10/300 increase column (GE) with PBS serving as mobile phase.

[423] *Methods of determination of IgG EC50 binding to human and cynomolgus monkey TSLP*

[424] Plates (Greiner Bio-One Cat:655081) were coated with 45.5ng/well human or cynomolgus monkey (cyno) TSLP antigen, then washed and blocked with 3% skim milk in PBS with 0.05% tween. Post blocking the tested IgG was added to the wells in a concentration range of 1nM-1000nM and incubated for 1 hour at room temperature (RT). The plates were washed and goat anti-human Fc- HRP conjugated secondary antibody (Jackson cat:109-035-008) diluted 1:20000 in PBS, was added. The reaction was developed and stopped using TMB (Southern-Biotech cat:0410-01) and stop solution (Southern-Biotech cat:0412-01) respectively and read at 450nm.

[425] *Methods of determination of IC50 competition between IL-13 and TSLP*

[426] Plates were coated with 1ng/ul hTSLP washed with TBS 0.05% tween (TBS-T) and blocked with TBS-T 2% BSA. 20nM of tested IgG was incubated with rhIL-13 at a concentration range of 0.78nM to 200nM for 1 hour, then the mixture was loaded on the plates for 10 minutes and the wells were washed and a bound IgG was detected using anti human Fc-HRP conjugate as described for the ELISA EC50 experiment above. A reciprocal competition experiment was conducted using the same conditions except this time IL-13 was coated on the wells, and TSLP served as free competing ligand at the same concentration range.

[427] *Methods of determination of IgG IC50 inhibition constant of blocking TSLP from binding to TSLP-R:*

[428] 150ng/well of TSLP-R-Fc tag (ACRO biosystems TSR-H525a) was diluted in 0.015M NaHCO₃, pH=9.5, and was then used to coat the wells of a 96 well plate (Greiner Bio-One

Cat:655081). Wells were then washed three times with TBS 0.05% tween (TBS-T) and blocked with TBS-T containing 2% BSA (w/v). Competitor IgG at a concentration range of 0.11nM to 300nM was mixed with 3nM hTSLP-His (ACRO biosystems cat: TSP-H52Hb) for one hour, then the mixture was loaded into the wells of the 96 well plate, incubated for 10 minutes, followed by washing the plate three times with TBS-T. Subsequently 1:200 anti-His-HRP conjugated secondary antibody was added (Santa Cruz Biothechnology cat SC-8036). The reaction was developed and stopped using TMB (Southern-Biotech cat:0410-01) and stop solution (Southern-Biotech cat:0412-01) respectively, and read at 450nm.

[429] *Methods of specificity determination by ELISA*

[430] 96 well plates (Greiner Bio-One Cat:655081) were coated with a total of 250ng ligand, blocked with PBS-T containing 0.5% (w/v) BSA, and incubated with 100nM IgG. Plates were developed using the same reagents and conditions as in the TSLP EC50 experiment described herein.

[431] *Methods of Calibration of MUTZ5 TSLP reporter cell line:*

[432] The *in-vitro* activity of anti-TSLP blockade of TSLP binding to its cognate receptor is based on detection of pSTAT5 activation by human TSLP in MUTZ5 human leukemia cell line (Francis et al., (2016) Hematopoiesis, 101(4):417-426). In order to determine the EC50 value of hTSLP STAT5 activation of MUTZ5 cell line, cells were inoculated in a total volume of 150µl, 250×10^5 cells/well and incubated for 1hr at 37°C 5% CO₂ in a 96 well plate. Then TSLP at concentration range of 0.1pg/ml to 1000pg/ml was added for 30 minutes. Subsequently cells were washed, blocked with Fc blocker (BD bioscience FC Blocker-MIX cat#BD564220), and fixated with cytofix fixation buffer (BD bioscience Cat# 554655). The cells were permeabilized with 90% methanol, washed and labeled with anti- pSTAT5-PE (BD bioscience cat#562077). Treated MUTZ5 cells were analyzed for pSTAT5 activation on a CytoFLEX S flow cytometer (Beckman). Cells gated for singlets and pSTAT5 were marked as pSTAT5 positive.

[433] *Methods of determination of IgG IC50 inhibition of MUTZ5 pSTAT5 activation by hTSLP*

[434] To test functional blocking of pSTAT5 activation, IgG at a concentration range of 0.48pM to 500pM, was mixed with 14pM hTSLP (ACROBiosystems,cat# TSP-H52Hb) and incubated for 30 minutes, then added to the cells for another 60 minutes. Subsequently the cells were washed, fixated, labeled, and analyzed as described for the calibration of MUTZ5 cells.

[435] *Methods of Surface Plasmon Resonance (SPR) analysis*

[436] Measurements of IgG binding to human IL-13: The SPR analysis was done on Biacore 200 (GE) on CM5 chips cat:br10005-30 (GE), the chip was crosslinked with primary capture Ab

(Cat: br-1008-39 GE) to a target of 8000RU, after cross linking of the primary Ab, the tested antibodies were immobilized on the primary Ab to a target of 500RU. The hIL-13 (Peprotech) analyte was streamed in HEB-EP buffer at concentrations ranging from 800nM to 1.6nM in a series of two-fold dilutions, one concentration for each cycle. Subsequent to a cycle, the analyte and tested antibody were stripped from the chip and new tested Ab was loaded on the chip as described above. KD was determined at a steady state condition.

[437] *Measurements of binding to cynomolgus monkey IL-13 (cIL-13, Sino biological, USA) and human TSLP*

[438] The SPR analysis was done on ProteOn™ XPR36 (BioRad) on a GLC chips cat:176-5011 (BioRad). The chip was crosslinked with primary capture Ab (Cat: br-1008-39 GE) to a target of 5500RU. After cross-linking of the primary Ab tested, antibodies 33.003 and 33.004 were immobilized on the primary Ab to a target of 2000RU. The cyno IL-13 analyte was streamed in HEB-EP buffer at concentrations ranging from 200nM to 12.5nM in a series of two-fold dilutions. KD was determined at a steady state condition. For measurements of binding kinetics to hTSLP, the same conditions were used but with TSLP serving as analyte at concentrations ranging from 3.2nM to 0.2nM in a series of two-fold dilutions.

[439] *Method of Dynamic Scanning Fluorescence (DSF)* Dynamic Scanning Fluorescence was measured as reported by (Niedziela-Majka et al., 2015) with minor modifications. Briefly: 0.3mg/ml tested antibody in sodium acetate pH 5.5 buffer was mixed 1:1 with 20x sypro orange (Thermo Fisher, USA cat#S6650) in the same buffer. Changes in fluorescence were monitored on a Bio-Rad cfx96 light cycler with setting of 0.5°C/min from 25°C-100°C. T_m was determined as the temperature corresponding to the maximum value of the first derivative of the DSF melting curve. Where mentioned, antibodies were diluted to 0.5 mg/ml in PBS and analyzed using NanoDSF Prometheus NT.48 (Nanotemper, Germany) in a temperature elevation rate of 1°C/min.

[440] *Methods of Cell based assays*

[441] HEK-Blue IL-4/IL-13 Cells (Invivogen, France Catalog # hkb-il413) were used to determine IL-13 inhibition. HEK-Blue cells were cultured in growth medium comprising of DMEM, 4.5 g/l glucose, 10% (v/v) fetal bovine serum (FBS), 50 U/ml penicillin, 50 mg/ml streptomycin, 100 mg/ml Normocin, 2 mM L-glutamine, 10 µg/ml of blasticidin and 100 µg/ml of Zeocin. HEK-Blue IL-4/IL-13 cells are specifically designed to monitor the activation of the STAT6 pathway induced by IL-4 and IL-13. These cells were generated by stably introducing the human STAT6 gene into HEK293 cells to obtain a fully active STAT6 signaling pathway. The other genes of the pathway are naturally expressed in sufficient amounts. HEK-BlueIL-4/dual cells stably express the reporter gene, secreted embryonic alkaline phosphatase (SEAP), under the

control of the IFN β minimal promoter fused to four STAT6 binding sites. Activation of the STAT6 pathway in HEK-Blue IL-4/IL-13 cells induces the expression of the reporter gene. SEAP, which is secreted in the supernatant is easily detectable when using QUANTI-Blue, a medium that turns purple/blue in the presence of SEAP.

[442] *Methods of Calibration of HEK-Blue IL-4/IL-13 system*

[443] In order to determine the EC₅₀ value for rh-IL-13 on HEK-Blue IL-4/IL-13 cells, 50000 cells (5×10^4 / ELISA well) were incubated with rh-IL-13 antibody (Peprotech, Israel) at concentration of 0nM to 8.13nM for 24 hrs at 37°C, 5% CO₂ in a 96 well plate. At the end of the incubation, 20ul of the cell's supernatant was incubated with 180μl of QUANTI-Blue reagent (Invivogen, France) for an additional 2 hrs, and the reaction was analyzed by measuring the absorbance at 620-655nm using a plate reader spectrophotometer (Synergy Neo2, BioTek Instruments, Inc. USA). Data shown is the mean of triplicate experiments, and error bars represent standard deviation.

[444] *IC₅₀ of antibody inhibition of IL-13 downstream signaling:*

[445] 0.4 nM of rh-IL-13 was incubated with antibodies at a range of concentrations for 1 hr at room temperature. After the incubation, the mixture of rh-IL-13-antibody was added to a total volume of 200μl, 50,000 cells/well and incubated for 24 hrs at 37C 5% CO₂ in a 96 well plate. At the end of the incubation, 20μl of the cell's supernatant was incubated with 180μl of QUANTI-Blue reagent for additional 2 hrs, and the reaction was analyzed by measuring the absorbance at 620-655nm using a plate reader spectrophotometer. Data shown is the mean of triplicate experiments, and error bars represent standard deviation.

Example 2: Screening And Selection of Dual Binding Antibodies

[446] *Objective:* Screen engineered dual binding antibodies to identify those with highest binding for IL-13 and TSLP.

[447] *Results:* Following screening and selection of the libraries to bind both IL-13 and TSLP, 45 clones were selected, isolated, and sequenced resulting in 26 unique Heavy chain (VH) – Light chain (VL) pair variable regions, wherein the amino acid sequences of the Heavy chain and Light chain pairs are presented in **Table 1** (antibodies 1-26), the nucleotide sequences of the Heavy chain and Light chain scFv for antibodies 1-26, including the encoded linker sequences, are presented in **Table 2**, and the nucleic acid sequences of the Heavy chain and Light chain pairs for antibodies 1-26, are presented in **Table 3**. The Clone ID number for antibodies 1-26, is provided as “C#”- of each “Name” provided, for example at row 2, Clone C2 variable region pair comprises C2 VL sequence SEQ ID NO: 3 and C2 VH sequence SEQ ID NO: 4.

[448] Subsequently, an affinity maturation library was screened and additional dual binding antibody clones identified that showed YSD tight binding to both hIL-13 and hTSLP (Ab clone #s: 33.023, 33.025, 38.014, 38.015, 38.018, 38.019, 38.021, 38.026, 38.040). The amino acid sequences of the VH/VL regions of clones 33.023, 33.025, 38.014, 38.015, 38.018, 38.019, 38.021, 38.026, 38.040, are presented in Table 1 and the nucleotide sequences encoding the VH/VL regions are presented in Table 3. The CDR regions of VH/VL pairs from Ab clone #s: 33.023, 33.025, 38.014, 38.015, 38.018, 38.019, 38.021, 38.026, 38.040, are provided in Table 4 and Table 5 below. These clones were selected for IgG production.

Table 1: Engineered dual binding antibodies: Variable Light chain (VL) and Variable Heavy chain (VH) amino acid sequences (See also Figures 1A and 1B)

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
1	3	C2-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDG VVFGGGTKLTVL	4	C2-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTAEAFDIWGQG TMVTVSS
2	5	C27-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDG VVFGGGTKLTVL	6	C27-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
3	7	C19-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDG VVFGGGTKLTVL	8	C19-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWQLVAEAFDIWGQG TMVTVSS
4	9	C43-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SAIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDG	10	C43-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWQLVAEAFDIWGQG TMVTVSS

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
			VVFGGGTKLTVL			
5	11	C5-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDTSSDG VVFGGGTKLTVL	12	C5- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
6	13	C8-VL	SYVLTQPPSVSVAP GQTARITCGGNLLG SKLVHWYQQKPGQ APVLVYDDSDRP SAIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	14	C8- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDIWGQG TMVTVSS
7	15	C6-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	16	C6- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWMLVAEAFDLWGQ GTMVTVSS
8	17	C3-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	18	C3- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWEWVAEAFDLWGQ GTMVTVSS
9	19	C37-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	20	C37- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDMWGQ GTMVTVSS
10	21	C32-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP	22	C32- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPGKGLEWVAVIWDGNS KHYADSVKGRFTITRDNSKN

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
			SKIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL			TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
11	23	C38-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SDIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	24	C38-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPKGLEWVAVIWDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWEYVAEAFDLWGQ GTMVTVSS
12	25	C26-VL	SYVLTQPPSVSVAP GQTARITCGGNIIGS KLVHWYQQKPGQ APVLVYDDGDRP SGIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	26	C26-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPKGLEWVAVIWDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTAEAFDLWGQG TMVTVSS
13	27	C13-VL	SYVLTQPPSVSVAP GQTARITCGGNLLG SKLVHWYQQKPGQ APVLVYDDGDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDH VVFGGGTKLTVL	28	C13-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPKGLEWVAVIWDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
14	29	C23-VL	SYVLTQPPSVSVAP GQTARITCGGNLLG SKLVHWYQQKPGQ APVLVYDDSDRP SEIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDSSSDG VVFGGGTKLTVL	30	C23-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPKGLEWVAVIWDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTAEAFDIWGQG TMVTVSS
15	31	C39-VL	SYVLTQPPSVSVAP GQTARITCGGNLLG SKLVHWYQQKPGQ APVLVYDDSDRP SWIPERFSGSNSGN TATLTISRVEAGDE ADYQCQVWDSSD HVVFGGGTKLTVL	32	C39-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPKGLEWVAVIWDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTAEAFDLWGQG TMVTVSS
16	33	C17-	SYVLTQPPSVSVAP	34	C17-	QMQLVESGGGVVQPGRSLR

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
		VL	GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SAIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDTSSDH VVFGGGTKLTVL		VH	LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTSEAFDLWGQG TMVTVSS
17	35	C45-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SEIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDSSSDG VVFGGGTKLTVL	36	C45-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELTSEAFDLWGQG TMVTVSS
18	37	C7-VL	SYVLTQPPSVSVAP GQTARITCGGNIIGS KLVHWYQQKPGQ APVLVYDDSDRP SGIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDSGSD HVVFGGGTKLTVL	38	C7- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWLLVAEAFDLWGQG TMVTVSS
19	39	C31-VL	SYVLTQPPSVSVAP GQTARITCGGNIIGS KLVHWYQQKPGQ APVLVYDDSDRP SDIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDSGSD GVVFGGGTKLTVL	40	C31-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
20	41	C15-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDGDRP SWIPERFSGSNSGN TATLTISRVEAGDE ADYYCQVWDSGS DGVVFGGGTKLTVL	42	C15-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
21	43	C35-VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA	44	C35-VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWVLVSEAFDLWGQG

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
			DYYCQVWDSGSD GVVFGGGTKLTVL			TMVTVSS
22	45	C12 - VL	SYVLTQPPSVSVAP GQTARITCGGNIIGS KLVHWYQQKPGQ APVLVYDDSDRP SAIPERFSGSNSGNT ATLTISRVEAGDEA DYYCQVWDSSSDG VVFGGGTKLTVL	46	C12 - VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWELVAEAFDLWGQG TMVTVSS
23	47	C4- VL	SYVLTQPPSVSVAP GQTARITCGGNNIG SKLVHWYQQKPGQ APVLVYDDSDRP SGIPERFSGSNSGNT ATLTISRVEAGDEA DYYCQVWDSSSDG VVFGGGTKLTVL	48	C4- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVISYDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARSPQWEWVHEAFDMWGQ GTMVTVSS
24	49	C41- VL	SYVLTQPPSVSVAP GQTARITCGGNILG SKLVHWYQQKPGQ APVLVYDDSDRP SEIPERFSGSNSGNT ATLTISRVEAGDEA DYYCQVWDTSSDG VVFGGGTKLTVL	50	C41- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVISYDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARSPQWEWVHEAFDLWGQ GTMVTVSS
25	51	C40- VL	SYVLTQPPSVSVAP GQTARITCGGNNIG SKLVHWYQQKPGQ APVLVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYYCQVWDTSSDG VVFGGGTKLTVL	52	C40- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARSPQWEWVHEAFDLWGQ GTMVTVSS
26	53	C9- VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVYDDGDRP SWIPERFSGSNSGN TATLTISRVEAGDE ADYYCQVWDSSSD GVVFGGGTKLTVL	54	C9- VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRITYGMHWVR QAPKGLEWVAVIWDGNSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARSPQWEWVHEAFDLWGQ GTMVTVSS
	156	33.02 3VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ	155	33.023 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRITYGMHWVR QAPKGLEWVAVIWDGNSN

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
			APVLVVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDHSSD HVVFGGGTKLTVL			THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWYLSAEAFDLWGQG TMVTVSS
	160	33.02 5VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SLIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDSSSDG VVFSGGGTKLTVL	159	33.025 VH	QMQLVESGGGVVQPGRSLR LSCAASGFTFRTYGMHWVR QAPGKGLEWVADIWYDGSN KHYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWYLVAEFPDLWGQG TMVTVSS
	170	38.01 4 VL	SYVLTQPPSVSVAP GQTARITCGGNLIG AKLVHWYQQKPG QAPVLVVYDDSDR PSRIPERFSGSNSGN TATLTISRVEAGDE ADYVCQVWDHSS DHVVFGGGTKLTV L	169	38.014 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRTYGMHWVR QAPGKGLEWVAVIWYDGSN THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC VRAPQWYLSAEAFDLWGQG TMVTVSS
	172	38.01 8 VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDHSSD HVVFGGGTKLTVL	171	38.018 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRTYGMHWVR QAPGKGLEWVAVIWDDGSN THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWYLSAEAFDLWGQG TMVTVSS
	174	38.01 9 VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDYSSN HVVFGGGTKLTVL	173	38.019 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRTYGMHWVR QAPGKGLEWVAVIWYDGSN THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC VRAPQWYLSAEAFDLWGQG TMVTVSS
	176	38.02 1 VL	SYVLTQPPSVSVAP GQTARITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYVCQVWDHSSD HYVFGGGTKLTVL	175	38.021 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFDYGMHWVR QAPGKGLEWVAVIWYDGSN TVYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC ARAPQWYLSAEAFDLWGQG TMVTVSS

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
	178	38.025 VL	SYVLTQPPSVSVAP GETATITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SRIPERFSGSNSGNT ATLTISRVEAGDEA DYQCQVWDHSSD HVVFGGGTKLTVL	177	38.025 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRTYGMHWVR QAPGKGLEWVAVIWDGSA THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC VRAPQWYLSAEAFDLWGQG TMVTVSS
	180	38.026 VL	SYVLTQPPSVSVAP GQTATITCGGNLIG SKLVHWYQQKPGQ APVLVVYDDSDRP SRIPERFSGSNIGNT ATLTISRVEAGDEA DYQCQVWDHSSD HVVFGGGTKLTVL	179	38.026 VH	QMQLVESGGGVVQPGRSLR LSCAASGFAFRTYGMHWVR QAPGKGLEWVAVIWDGSA THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC VRAPQWYLSAEAFDLWGQG TMVTVSS
	182	38.040 VL	SYVLTQPPSVSVAP GQTARITCGGNLIG TKLVHWYQQKPG QAPVLVVYDDSDR PSRIPERFSGSNSGN TATLTISRVEAGDE ADYQCQVWDHNE DEVVFGGGTKLTVL	181	38.040 VH	QMQLVESGGGVVQPGRSLR LSCAASGFDFRTYGMHWVR QAPGKGLEWVAVIWDGSI THYADSVKGRFTITRDNSKN TLNLQMNSLRAEDTAVYYC VRAPQWYLTAEAFDLWGQG TMVTVSS

Table 2: Nucleotide Sequences Encoding Engineered dual Binding scFv: Variable Heavy chain (VH)-Linker-Variable Light chain (VL)

Ab #	SEQ ID NO:	Name	VH-Linker-VL Sequence
	109	Template VH-L-VL	CAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCAGTTTATTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTACACGAAGCAT TCGATATCTGGGGTCAGGGTACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC AATCTGGGCTCTAAGTCTGTTCAGTGGTATCAGCAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATCCGATC

			GTCCAAGCTGGATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGGGGAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTTCTT CCGACCATGTAGTCTTTGGCGGGGGCACCAAAGTACCGTT TTG
1	110	C2- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACGGCGGAAGCAT TCGATATTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCCTGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTAGC TCCGACGGTGTAGTCTTTGGCGGGGGCACCAAAGTACCGT TTTG
2	111	C27- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCGGAAGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCCTGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTAGC TCCGACGGTGTAGTCTTTGGCGGGGGCACCAAAGTACCGT TTTG
3	112	C19- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGCAGTTAGTAGCGGAAGCAT

			TCGATATTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCTGGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTACTGGTTGTGTACGATGATAGCGAT CGTCCAAGCCTTATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
4	113	C43- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGCAGTTAGTAGCCGAAGCATT CGATATATGGGGCCAGGGCACTATGGTGACCGTTAGCTCTG GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACC TGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACCA GGCCAGGCACCACTACTGGTTGTGTACGATGATAGCGATC GTCCAAGCGCTATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACACTAGCT CCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACCGTT TTG
5	114	C5- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTGGCGGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGTGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACC TGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACCA GGCCAGGCACCACTACTGGTTGTGTACGATGATAGCGATC GTCCAAGCCTTATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAGC TCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACCGT TTTG
6	115	C8- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT

			ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTGGTAGCCGAAGCAT TCGATATCTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGCTGGGCTCTAAGCTGGTTCCTGTTATCAGCAAAAACC AGGCCAGGCACCACTGTTGTTGTACGATGATAGCGAT CGTCCAAGCGCAATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTAGC TCCGACCACGTAGTCTTTGGCGGGGGCACCAAACCTGACCGT TTTG
7	116	C6- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGATGTTAGTAGCGGAAGCATT CGATCTATGGGGCCAGGGCACTATGGTGACCGTTAGCTCTG GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACC TGATCGGCTCTAAGCTGGTTCCTGTTATCAGCAAAAACCA GGGCAGGCACCACTGTTGTTGTACGATGATAGCGATC GTCCAAGCCGGATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAGCT CCGACCATGTAGTCTTTGGCGGGGGCACCAAACCTGACCGT TTG
8	117	C3- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATGGGTAGCCGAAGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCCTGTTATCAGCAAAAACC AGGCCAGGCACCACTGTTGTTGTACGATGATAGCGAT CGTCCAAGCCGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGTAAGCCG

			GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAGCTCCGACCACGTAGTCTTTGGCGGGGGCACCAAACCTGACCGTTTTG
9	118	C37-VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCCGAGGCAT TCGATATGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAAC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCCGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTAGC TCCGACCATGTAGTCTTTGGCGGGGGCACCAAACCTGACCGT TTTG
10	119	C32-VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCCGAAGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAAC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCAAAATTCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAG CTCCGACCATGTAGTCTTTGGCGGGGGCACCAAACCTGACCG TTTTG
11	120	C38-VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATACGTAGCGGAAGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC

			GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTACTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGATATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAG CTCCGACCATGTAGTCTTTGGCGGGGGCACCAAACCTGACCG TTTTG
12	121	C26- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACGGCCGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC ATCATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTACTGGTTGTGTACGATGATGGCGAT CGTCCAAGCGGTATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACTAG CTCCGACCACGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
13	122	C13- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTGGCGGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGCTGGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTACTGGTTGTGTACGATGATGGCGAT CGTCCAAGCCTGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACTAG CTCCGACCACGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
14	123	C23- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT

			GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACGGCCGAAGCAT TCGATATTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGCTGGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGAAATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
15	124	C39- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACGGCCGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGCTGGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCTGGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACTAG CTCCGACCATGTAGTCTTTGGCGGGGGCACCAAACCTGACCG TTTTG
16	125	C17- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACGTCGGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGCAATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAG CTCCGACACGTAGTCTTTGGCGGGGGCACCAAACCTGACC

			GTTTTG
17	126	C45- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAACCTCCGAAGCATT CGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCTG GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACC TGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACCA GGCCAGGCACCAGTACTGGTTGTGTACGATGATAGCGATC GTCCAAGCGAAATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTGAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAGCT CCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACCGTT TTG
18	127	C7- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGCTGTTAGTAGCGGAAGCATT CGATCTCTGGGGCCAGGGCACTATGGTGACCGTTAGCTCTG GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACA TCATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACCA GGCCAGGCACCAGTACTGGTTGTGTACGATGATAGCGATC GTCCAAGCGGTAATCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTGAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTGGCT CCGACCACGTAGTCTTTGGCGGGGGCACCAAACCTGACCGTT TTG
19	128	C31- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTGGTATGAT GGTAAGCAATAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCGGAAGCAT TCGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC

			ATCATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGATATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCGG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
20	129	C15- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTTCGAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCGGAAGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATGGCGAT CGTCCAAGCTGGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTGGC TCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACCGT TTTG
21	130	C35- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTTCGAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGTCTTAGTATCCGAAGCATT CGATCTTTGGGGCCAGGGCACTATGGTGACCGTTAGCTCTG GCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGCG GCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGTA GCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAACC TGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACCA GGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGATC GTCCAAGCCGTATTCCAGAGCGTTTCAGCGGCTCTAATTCC GGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCGG CGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCGGCT CCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACCGTT TTG
22	131	C12 – VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTTCGAGTAATCTGGTATGAT GGTAGCAATAAACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACCTCTTAACCTAC

			AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTGCACCACAGTGGGAATTAGTAGCGGAGGCAT TCGATCTGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC ATCATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCAGTACTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGCAATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACGACC GTTTTG
23	132	C4- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTCTTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTAGCCACAGTGGGAATGGGTACACGAAGCAT TCGATATGTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC AACATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCAGTACTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGGTATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCTAGC TCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACGACCCT TTTG
24	133	C41- VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTGCGCAGTAATCTCTTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACACTCTTAACCTTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTTGCCACAGTGGGAATGGGTACACGAAGCAT TCGATCTCTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC ATCCTGGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCAGTACTGGTTGTGTACGATGATAGCGAT CGTCCAAGCGAAATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTTGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACCAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACGACC GTTTTG

25	134	C40-VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTTCGCAGTAATCTGGTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTAGCCACAGTGGGAATGGGTACACGAAGCAT TCGATCTATGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC AACATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATAGCGAT CGTCCAAGCCGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACACTAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG
26	135	C9-VH-L-VL	CAAATGCAGCTGGTCGAGTCTGGCGGTGGGGTAGTGCAAC CAGGCCGTTCTCTGCGTCTTAGCTGCGCCGCATCTGGTTTT ACCTTTTCGTACCTACGGTATGCACTGGGTGCGTCAGGCACC AGGCAAAGGTCTGGAATGGGTTCGCAGTAATCTGGTATGAT GGTAGCAATAAAACACTATGCTGACTCAGTCAAAGGCCGTTT CACCATCACCCGTGATAACAGCAAGAACAACACTCTTAACCTAC AGATGAACTCTCTGCGTGCCGAAGACACCGCCGTTTACTAC TGTGCCCCGTTCGCCACAGTGGGAATGGGTACACGAAGCAT TCGATCTCTGGGGCCAGGGCACTATGGTGACCGTTAGCTCT GGCGGTGGTGGTAGCGGAGGCGGAGGATCAGGTGGAGGC GGCAGTTCTTACGTGCTGACTCAACCACCATCAGTGTCTGT AGCACCAGGCCAGACCGCACGTATTACCTGTGGCGGTAAC CTGATCGGCTCTAAGCTGGTTCAGTGGTATCAGCAAAAACC AGGCCAGGCACCACTAGTGGTTGTGTACGATGATGGCGAT CGTCCAAGCTGGATTCCAGAGCGTTTCAGCGGCTCTAATTC CGGCAACACCGCTACTCTGACTATTTCCCGTGTGGAAGCCG GCGATGAAGCCGACTACTATTGCCAGGTCTGGGACTCCAG CTCCGACGGTGTAGTCTTTGGCGGGGGCACCAAACCTGACC GTTTTG

Table 3: Nucleotide Sequences Encoding Engineered dual binding antibodies: Variable Light chain (VL) and Variable Heavy chain (VH) nucleic acid sequences

Ab #	SEQ ID NO:	Name	VL Sequence	SEQ ID NO:	Name	VH Sequence
	56	Template-VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACAATCTGGG	55	Template-VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT

			CTCTAAGTCTGTTCACT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TTCCGATCGTCCAAGCT GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGGGGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTTCTTCCGA CCATGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG			TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGC GTGCCGAAGACACC GCAGTTTATTACTGTGCC CGTGACCACAGTGGGA ATTAGTACACGAAGCAT TCGATATCTGGGGTCAG GGTACTATGGTGACCGT TAGCTCT
1	58	C2- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC TGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTAGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG	57	C2- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGC GTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACCACAGTGGGA ATTAACGGCGGAAGCAT TCGATATTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
2	60	C27- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC TGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA	59	C27- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT GGTATGATGGTAGCAAT AAACACTAT GCTGACTCAGTCAAAGG CCGTTTCACCATCACCC

			CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTAGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG			GTGATAACAGCAAGAAC ACTCTTAACCTACAGAT GAACTCTCTGCGTGCCG AAGACACCGCCGTTTAC TACTGTGCCCCGTGCACC ACAGTGGAATTAGTAG CGGAAGCATTTCGATCTG TGGGGCCAGGGCACTAT GGTGACCGTTAGCTCT
3	62	C19- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC TTATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACACCAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG	61	C19- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGCA GTTAGTAGCGGAAGCAT TCGATATTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
4	64	C43- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GCTATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACTAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG	63	C43- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGCA GTTAGTAGCCGAAGCAT TCGATATATGGGGCCAG GGCACTATGGTGACCGT

						TAGCTCT
5	66	C5- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC TTATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACACCAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	65	C5- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTGGCGGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
6	68	C8- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGCTGGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GCAATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCTAGCTCCG ACCACGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	67	C8- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTGGTAGCCGAAGCAT TCGATATCTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
7	70	C6- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG	69	C6- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT

			CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGGCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCCAGCTCCGA CCATGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG			TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCTGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCCTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGAT GTTAGTAGCGGAAGCAT TCGATCTATGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
8	72	C3- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCCAGCTCCGA CCACGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG	71	C3- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCTGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCCTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGGA ATGGGTAGCCGAAGCAT TCGATCTGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
9	74	C37- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA	73	C37- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCTGTGATAAC

			CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTAGCTCCGA CCATGTAGTCTTTGGCG GGGGCACCAAACTGAC CGTTTTG			AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTAGCCGAGGCAT TCGATATGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
10	76	C32- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC AAAATTCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCCAGCTCCG ACCATGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	75	C32- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTAGCCGAAGCAT TCGATCTGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
11	78	C38- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GATATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACCAGCTCC GACCATGTAGTCTTTGG CGGGGGCACCAAACTG ACCGTTTTG	77	C38- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATACGTAGCGGAAGCAT TCGATCTGTGGGGCCAG GGCACTATGGTGACCGT

						TAGCTCT
12	80	C26- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACATCATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TGGCGATCGTCCAAGC GGTATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACTAGCTCCG ACCACGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	79	C26- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAACGGCCGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
13	82	C13- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGCTGGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TGGCGATCGTCCAAGCC TGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACACTAGCTCCG ACCACGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	81	C13- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTGGCGGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
14	84	C23- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGCTGGG	83	C23- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT

			CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GAAATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACCAGCTCC GACGGTGTAGTCTTTGG CGGGGGCACCAAACCTG ACCGTTTTG			TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACCACAGTGGGA ATTAACGGCGGAAGCAT TCGATATTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
15	86	C39- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGCTGGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCT GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACACTAGCTCCG ACCATGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG	85	C39- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACCACAGTGGGA ATTAACGGCGGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
16	88	C17- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCAC GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GCAATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG	87	C17- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC

			ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACCAGCTCC GACCACGTAGTCTTTGG CGGGGGCACAAACTG ACCGTTTTG			AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAACGTCCGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
17	90	C45- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GAAATCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCCAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	89	C45- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAACCTCCGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
18	92	C7- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACATCATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GGTATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCTGGCTCCG ACCACGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	91	C7- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGCT GTTAGTAGCGGAAGCAT TCGATCTCTGGGGCCAG GGCACTATGGTGACCGT

						TAGCTCT
19	94	C31- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACATCATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GATATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCCGGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	93	C31- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTAGCGGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
20	96	C15- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TGGCGATCGTCCAAGCT GGATTCCAGAGCGTTT AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTGGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACTGAC CGTTTTG	95	C15- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGGA ATTAGTAGCGGAAGCAT TCGATCTGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
21	98	C35- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCCT	97	C35- VH	CAAATGCAGCTGGTCGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC

			GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GTATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCCGGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG			ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCCTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGGT CTTAGTATCCGAAGCAT TCGATCTTTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
22	100	C12 - VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACATCATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GCAATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCCAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG	99	C12 - VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCCTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACACACAGTGGGA ATTAGTAGCGGAGGCAT TCGATCTGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
23	102	C4- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACAACATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GGTATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA	101	C4- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGCAGTAATCT CTTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA

			AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACTCTAGCTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG			CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTAGCCCACAGTGGA ATGGGTACACGAAGCAT TCGATATGTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
24	104	C41- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACATCCTGGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGC GAAATTCCAGAGCGTTT CAGCGGCTCTAATTCCG GCAACACCGCTACTCTG ACTATTTCCCGTGTTGA AGCCGGCGATGAAGCC GACTACTATTGCCAGGT CTGGGACACCAGCTCC GACGGTGTAGTCTTTGG CGGGGGCACCAAACCTG ACCGTTTTG	103	C41- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT CTTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTTCGCCACAGTGGA ATGGGTACACGAAGCAT TCGATCTCTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
25	106	C40- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACAACATCGG CTCTAAGCTGGTTCCT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GGATTCCAGAGCGTTT AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACACTAGTCCG ACGGTGTAGTCTTTGGC GGGGGCACCAAACCTGA CCGTTTTG	105	C40- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTCCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCGTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTAGCCCACAGTGGA ATGGGTACACGAAGCAT TCGATCTATGGGGCCAG GGCACTATGGTGACCGT TAGCTCT

26	108	C9- VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCACT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TGGCGATCGTCCAAGCT GGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCCAGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACTGAC CGTTTTG	107	C9- VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGC GTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTTTCGCCACAGTGGGA ATGGGTACACGAAGCAT TCGATCTCTGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
	162	33.023 VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCACT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC GTATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACCACAGCTCCG ACCATGTAGTCTTTGGC GGGGGCACCAAACTGA CCGTTTTG	161	33.023 VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTGCGTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA ATGGGTTCGAGTAATCT GGTATGATGGTAGCAAT ACCCACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGC GTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGACCACAGTGGTA CTTAAGCGCGGAAGCAT TCGATCTATGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
	164	33.025 VL	TCTTACGTGCTGACTCA ACCACCATCAGTGTCTG TAGCACCAGGCCAGAC CGCACGTATTACCTGTG GCGGTAACCTGATCGG CTCTAAGCTGGTTCACT GGTATCAGCAAAAACC AGGCCAGGCACCAGTA	163	33.025 VH	CAAATGCAGCTGGTCTGA GTCTGGCGGTGGGGTAG TGCAACCAGGCCGTTCT CTGCGTCTTAGCTGCGC CGCATCTGGTTTTACCTT TCGTACCTACGGTATGC ACTGGGTGCGTCAGGCA CCAGGCAAAGGTCTGGA

			CTGGTTGTGTACGATGA TAGCGATCGTCCAAGCC TGATTCCAGAGCGTTTC AGCGGCTCTAATTCCGG CAACACCGCTACTCTGA CTATTTCCCGTGTTGAA GCCGGCGATGAAGCCG ACTACTATTGCCAGGTC TGGGACTCTAGCTCCGA CGGTGTAGTCTTTGGCG GGGGCACCAAACCTGAC CGTTTTG			ATGGGTTCGCAGACATCT GGTATGATGGTAGCAAT AAACACTATGCTGACTC AGTCAAAGGCCGTTTCA CCATCACCCGTGATAAC AGCAAGAACACTCTTAA CTTACAGATGAACTCTC TGCCTGCCGAAGACACC GCCGTTTACTACTGTGCC CGTGCACCACAGTGGTA CTTAGTAGCGGAACCGT TCGATCTATGGGGCCAG GGCACTATGGTGACCGT TAGCTCT
	184	38.014 VL	AGTTACGTGCTGACACA ACCTCCAAGTGTTAGTG TCGCACCAGGACAAAC AGCACGTATTACATGTG GAGGAAATCTTATCGGT GCCAAGCTGGTGCAC GGTACCAGCAGAAACC TGGTCAGGCCCCAGTAC TGGTTGTGTATGATGAC AGCGACCGTCCAAGCC GTATCCCAGAACGTTTT TCTGGGAGCAACTCAG GTAATACAGCCACTCTG ACCATTTACGTGTTGA GGCAGGAGATGAGGCC GATTATTATTGCCAAGT ATGGGACCACAGCTCT GACCATGTTGTTTTTGG CGGAGGGACTAAGCTG ACCGTGCTT	183	38.014 VH	CAGATGCAGTTGGTGGA GTCCGGAGGTGGAGTGG TGCAACCAGGGCGTTCC TTGCGTTTGTCTTGTGCT GCTTCCGGATTGCGCTTT CGTACATATGGCATGCA TTGGGTGCGTCAAGCAC CTGGTAAGGGCCTGGAG TGGGTTGCCGTTATTTGG TACGACGGCTCCAACAC CCACTACGCAGATAGCG TGAAAGGACGTTTCACT ATTACCCGTGATAACTC CAAGAATACCCTTAACC TGCAGATGAATAGCTTG CGTGCTGAGGACACAGC AGTATATTACTGCGTCC GTGCACCACAATGGTAC CTGAGCGCCGAGGCCTT TGATCTGTGGGGGCAGG GCACAATGGTGACCGTT TCCTCA
	186	38.018 VL	TCCTATGTGCTGACACA GCCACCTAGCGTGAGC GTCGCCCCAGGTCAGA CCGCTCGTATCACTTGT GGCGGGAACCTTATCG GCAGCAAGCTGGTGCA CTGGTACCAGCAGAAG CCTGGCCAAGCACCTGT GCTGGTCGTTTATGACG ACTCTGACCGTCCATCC CGTATCCCAGAACGTTT CTCTGGCTCTAACTCTG GGAATACCGCTACCCCTG ACAATCTCACGTGTTGA AGCTGGCGACGAGGCA GATTATTATTGCCAAGT	185	38.018 VH	CAGATGCAACTGGTGGA GTCAGGAGGCGGCGTGG TGCAGCCAGGACGTTCT CTGCGTCTGTCTTGCGCA GCTTCCGGGTTGCGCTTT CGTACCTATGGGATGCA TTGGGTGCGTCAGGCTC CAGGTAAGGGACTGGAG TGGGTGCGTGTTATTTGG GACGACGGAAGTAACAC TCATTACGCCGACAGCG TGAAGGGCCGTTTCACA ATTACCCGTGACAATTC CAAGAATACCTTGAACC TGCAGATGAACTCTCTT CGTGCTGAAGATACCGC

			CTGGGATCACTCCAGCG ATCACGTCGTGTTCCGA GGCGGAACAAAATTGA CTGTCCTG			CGTGTACTATTGCGCCC GTGCTCCACAGTGGTAT CTGTCAGCAGAGGCCTT CGACCTGTGGGGACAGG GAACAATGGTGACCGTA TCTTCA
	188	38.019 VL	TCTTACGTGTTGACACA ACCACCAAGTGTTAGTG TCGCACCTGGCCAAACC GCTCGTATCACCTGTGG TGGGAATCTTATTGGCT CTAAGCTGGTGCCTGG TATCAGCAGAAACCAG GCCAGGCTCCAGTACTG GTGGTGTACGACGACTC TGACCGTCCAAGCCGTA TCCCAGAGCGTTTCAGT GGCTCTAACTCCGGGA ACACAGCAACTCTTACA ATTTACGTGTGGAGGC CGGTGATGAAGCCGAC TACTATTGCCAGGTTTG GGACTACAGTAGTAAT CACGTGGTTTTCGGTGG TGGTACCAAGCTGACTG TGTTG	187	38.019 VH	CAAATGCAGCTGGTGGA ATCCGGGGGTGGGGTCG TCCAGCCTGGCCGTAGT CTGCGTCTTTCTGTGCC GCATCAGGCTTTGCTTTC CGTACCTACGGGATGCA CTGGGTGCGTCAGGCCC CAGGAAAGGGACTTGAA TGGGTGGCTGTCATCTG GTACGATGGTTCCAACA CACACTATGCCGATTCA GTGAAAGGGCGTTTCAC CATTACTCGTGACAATA GTAAGAATACTCTGAAT CTGCAAATGAATTCCT GCGTGCTGAGGACACCG CTGTTTATTACTGTGTGC GTGCTCCTCAGTGGTAC CTGAGTGCCGAAGCTTT CGATTTGTGGGGACAGG GCACAATGGTGACAGTC AGTTCT
	190	38.021 VL	AGCTATGTTCTGACTCA ACCACCTAGTGTGAGTG TGGCCCCTGGTCAGACT GCACGTATTACCTGTGG CGGAAACCTTATCGGC AGTAAGCTGGTTCATTG GTATCAGCAGAAGCCA GGACAGGCACCAAGTGC TGGTCGTTTACGACGAT AGTGACCGTCCATCACG TATCCCAGAGCGTTTTA GCGGGTCCAATTCCGG AAATACAGCAACCTTG ACCATTAGCCGTGTGGA AGCCGGCGATGAAGCT GATTATTACTGCCAGGT ATGGGACCATTCTCCG ACCACTACGTTTTTGGT GGCGGAACCTAAGCTGA CAGTCTTG	189	38.021 VH	CAGATGCAGCTGGTTGA ATCTGGCGGCGGTGTGG TCCAGCCTGGTCGTAGC CTGCGTCTGTCTGTGCT GCAAGCGGATTTGCCTT TGACACCTATGGGATGC ACTGGGTACGTCAGGCC CCAGGAAAGGGCCTGGA ATGGGTGGCAGTTATCT GGTATGATGGTTCTAAT ACCGTGTATGCCGACTC CGTTAAAGGCCGTTTCA CTATCACCCGTGATAAT AGTAAAAACACACTGAA CCTGCAGATGAATAGCT TGCCTGCTGAGGACACC GCAGTGTAATACTGTGC CCGTGCTCCTCAGTGGT ATCTGTGAGCAGAGGCC TTCGATCTGTGGGGCCA AGGGACAATGGTGACCG TGTCTTCC
	192	38.025	TCTTACGTGCTTACTCA	191	38.025	CAGATGCAGCTTGTGGA

		VL	GCCTCCTAGCGTCTCAG TGGCCCCAGGCGAGAC AGCAACCATTACATGC GGGGGTAATTTGATCG GTAGCAAGCTGGTGCA TTGGTATCAGCAGAAG CCTGGCCAGGCCCCAGT GCTGGTTGTATATGACG ATAGTGATCGTCCAAGT CGTATCCCTGAGCGTTT TAGCGGATCTAACTCCG GCAACACAGCCACATT GACAATCAGCCGTGTG GAGGCAGGCGATGAGG CCGACTACTACTGCCAA GTTTGGGACCACTCCTC TGACCACGTGGTATTTG GCGGAGGAACAAAGCT TACAGTTTTG		VH	GAGCGGCGGAGGCGTG TGCAACCAGGCCGTTCA TTGCGTCTGTCTGCGCC GCCAGCGGCTTTGCTTTT CGTACATACGGCATGCA CTGGGTGCGTCAGGCC CTGGCAAGGGGCTGGAA TGGGTGCGCGTGATTTG GTATGACGGTAGTAACA CCCATTATGCTGATTCCG TCAAGGGACGTTTCACT ATCACCCGTGACAATAG CAAAAATACACTGAATC TGCAATGAATTCATTG CGTGCCGAAGACACCGC CGTATATTACTGTGTCCG TGCCCCACAGTGGTACC TGAGCGCTGAGGCCTTC GATCTGTGGGTCAGGG GACTATGGTGACCGTAT CATCC
	194	38.026 VL	TCTTATGTTTTGACCCA ACCTCCATCCGTTAGCG TGGCTCCAGGTCAAAC AGCTACCATCACATGTG GCGGTAACCTTATTGGC TCAAAGCTGGTTCATTG GTATCAACAGAAACCA GGCCAAGCCCCAGTGC TGGTGGTGTATGACGAC AGTGACCGTCCTTCTCG TATTCCTGAGCGTTTTT CCGGCTCTAATATTGGC AACACTGCCACCCTGAC CATTTCTCGTGTGGAAG CAGGAGATGAGGCAGA CTATTATTGTCAGGTTT GGGATCACTCCAGCGA TCATGTGGTATTGCGAG GTGGGACAAAACCTTAC TGTTCTT	193	38.026 VH	CAGATGCAGCTGGTGGA GAGTGGAGGTGGTGTGG TGCAACCTGGGCGTAGC CTGCGTTTTGAGCTGCGC TGCCTCTGGATTTGCCTT CCGTACCTATGGCATGC ACTGGGTGCGTCAGGCT CCAGGAAAGGGGTTGGA ATGGGTGGCTGTGATTT GGTACGACGGGAGCGCC ACACATTACGCAGACAG CGTTAAGGGCCGTTTCA CAATTACCCGTGACAAT AGCAAAAATACATTGAA CCTGCAGATGAATTCCC TGCGTGACAGAGGATACT GCAGTGTACTATTGCGT CCGTGCCCCACAGTGGT ATCTGTGACCCGAAGCC TTCGATCTGTGGGGGCA GGGTACTATGGTCACCG TAAGTTCC
	196	38.040 VL	AGCTACGTGCTTACCCA GCCACCATCAGTCAGTG TGGCTCCAGGCCAAACT GCCCCGTATCACCTGCGG CGGCAATTTGATTGGCA CCAAGCTTGTGCACTGG TACCAACAGAAGCCAG GGCAGGCCCTGTGCTG GTTGTCTACGACGATAG	195	38.040 VH	CAGATGCAGCTGGTGGA AAGTGGTGGGGGAGTCG TGCAACCAGGACGTTCC TTGCGTCTGTGATGCGCT GCTTCAGGTTTCGACTTT CGTACCTACGGCATGCA TTGGGTGCGTCAGGCTC CAGGTAAAGGACTTGAG TGGGTGCGCAGTGATCTG

			TGATCGTCCTTCCCGTA TTCCTGAACGTTTCTCT GGAAGCAATTCCGGAA ACACAGCCACACTTACC ATTTCTCGTGTTGAGGC TGGGGATGAAGCCGAC TACTATTGCCAGGTTTG GGACCACAATGAAGAC GAAGTTGTTTTTGGAGG AGGAACTAAGCTGACA GTTCTG			GTACGACGGATCAATTA CTCACTACGCCGATAGC GTGAAAGGCCGTTTCAC CATCACCCGTGACAACT CCAAGAACACCCTGAAC TTGCAGATGAACAGTCT GCGTGCAGAAGACACTG CAGTATATTATTGTGTCC GTGCCCCACAGTGGTAC TTGACCGCCGAGGCTTT TGATCTGTGGGGACAGG GCACAATGGTGACCGTA TCTAGC
--	--	--	--	--	--	---

Table 4: Amino Acid Sequences of Heavy-chain CDR Regions for Antibody Clone #s. 33.023, 33.025, 38.014, 38.015, 38.018, 38.019, 38.021, 38.026, 38.040

CLONE	HCDR1	SEQ ID	HCDR2	SEQ ID	HCDR3	SEQ ID
BDG33_023_VH	GFAFRTYG	149	IWYDGSNT	150	ARAPQWYLSAEAFDL	151
BDG33_025_VH	GFTFRTYG	165	IWYDGSNK	166	ARAPQWYLVAEPFDL	167
BDG38_014_VH	GFAFRTYG	149	IWYDGSNT	150	VRAPQWYLSAEAFDL	201
BDG38_018_VH	GFAFRTYG	149	IWDDGSNT	208	ARAPQWYLSAEAFDL	151
BDG38_019_VH	GFAFRTYG	149	IWYDGSNT	150	VRAPQWYLSAEAFDL	201
BDG38_021_VH	GFAFDTYG	197	IWYDGSNT	150	ARAPQWYLSAEAFDL	151
BDG38_025_VH	GFAFRTYG	149	IWYDGSNT	150	VRAPQWYLSAEAFDL	201
BDG38_026_VH	GFAFRTYG	149	IWYDGSAT	199	VRAPQWYLSAEAFDL	201
BDG38_040_VH	GFDFTYG	198	IWYDGSIT	200	VRAPQWYLTAEAFDL	202

Table 5: Amino Acid Sequences of Light-chain CDR Regions for Antibody Clone #s. 33.023, 33.025, 38.014, 38.015, 38.018, 38.019, 38.021, 38.026, 38.040

CLONE	LCDR1	SEQ ID	LCDR2	SEQ ID	LCDR3	SEQ ID
BDG33_023_VL	LIGSKL	152	DDS	153	QVWDHSSDHVV	154
BDG33_025_VL	LIGSKL	152	DDS	153	QVWDSSSDGVV	168
BDG38_014_VL	LIGAKL	203	DDS	153	QVWDHSSDHVV	154
BDG38_018_VL	LIGSKL	152	DDS	153	QVWDHSSDHVV	154
BDG38_019_VL	LIGSKL	152	DDS	153	QVWDYSSNHVV	205
BDG38_021_VL	LIGSKL	152	DDS	153	QVWDHSSDHVV	206
BDG38_025_VL	LIGSKL	152	DDS	153	QVWDHSSDHVV	154
BDG38_026_VL	LIGSKL	152	DDS	153	QVWDHSSDHVV	154
BDG38_040_VL	LIGTKL	204	DDS	153	QVWDHNEDEVV	207

[449] The clones were tested for their binding to 10nM rh-IL-13 in yeast scFv format and were compared to a positive rh-IL-13 binder displayed on yeast as well. The affinity was normalized based on the mean fluorescence values of the positive control (normalized MFI).

Relative affinity of the isolated clones for rh-IL-13 was between 3% and 30% of the affinity displayed by the positive control (Figure 2A). C2, C6, C9, and C40 clones that exhibited above 20% of the relative affinity for rh-IL-13 and were shown to bind 10nM TSLP in YSD (Figure 2B) were chosen to be expressed as human IgG1.

Example 3: Ab Production And Biochemical Characterization

[450] *Objective:* To reformat the selected clones to a human IgG1 format and analyze the IgG1 antibodies for dual IL-13 and TSLP binding.

[451] *Results:* Subsequent to characterization in the yeast surface display format described in Example 2, the selected clones C2, C6, and C9, were reformatted to human IgG1 by subcloning the variable domain into two separate expression vectors, pSF-CMV-HuIgG1_HC and pSF-CMV-HuLambda_LC, as described in Example 1 (*Methods*).

[452] Clones BDG 33.003, BDG 33.004, and BDG 33.005 (Clones C2, C6, and C9, respectively) were expressed and purified as described in Example 1 (*Methods*), following protein A purification, the IgGs were >95% pure as evident from an SDS PAGE analysis (data not shown). Size exclusion chromatography of BDG 33.003 (clone C2), BDG 33.004 (clone C6), and BDG 33.005 (clone C9) on Superdex®200 10/300, showed two main peaks the first with a retention time of 9.2ml (0.36CV), typical of large aggregate and a second peak with retention of approximately 13.2ml (0.528CV), typical of an ordinary human IgG1 (hIgG1). The integrated area under the curve of these two peaks showed a ratio of 22% and 78% respectively (**Figures 3A-3D**).

[453] Both BDG33.0023 and BDG33.025 migrated on Superdex®200 10/300 with small leading peak corresponds to (0.36CV) that is typical of a large diameter aggregate, and a second peak with retention of approximately 13.8ml (0.55CV) that is typical of an ordinary human IgG. Area Under the Curve (AUC) peak ratio is 97.3% folded/2.8% misfolded and 98.5% folded/1.5% misfolded for BDG33.023 and BDG33.025 respectively (**Figures 3E-3F**).

[454] To test the thermostability of clones BDG 33.003, BDG 30.004, and BDG 30.005, the clones thermal melting was monitored by differential scanning fluorescence (DSF) as described in Example 1. As was evident from the first derivative of the fluorescence thermal shift graph, BDG 33.004 had one distinct transition at point at 62°C which could possibly correspond to both Tm1 and Tm2. (Data not shown). BDG 33.003 and BDG33.005 each had two transition points, a major one at 62°C (BDG 33.003) and 64.5°C (BDG 33.005), respectively, and a minor one at 73°C (BDG 33.003) and 74.5°C (BDG 33.005), respectively.

[455] BDG 33.023 and BDG33.025 were tested using NanoDSF Prometheus NT.48

(NanoTemper Technologies, Germany). BDG33.0023 had a T-onset of 64.2°C and first transition point at 67.7°C, BDG33.0025 had a T-onset of 56.4°C and first transition point at 60.9°C and second transition point at 67.4°C (**Figures 4A-4B**)

[456] The affinities of the IgGs to human TSLP, human IL-13, and cynomolgus monkey IL-13 were tested. Binding kinetics of hIL-13 to BDG33.023 and BDG33.025 was tested on BiacoreT200 as described herein, (**Figure 5G-H**). BDG33.003 and BDG33.004 clones were tested by SPR analysis on BiacoreT200 and ProteOn™ XPR36, respectively, using the GE capture antibody kit. While it was not possible to obtain kinetics parameters for binding the human IL-13, steady-state binding measurement resulted in an apparent K_D of 21.6 nM and 57.4 nM for BDG33.003 and BDG33.004, respectively (**Figures 5A-5B**). For all other measurements of BDG33.003, BDG 33.004, BDG33.023, and BDG33.025, kinetics of binding to hTSLP and hIL-13 are presented in **Tables 6A and 6B**, and **Figure 5E and 5H**.

[457] The antibodies were also tested for binding of recombinant cynomolgus monkey IL-13 (rc-IL-13), which shares 85% identity and 88% homology with the human IL-13, as can be seen in **Figures 5C-5D**, injection of cIL-13 as analyte, resulted in strong, dose dependent response indicating that BDG33.003 and BDG33.004 bind to rc-IL-13. Although kinetics for rc-IL-13 could not be obtained, the binding and dissociation slopes had similar profile for rh-IL-13 and rc-IL-13, suggesting that the binding mode for recombinant h-IL-13 and rc-IL-13 is likely similar (**Figures 5A-5B** for human IL-13 and **Figures 5C-5D** for cyno IL-13).

[458] To further test the IgGs affinity to human TSLP, cynomolgus monkey TSLP and cynomolgus monkey IL-13, an ELISA EC50 experiment was done as described herein. Briefly, wells were coated with the respective ligand, then incubated with clones BDG33.003, BDG33.004, BDG33.023, or BDG33.025 at a concentration range of 1nM to 1000nM, washed and developed using HRP conjugated secondary antibody. EC50 values are presented in **Table 7**. Since the IgGs mentioned in the above sections are symmetrical IgGs, and since these same IgGs bind both hIL-13 and hTSLP this data demonstrates that BGD33.003, BGD33.004, BGD33.023, and BGD33.025 antibodies bind the two unrelated targets -TSLP and/or IL-13 from the same standard IgG CDRs, as appose to bi-specific antibody where the Light chain variable domain binds one target and heavy chain binds the other target (**Figures 6A-6E**).

[459] To test whether the IgGs are binding IL-13 and TSLP with overlapping paratopes, a competition assay was done as described herein. Briefly, BDG 33.023 or BDG33.025 were incubated with hIL-13 or hTSLP in concentration range of 0.78nM to 200nM and then tested for binding to either IL-13 or TSLP that were pre-coated on an ELISA plate. As can be seen in **Figures 7A-7D**, IL-13 blocks BDG33.023 from binding to IL-13 coated wells, and TSLP blocks

BDG33.023 and BDG33.025 from binding to TSLP coated wells (**Figures 7A-7B**). In addition, IL-13 blocked binding of BDG33.023 from binding to TSLP coated wells and reciprocally TSLP blocked binding of BDG33.023 from binding to IL-13 coated wells (**Figures 7C-7D**). This experiment indicates that each of IL-13 and TSLP share at least partially BDG33.023 binding paratope.

[460] IL-13 and TSLP are sequence and structurally unrelated. To test whether binding to these ligands by BDG33.023 and BDG33.025 is specific and not a result of non-specific binding or “stickiness”, BDG33.0023 and BDG33.025 binding to IL-4, IL-2, IL-17, BSA IL-13, and TSLP was tested by ELISA as described herein. As can be seen in **Figure 8**, while BDG33.025 shows strong binding to IL-13, TSLP and IL-4, but not to IL-2 and IL-17. BDG33.023 bind strongly to TSLP and IL-13 but shows no binding to the other ligands, indicating that its binding to IL-13 and TSLP is specific.

[461] To test if BDG33.023 binds TSLP at a functional epitope, the ability of BDG33.023 to cross-block TSLP from binding to a TSLP receptor was tested. Briefly TSLP-R was coated on ELISA plate wells, and its ability to bind hTSLP in the presence of 0nM to 500nM BDG33.023 was tested. As can be seen in **Figure 9**, BDG 33.023 can cross block TSLP binding to TSLP-R with an IC₅₀ of 0.41nM indicating that BDG33.023 binds tightly TSLP at a biologically functional site.

Table 6A: KD values of antibody clones for human IL-13 and TSLP

	Human IL-13		
Antibody	K _D (Steady state) (M)	SE(K _D) (M)	
BDG33.003	2.16E-08	2.5E-09	
BDG33.004	5.74E-08	4.1E-09	
	Human TSLP		
	ka (1/Ms)	kd (1/s)	KD (M)
BDG33.003	2.87E+05	1.29E-3	4.5E-9
BDG33.004	3.44E+05	1.14E-3	3.33E-9

Table 6B: KD values of antibody clones for human IL-13 and TSLP

	Human IL-13		
Antibody	ka (1/Ms)	kd (1/s)	KD (M)
BDG33.023	4.31E+06	0.01573	3.65E-09
BDG33.025	4.07E+6	0.0036	9.03E-10

Table 7. EC₅₀ values for human and cyno TSLP, and cyno IL-13

<i>IgG</i>	<i>EC50 value for ligand</i>		
	<i>hTSLP</i>	<i>cTSLP</i>	<i>cIL-13</i>
<i>BDG33.023</i>	<i>2.5nM</i>	<i>8.2nM</i>	<i>7.1nM</i>
<i>BDG33.025</i>	<i>12.8nM</i>	<i>148nM</i>	<i>1.5nM</i>

Example 4: Cell Based Assays For The Inhibitor Antibody BDG33.003 (Clone C2)

[462] *Objective:* Analyze the IgG1 antibodies for the ability to inhibit IL-13 activity.

[463] *Results:* To evaluate the capability of the antibody to inhibit rh-IL-13, the HEK-Blue IL-4/IL-13 system was used. The system uses HEK293 cells, which were stably transfected with human STAT6 gene and the reporter gene secreted embryonic alkaline phosphatase (SEAP) under the control of the IFN β minimal promoter fused to four STAT6 binding sites (Example 1 (Methods), and **Figure 10**). The system was initially tested by introducing rh-IL-13 to the cells and following the cell signaling cascade resulting in IL-13R (IL-13 receptor) activation by rh-IL-13. The results showed that IL-13 had an EC50 of about 0.12nM to the cells (**Figure 11**). Next, the engineered BDG33.003 (clone C2), BDG33.023 and BDG33.025 antibodies were tested to determine if they could inhibit IL-13 mediated activation of the cell's signaling cascade. The antibody was incubated with 0.4nM rh-IL-13, which was shown to activate IL-13R to approximately 70% of the saturation level, and the IgG/IL-13mixture was introduced to the cells for 24 hrs. The results obtained showed that the antibodies were able to inhibit IL-13 from binding to the IL-13R/IL-4R receptor complex, thus interfering with the signaling cascade. While for BDG33.003 the exact IC50 value was hard to determine, it is clear that BDG33.003 is inhibiting IL-13 signaling cascade. In addition, BDG33.023 and BDG33.025 inhibited IL-13 signaling cascade with an IC50 of 1.3nM and 25nM respectively, indicating that the IgGs are functionally blocking IL-13 in a biologically relevant setting. (**Figures 11B-11D**).

[464] To evaluate the capability of the antibodies to inhibit human TSLP in cells, MUTZ5 cells were used to test pSTAT5 TSLP dependent activation in a similar manner reported by Francis OL et al (Hematopoiesis 2016). TSLP induced phospho-STAT5 (pSTAT5) cellular activation cascade requires IL-7 receptor and TSLP-R receptor to function, as can be seen in figure 12A both are expressed on the MUTZ5 cell line surface indicating that these cells have the necessary receptors for this assay. To establish the cellular response to TSLP cells were incubated with TSLP at a concentration range of 0.0001 to 1000pg/ml and their pSTAT5 activation levels were determined by flow cytometry. As can be seen in figure 12B treatment of MUTZ 5 cells with TSLP activates pSTAT5 in a dose dependent manner, indicating that these cells respond to TSLP via the pSTAT5 pathway. To test BDG33.023 inhibition of TSLP dependent STAT5 activation,

TSLP at a concentration of 14pM was mixed with 0.48pM to 500pM BDG33.023 and incubated with MUTZ cells, as can be seen in **Figure 12C**. BDG33.023 inhibit TSLP pSTAT5 activation with an IC₅₀ value of 13pM. These experiments demonstrate that BDG33.023 is functionally blocking TSLP in a biologically relevant cell-based setting.

[465] *Summary:* The “re-epitoped” engineered BDG33.003 (clone 2), BDG33.023, and BDG33.025 antibodies were shown to bind both TSLP and IL-13. In contrast to the bispecific antibody format where each Fv has a specificity to a single antigen, these three antibodies are a standard IgG format, and each Fv has specificity to both IL-13 and TSLP. In addition, BDG33.023’s paratopes for IL-13 and TSLP was shown to be at least partly overlapping. All three IgGs interfere with the IL-13R/IL4R and TSLPR/IL-7R signaling cascade. Such antibodies could be used as a component of a therapeutic treatment, for example but not limited to, severe asthma, atopic dermatitis, and other allergic and respiratory conditions.

[466] While certain features of the engineered dual antibodies disclosed have been illustrated and described herein, many modifications, substitutions, changes, and equivalents will now occur to those of ordinary skill in the art. It is, therefore, to be understood that the appended claims are intended to cover all such modifications and changes as fall within the true spirit of what has been described herein.

Example 5: Biochemical Characterization of Dual Binding Antibodies

[467] *Objective:* To examine the biochemical and functional properties of dual binding antibodies BDG38.074 to BDG38.143.

[468] *Methods:*

[469] Peripheral blood mononuclear cells (PBMCs) (Cell Generation, CAT:101061021) were used to determine IL-13 and hTSLP inhibition. PBMCs were thawed and cultured in growth medium comprising of RPMI – 1640, 10% FBS, 1% Glutamax, 1% Sodium – Pyruvate, 0.1% 2-ME, 1% Pen-Strep and 1% nonessential AA. The cells were seeded in 96 well plate at 5x10⁵ cells/well. Fifteen ng/mL hTSLP and 1.25ng/mL IL-13 were incubated with antibodies for half an hour in 37°C, then added to the cells, to a total of 200uL. Cells were incubated for 48 hours at 37°C, 5% CO₂.

[470] CD23 upregulates in human monocytes in the presence of IL-13 (RD May et.al, 2011). IC₅₀ of antibody inhibition of IL-13 was determined by measuring CD23 expression level in monocytes. At the end of 48 hours incubation of the cells with different concentrations of antibodies, monocytes were detached from the bottom of the wells using cold PBS and scraping. Cells were marked using CD3 (Bio Legend, CAT: 300450), CD14 (Bio Legend, CAT: 301814),

CD19 (Bio Legend, CAT: 302212) and CD23 (Bio Legend, CAT: 338506) antibodies. CD23 percentage of CD14+ population was measured using CytoFLEX flow cytometer (Beckman Coulter).

[471] IC₅₀ of antibody inhibition of hTSLP was determined by TARC inhibition. TARC levels were determined using TARC DUOSET ELISA kit DY364 (R&D systems) according to kit instructions. Briefly, ELISA high bonding protein plates were plated with capture (non-biotinylated) antibody, diluted in PBSX1. Plates were sealed and incubated overnight. The following day plates were washed and blocked using PBST 2%BSA in room temperature, shaking, for two hours. Supernatant from the PBMCs plates was transferred to the wells. Detection was preformed using the kit's detection antibodies (Biotinylated) in PBS 1%BSA and Streptavidin – HRP in PBST 2%BSA. After adding TMB stop solution, ELISA plates were read at 450nm. Values were analyzed using standard sample curve.

[472] Other methods used to provide the results described and presented in this Example have been described in Example 1 above.

[473] *Results:* The tables below present the amino acid sequences for antibodies BDG38.074 to BDG38.143. The VH and VL sequences are shown in Table 10, whereas the heavy and light chain CDRs are shown in Table 8 and 9 respectively.

Table 8: Amino Acid Sequences of Heavy-Chain CDR Regions for Antibodies BDG38.074 to BDG38.143

Antibodies	HCDR1	SEQ ID NO	HCDR2	SEQ ID NO	HCDR3	SEQ ID NO
BDG38_074_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_079_VH	GFAFRTYG	349	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_094_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_138_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_075_VH	GFAFRTYG	349	IWYDGSAT	352	VRAPQWYLSAEAFDL	353
BDG38_076_VH	GFAFRTYG	349	IWDDGSAT	354	VRAPQWYLTAEAFDL	351
BDG38_077_VH	GFAFRTYG	349	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_078_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_080_VH	GFAFRTYG	349	IWYDGSAT	352	VRAPQWYLSAEAFDL	353
BDG38_081_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_082_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_083_VH	GFEFRTYG	355	IWDDGSNT	350	ARAPQWYLTAEAFDL	357
BDG38_084_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_085_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_086_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_087_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_088_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351

BDG38_089_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_090_VH	GFEFRTYG	355	IWDDGSNT	350	ARAPQWYLTAEAFDL	357
BDG38_091_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_092_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_093_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_095_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_096_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_097_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_098_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_099_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_100_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_101_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_102_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_103_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_104_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_105_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_106_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_107_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_108_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_109_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_110_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_111_VH	GFEFRTYG	355	IWDDGSAT	354	VRAPQWYLTAEAFDL	351
BDG38_112_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_113_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_114_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_115_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_116_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_117_VH	GFEFRTYG	355	IWDDGSNT	350	ARAPQWYLTAEAFDL	357
BDG38_118_VH	GFEFRTYG	355	IWYDGSAT	352	VRAPQWYLTAEAFDL	351
BDG38_119_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_120_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_121_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYLTAEAFDL	351
BDG38_122_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_123_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_124_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_125_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_126_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_127_VH	GFEFRTYG	355	IWYDGSAT	352	ARAPQWYLTAEAFDL	357
BDG38_128_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_129_VH	GFEFRTYG	355	IWDDGSNT	350	ARAPQWYLTAEAFDL	357
BDG38_130_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_131_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_132_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_133_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_134_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYNSAEAFDL	358
BDG38_135_VH	GFAFRTYG	349	IWYDGSNT	356	VRAPQWYNSAEAFDL	358

BDG38_136_VH	GFEFRTYG	355	IWYDGSNT	356	VRAPQWYNSAEAFDL	358
BDG38_137_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_139_VH	GFAFRTYG	349	IWDDGSNT	350	VRAPQWYLSAEAFDL	353
BDG38_140_VH	GFEFRTYG	355	IWDDGSNT	350	ARAPQWYLTAEAFDL	357
BDG38_141_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_142_VH	GFEFRTYG	355	IWDDGSNT	350	VRAPQWYLTAEAFDL	351
BDG38_143_VH	GFEFRTYG	355	IWDDGSAT	354	VRAPQWYLTAEAFDL	351

Table 9: Amino Acid Sequences of Light-Chain CDR Regions for Antibodies BDG38.074 to BDG38.143

Antibodies	LCDR1	SEQ ID NO	LCDR2	SEQ ID NO	LCDR3	SEQ ID NO
BDG38_074_VL	MIGAYL	359	DDV	360	QVWDHNTDKMV	361
BDG38_079_VL	MIGGYL	364	DDV	360	QVWDHDSNTMV	371
BDG38_094_VL	LIGAYL	362	DDV	360	QVWDHNSNHMV	384
BDG38_138_VL	MIGGYL	364	DDV	360	QVWDHNSNHMV	384
BDG38_075_VL	LIGAYL	362	DDV	360	QVWDHDTNTMV	363
BDG38_076_VL	MIGGYL	364	DDV	360	QVWDHDTNHMV	365
BDG38_077_VL	LIGSRL	366	DDS	367	QVWDHSSNTYV	368
BDG38_078_VL	LIGARL	369	DDS	367	QVWDYYS DHMV	370
BDG38_080_VL	LIGAYL	362	DDS	367	QVWDHNTNHMV	372
BDG38_081_VL	MIGAYL	359	DDV	360	QVWDHNTQQMV	373
BDG38_082_VL	MIGGYL	364	DDV	360	QVWDHDTNQVV	374
BDG38_083_VL	LIGAKL	375	DDS	367	QVWDYSSDTMV	376
BDG38_084_VL	LIGAYL	362	DDV	360	QVWDHSTNTMV	377
BDG38_085_VL	LIGARL	369	DDS	367	QVWDYSSNSYV	378
BDG38_086_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_087_VL	MIGAYL	359	DDV	360	QVWDHNSNQMV	380
BDG38_088_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_089_VL	LIGARL	369	DDS	367	QVWDHSSNHYV	381
BDG38_090_VL	LIGARL	369	DDS	367	QVWDYYS DHMV	370
BDG38_091_VL	LIGSRL	366	DDS	367	QVWDHYSNHYV	382
BDG38_092_VL	LIGARL	369	DDS	367	QVWDYYS DHMV	370
BDG38_093_VL	LIGARL	369	DDS	367	QVWDYSADSYV	383
BDG38_095_VL	LIGSRL	366	DDS	367	QVWDYYS DSYV	385
BDG38_096_VL	LIGARL	369	DDS	367	QVWDYSSDSMV	386
BDG38_097_VL	LIGARL	369	DDS	367	QVWDYYS DHYV	387
BDG38_098_VL	LIGARL	369	DDS	367	QVWDYSSDSMV	386
BDG38_099_VL	LIGARL	369	DDS	367	QVWDYSSDSYV	388
BDG38_100_VL	LIGARL	369	DDS	367	QVWDYYSNSYV	389
BDG38_101_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_102_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_103_VL	LIGARL	369	DDS	367	QVWDYYSNSYV	389
BDG38_104_VL	LIGARL	369	DDS	367	QVWDYYS DSYV	390
BDG38_105_VL	LIGARL	369	DDS	367	QVWDYYSNSYV	389
BDG38_106_VL	LIGSRL	366	DDS	367	QVWDHYS DHMV	391

BDG38_107_VL	LIGARL	369	DDS	367	QVWDYSSDSYV	388
BDG38_108_VL	LIGARL	369	DDS	367	QVWDYSSNSYV	389
BDG38_109_VL	LIGARL	369	DDS	367	QVWDYSSNSYV	378
BDG38_110_VL	LIGAKL	375	DDS	367	QVWDYSSNHMV	392
BDG38_111_VL	LIGARL	369	DDS	367	QVWDYYSANSYV	393
BDG38_112_VL	LIGARL	369	DDS	367	QVWDYSSDTYV	394
BDG38_113_VL	LIGARL	369	DDS	367	QVWDYSSDSYV	388
BDG38_114_VL	LIGARL	369	DDS	367	QVWDYSANSYV	395
BDG38_115_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_116_VL	LIGARL	369	DDS	367	QVWDYYSDTMV	396
BDG38_117_VL	LIGAKL	375	DDS	367	QVWDYSSDTMV	376
BDG38_118_VL	LIGARL	369	DDS	367	QVWDYSSDHVYV	397
BDG38_119_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_120_VL	LIGARL	369	DDS	367	QVWDYSSDHMV	398
BDG38_121_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379
BDG38_122_VL	LIGARL	369	DDS	367	QVWDYSSDSYV	388
BDG38_123_VL	LIGARL	369	DDS	367	QVWDYSSDTYV	394
BDG38_124_VL	LIGARL	369	DDS	367	QVWDYSSDTYV	394
BDG38_125_VL	LIGARL	369	DDS	367	QVWDYSSDHMV	398
BDG38_126_VL	LIGAKL	375	DDS	367	QVWDYSSDHMV	398
BDG38_127_VL	LIGAKL	375	DDS	367	QVWDYYSDTYV	399
BDG38_128_VL	LIGAKL	375	DDS	367	QVWDYYADTMV	400
BDG38_129_VL	LIGAKL	375	DDS	367	QVWDYSSDHMV	398
BDG38_130_VL	LIGARL	369	DDS	367	QVWDYSSNSYV	378
BDG38_131_VL	LIGAKL	375	DDS	367	QVWDYYSNTMV	401
BDG38_132_VL	LIGARL	369	DDS	367	QVWDYSSDHMV	398
BDG38_133_VL	LIGARL	369	DDS	367	QVWDYSSDTYV	394
BDG38_134_VL	LIGARL	369	DDS	367	QVWDYSADTMV	402
BDG38_135_VL	LIGARL	369	DDS	367	QVWDHSADTMV	403
BDG38_136_VL	LIGAKL	375	DDS	367	QVWDYSSDTMV	376
BDG38_137_VL	LIGARL	369	DDS	367	QVWDYSADTMV	402
BDG38_139_VL	MIGAYL	359	DDV	360	QVWDHNSDHMV	404
BDG38_140_VL	LIGAKL	375	DDS	367	QVWDYSANHMV	405
BDG38_141_VL	LIGSRL	366	DDS	367	QVWDYYSNHMV	406
BDG38_142_VL	LIGAKL	375	DDS	367	QVWDYYSHTMV	407
BDG38_143_VL	LIGARL	369	DDS	367	QVWDYSSNTYV	379

Table 10: Amino Acid Sequences of VH and VL Regions for Antibodies BDG38.074 to BDG38.143

Antibodies	Amino Acid Sequence	SEQ ID NO:
BDG38_074_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAIVWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	209

BDG38_074_VL	SYVLTQPPSVSVAPGQTATITCGGNMIGAYLVHWYQQKPGQAPLLV VYDDVDRPNRIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWD HNTDKMVFGGGTKLTVL	210
BDG38_079_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	219
BDG38_079_VL	SYVLTQPPSVSVAPGETATITCGGNMIGGYLVHWYQQKPGQAPLLV IYDDVDRPDRIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH DSNTMVFGGGTKLTVL	220
BDG38_094_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	249
BDG38_094_VL	SYVLTQPPSVSVAPGETASITCGGNLIGAYLVHWYQQKPGQAPLLVI YDDVDRPARIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDHN SNHMFVFGGGTKLTVL	250
BDG38_138_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	337
BDG38_138_VL	SYVLTQPPSVSVAPGETASITCGGNMIGGYLVHWYQQKPGQAPVLV IYDDVDRPSRIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH NSNHMFVFGGGTKLTVL	338
BDG38_075_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLSAEAFDLWGQGTMTVSS	211
BDG38_075_VL	SYVLTQPPSVSVAPGETATITCGGNLIGAYLVHWYQQKPGQAPVLV IYDDVDRPARIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH DTNTMVFGGGTKLTVL	212
BDG38_076_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	213
BDG38_076_VL	SYVLTQPPSVSVAPGETASITCGGNMIGGYLVHWYQQKPGQAPLLV IYDDVDRPARIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH DTNHMFVFGGGTKLTVL	214
BDG38_077_VH	QMQLVESGGGVVQPGRSLTLSCAASGFAFRITYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	215
BDG38_077_VL	SYVLTQPPSVSVAPGETATITCGGALIGSRLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDHSS NTYVFGGGTKLTVL	216
BDG38_078_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRITYGMHWVRQAPGKG LEWLGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	217
BDG38_078_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEAGDEADYYCQVWDYYS DHMFVFGGGTKLTVL	218

BDG38_080_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLSAEAFDLWGQGTMTVSS	221
BDG38_080_VL	SYVLTQPPSVSVAPGETATITCGGNLIGAYLVHWYQQKPGQAPVLV IYDDSDRPDRIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH NTNHMVFGGGTKLTVL	222
BDG38_081_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	223
BDG38_081_VL	SYVLTQPPSVSVAPGQTARITCGGNMIGAYLVHWYQQKPGQAPLL VIYDDVDRPDRIPIERFSGSNSGNTATLTISRVEAGDEADYYCQVWD HNTQQMVFGGGTKLTVL	224
BDG38_082_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	225
BDG38_082_VL	SYVLTQPPSVSVAPGETATITCGGNMIGGYLVHWYQQKPGQAPVL VIYDDVDRPDRIPIERFSGSNSGNTATLTISRVEAGDEADYYCQVWD HDTNQVVFGGGTKLTVL	226
BDG38_083_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRITYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	227
BDG38_083_VL	SYILTQPPSVSVAPGQTATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPIERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYSS DTMVFGGGTKLTVL	228
BDG38_084_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	229
BDG38_084_VL	SYVLTQPPSVSVAPGETATITCGGNLIGAYLVHWYQQKPGQAPVLV VYDDVDRPDRIPIERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH STNTMVFGGGTKLTVL	230
BDG38_085_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRITYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	231
BDG38_085_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPIERFSGSNIGNTATLTIERVEAGDEADYYCQVWDYS SNSYVFGGGTKLTVL	232
BDG38_086_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRITYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	233
BDG38_086_VL	SYVLTQPPSVSVAPGQTARITCGGNLIGARLVHWYQQKPGQAPVLV VYDDSDRPSRIPIERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYS SNTYVFGGGTKLTVL	234
BDG38_087_VH	QMQLVESGGGVVQPGRSLRLSCAASGFAFRITYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	235

BDG38_087_VL	SYVLTQPPSVSVAPGQTATITCGGNMIGAYLVHWYQQKPGQAPVL VIYDDVDRPDRIPEFSGSNSGNTATLTISRVEAGDEADYYCQVWD HNSNQMVFGGGTKLTVL	236
BDG38_088_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	237
BDG38_088_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDYS SNTYVFGGGTKLTVL	238
BDG38_089_VH	QMQLVESGGGVVQPGRSLRSLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	239
BDG38_089_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDHSS NHVYVFGGGTKLTVL	240
BDG38_090_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWLGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	241
BDG38_090_VL	SYILTQPPSVSVAPGETARITCGGNLIGARLVHWYQQKPGQAPVLV VYDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDY YSDHMFVFGGGTKLTVL	242
BDG38_091_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWLGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	243
BDG38_091_VL	SYILTQPPSVSVAPGETATITCGGNLIGSLRVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDHYS NHVYVFGGGTKLTVL	244
BDG38_092_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWLAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	245
BDG38_092_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYY SDHMFVFGGGTKLTVL	246
BDG38_093_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	247
BDG38_093_VL	SYVLTQPPSVSVAPGQTARITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYS ADSYVFGGGTKLTVL	248
BDG38_095_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	251
BDG38_095_VL	SYVLTQPPSVSVAPGQTATITCGGALIGSLRVHWYQQKPGQAPVLV VYDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYY SDSYVFGGGTKLTVL	252

BDG38_096_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAWIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	253
BDG38_096_VL	SYILTQPPSVSVAPGETARITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYSS DSMVFGGGTKLTVL	254
BDG38_097_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWLGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	255
BDG38_097_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYSS DHYVFGGGTKLTVL	256
BDG38_098_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVAWIWDDGSNTVYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	257
BDG38_098_VL	SYVLTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTISRVEAGDEADYYCQVWDYS SDSMVFGGGTKLTVL	258
BDG38_099_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	259
BDG38_099_VL	SYILTQPPSVSVAPGETARITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDYSS DSYVFGGGTKLTVL	260
BDG38_100_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	261
BDG38_100_VL	SYVLTQPPSVSVAPGETARITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYY SNSYVFGGGTKLTVL	262
BDG38_101_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	263
BDG38_101_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYSS NTYVFGGGTKLTVL	264
BDG38_102_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	265
BDG38_102_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV VYDDSDRPSRIPERFSGSNIGNTATLTIEDVEAGDEADYYCQVWDY SSNTYVFGGGTKLTVL	266
BDG38_103_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	267

BDG38_103_VL	SYVLTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLV VYDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDY YSNSYVFGGGTKLTVL	268
BDG38_104_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	269
BDG38_104_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDYYS DSYVFGGGTKLTVL	270
BDG38_105_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	271
BDG38_105_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIERVEAGDEADYYCQVWDYYS SNSYVFGGGTKLTVL	272
BDG38_106_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTVYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	273
BDG38_106_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDHYS DHMVFGGGTKLTVL	274
BDG38_107_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	275
BDG38_107_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYSS DSYVFGGGTKLTVL	276
BDG38_108_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	277
BDG38_108_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYYS NSYVFGGGTKLTVL	278
BDG38_109_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	279
BDG38_109_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYSS NSYVFGGGTKLTVL	280
BDG38_110_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	281
BDG38_110_VL	SYILTQPPSVSVAPGETTRITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEAGDEADYYCQVWDYSS NHMVFGGGTKLTVL	282

BDG38_111_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	283
BDG38_111_VL	SYVLTQPPSVSVAPGETATITCGGALIGARLVHWYQQKPGQAPVLV YDDSDRPSRIPERFSGSNIGNTATLTISRVEAGDEADYYCQVWDY YANSYVFGGGTKLTVL	284
BDG38_112_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWLGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	285
BDG38_112_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYSS DTYVFGGGTKLTVL	286
BDG38_113_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	287
BDG38_113_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDYS SDSYVFGGGTKLTVL	288
BDG38_114_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	289
BDG38_114_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEEGDEADYYCQVWDYSA NSYVFGGGTKLTVL	290
BDG38_115_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWLGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	291
BDG38_115_VL	SYILTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNSGNTATLTISDVEEGDEADYYCQVWDYSS NTYVFGGGTKLTVL	292
BDG38_116_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	293
BDG38_116_VL	SYVLTQPPSVSVAPGQTATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIERVEAGDEADYYCQVWDYY SDTMVFGGGTKLTVL	294
BDG38_117_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	295
BDG38_117_VL	SYVLTQPPSVSVAPGETATITCGGNLIGAKLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIEDVEAGDEADYYCQVWDYS SDTMVFGGGTKLTVL	296
BDG38_118_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	297

BDG38_118_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYSS DHYVFGGGTKLTVL	298
BDG38_119_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	299
BDG38_119_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYSS NTYVFGGGTKLTVL	300
BDG38_120_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	301
BDG38_120_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNSGNTATLTISDVEAGDEADYYCQVWDYS SDHMFVFGGGTKLTVL	302
BDG38_121_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	303
BDG38_121_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEAGDEADYYCQVWDYSS NTYVFGGGTKLTVL	304
BDG38_122_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	305
BDG38_122_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYSS DSYVFGGGTKLTVL	306
BDG38_123_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	307
BDG38_123_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDYSS DTYVFGGGTKLTVL	308
BDG38_124_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	309
BDG38_124_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDYSS DTYVFGGGTKLTVL	310
BDG38_125_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	311
BDG38_125_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYS SDHMFVFGGGTKLTVL	312

BDG38_126_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	313
BDG38_126_VL	SYILTQPPSVSVAPGQTARITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDYSS DHMVFGGGTKLTVL	314
BDG38_127_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWYDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	315
BDG38_127_VL	SYILTQPPSVSVAPGETATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEAGDEADYYCQVWDYSS DTYVFGGGTKLTVL	316
BDG38_128_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	317
BDG38_128_VL	SYILTQPPSVSVAPGETATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNSGNTATLTISDVEEGDEADYYCQVWDYY ADTMVFGGGTKLTVL	318
BDG38_129_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	319
BDG38_129_VL	SYILTQPPSVSVAPGQTATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYSS DHMVFGGGTKLTVL	320
BDG38_130_VH	QMQLVESGGGVVQPGRSLTLSCAASGFAFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	321
BDG38_130_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIERVEAGDEADYYCQVWDYSS NSYVFGGGTKLTVL	322
BDG38_131_VH	QMQLVESGGGVVQPGRSLTLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	323
BDG38_131_VL	SYILTQPPSVSVAPGETATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYSS NTMVFGGGTKLTVL	324
BDG38_132_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	325
BDG38_132_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEAGDEADYYCQVWDYSS DHMVFGGGTKLTVL	326
BDG38_133_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	327

BDG38_133_VL	SYILTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEAGDEADYYCQVWDYSS DTYVFGGGTKLTVL	328
BDG38_134_VH	QMQLVESGGGVVQPGQSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWLAVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYNSAEAFDLWGQGTMTVSS	329
BDG38_134_VL	SYVLTQPPSVSVAPGQTARITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSHIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDYS ADTMVFGGGTKLTVL	330
BDG38_135_VH	QMQLVESGGGVVQPGQSLRLSCAASGFAFRTYGMHWVRQAPGKG LEWLAVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYNSAEAFDLWGQGTMTVSS	331
BDG38_135_VL	SYVLTQPPSVSVAPGQTARITCGGNLIGARLVHWYQQKPGQAPVLV VYDDSDRPSHIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH SADTMVFGGGTKLTVL	332
BDG38_136_VH	QMQLVESGGGVVQPGQSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWLAVIWYDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYNSAEAFDLWGQGTMTVSS	333
BDG38_136_VL	SYVLTQPPSVSVAPGQTARITCGGNLIGAKLVHWYQQKPGQAPVLV IYDDSDRPSHIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDYS SDTMVFGGGTKLTVL	334
BDG38_137_VH	QMQLVESGGGVVQPGQSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	335
BDG38_137_VL	SYVLTQPPSVSVAPGETATITCGGNLIGARLVHWYQQKPGQAPVLV IYDDSDRPSHIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDYS ADTMVFGGGTKLTVL	336
BDG38_139_VH	QMQLVESGGGVVQPGQSLRLSCAASGFAFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLSAEAFDLWGQGTMTVSS	339
BDG38_139_VL	SYVLTQPPSVSVAPGETATITCGGNMIGAYLVHWYQQKPGQAPLLV IYDDVDRPARIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDH NSDHMVFGGGTKLTVL	340
BDG38_140_VH	QMQLVESGGGVVQPGQSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCARAPQWYLTAEAFDLWGQGTMTVSS	341
BDG38_140_VL	SYILTQPPSVSVAPGETATITCGGNLIGAKLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTIEDVEAGDEADYYCQVWDYS ANHMOVFGGGTKLTVL	342
BDG38_141_VH	QMQLVESGGGVVQPGQSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVSS	343
BDG38_141_VL	SYILTQPPSVSVAPGETATITCGGALIGSRLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISDVEEGDEADYYCQVWDYYS NHMVFGGGTKLTVL	344

BDG38_142_VH	QMQLVESGGGVVQPGRSLTSCAASGFEFRTYGMHWVRQAPGKG LEWVAVIWDDGSNTHYADSVKGRFTITRDNSKNTLNLQMNSLRAE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVTVSS	345
BDG38_142_VL	SYILTQPPSVSVAPGQTATITCGGNLIGAKLVHWYQQKPGQAPVLV VYDDSDRPSRIPERFSGSNIGNTATLTIERVEEGDEADYYCQVWDY YSHTMVFGGGTKLTVL	346
BDG38_143_VH	QMQLVESGGGVVQPGRSLRLSCAASGFEFRTYGMHWVRQAPGKG LEWVGVIWDDGSATHYADSVKGRFTITRDNSKNTLNLQMNSLRVE DTAVYYCVRAPQWYLTAEAFDLWGQGTMTVTVSS	347
BDG38_143_VL	SYILTQPPSVSVAPGQTATITCGGALIGARLVHWYQQKPGQAPVLVI YDDSDRPSRIPERFSGSNIGNTATLTISRVEEGDEADYYCQVWDYSS NTYVFGGGTKLTVL	348

[474] Data presented below demonstrate biochemical and functional properties for some of the dual binding antibodies disclosed herein. **Figure 13** shows retention time and calculated pI for some of the dual binding antibodies disclosed herein, demonstrating that BDG antibodies are monodispersed and highly homogenic.

[475] **Figures 14A-14C** show competitive ELISA of some of the dual binding antibodies and Tezepelumab with TSLP. ELISA plates were coated overnight at 4°C with 50 ng/well Tezepelumab. BDG antibodies were double diluted and pre-incubated with a constant concentration of 7nM TSLP-His for 1 hour at room temperature. After blocking and washing steps, BDG antibodies-TSLP mix were added to the plates, incubated for 10 minutes and washed again, followed by a 30 minute incubation with anti-HIS. BDG antibodies exhibited similar IC50 values and complete competition with Tezepelumab at the TSLP epitope.

[476] **Figure 15** shows nanoscale differential scanning fluorimetry (nanoDSF) analysis of some of the dual binding antibodies disclosed herein. All BDG antibodies were analyzed to test stability over a range of thermal changes using nanoDSF (See, Table 11 below). Antibodies with a $T_m > 60^\circ\text{C}$ were selected to further characterizations.

[477] **Table 11: SEC and DSF summary table. SEC column presents Area Under the Curve (AUC) peak ratio of monodispersed BDG antibodies fraction. DSF columns present T-onset and T_m values of the different BDG antibodies.**

Antibody#	SEC (%)	DSF 1st repetition		DSF 2nd repetition	
		T-onset	T_m -1	T-onset	T_m -1
BDG38.074	100	60.3	62.2		
BDG38.075	88.65	61	65.5	60.7	65.4
BDG38.076	91.98				
BDG38.077	97.43	66.4	70.4	66	70.3
BDG38.078	89.12	64.3	67.4	64.4	67.4

BDG38.079	96.28	61.4	64.2	61.7	64.3
BDG38.080	97.4	63.7	66.8	63.8	66.8
BDG38.081	97.54	63.3	65.9	63.4	65.9
BDG38.082	99.35	65.6	69.3	65.4	69.3
BDG38.083	92.88	66.6	70.1	66.8	70.2
BDG38.084	96.7	20.0°C	49.6°C	23.7°C	48.4°C
BDG38.085	90		51.1°C		51.5°C
BDG38.086	68		51.3°C		51.1°C
BDG38.087					
BDG38.088	84.3	65.2°C	71.5°C		51.6°C
BDG38.089	92.34	61.0°C	71.2°C	60.8°C	71.3°C
BDG38.090	96.6	63.5°C	69.7°C	63.5°C	69.7°C
BDG38.091	86.78	57.6°C	72.0°C	58.0°C	72.1°C
BDG38.092	95.8	64.5°C	68.4°C	63.9°C	68.2°C
BDG38.093	82.33		49.6°C		49.6°C
BDG38.094	96.15	21.4°C	49.7°C	62.4°C	65.3°C
BDG38.095	98.01	64.6°C	68.2°C	53.0°C	53.5°C
BDG38.096	97	63.5°C	66.9°C	64.2°C	67.0°C
BDG38.097	68	65.0°C	68.5°C	64.7°C	68.5°C
BDG38.098	95.8	63.4°C	66.1°C	62.8°C	66.2°C
BDG38.099	75.2		49.4°C		58.3°C
BDG38.100	84		50.8°C		51.2°C
BDG38.101	97	63.6°C	70.6°C	64.5°C	70.5°C
BDG38.102	70.6		58.9°C	65.4°C	68.8°C
BDG38.103	78.26	51.0°C	51.2°C	65.5°C	69.2°C
BDG38.104	76.36	65.0°C	69.3°C	65.4°C	69.3°C
BDG38.105	74.87		51.8°C	64.7°C	71.2°C
BDG38.106	96.8	63.9°C	67.5°C	64.3°C	67.6°C
BDG38.107	69.19		49.3°C	49.1°C	49.2°C
BDG38.108	75.47	42.2°C	42.9°C		51.5°C
BDG38.109	56		51.2°C		51.4°C
BDG38.110	77.15	27.7°C	56.8°C	26.0°C	57.2°C
BDG38.111	92	23.4°C	55.4°C	43.9°C	54.9°C
BDG38.112	71.97		49.5°C		49.3°C
BDG38.113	71		49.6°C		50.0°C
BDG38.114	67.9		59.1°C		52.0°C
BDG38.115	72.41		56.8°C	57.4°C	71.5°C
BDG38.116	88	63.8°C	67.6°C	64.3°C	67.9°C
BDG38.117	96	64.6°C	69.7°C	63.9°C	69.7°C
BDG38.118	90.5	64.5°C	69.6°C	64.8°C	69.7°C
BDG38.119	63.19		51.7°C		59.9°C
BDG38.120	82	28.3°C	55.8°C	62.9°C	66.8°C
BDG38.121	51		49.9°C		50.1°C
BDG38.122	67	58.5°C	58.9°C		59.0°C
BDG38.123	59		50.6°C		50.5°C

BDG38.124	69	57.5°C	58.5°C	56.2°C	58.3°C
BDG38.125	85.47	63.9°C	67.2°C	63.6°C	67.1°C
BDG38.126	93	63.9°C	67.3°C	64.0°C	67.3°C
BDG38.127	89	62.9°C	69.3°C	62.8°C	69.3°C
BDG38.128	81	63.4°C	67.4°C	63.2°C	67.4°C
BDG38.129	95.98	63.8°C	70.1°C	63.4°C	70.1°C
BDG38.130	38.5		60.2°C		60.5°C
BDG38.131	71.5	64.2°C	68.8°C	64.4°C	68.8°C
BDG38.132	94.35	63.3°C	67.1°C	63.6°C	67.1°C
BDG38.133	61.3		50.8°C		50.4°C
BDG38.134	93.57	63.5°C	67.4°C	63.6°C	67.4°C
BDG38.135	92.78	62.6°C	65.6°C	62.6°C	65.6°C
BDG38.136	93.38	63.8°C	67.3°C	63.6°C	67.3°C
BDG38.137	77.8	63.4°C	67.9°C	62.7°C	67.8°C
BDG38.138	94.8	20.0°C	50.4°C	62.8°C	66.0°C
BDG38.139	90	61.6°C	64.8°C	62.0°C	64.8°C
BDG38.140	82.2	63.6°C	70.6°C	20.0°C	50.9°C
BDG38.141	73.25	20.0°C	55.2°C	20.0°C	56.1°C
BDG38.142	80	63.8°C	67.8°C	63.8°C	67.7°C
BDG38.143	74	21.3°C	57.5°C		56.9°C

[478] **Figures 16A-16F** show the results of SPR (Surface Plasmon Resonance) analysis for some of the dual binding antibodies disclosed herein for human/cyno IL-13 and TSLP. SPR analysis was performed to assess the kinetics of BDG antibodies binding to TSLP and IL-13. SPR data indicates dissociation constants of double digit pM for BDG antibodies binding to IL-13 and TSLP.

[479] Table 12 presents IC₅₀ inhibition values of CD23 expression on monocytes and IC₅₀ inhibition values of TARC secretion by PBMCs. This data demonstrates that each BDG dual antibody can inhibit both IL-13 and TSLP functions in human PBMCs.

[480] **Table 12: PBMC Functional Assay Summary Table.**

Antibody#	PBMC Functional Assay	
	IC ₅₀ CD23 [nM]	IC ₅₀ TARC [nM]
BDG38.074	0.1578	2.771
BDG38.075	0.09752	4.168
BDG38.076	0.3078	3.206
BDG38.077	0.01245	2.466
BDG38.078	0.1252	6.516
BDG38.079	0.07179	1.961
BDG38.080	0.2011	0.7904
BDG38.081	0.03386	5.929
BDG38.082	0.03908	1.577

BDG38.083	0.02146	11.37
BDG38.090	0.006556	0.007555
BDG38.092	0.06747	5.876
BDG38.094	0.1724	0.3556
BDG38.096	0.2504	23.41
BDG38.106	0.1616	NaN
BDG38.112	0.05937	0.1039
BDG38.116	0.08366	59.36
BDG38.117	0.1207	5.40E+15
BDG38.125	0.0527	NaN
BDG38.126	0.0868	2.575
BDG38.128	0.01492	NaN
BDG38.129	0.029	NaN
BDG38.131	0.07	NaN
BDG38.132	0.43	22.39
BDG38.135	0.4075	0.8042
BDG38.137	0.3728	0.8233
BDG38.138	0.1638	2.358
BDG38.139	0.012	-
BDG38.140	0.146	-

*NaN- stands for values that could not be determined.

[481] Table 13 presents the IC50 values obtained from competitive ELISA with Tezepelumab. The data demonstrates that BDG dual antibodies compete with Tezepelumab for TSLP binding.

[482] **Table 13: IC50 Competition ELISA Values Summary Table.**

Antibody#	IC50(nM) vs anti-TSLP benchmark
BDG38.074	3.66
BDG38.075	3.94
BDG38.076	5.382
BDG38.077	8.47
BDG38.078	4.3
BDG38.079	3.63
BDG38.080	3.27
BDG38.081	2.2
BDG38.082	2.7
BDG38.083	3.33
BDG38.090	7.31
BDG38.092	7.382
BDG38.096	5.871
BDG38.106	10.09
BDG38.112	29.14
BDG38.116	5.123
BDG38.117	6.137

BDG38.125	8.442
BDG38.126	5.953
BDG38.128	5.378
BDG38.129	5.494
BDG38.132	9.203
BDG38.137	10.89
BDG38.139	9.571
BDG38.140	5.628

[483] Table 14 presents data from a TSLP functional inhibition assay using MUTZ5 cells (human B cell precursor leukemia cells). IC50 values of STAT5 phosphorylation inhibition are presented in pM, and the R² column presents goodness of fit values. These results demonstrate that BDG dual antibodies can inhibit TSLP function in the MUTZ5 cell line expressing the native TSLP-receptor heterocomplex (TLSP-R and IL-7R).

[484] **Table 14: TSLP Functional Inhibition Assay in MUTZ5 cell line.**

Antibody#	IC50 [pM]	R ²
BDG38.074	46.74	0.925
BDG38.075	43.86	0.9919
BDG38.076	30.14	0.9322
BDG38.077	304.1	0.7798
BDG38.078	337.5	0.9148
BDG38.079	21.84	0.8359
BDG38.080	34.33	0.9534
BDG38.081	38.74	0.9556
BDG38.086	830.1	0.01605
BDG38.090	43.35	0.2951
BDG38.091	0.5069	0.43
BDG38.092	33.05	0.5048
BDG38.094	27.16	0.9719
BDG38.096	5.44E-05	0.6063
BDG38.098	68.84	0.07219
BDG38.107	0.5081	0.3199
BDG38.110	0.2951	0.8273
BDG38.112	11383	0.05737
BDG38.115	300.3	0.1132
BDG38.117	0.3628	0.5132
BDG38.118	0.6884	0.4463
BDG38.120	0.2315	0.7447
BDG38.125	8.74E-05	0.7192
BDG38.126	0.04215	0.7997
BDG38.127	0.1433	0.5647
BDG38.128	0.004796	0.639
BDG38.129	0.6765	0.5382

BDG38.132	0.3858	0.737
BDG38.134	32.04	0.7965
BDG38.137	1.855	0.08069
BDG38.138	45.24	0.9313
BDG38.139	88.95	0.8952
BDG38.140	2.211	0.5831

[485] Table 15 shows the results of a functional inhibition assay in IL-4/IL-13 reporter HEK 293 cells. IC₅₀ values of STAT6 phosphorylation inhibition is presented in nM. This data demonstrated that BDG dual antibodies can inhibit IL-13 function in reporter HEK cell line cells expressing the IL-13-receptor heterocomplex (IL-13R α 1 and IL-4R α).

[486] **Table 15: IL-13 Functional Inhibition Assay in IL-4/IL-13 Reporter HEK 293 cell line.**

Antibody#	HEK-IL-13 [nM]
BDG38.074	0.02407
BDG38.075	0.06912
BDG38.076	0.04938
BDG38.077	0.05055
BDG38.078	0.05079
BDG38.079	0.0253
BDG38.080	0.05923
BDG38.081	0.04828
BDG38.082	0.04295
BDG38.083	0.04194
BDG38.084	0.0419
BDG38.085	0.02355
BDG38.086	0.0174
BDG38.087	-----
BDG38.088	0.04861
BDG38.089	0.07497
BDG38.090	0.1035
BDG38.091	0.08321
BDG38.092	0.1197
BDG38.093	0.04018
BDG38.094	0.04873
BDG38.095	0.08224
BDG38.096	0.1505
BDG38.097	0.04645
BDG38.098	0.08838
BDG38.099	0.06009
BDG38.100	0.08009
BDG38.101	0.04035
BDG38.102	0.04211

BDG38.103	0.084
BDG38.104	0.03805
BDG38.105	0.1341
BDG38.106	0.01134
BDG38.107	0.09427
BDG38.108	0.06595
BDG38.109	0.06295
BDG38.110	0.02988
BDG38.111	0.05454
BDG38.112	0.03665
BDG38.113	0.02843
BDG38.114	0.09887
BDG38.115	0.06638
BDG38.116	0.1799
BDG38.117	0.08006
BDG38.118	0.08044
BDG38.119	0.06617
BDG38.120	0.2649
BDG38.121	0.05618
BDG38.122	0.07895
BDG38.123	0.04405
BDG38.124	0.05233
BDG38.125	0.05618
BDG38.126	0.07895
BDG38.127	0.04405
BDG38.128	0.05233
BDG38.129	0.07253
BDG38.130	0.06463
BDG38.131	0.0443
BDG38.132	0.08986
BDG38.133	0.02895
BDG38.134	0.2711
BDG38.135	0.4058
BDG38.136	0.1509
BDG38.137	0.1951
BDG38.138	0.02862
BDG38.139	0.04644
BDG38.140	0.04303
BDG38.141	0.08244
BDG38.142	0.05843
BDG38.143	0.04382

[487] *Antibody BDG38.074*

[488] The size exclusion chromatography (SEC) scans and nano-differential scanning fluorimetry (DSF) analysis of the melting point for antibody BDG38.074 are shown in **Figures**

17A and **17B**. SEC analysis (**Figures 17A**) was performed using BioResolve SEC mAb Column, 200Å, 2.5 µm, 4.6 x 300 mm at 0.5ml/min in PBS as a mobile phase and analyzed at 280nm. BDG38.074 shows a predominant monodisperse peak with undetectable aggregates. DSF analysis (**Figures 17B**) was performed using nanoDSF at a 1°C/min from 20-95°C.

[489] NanoDSF is monitoring the thermal unfolding of BDG38.074 according to the intrinsic fluorescence change at 350 and 330 nm. The top half of the graph in **Figure 17B** shows the fluorescence ratio of 350nm/330nm as a function of temperature and the bottom half shows the first derivative as a function of temperature. BDG38.074 was analyzed at 0.5 mg/ml in PBS showed to have a T-onset of 60.3°C and T_m of 62.2°C suggesting a relatively stable fold.

[490] The binding affinities of antibody BDG38.074 to human/cyno IL-13 and human/cyno TSLP are shown in **Figures 18A-18D**. Surface Plasmon Resonance (SPR) analysis of BDG38.074 binding to human IL-13 (**Figure 18A**), human TSLP (**Figure 18B**), cyno IL-13 (**Figure 18C**) and cyno TSLP (**Figure 18D**) using BiacoreT200. CM5 chip was coated with human antibody capture kit to obtain 3000-5000 RU and the antibody, served as the capture, was injected at a flow rate of 10 ul/ml to obtain 250-350 RU. Human/cynoTSLP and human/cynoIL-13 served as analytes in a concentration range of 30-0.153 nM. Contact time 300 sec and dissociation time 600 sec at a flow rate of 30 ul/min. BDG38.074 shows a high affinity to all cytokines with K_D values for hIL-13: 2.95E-11M, for hTSLP: 3.34E-12 M, for cyno IL-13: 7.33E-10 M and for cyno TSLP: 4.16E-12 M.

[491] The results of SPR (Surface Plasmon Resonance) analysis of antibody BDG38.074 for human or cyno IL-13 or TSLP are shown in **Figures 21A and 21B**. Based on the SPR analysis, the affinities of BDG 38.074 to human and cynomolgus IL-13 are at double and triple digit picomolar, respectively, and the affinities to human and cynomolgus TSLP at a single digit picomolar.

[492] **Figure 22A** shows antibody BDG38.074 exhibits inhibition of CD23 expression similar to the anti-IL-13 benchmark (Tralokinumab). **Figure 22B** shows antibody BDG38.074 inhibits TARC expression similar to anti-TSLP benchmark (Tezepelumab). These data demonstrate that while the anti-TSLP benchmark has only limited effect at inhibiting CD23 expression in monocytes, and the anti-IL-13 benchmark has only limited effect at inhibiting TARC levels, BDG38.074 inhibits both CD23 and TARC expression, demonstrating the unique ability of BDG dual antibodies to exert two distinct functions as a single standard IgG1 (LALA PG) antibody.

[493] **Figure 23** shows antibody BDG38.074 exhibits functional inhibition similar to anti-TSLP benchmarks in MUTZ-5 cell line, demonstrating that BDG38.074 inhibits TSLP function

with an IC₅₀ of about 35pM in cells expressing the native TSLP receptor subunits.

[494] **Figure 24** shows antibody BDG38.074 inhibits IL-13 function in HEK reporter cell line with double digit picomolar affinity, demonstrating that BDG 38.074 inhibits IL-13 function in cells expressing the IL-13 receptor heterocomplex IL-4R α and IL-13R α 1.

Antibody BDG38.079

[495] The size exclusion chromatography (SEC) scans and nano-differential scanning fluorimetry (DSF) analysis of the melting point for antibody BDG38.074 are shown in **Figures 19A and 19B**.

[496] SEC analysis (**Figure 19A**) was performed using BioResolve SEC mAb Column, 200Å, 2.5 μ m, 4.6 x 300 mm at 0.5ml/min in PBS as a mobile phase and analyzed at 280nm. BDG38.079 shows a predominant monodisperse peak with a minor percentage of aggregates (1.34%). DSF analysis (**Figure 19B**) was performed using nanoDSF at a 1°C/min from 20-95°C. NanoDSF is monitoring the thermal unfolding of BDG38.079 according to the intrinsic fluorescence change at 350 and 330 nm. The top half of the graph shows the fluorescence ratio of 350nm/330nm as a function of temperature and the bottom half shows the first derivative as a function of temperature. BDG38.079 was analyzed at 0.5 mg/ml in PBS showed to have a T-onset of 61.4°C and T_m of 64.2°C suggesting a relatively stable fold.

[497] The binding affinities of antibody BDG38.079 to human/cyno IL-13 and human/cyno TSLP are shown in **Figures 20A-20D**. Surface Plasmon Resonance (SPR) analysis of BDG38.079 binding to human IL-13 (**Figure 20A**), human TSLP (**Figure 20B**), cyno IL-13 (**Figure 20C**) and cyno TSLP (**Figure 20D**) using BiacoreT200. CM5 chip was coated with human antibody capture kit to obtain 3000-5000 RU and the antibody, served as the capture, was injected at a flow rate of 10 ul/ml to obtain 250-350 RU. Human/cynoTSLP and human/cynoIL-13 served as analytes in a concentration range of 30-0.153 nM. Contact time is 300 sec and dissociation time is 600 sec at a flow rate of 30 ul/min. BDG38.079 exhibits high affinity binding to all cytokines tested, with K_D values for hIL-13: 3.17E-11 M, for hTSLP: 2.21E-12 M, for cyno IL-13 9.01E-10 M and for cyno TSLP: 3.97E-12M.

[498] Results of SPR (Surface Plasmon Resonance) analysis of antibody BDG38.079 for human or cyno IL-13 or TSLP are shown in **Figure 21**. From the SPR analysis, the affinities of BDG 38.079 to human and cynomolgus IL-13 are double and triple digit picomolar respectively and the affinities to human and cynomolgus TSLP are single digit picomolar .

[499] **Figure 22** shows antibody BDG38.079 inhibits IL-13 function in HEK reporter cell

line with double digit picomolar affinity, demonstrating that BDG 38.079 inhibits IL-13 function in cells expressing the IL-13 receptor heterocomplex IL-4R α and IL-13R α 1.

[500] **Figure 23** shows antibody BDG38.079 exhibits similar functional inhibition to anti-TSLP benchmarks in MUTZ-5 cell line, demonstrating that BDG38.079 inhibits TSLP function with an IC50 of about 13pM in cells expressing the native TSLP receptor subunits.

[501] **Figure 24A** shows antibody BDG38.079 exhibits similar inhibition of CD23 expression as the anti-IL-13 benchmark (Tralokinumab). **Figure 24B** shows antibody BDG38.079 inhibits TARC expression similarly to anti-TSLP benchmark (Tezepelumab).

[502] These data demonstrate that while the anti-TSLP benchmark has only limited effect at inhibiting CD23 expression in monocytes, and the anti-IL-13 benchmark has only limited effect at inhibiting TARC levels, BDG38.079 inhibits both CD23 and TARC expression, demonstrating the unique ability of BDG dual antibodies to exert two distinct functions as a single standard IgG1 (LALA PG) antibody.

Antibody BDG38.094

[503] The size exclusion chromatography (SEC) scans and nano-differential scanning fluorimetry (DSF) analysis of the melting point for antibody BDG38.09 are shown in **Figures 25A and 25B**.

[504] The size exclusion chromatography (SEC) scans and nano-differential scanning fluorimetry (DSF) analysis of the melting point for antibody BDG38.094 are shown in **Figures 25A and 25B**. SEC analysis (**Figure 25A**) was performed using BioResolve SEC mAb Column, 200Å, 2.5 μ m, 4.6 x 300 mm at 0.4ml/min in PBS as a mobile phase and analyzed at 280nm. BDG38.094 shows a predominant monodisperse peak with undetectable aggregates. DSF analysis (**Figure 25B**) was performed using nanoDSF at a 1°C/min from 20-95°C. NanoDSF is monitoring the thermal unfolding of BDG38.094 according to the intrinsic fluorescence change at 350 and 330 nm. The top half of the graph shows the fluorescence ratio of 350nm/330nm as a function of temperature and the bottom half shows the first derivative as a function of temperature. BDG38.094 was analyzed at 0.5 mg/ml in PBS showed to have a T-onset of 62.4°C and Tm of 65.3°C suggesting a relatively stable fold.

[505] The binding affinities of antibody BDG38.094 to human/cyno IL-13 and human/cyno TSLP are shown in **Figures 26A-26D**. Surface Plasmon Resonance (SPR) analysis of BDG38.094 binding to human IL-13 (**Figure 26A**), human TSLP (**Figure 26B**), cyno IL-13 (**Figure 26C**) and cyno TSLP (**Figure 26D**) using BiacoreT200. CM5 chip was coated with human antibody capture

kit to obtain 3000-5000 RU and the antibody, served as the capture, was injected at a flow rate of 10 ul/ml to obtain 250-350 RU. Human/cyno TSLP and human/cyno IL-13 served as analytes in a concentration range of 30-0.153 nM. Contact time was 300 sec and dissociation time was 600 sec (for cyno IL-13 dissociation time was 200 sec) at a flow rate of 30 ul/min. BDG38.094 shows a high affinity to all cytokines with K_D values for hIL-13: $1.38E-11$ M for hTSLP: $<1E-12$ M (below limit of detection), for cyno IL-13: $2.70E-10$ M and for cyno TSLP: $2.75E-12$ M.

Antibody BDG38.138

[506] The size exclusion chromatography (SEC) scans and nano-differential scanning fluorimetry (DSF) analysis of the melting point for antibody BDG38.138 are shown in **Figures 27A and 27B**.

[507] SEC was performed using BioResolve SEC mAb Column, $200\text{\AA}$, $2.5\text{ }\mu\text{m}$, 4.6×300 mm at 0.4ml/min in PBS as a mobile phase and analyzed at 280nm . BDG38.138 shows a predominant monodisperse peak with 0.8% aggregates. DSF analysis was performed using nanoDSF at a 1°C/min from $20-95^\circ\text{C}$. NanoDSF is monitoring the thermal unfolding of BDG38.138 according to the intrinsic fluorescence change at 350 and 330 nm. The top half of the graph shows the fluorescence ratio of $350\text{nm}/330\text{nm}$ as a function of temperature and the bottom half shows the first derivative as a function of temperature. BDG38.138 was analyzed at 0.5 mg/ml in PBS showed to have a T-onset of 62.8°C and T_m of 66°C suggesting a relatively stable fold.

[508] The binding affinities of antibody BDG38.138 to human/cyno IL-13 and human/cyno TSLP are shown in **Figures 28A-28D**. (Surface Plasmon Resonance (SPR) analysis of BDG38.138 binding to human IL-13 (**Figure 28A**), human TSLP (**Figure 28B**), cyno IL-13 (**Figure 28C**) and cyno TSLP (**Figure 28D**) using BiacoreT200. CM5 chip was coated with human antibody capture kit to obtain 3000-5000 RU and the antibody, served as the capture, was injected at a flow rate of 10 ul/ml to obtain 250-350 RU. Human/cyno TSLP and h/cyno IL-13 served as analytes in a concentration range of 30-0.153 nM. Contact time was 300 sec and dissociation time was 600 sec (for cyno IL-13 dissociation time was 200 sec) at a flow rate of 30 ul/min. BDG38.138 shows a high affinity to all cytokines with K_D values for hIL-13: $1.34E-11$ M, for hTSLP: $2.85E-12$ M, for cynoIL-13: $8.49E-10$ M and for cyno TSLP: $1.35E-12$ M.

[509] *Summary:*

[510] The presented data demonstrate that BDG dual antibodies are highly monomeric and stable molecules (Table 11). Those antibodies can bind human IL-13, cynomolgus IL-13, human TSLP and cynomolgus TSLP with picomolar affinities (**Figures 16A-16F**). Unlike different bi-specific antibodies, each BDG dual antibody mentioned in the patent is a standard IgG1 (LALA,

PG) that can bind the above-mentioned antigens.

[511] Since some of the antibody's binding was above the limit of detection IC50 competition ELISA against Tezepelumab was done (Table 13, **Figures 14A-14C**). This data supports the SPR analysis and demonstrate that BDG dual antibodies can compete with Tezepelumab over binding to TSLP.

[512] BDG antibodies demonstrated functional inhibition of IL-13 signaling in HEK reporter cells, by blocking signal transduction of IL-13 through STAT6 (Table 15, **Figure 22**). Similarly, BDG antibodies inhibited TSLP mediated STAT5 phosphorylation in MUTZ cells, expressing the natural TSLP-receptor heterocomplex comprising TSLP-R and IL-7R α .

[513] To test the effect of a dual antibody, hP BMC was treated with both hTSLP and hIL-13 for 48 hours in the presence of rising concentration of BDG antibodies or benchmark antibodies (Table 12, **Figures 24A and 24B**). In this setup TSLP treatment led to increase in TARC level and IL-13 treatment led to increase in CD23 levels. While BDG dual antibodies inhibited CD23 expression similarly to the anti-IL-13 benchmark (Tralokinumab), anti-TSLP benchmark had only minor effect. Similarly, while BDG dual antibodies inhibited TARC secretion similarly to Tezepelumab, anti-IL-13 benchmark had only minor effect on the level of TARC. This demonstrates that BDG dual antibodies inhibit as a single IgG1 molecule having both functions of TSLP and IL-13.

CLAIMS

What is claimed is:

1. An isolated dual binding antibody comprising three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein
 - (i) the HCDR1 comprises the amino acid sequence of SEQ ID NO:349 or 355, or the amino acid sequence of SEQ ID NO:149 or SEQ ID NO:136;
 - (ii) the HCDR2 comprises the amino acid sequence of one of SEQ ID NOs:350, 352, 354 and 356, or the amino acid sequence of SEQ ID NO:150 or the sequence set forth as: I HX1 Y D G S N K (SEQ ID NO:142), wherein HX1 is any amino acid;
 - (iii) the HCDR3 comprises the amino acid sequence of one of SEQ ID NOs:351, 353, 357, and 358, or the amino acid sequence of SEQ ID NO:151 or the sequence set forth as: A R HX2 HX3 HX4 HX5 HX6 HX7 HX8 HX9 HX10 HX11 F D HX12 (SEQ ID NO:143), wherein HX2, HX3, HX4, HX5, HX6, HX7, HX8, HX9, HX10, HX11, and HX12 are any amino acid;
 - (iv) the LCDR1 comprises the amino acid sequence of one of SEQ ID NOs:359, 362, 364, 366, 369, and 375, or the amino acid sequence of SEQ ID NO:152 or the sequence set forth as LX1, LX2, G S K LX3 V (SEQ ID NO:144), wherein LX1, LX2, and LX3 are any amino acid;
 - (v) the LCDR2 comprises the amino acid sequence of SEQ ID NO:360 or 367, or the amino acid sequence of SEQ ID NO:153 or the sequence set forth as D D LX4 (SEQ ID NO:145), wherein LX4 is any amino acid; and
 - (vi) the LCDR3 comprises the amino acid sequence of one of SEQ ID NOs:361, 363, 365, 368, 370-374, 376-407, or the amino acid sequence of SEQ ID NO:154 or the sequence set forth as Q V W D LX5 LX6 S D LX7 V V (SEQ ID NO:146), wherein LX5, LX6, and LX7 are any amino acid.

2. The isolated dual binding antibody of claim 1, wherein

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences of SEQ ID NOs:359, 360 and 361 respectively; or

the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise

the amino acid sequences of SEQ ID NOs:364, 360 and 371 respectively; or
 the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID
 NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise
 the amino acid sequences of SEQ ID NOs:362, 360 and 384 respectively; or
 the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences of SEQ ID
 NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise
 the amino acid sequences of SEQ ID NOs:364, 360 and 384 respectively.

3. The isolated dual binding antibody of claim 1, wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.
4. The isolated dual binding antibody of claim 1,
 wherein HX1 is W or S; HX2 is A or S; HX3 is P; HX4 is Q; HX5 is W; HX6 is E, Q, M, L, or V; HX7 is L, W, or Y; HX8 is V or T; HX9 is H, A, or S; HX10 is E; HX11 is A; HX12 is I, L, or M;
 wherein LX1 is N, L, or I; LX2 is L or I; LX3 is S or L; LX4 is S or G; LX5 is S or T; LX6 is S or G; LX7 is H or G.
5. The isolated dual binding antibody of claim 4, wherein HX1 is W, HX2 is A or S, HX6 is E or M, HX7 is L or W, HX8 is V or T, HX9 is H or A, HX12 is I or L, LX1 is L, LX2 is I, LX3 is L, LX4 is S or G, LX5 is S, LX6 is S, LX7 is H or G.
6. The isolated dual binding antibody of claim 5, wherein
 - a. HX1 is W, HX2 is A, HX6 is E, HX7 is L, HX8 is T, HX9 is A, HX12 is I, LX4 is S, and LX7 is G; or
 - b. HX1 is W, HX2 is A, HX6 is M, HX7 is L, HX8 is V, HX9 is A, HX12 is L, LX4 is S, and LX7 is H; or
 - c. HX1 is W, HX2 is S, HX6 is E, HX7 is W, HX8 is V, HX9 is H, HX12 is L, LX4 is G, and LX7 is G.
7. The isolated dual binding antibody of claim 1, wherein the antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), said VH and VL

comprise the amino acid sequences of SEQ ID Nos:209 and 210, SEQ ID Nos:219 and 220, SEQ ID Nos:249 and 250, SEQ ID Nos:337 and 338, SEQ ID Nos:155 and 156, SEQ ID Nos:157 and 158, SEQ ID Nos:4 and 3, SEQ ID Nos:6 and 5, SEQ ID Nos:8 and 7, SEQ ID Nos:10 and 9, SEQ ID Nos:12 and 11, SEQ ID Nos:14 and 13, SEQ ID Nos:16 and 15, SEQ ID Nos:18 and 17, SEQ ID Nos:20 and 19, SEQ ID Nos:22 and 21, SEQ ID Nos:24 and 23, SEQ ID Nos:26 and 25, SEQ ID Nos:28 and 27, SEQ ID Nos:30 and 29, SEQ ID Nos:32 and 31, SEQ ID Nos:34 and 33, SEQ ID Nos:36 and 35, SEQ ID Nos:38 and 37, SEQ ID Nos:40 and 39, SEQ ID Nos:42 and 41, SEQ ID Nos:44 and 43, SEQ ID Nos:46 and 45, SEQ ID Nos:48 and 47, SEQ ID Nos:50 and 49, SEQ ID Nos:52 and 51, or SEQ ID Nos:54 and 53.

8. The isolated dual binding antibody of claim 1, wherein the antibody comprises a heavy chain variable domain (VH) and a light chain variable domain (VL), said VH and VL comprise the amino acid sequences as shown in Table 10 or Table 1.
9. The isolated dual binding antibody of claim 1, comprising a heavy chain variable domain (VH) and a light chain variable domain (VL), wherein
 - (c) said VH domain comprises the amino acid sequence set forth in SEQ ID NO: 1 with amino acid variants at two or more of positions 52, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, or 111, or any combination thereof (IMGT positions: 57, 107, 108, 109, 110, 111, 111A, 112A, 112, 113, 114, or 117, or a combination thereof); and
 - (d) said VL domain comprises the amino acid sequence set forth in SEQ ID NO: 2 with amino acid variants at two or more of positions 26, 27, 31, 51, 56, 77, 92, 93, or 96, or any combination thereof (IMGT positions: 27, 28, 38, 65, 70, 94, 109, 110, or 115, or a combination thereof).
10. The isolated dual binding antibody of claim 1, wherein said antibody comprises an IgG, an Fv, an scFv, an Fab, an F(ab')₂, a minibody, a diabody, or a triabody.
11. The isolated dual binding antibody of claim 10, wherein said IgG is IgG1, IgG2, IgG3, or IgG4.
12. The isolated dual binding antibody of claim 10, wherein said IgG comprises a mutated

IgG, said mutant IgG is unable to bind to antibody-dependent cellular cytotoxicity components.

13. A composition comprising the isolated dual binding antibody of claim 1 and a pharmaceutically acceptable carrier.
14. An isolated polynucleotide construct encoding the isolated dual binding antibody of claim 1.
15. An expression vector comprising the polynucleotide construct of claim 14.
16. A host cell comprising the expression vector of claim 15.
17. A method of treating a subject suffering from a disease or condition, said method comprises administering to said subject a composition comprising the isolated dual binding antibody of claim 1.
18. The method of claim 17, wherein said disease or condition is an allergic or respiratory condition, an inflammatory or autoimmune condition, or tumors or cancers.
19. The method of claim 17, wherein said disease or condition is asthma, allergic asthma, nonallergic asthma, severe asthma, mild asthma, chronic obstructive pulmonary disease (COPD), a condition involving airway inflammation, cystic fibrosis, allergic lung disease, airway hyperresponsiveness, goblet cell metaplasia, mucus hypersecretion, airway remodeling, pulmonary fibrosis, atopic dermatitis, urticaria, eczema, allergic enterogastritis, allergic rhinitis, inflammatory bowel diseases, liver cirrhosis or fibrosis, or a combination thereof.
20. The method of claim 17, wherein said dual binding antibody comprises three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:359, 360 and 361 respectively; or

- the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 356 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 371 respectively; or
- the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:362, 360 and 384 respectively; or
- the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequence of SEQ ID NOs:349, 350 and 351 respectively, and the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequence of SEQ ID NOs:364, 360 and 384 respectively; or
- the CDRs having the sequences of SEQ ID NOs:149-154.
21. The method of claim 17, wherein said dual binding antibody comprises three complementarity determining regions (CDRs) on a heavy chain (HCDR1, HCDR2, and HCDR3) and three CDRs on a light chain (LCDR1, LCDR2, and LCDR3), wherein the HCDR1, HCDR2 and HCDR3 comprise the amino acid sequences as shown in Table 8 or Table 4, wherein the LCDR1, LCDR2 and LCDR3 comprise the amino acid sequences as shown in Table 9 or Table 5.
22. The method of claim 17, wherein said dual binding antibody comprises VH and VL having the sequences of SEQ ID Nos:209 and 210, SEQ ID Nos:219 and 220, SEQ ID Nos:249 and 250, SEQ ID Nos:337 and 338, SEQ ID NOs:155 and 156, SEQ ID NOs:157 and 158 .
23. The method of claim 17, wherein said dual binding antibody comprises VH and VL having the sequences as shown in Table 10 or Table 1.