

MULTISPECIFIC ANTIBODIES AGAINST CD40 AND CD137

FIELD OF THE INVENTION

The present invention relates to multispecific antibodies binding to CD40 and CD137, and to uses of such multispecific antibodies, in particular to the use for
5 treatment of cancer.

BACKGROUND OF THE INVENTION

CD40 is a member of the tumor necrosis factor (TNF) receptor (TNFR) family and is known as a co-stimulatory protein found on a diversity of cell types. CD40 is constitutively expressed by antigen-presenting cells (APCs), including dendritic cells
10 (DCs), B cells and macrophages. It can also be expressed by endothelial cells, platelets, smooth muscle cells, fibroblasts and epithelial cells. Consistent with its widespread expression on normal cells, CD40 is also expressed on a wide range of tumor cells.

The presentation of peptide antigens in the context of MHC class II molecules to antigen-specific CD4⁺ T cells, together with co-stimulatory signals (from CD80 and/or
15 CD86), results in the activation of CD4⁺ T cells and the up-regulation of the DC licensing factors CD40 ligand (CD40L) and lymphotoxin- α 1 β 2 (LT α 1 β 2). Expression of CD40L and LT α 1 β 2 on activated antigen-specific CD4⁺ T cells induces signaling through CD40 and the LT β receptor (LT β R), and this licenses DCs to induce CD8⁺ T-cell responses. CD40 signaling results in the production of interleukin-12 (IL-12) and the up-regulation of
20 CD70, CD86, 4-1BB ligand (4-1BBL), OX40 ligand (OX40L) and GITR ligand (GITRL), whereas LT β R signaling leads to the production of type I interferons (IFNs). The signaling system that controls the activity of nuclear factor kappaB (NF- κ B) is responsive to virtually all TNFR superfamily members. Pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs) also contribute to these
25 events. Priming of CD8⁺ T cells by MHC class I-restricted peptides results in the up-regulation of CD27, 4-1BB, OX40 and glucocorticoid-induced TNFR-related protein (GITR). Stimulation of these receptors on CD8⁺ T cells by their cognate TNF superfamily ligands, in combination with IL-12 and type I IFNs, results in robust CD8⁺ T cell activation, proliferation and effector function, as well as the formation and maintenance
30 of CD8⁺ T cell memory. CD40 antibodies can exert different actions, CD40-expressing tumor cell kill by induction of antibody-dependent cell-mediated cytotoxicity (ADCC), complement-dependent cytotoxicity (CDC) or antibody-dependent cell-mediated

phagocytosis (ADCP), induction of cell signaling to induce direct apoptosis or growth arrest, but also, independent of CD40 expression on the tumor cells, through licensing of APCs to stimulate an anti-cancer immune response. Antibodies binding to CD40 can trigger CD40 on APCs to prime effector cytotoxic T lymphocytes (CTLs) and induce release of IL-2 by these cells, and indirectly activate NK cells. Antibodies stimulating CD40 have been disclosed in the prior art, and include CP-870,893, a human IgG2 antibody (WO 2003/040170); dacetuzumab, a humanized IgG1 antibody (WO 2000/075348) and Chi Lob 7/4, a chimeric IgG1 antibody (US 2009/0074711). Furthermore, an antagonistic CD40 antibody has been disclosed, lucatumumab, a human IgG1 antibody (WO 2002/028481).

CD137 (4-1BB) is also a member of the TNFR family. CD137 is a co-stimulatory molecule on CD8⁺ and CD4⁺ T cells, regulatory T cells (Tregs), Natural Killer T cells (NK(T) cells), B cells and neutrophils. On T cells, CD137 is not constitutively expressed, but induced upon T-cell receptor (TCR) activation (for example, on tumor infiltrating lymphocytes (TILs) (Gros et al., J. Clin Invest 2014;124(5):2246-59)). Stimulation via its natural ligand 4-1BBL or agonist antibodies leads to signaling using TRAF-2 and TRAF-1 as adaptors. Early signaling by CD137 involves K-63 poly-ubiquitination reactions that ultimately result in activation of the nuclear factor(NF)-kB and mitogen-activated protein(MAP)-kinase pathways. Signaling leads to increased T cell co-stimulation, proliferation, cytokine production, maturation and prolonged CD8⁺ T-cell survival. Agonistic antibodies against CD137 have been shown to promote anti-tumor control by T cells in various pre-clinical models (Murillo et al., Clin Cancer Res 2008;14(21):6895-906). Antibodies stimulating CD137 can induce survival and proliferation of T cells, thereby enhancing the anti-tumor immune response. Antibodies stimulating CD137 have been disclosed in the prior art, and include urelumab, a human IgG4 antibody (AU2004279877) and utomilumab, a human IgG2 antibody (Fisher et al. 2012 Cancer Immunol. Immunother. 61: 1721-1733).

Westwood JA, et al., Leukemia Research 38 (2014), 948-954 discloses "Combination anti-CD137 and anti-CD40 antibody therapy in murine myc-driven hematological cancers".

US20090074711 discloses "Human therapies using chimeric agonistic anti-human CD40 antibody".

However, despite these and other advances in the art, there is a need for multispecific antibodies that can bind both CD40 and CD137, simultaneously binding to CD40-expressing APCs and CD137-expressing T cells, thereby bringing these cell types in close contact. This, in turn, can lead to activation of both cell types and efficient induction of anti-tumor immunity.

SUMMARY OF THE INVENTION

The present inventors have identified multispecific antibodies that can bind both CD40 and CD137 and elicit T cell and APC activation.

So, in one aspect, the invention relates to a multispecific antibody comprising

- 5 (i) a first antigen-binding region binding to human CD40, and (ii) a second antigen-binding region binding to human CD137.

In some embodiments, the invention relates to such a multispecific antibody wherein the first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 which comprise specific amino acid sequences, optionally with
10 mutations, or the amino acid sequences of an antibody which competes with or has the specificity of an antibody comprising such specific amino acid sequences. In specific embodiments, the first antigen-binding region comprises heavy and light chain variable sequences comprising the CDR1, CDR2 and CDR3 of anti-CD40 antibody 001, or competes with or has the specificity of such an antibody.

15 In some embodiments, the invention relates to such a multispecific antibody wherein the second antigen-binding region comprises heavy and light chain variable sequences wherein the CDR1, CDR2 and CDR3 comprise specific amino acid sequences or provide specific amino acid sequences, optionally with mutations, or comprise the amino acid sequences of an antibody which competes with or has the specificity of an
20 antibody comprising such specific amino acid sequences. In specific embodiments, the second antigen-binding region comprises heavy and light chain variable sequences comprising the CDR1, CDR2 and CDR3 of anti-CD137 antibody 001, 002, 003, 004, 005, 006, 007, 008, 009, 010, 011 or 012, or competes with or has the specificity of any such antibody.

25 These and other aspects and embodiments, including nucleic acids encoding the amino acid sequences of such multispecific antibodies; expression vectors comprising such nucleic acids; host cells comprising such nucleic acids or expression vectors; compositions comprising such multispecific antibodies; such compositions for use in treating cancer or other diseases; methods for producing such multispecific antibodies;
30 and diagnostic methods and kits based on such multispecific antibodies, are described in further detail below.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Sequence alignments for human, African elephant and wild boar CD137. Amino acids in African elephant or wild boar CD137 that differ from those in the
35 human sequence are highlighted in black.

Figure 2: CD137 shuffle constructs, containing African elephant (shuffle 5) or wild boar (shuffle 1-4, 6) CD137 domains.

Figure 3: Expression of CD137 shuffle constructs on HEK293-T17 cells. HEK293-T17 cells were transfected with the CD137 shuffle constructs. Cell surface expression of the constructs was measured by flow cytometry, using a polyclonal anti-CD137 antibody that recognizes human, wild boar and African elephant CD137.

Figure 4: Binding of CD137 antibody clones to CD137 shuffle constructs expressed on HEK293-T17 cells. HEK293-T17 cells were transfected with the CD137 shuffle constructs, and with human CD137 (hCD137 wt), African elephant or wild boar CD137, as indicated. Binding of the different CD137 antibody clones to these constructs expressed on HEK293-T17 cells was measured by flow cytometry. Staining with polyclonal anti-CD137 antibody is shown as a control.

Figure 5: Matrix-like mixing grids used for automated bispecific antibody discovery. Parental antibodies can be plated as indicated. Parental antibodies can then be combined to obtain bispecific antibodies using a simple matrix-like mixing grid. Controlled Fab-arm exchange can then be performed to obtain bispecific antibodies.

Figure 6: Expression of CD40 and CD137 on the cell surface of stably transduced HEK293-NFK-gfp-luc and K562 cells. NF- κ B/293/GFP-Luc (A) and K562 (B) cells were stably transduced with CD40 or CD137. Surface expression of CD40 (left panels) and CD137 (right panels) was determined by flow cytometry (white curves: control without antibody; grey curves: antibody staining).

Figure 7: Analysis of bispecific antibodies simultaneously targeting CD40 and CD137 (CD40xCD137). Bispecific antibodies targeting CD40 and CD137 (CD40-FEALxCD137-FEAR) were tested in the reporter assay in duplicate (A-L: CD40-001xCD137-001 until CD40-001xCD137-012). Activation of CD137 was measured by luciferase activity (relative luminescence units, RLU) of NF- κ B/293/GFP-Luc transduced with CD137 (HEK293_NFK_CD137_gfp_luc) upon incubation with the indicated bispecific antibodies and K562 cells transduced with CD40 (K562_CD40) for trans-activation or wildtype K562 cells (K562_wt) as a control. Activation of CD40 was measured by luciferase activity (RLU) of NF- κ B/293/GFP-Luc transfected with CD40 (HEK293_NFK_CD40_gfp_luc) upon incubation with the indicated bispecific antibodies and K562 transduced with CD137 (K562_CD137) for trans-activation or wildtype K562 cells (K562_wt) as a control. The two monospecific, monovalent antibodies with one irrelevant arm (b12-FEALxCD137-FEAR, b12-FEALxCD40-FEAR) were used as control for the bispecific CD40xCD137 antibodies.

Figure 8: Induction of CD8⁺ T-cell proliferation by CD40xCD137 bispecific antibodies in a non-antigen-specific T cell assay. CFSE-labeled PBMCs were incubated with CD40xCD137 bispecific antibodies or monospecific, monovalent control antibodies,

for four days. Proliferation of CD8⁺ cells was measured by flow cytometry. Data shown are CFSE plots showing CD8⁺ T-cell proliferation induced by the indicated bispecific and control antibodies at 0.02 µg/mL (A), percentages of divided cells and proliferation indices for CD40-001-FEALxCD137-005-FEAR (B), CD40-001-FEALxCD137-009-FEAR (C), CD40-001-FEALxCD137-003-FEAR (D) and CD40-001-FEALxCD137-011-FEAR (E), as calculated using FlowJo software.

Figure 9: Enhancement of CD8⁺ T-cell proliferation by CD40xCD137 bispecific antibodies in an antigen-specific T-cell assay. T cells transfected with a claudin-6-specific TCR and labeled with CFSE were incubated with claudin-6 IVT-RNA-electroporated immature DCs in the presence or absence of CD40xCD137 bispecific antibodies or control antibodies for five days. CD8⁺ T-cell proliferation was measured by flow cytometry. Data shown are CFSE plots showing CD8⁺ T-cell proliferation induced by the indicated bispecific antibodies and controls at 0.02 µg/mL (A), percentages divided cells and proliferation indices for the indicated bispecific antibodies (B) and for CD40-001-FEALxCD137-005-FEAR and control antibodies (C) and CD40-001-FEALxCD137-009-FEAR and control antibodies (D) at the indicated concentrations, as calculated using FlowJo software. Proliferation index curves for the indicated bispecific antibodies at serial dilutions ranging from 6.4×10^{-5} to 5 µg/mL are also shown (E). Curves were analyzed by non-linear regression (sigmoidal dose-response with variable slope) using GraphPad Prism 5 software (GraphPad Software, San Diego, CA, USA). The EC₅₀ values for induction of T-cell proliferation for CD40-001-FEALxCD137-005-FEAR and CD40-001-FEALxCD137-009-FEAR were 0.005 and 0.030 µg/mL, respectively.

Figure 10: Induction of CD8⁺ T-cell proliferation by the humanized CD40xCD137 bispecific antibody in a non-antigen-specific T cell assay. CFSE-labeled PBMCs were incubated with humanized CD40xCD137 bispecific antibody, the parental bispecific antibody or IgG1 control antibody for four days. Proliferation of CD8⁺ T cells was measured by flow cytometry. Data shown are percentages of divided cells and proliferation index, as calculated by FlowJo software. (n.d. = not determined)

Figure 11: Enhancement of CD8⁺ T-cell proliferation by a humanized CD40xCD137 bispecific antibody in an antigen-specific T-cell assay. T cells transfected with a claudin-6-specific TCR and labeled with CFSE were incubated with claudin-6 IVT-RNA-electroporated immature DCs in the presence or absence of a humanized CD40xCD137 bispecific antibody (BisG1-CD40-001-H6LC1-FEALxCD137-009-HC7LC2-FEAR), the parental bispecific antibody or an IgG1 control antibody for four days. CD8⁺ T-cell proliferation was measured by flow cytometry. Data shown are percentages divided cells and proliferation indices for the indicated antibodies, as calculated using FlowJo software. (n.d. = not determined)

Figure 12: Ex vivo expansion of TILs from a human melanoma tissue resection by a CD40xCD137 bispecific antibody. Tumor pieces from the resected tissue were cultured with 100 U/mL IL-2 and the indicated concentration of a CD40xCD137 bispecific antibody (BisG1-CD40-001-FEALxCD137-009-FEAR). After 14 days of culture, cells were harvested and analyzed by flow cytometry. Relative viable TIL count per sample (normalized to 1,000 measured counting beads) is shown. Each data point refers to a single well, representing the expansion of TILs out of two tumor pieces analyzed in one FACS tube. The line indicates the mean of five measured samples.

Figure 13: Ex vivo expansion of TILs from a human non-small cell lung cancer (NSCLC) tissue resection by a CD40xCD137 bispecific antibody. Tumor pieces from resected NSCLC tissue were cultured with 10 U/mL IL-2 and the indicated concentration of CD40xCD137 bispecific antibody (BisG1-CD40-001-FEALxCD137-009-FEAR). After 10 days of culture, cells were harvested and analyzed by flow cytometry. Relative viable TIL count per sample (normalized to 1,000 measured counting beads) is shown. Each data point refers to a single well, representing the expansion of TILs out of two tumor pieces analyzed in one FACS tube. The line indicates the mean of five measured samples.

Table 1 - Sequences

Sequence name	Type of sequence	Sequence	Sequence identifier
CD40-001 antibody (mouse)	VH CDR1	GYTFTEYI	SEQ ID NO:1
	VH CDR2	IIPNNGGT	SEQ ID NO:2
	VH CDR3	TRREVYGRNYYALDY	SEQ ID NO:3
	VL CDR1	QGINNY	SEQ ID NO:4
	VL CDR2	YTS	
	VL CDR3	QQYSNLPYT	SEQ ID NO:5
	VH	EVQLQQSGPDLVKPGASVKISCKTS GYTFTEYIMHWVKQSHGKSLEWIG GIIPNNGGTSYNQKFKDKATMTVDK SSSTGYMELRSLTSEDSAVYYCTRRE VYGRNYYALDYWGQGLTVTVSS	SEQ ID NO:6
	VL	DIQMTQTTSSLSASLGDRVITITCSA SQGINNYLNWYQQKPDGTVKLLIYY	SEQ ID NO:7

		TSSLHSGVPSRFSGSGSGTDYSLTIS NLEPEDIATYYCQQYSNLPYTFGGGT KLEIK	
CD137 antibody clone 001 (rabbit)	VH CDR1	GFSLSYYA	SEQ ID NO:8
	VH CDR2	IWNNGAT	SEQ ID NO:9
	VH CDR3	ARSANDAWSTYSDL	SEQ ID NO:10
	VL CDR1	QTITNY	SEQ ID NO:11
	VL CDR2	KAS	
	VL CDR3	QNYYYGSSSGYGFV	SEQ ID NO:12
	VH	QSVEESGGRLVTPGTPLTLTCTVSGFS LSSYAVSWVRQAPGKGLEWIGVIWN NGATHYATWAKGRFTISKASTTVDLK VTSPPTEDTATYFCARSANDAWSTYS DLWGQGTLTVSS	SEQ ID NO:13
	VL	DIVMTQTPASVEAAVGGTVTIKCQASQ TITNYLSWYQQKPGQPPKLLIYKASTLT SGVSSRFKSGSGTEFTLTISDLESDDA ATYYCQNYYYGSSSGYGFVFGGGTEVVV K	SEQ ID NO:14
CD137 antibody clone 002 (rabbit)	VH CDR1	GFSLTYYA	SEQ ID NO:15
	VH CDR2	IYDNGAT	SEQ ID NO:16
	VH CDR3	ARSANNAWSTYSNL	SEQ ID NO:17
	VL CDR1	EDISSY	SEQ ID NO:18
	VL CDR2	KAS	
	VL CDR3	QSYYSGISISGYGFV	SEQ ID NO:19
	VH	QSVEESGGRLVTPGTPLTLTCTVSGFS LTYAVTWVRQPPGKGLEWIGVIYDN GATAFATWAKGRFTMSKNSTTVALKV	SEQ ID NO:20

		TSPTTEDTATYFCARSANNAWSTYSN LWGQGLTVTVSS	
	VL	DIVMTQTPSSVSAAVGGTVTINCQAS EDISSYLSWYQQKLGQPPKLLIYKAST LESGVPSRFGSGSGTEYTLTISDLES DDAATYYCQSYSGSISGYGFVFGGGT GVVVK	SEQ ID NO:21
CD137 antibody clone 003 (rabbit)	VH CDR1	GFTISSYH	SEQ ID NO:22
	VH CDR2	IYGGTATT	SEQ ID NO:23
	VH CDR3	ARARYSGGSYANYVFNL	SEQ ID NO:24
	VL CDR1	QSISSY	SEQ ID NO:25
	VL CDR2	RTS	
	VL CDR3	QGYDWSSSNRYDNT	SEQ ID NO:26
	VH	QSVEESGGRLVTPGTPLTLTCTAS GFTISSYHMIWVRQAPGEGLEWI GDIYGGTATTDYASWAKGRFTIS KTSTTVDLKMTSLTTEDTATYFCA RARYSGGSYANYVFNLWGQGLTV TVSS	SEQ ID NO:27
	VL	DIVMTQTPASVEAAVGGTVTIKCQ ASQSISSYLAWYQQKPGQPPKLLI YRTSTLESGVPSRFGSGSGTEFTL TISDLESADAATYYCQGYDWSSSN RYDNTFGGGTEVVVK	SEQ ID NO:28
CD137 antibody clone 004 (rabbit)	VH CDR1	GFSLSRDYD	SEQ ID NO:29
	VH CDR2	ISSSGGT	SEQ ID NO:30
	VH CDR3	AREGDYWDFNL	SEQ ID NO:31
	VL CDR1	QISISNL	SEQ ID NO:32

	VL CDR2	GAS	
	VL CDR3	AGGFPGLDTVAA	SEQ ID NO:33
	VH	QSLEESGGRLVTPGTPLTLTCTASGF SLSRYDMSWVRQAPGKGLEYGIVIS SSGGTNYANWAKGRFTISKSTTV LKITSPTTEDTATYFCAREGDYWDFN LWGPGTLTVSS	SEQ ID NO:34
	VL	AQVLTQTPSSVSAAVGGTVTINCQA SQSISNLLAWYQQKPGQPPKLLIYG ASTLASGVPSRFSGSGSGTEFTLTIS DLESDDAATYYCAGGFPGLDTVAAF GGGTEAVVT	SEQ ID NO:35
CD137 antibody clone 005 (rabbit)	VH CDR1	GFTISDFH	SEQ ID NO:36
	VH CDR2	IITSASTT	SEQ ID NO:37
	VH CDR3	ARSTYTDTS GYYFDF	SEQ ID NO:38
	VL CDR1	QSIYNGNR	SEQ ID NO:39
	VL CDR2	SAS	
	VL CDR3	LGSYDCDSADCFA	SEQ ID NO:40
	VH	QSVEESGGRLVTPGTPLTLTCTASG FTISDFHVTWVRQAPGKLEWIGTI ITSASTTAYATWARGRFTISKSTTV NLKIVSPTTEDTATYFCARSTYDTS GYYFDFWGQGTLTVSS	SEQ ID NO:41
	VL	AQVLTQTASPVSAAVGGTVIINCQSS QSIYNGNRLSWYQQKPGQPPKLLIYS ASTLASGVSSRFKSGSGTQFTLAISD VQSDDAATYYCLGSYDCDSADCFAFG GGTEVVVE	SEQ ID NO:42
CD137 antibody clone 006 (rabbit)	VH CDR1	GFSLSYYA	SEQ ID NO:43

	VH CDR2	ISTSGIT	SEQ ID NO:44
	VH CDR3	ARLNGFDDYVRYFDF	SEQ ID NO:45
	VL CDR1	ESIASN	SEQ ID NO:46
	VL CDR2	AAS	
	VL CDR3	QSAFYVSSSDNA	SEQ ID NO:47
	VH	QSVEESGGRLVTPGTPLTLTCTVSGFS LSSYAMSWVRQAPGKGLEWIGIISTS GITYYASWAKGRFTISKSTSTMVDLKIT SPTTEDTATYFCARLNGFDDYVRYFDF WGLGTLTVSS	SEQ ID NO:48
	VL	AIELTQTPSSVSAAVGGTVTINCQASE SIASNLAWYQQKPGQPPKLLIYAASYL ASGVPSRFKSGSGSTEYTLTISGVQSA DAATYYCQSAFYVSSSDNAFGGGTEVV VK	SEQ ID NO:49
CD137 antibody clone 007 (rabbit)	VH CDR1	GFSLSYD	SEQ ID NO:50
	VH CDR2	IGSDGSA	SEQ ID NO:51
	VH CDR3	ARDWNDYWAHDL	SEQ ID NO:52
	VL CDR1	QIVTSKSA	SEQ ID NO:53
	VL CDR2	KAS	
	VL CDR3	AGGYNSGDLNP	SEQ ID NO:54
	VH	QSLEESGGRLVTPGTPLTLTCTASGFS LSSYDVSWVRQAPGKLEYIGFIGSD GSAHYATWVKGRFTISKSTTVDLKIT SPTTEDTATYFCARDWNDYWAHDLW GPGTLTVSS	SEQ ID NO:55
	VL	AQVLTQTTSPVSAAVGGTVTINCQAS QIVTSKSALSWYQQKPGQPPRLIYK ASTLASGVPSRFSGSGGTQFTLTIS DLESDDAATYYCAGGYNSGDLNPF	SEQ ID NO:56

		GGGTEVVVK	
CD137 antibody clone 008 (rabbit)	VH CDR1	GFSLSSTYD	SEQ ID NO:57
	VH CDR2	ISSSGNT	SEQ ID NO:58
	VH CDR3	AREGDYWDFNL	SEQ ID NO:59
	VL CDR1	QISISNL	SEQ ID NO:60
	VL CDR2	RAS	
	VL CDR3	AGGFPGLDTGAT	SEQ ID NO:61
	VH	QSLEESGGRLVTPGTPLTLTCTASGFSL SSYDMSWVRQAPGKLEYIGYISSSG NTYYASWAKSRFTISKSTTVDLKITS PTTEDTATYFCAREGDYWDFNLWGPG TLVTVSS	SEQ ID NO:62
	VL	AQVLTQTPSSVSAAVGGTVTINCQAS QISNLLAWYQQKPGQRPRLIYRAS TLASGVPSRFKSGSGTEFTLTISDLE SEDAATYYCAGGFPGLDTGATFGGGT EAVVT	SEQ ID NO:63
CD137 antibody clone 009 (rabbit)	VH CDR1	GFSLNNDYW	SEQ ID NO:64
	VH CDR2	IDVGGSL	SEQ ID NO:65
	VH CDR3	ARGGLTYGFDL	SEQ ID NO:66
	VL CDR1	EDISSY	SEQ ID NO:67
	VL CDR2	GAS	
	VL CDR3	HYYATISGLGVA	SEQ ID NO:68
	VH	QSLEESGGRLVTPGTPLTLTCTVSG FSLNDYWMSWVRQAPGKGLEWIG YIDVGGSLYYASWAKGRFTISRTST TVDLKMTSLTTEDTATYFCARGGLT YGFDLWGPGTLVTVSS	SEQ ID NO:69

	VL	DIVMTQTPASVSEPVGGTVTINCQA SEDISSYLAWYQQKPGQRPKRLIYG ASDLASGVPSRFSASGSGTEYALTIS DLESADAATYYCHYYATISGLGVAFG GGTEVVVK	SEQ ID NO:70
CD137 antibody clone 010 (rabbit)	VH CDR1	GFSLSTYA	SEQ ID NO:71
	VH CDR2	VYDNGYI	SEQ ID NO:72
	VH CDR3	ARSADGSWSTYFNL	SEQ ID NO:73
	VL CDR1	ESISNY	SEQ ID NO:74
	VL CDR2	KAS	
	VL CDR3	QTNYYCCSSSDNGFA	SEQ ID NO:75
	VH	QSVEESGGRLVTPGTPLTLTCTVSGFSL STYAMIWVRQAPGKGLEWIGVVYDNG YISHATWVKGRFTISKSTTTVGLEITSP TTEDTATYFCARSADGSWSTYFNLWG QGTLVTVSS	SEQ ID NO:76
	VL	DIVMTQTPASVEAAVGGTVTIKCQAS ESISNYLAWYQQKPGQPPKLLIYKAS TLASGVSSRFKSGSGTEFTLTISDL ESADAATYYCQTNYYCCSSSDNGFAF GGGTEVVVK	SEQ ID NO:77
CD137 antibody clone 011 (rabbit)	VH CDR1	GIDLSSYH	SEQ ID NO:78
	VH CDR2	IAYGGNT	SEQ ID NO:79
	VH CDR3	ARGYSEDSYWGL	SEQ ID NO:80
	VL CDR1	QNIENY	SEQ ID NO:81
	VL CDR2	DTS	
	VL CDR3	QQDYGIIFVDNI	SEQ ID NO:82
	VH	QSLEESGGRLVTPGTPLTLTCTVSGIDL	SEQ ID NO:83

		SSYHMCWVRQAPGKGLEYIGYIAYGG NTYYANWAKGRFTISKSTTVDLRITS PTTEDTATYFCARGYSEDSYWGLWGP GTLVTVSS	
	VL	AYDMTQTPASVEAAVGGTVTIKCQAS QNIENYLAWYQQKPGQPPKLLIYDTS KLTSGVPSRFSGSGSGTDFLTISGVQ SDDAATYYCQQDYGIIFVDNIFGGGTE VVK	SEQ ID NO:84
CD137 antibody clone 012 (rabbit)	VH CDR1	GFSLSDY	SEQ ID NO:85
	VH CDR2	MSGSGST	SEQ ID NO:86
	VH CDR3	ARDGDYAGWGYATGAFFD	SEQ ID NO:87
	VL CDR1	QSVVGNL	SEQ ID NO:88
	VL CDR2	SAS	
	VL CDR3	TGRYNSDTDFV	SEQ ID NO:89
	VH	QSVEESGGRLVTPGTPLTCTVSGFSL SDYYMTWVRQAPGKGLEYIGIMSGSG STYYASWAKGRFTISKSTTLELKITS PTTEDTAIYFCARDGDYAGWGYATGAF DPWGPGLTVTVSS	SEQ ID NO:90
	VL	AAVLTQTPSPVSAAVGGTVTISCQASQ SVVGNLSSWFQKKPGQPPKLLIYSAS SLASGVPSRFGSGSGTQFTLTISDLES DDAATYYCTGRYNSDTDFVFGGGTEV VVK	SEQ ID NO:91
Human CD137 (TNR9_Human)		MGNSCYNIVATLLLVNFERTRSLQDPCS NCPAGTFCDNNRNQICSPCPPNSFSSAG GQRTCDICRQCKGVFRTRKECSSTNAE CDCTPGFHCLGAGCSMCEQDCKQGQELT KKGCKDCCFGTFNDQKRGICRPWTNCSL DGKSVLVNGTKERDVVCGPSPADLSPGA SSVTPPAPAREPGHSPQIISFFLALTSTALL FLLFFLTLRFSVVKRGRKKLLYIFKQPFMRP	SEQ ID NO:92

		VQTTQEEDGCSCRFPFFFFFFGGCEL	
Human CD137 shuffle 6 (amino acids 24-47 replaced by wild boar CD137)		MGNSCYNIVATLLLVNLFERTRSVDPDPCS NCSAGTFCGKNIQELCMPCPPNSFSSAG GQRTCDICRQCKGVFRTRKECSSTSNAE CDCTPGFHCLGAGCSMCEQDCKQGQELT KKGCKDCCFGTFNDQKRGICRPWTNCSL DGKSVLVNGTKERDVVCGPSPADLSPGA SSVTPPAPAREPGHSPQIISFFLALTSTALL GGCEL	SEQ ID NO:93
Human CD137 shuffle 5 (amino acids 48-88 replaced by African elephant CD137)		MGNSCYNIVATLLLVNLFERTRSLQDPCS NCPAGTFCDNRRNQICSPCLNSFSSTGG QMNCDCMRKCEGVFKTKRACSPTRDAEC ECTPGFHCLGAGCSMCEQDCKQGQELTK KGCKDCCFGTFNDQKRGICRPWTNCSLD GKSVLVNGTKERDVVCGPSPADLSPGAS SVTPPAPAREPGHSPQIISFFLALTSTALLF LLFFLTLRFSVVKRGRKKLLYIFKQPFMRP VQTTQEEDGCSCRFPFFFFFFGGCEL	SEQ ID NO:94
Human CD137 shuffle 4 (amino acids 89-114 replaced by wild boar CD137)		MGNSCYNIVATLLLVNLFERTRSLQDPCS NCPAGTFCDNRRNQICSPCPNSFSSAG GQRTCDICRQCKGVFRTRKECSSTSNAE CDCVPGFRCLGAGCAMCEEYCQOQELT QKGCKDCCFGTFNDQKRGICRPWTNCSL DGKSVLVNGTKERDVVCGPSPADLSPGA SSVTPPAPAREPGHSPQIISFFLALTSTALL FLLFFLTLRFSVVKRGRKKLLYIFKQPFMRP VQTTQEEDGCSCRFPFFFFFFGGCEL	SEQ ID NO:95
Human CD137 shuffle 3 (amino acids 115-138 replaced by wild boar CD137)		MGNSCYNIVATLLLVNLFERTRSLQDPCS NCPAGTFCDNRRNQICSPCPNSFSSAG GQRTCDICRQCKGVFRTRKECSSTSNAE CDCTPGFHCLGAGCSMCEQDCKQGQELT KEGCKDCSFGTFNDEEHGVCRPWTDCSL DGKSVLVNGTKERDVVCGPSPADLSPGA SSVTPPAPAREPGHSPQIISFFLALTSTALL FLLFFLTLRFSVVKRGRKKLLYIFKQPFMRP VQTTQEEDGCSCRFPFFFFFFGGCEL	SEQ ID NO:96
Human CD137 shuffle 2 (amino acids 139-161 replaced by wild boar CD137)		MGNSCYNIVATLLLVNLFERTRSLQDPCSNCPA GTFCDNRRNQICSPCPNSFSSAGGQRTCDICR QCKGVFRTRKECSSTSNAECDCTPGFHCLGAGC SMCEQDCKQGQELTKKGCKDCCFGTFND QKRGICRPWTNCSLAGKPVLMTNGTKARD VVCGPRPADLSPGASSVTPPAPAREPGHS PQIISFFLALTSTALLFLLFFLTLRFSVVKRGR RKLLYIFKQPFMRPVQTTQEEDGCSCRFP FFFFFFGGCEL	SEQ ID NO:97
Human CD137 shuffle 1 (amino acids 162-186)		MGNSCYNIVATLLLVNLFERTRSLQDPCSNCPA GTFCDNRRNQICSPCPNSFSSAGGQRTCDICR	SEQ ID NO:98

replaced by wild boar CD137)		QCKGVFRTRKECSSTSNAECDCTPGFHCLGAGC SMCEQDCKQGQELTKKGCKDCCFGTFNDQKR GICRPWTNCSLDGKSVLVNGTKERDVVCGPSPT DFSPGTPSTTMPVPGGEPGHTSHIISFFLALTST ALLFLLFFLTFRFSVVKRGRKKLLYIFKQPFMRPV QTTQEEDGCSCRFPEEEEGGCEL	
b12	VH CDR1	GYRFSNFV	SEQ ID NO:99
	VH CDR2	INPYNGNK	SEQ ID NO:100
	VH CDR3	ARVGPYSWDDSPQDNYYMDV	SEQ ID NO:101
	VL CDR 1	HSIRSRR	SEQ ID NO:102
	VL CDR 2	GVS	
	VL CDR 3	QVYGASSYT	SEQ ID NO:103
	VH	QVQLVQSGAEVKKPGASVKVSCQASGYR FSNFVIHWVRQAPGQRFWMGWINPYN GNKEFSAKFQDRVTFADTSANTAYMELR SLRSADTAVYYCARVGPYSWDDSPQDN YMDVWGKGTTIVSS	SEQ ID NO:104
	VL	EIVLTQSPGTLSPGERATFSCRSSHSIR SRRVAWYQHKGQAPRLVIHGVSNRASG ISDRFSGSGSGTDFLTITRVEPEDFALYY CQVYGASSYTFGQGTKLERK	SEQ ID NO:105
IgG1m(a) CH3 region		GQPREPQVYTLPPSRDELTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVL DSDGSFFLYSKLTVDKSRWQQGNVFSCS VMHEALHNHYTQKSLSLSPGK	SEQ ID NO:106
IgG1m(f) CH3 region		GQPREPQVYTLPPSREEMTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVL DSDGSFFLYSKLTVDKSRWQQGNVFSCS VMHEALHNHYTQKSLSLSPGK	SEQ ID NO:107
IgG1m(ax) CH3 region		GQPREPQVYTLPPSRDELTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVL DSDGSFFLYSKLTVDKSRWQQGNVFSCS VMHEGLHNHYTQKSLSLSPGK	SEQ ID NO:108
IgG1 heavy chain constant region – WT*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE	SEQ ID NO:109

		LLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSFFLY SKLTVDKSRWQQGNVFSCSVMHEALHN HYTQKSLSLSPGK	
IgG1 heavy chain constant region – F405L*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTWSWNSGALTSGVHTFPAVLQ SSGLYSLSVVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE LLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSFLL YSKLTVDKSRWQQGNVFSCSVMHEALH NHYTQKSLSLSPGK	SEQ ID NO:110
IgG1 heavy chain constant region – K409R*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTWSWNSGALTSGVHTFPAVLQ SSGLYSLSVVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE LLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSFFLY SRLTVDKSRWQQGNVFSCSVMHEALHN HYTQKSLSLSPGK	SEQ ID NO:111
Human IgG1 heavy chain constant sequence with FEAR*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTWSWNSGALTSGVHTFPAVLQ SSGLYSLSVVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE FEGGPSVFLFPPKPKDTLMISRTPEVTCVV VAVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSDGSFFLY SRLTVDKSRWQQGNVFSCSVMHEALHN HYTQKSLSLSPGK	SEQ ID NO:112
Human IgG1 heavy chain constant sequence with FEAL*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTWSWNSGALTSGVHTFPAVLQ SSGLYSLSVVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE FEGGPSVFLFPPKPKDTLMISRTPEVTCVV VAVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV	SEQ ID NO:113

		YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSGDSFLL YSKLTVDKSRWQQGNVFSCSVMHEALH NHYTQKSLSLSPGK	
Human Kappa light chain constant sequence		RTVAAPSVFIFPPSDEQLKSGTASVVCCLN NFYPREAKVQWKVDNALQSGNSQESVTE QDSKDSSTYSLSSTLTLSKADYEKHKVYAC EVTHQGLSSPVTKSFNRGEC	SEQ ID NO:114
Human CD40		MVRLPLQCVLWGCCLLTAVHPEPPTACREK QYLINSQCCSLCQPGQKLVSDCTEFTETE CLPCGESEFLDTWNRETHCHQHKYCDPN LGLRVQQKGTSETDTICTCEEGWHCTSE ACESCVLHRSCSPGFGVKQIATGVSDTIC EPCPVGFFSNVSSAFEKCHPWTSCETKDL VVQQAGTNKTDVVCGPQDRLRALVVIPII FGILFAILLVLVFIKKVAKKPTNKAPHPKQE PQEIFPDDLPGSNTAAPVQETLHGCQPV TQEDGKESRISVQERQ	SEQ ID NO:115
Human IgG1 heavy chain constant sequence with FEA*		ASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTWSWNSGALTSGVHTFPAVLQ SSGLYSLSVVTVPSSSLGTQTYICNVNH KPSNTKVDKRVEPKSCDKTHTCPPCPAPE FEAGGPSVFLFPPKPKDTLMISRTPEVTCVV VAVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQV YTLPPSREEMTKNQVSLTCLVKGFYPSDIA VEWESNGQPENNYKTTTPVLDSGDSFFLY SKLTVDKSRWQQGNVFSCSVMHEALHN HYTQKSLSLSPGK	SEQ ID NO: 116
CD40-001 humanized antibody (HC6 and LC1):			
VH-CD40-001- HC6	VH	EVQLVQSGAEVKKPGASVKVSCKTSGYT FTEYIMHWVRQAPGQGLEWMGGIIPNNG GTSYNQKFQGRVTMTVDKSTSTGYMELS SLRSEDTAVYYCTRREYVGRNYYALDYW GQGTLTVTVSS	SEQ ID NO:117
CD40-001-HC6	HC, IgG1	EVQLVQSGAEVKKPGASVKVSCKTSGYT FTEYIMHWVRQAPGQGLEWMGGIIPNNG GTSYNQKFQGRVTMTVDKSTSTGYMELS SLRSEDTAVYYCTRREYVGRNYYALDYW GQGTLTVTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVTWSWNSGALTS GVHTFPAVLQSSGLYSLSVVTVPSSSLG TQTYICNVNHKPSNTKVDKRVEPKSCDKT HTCPPCPAPPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTTTPV	SEQ ID NO:118

		LDSDGSFFLYSKLTVDKSRWQQGNVFSC SVMHEALHNHYTQKSLSLSPGK	
CD40-001-HC6- FEAL	HC, IgG1 FEAL	EVQLVQSGAEVKKPGASVKVSCKTSGYT FTEYIMHWVRQAPGQGLEWMGGIIPNNG GTSYNQKFQGRVTMTVDKSTSTGYMELS SLRSED TAVYYCTRREVYGRNYYALDYW GQGT LTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVT VSWNSGALTS GVHTFPAVLQSSGLYSLSSV VTPSSSLG TQTYICNVNHKPSNTKVDKRVEPKSCDKT HTCPPCPAPEFEGGPSVFLFPPKPKDTLMI SRTPEVTCVVVAVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFLLYSKLTVDKSRWQQGNVFSC SVMHEALHNHYTQKSLSLSPGK	SEQ ID NO:119
CD40-001-HC6- FEAR	HC, IgG1 FEAR	EVQLVQSGAEVKKPGASVKVSCKTSGYT FTEYIMHWVRQAPGQGLEWMGGIIPNNG GTSYNQKFQGRVTMTVDKSTSTGYMELS SLRSED TAVYYCTRREVYGRNYYALDYW GQGT LTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVT VSWNSGALTS GVHTFPAVLQSSGLYSLSSV VTPSSSLG TQTYICNVNHKPSNTKVDKRVEPKSCDKT HTCPPCPAPEFEGGPSVFLFPPKPKDTLMI SRTPEVTCVVVAVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTTPPV LDSDGSFFLYSR LTVDKSRWQQGNVFSC SVMHEALHNHYTQKSLSLSPGK	SEQ ID NO:120
VL-CD40-001-LC1	VL	DIQMTQSPSSLSASVGDRVTITCSASQGI NNYLNWYQQKPGKAVKLLIYYTSSLHSGV PSRFSGSGSGTDYFTFTISLQPEDATYYC QQYSNLPYTFGGGTKVEIK	SEQ ID NO:121
CD40-001-LC1	LC, kappa	DIQMTQSPSSLSASVGDRVTITCSASQGI NNYLNWYQQKPGKAVKLLIYYTSSLHSGV PSRFSGSGSGTDYFTFTISLQPEDATYYC QQYSNLPYTFGGGTKVEIKRTVAAPSVFIF PPSDEQLKSGTASVVCLLNNFYPREAKVQ WKVDNALQSGNSQESVTEQDSKDSSTYS SSTLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGEC	SEQ ID NO:122
CD137-009 humanized antibody (HC7 and LC2):			
VH-CD137-009- HC7	VH	EVQLVESGGGLVQPGRSLRLSCTASGFSL NDYWMSSWVRQAPGKGLEWVG YIDVGG LYYAASVKGRFTISRDDSKSIAYLQMNSL	SEQ ID NO:123

		KTEDTAVYYCARGGLTYGFDLWGQGTLV TVSS	
CD137-009-HC7	HC, IgG1	EVQLVESGGGLVQPGRSLRLSCTASGFSL NDYWMSWVRQAPGKGLEWVGIDVGGS LYYAASVKGRFTISRDDSKSIAYLQMNSL KTEDTAVYYCARGGLTYGFDLWGQGTLV TVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPP CPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQP REPQVYTLPPSREEMTKNQVSLTCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK	SEQ ID NO:124
CD137-009-HC7- FEAR	HC, IgG1 FEAR	EVQLVESGGGLVQPGRSLRLSCTASGFSL NDYWMSWVRQAPGKGLEWVGIDVGGS LYYAASVKGRFTISRDDSKSIAYLQMNSL KTEDTAVYYCARGGLTYGFDLWGQGTLV TVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPP CPAPE FE GGPSVFLFPPKPKDTLMISRTPE VTCVV AV SHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQP REPQVYTLPPSREEMTKNQVSLTCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYS R LTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK	SEQ ID NO:125
CD137-009-HC7- FEAL	HC, IgG1 FEAL	EVQLVESGGGLVQPGRSLRLSCTASGFSL NDYWMSWVRQAPGKGLEWVGIDVGGS LYYAASVKGRFTISRDDSKSIAYLQMNSL KTEDTAVYYCARGGLTYGFDLWGQGTLV TVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVPSSSLGTQTYIC NVNHKPSNTKVDKRVEPKSCDKTHTCPP CPAPE FE GGPSVFLFPPKPKDTLMISRTPE VTCVV AV SHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQP REPQVYTLPPSREEMTKNQVSLTCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSD GS F LLYSKLTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK	SEQ ID NO:126
VL-CD137-009- LC2	VL	DIVMTQSPSSLSASVGDRVTITCQASEDI SSYLAWYQQKPGKAPKRLIYGASDLASG	SEQ ID NO:127

		VPSRFSASGSGTDYFTFTISLQPEDATYY CHYYATISGLGVAFGGGTKVEIK	
CD137-009-LC2	LC, kappa	DIVMTQSPSSLSASVGDRTITCQASEDI SSYLAWYQQKPGKAPKRLIYGASDLASG VPSRFSASGSGTDYFTFTISLQPEDATYY CHYYATISGLGVAFGGGTKVEIKRTVAAP SVFIFPPSDEQLKSGTASVCLLNNFYPRE AKVQWKVDNALQSGNSQESVTEQDSKD STYLSSTLTLSKADYEKHKVYACEVTHQ GLSSPVTKSFNRGEC	SEQ ID NO:128
Human CD137 (shuffle 6)	Amino acids 24-47 of human CD137	LQDPCSNCPAGTFCDNNRNQICSP	SEQ ID NO:129
Human CD137 (shuffle 5)	Amino acids 48-88 of human CD137	CPPNSFSSAGGQRTCDICRQCKGVFRTR KECSSTSNAECDC	SEQ ID NO:130
Human CD137 (shuffle 4)	Amino acids 89-114 of human CD137	TPGFHCLGAGCSMCEQDCKQGQELTK	SEQ ID NO:131
Human CD137 (shuffle 3)	Amino acids 115-138 of human CD137	KGCKDCCFGTFNDQKRGICRPWTN	SEQ ID NO:132
Human CD137 (shuffle 2)	Amino acids 139-161 of human CD137	CSLDGKSVLVNGTKERDVVCGPS	SEQ ID NO:133
Human CD137 (shuffle 1)	Amino acids 162-186 of human CD137	PADLSPGASSVTPAPAREPGHSPQ	SEQ ID NO:134
Wild Boar CD137		MGNNGYNNIVATVLLVMNFERTRSVPDPCS NCSAGTFCGKNIQELCMPCPSNSFSSTSG QKACNVCRKCEGVFRTKKECSSTSNAVC ECVPGFRCLGAGCAMCEEYCQQGQELTQ EGCKDCSFGTFNDEEHGVCRPWTDCLSLA GKPVLMNGTKARDVVCGPRPTDFSPGTP STTMPVPGGEPGHTSHVIFFLALMSTAVF VLVSYLALRFSVVQQGRKKLLYIVKQPFLK PAQTVQEEDACSRFPEEEEGECEL	SEQ ID NO:135
African Elephant CD137		MGNNGYNNMIVATVLLVMNFERTGAVQDSC RDCLAGTYCVKNESQICSPCLNSFSSTG GQMNCDCMRKCEGVFKTKRACSPTRDAE CECVSGFHCLGAGCTMCQQDCKQGQEL TKEGCKDCCLGTFNDQKNGICRPWTNCS LEGKSVLANGTKKRDVVCGPPAADSFPDT SSVTVPAPERKPDHHPQIITFFLALISAALL FLVFFLVVRFSAKVGWRKKLLYIFKQPFIK	SEQ ID NO:136

		PVQTAQEEDGCSCRFPEEEEGDCEL	
--	--	---------------------------	--

*amino acids positions 118-447 according to EU numbering

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The term "CD40" as used herein, refers to CD40, also referred to as tumor necrosis factor receptor superfamily member 5 (TNFRSF5), which is the receptor for the ligand TNFSF5/CD40L. CD40 is known to transduce TRAF6- and MAP3K8-mediated signals that activate ERK in macrophages and B cells, leading to induction of immunoglobulin secretion by the B cells. Other synonyms used for CD40 include, but are not limited to, B-cell surface antigen CD40, Bp50, CD40L receptor and CDw40. In one embodiment, CD40 is human CD40, having UniProt accession number P25942. The sequence of human CD40 is also shown in SEQ ID NO:115. Amino acids 1-20 of SEQ ID NO:115 correspond to the signal peptide of human CD40; while amino acids 21-193 of SEQ ID NO:115 correspond to the extracellular domain of human CD40; and the remainder of the protein; i.e. from amino acids 194-215 and 216-277 of SEQ ID NO:115 is transmembrane and cytoplasmic domain, respectively.

The term "CD137" as used herein, refers to CD137 (4-1BB), also referred to as tumor necrosis factor receptor superfamily member 9 (TNFRSF9), which is the receptor for the ligand TNFSF9/4-1BBL. CD137 (4-1BB) is believed to be involved in T-cell activation. Other synonyms for CD137 include, but are not limited to, 4-1BB ligand receptor, CDw137, T-cell antigen 4-1BB homolog and T-cell antigen ILA. In one embodiment, CD137 (4-1BB) is human CD137 (4-1BB), having UniProt accession number Q07011. The sequence of human CD137 is also shown in SEQ ID NO:92. Amino acids 1-23 of SEQ ID NO:92 correspond to the signal peptide of human CD137; while amino acids 24-186 of SEQ ID NO:92 correspond to the extracellular domain of human CD137; and the remainder of the protein, i.e. from amino acids 187-213 and 214-255 of SEQ ID NO:92 are transmembrane and cytoplasmic domain, respectively.

The term "chimeric antibody" as used herein, refers to an antibody wherein the variable region is derived from a non-human species (e.g. derived from rodents) and the constant region is derived from a different species, such as human. Chimeric antibodies may be generated by antibody engineering. "Antibody engineering" is a term used generically for different kinds of modifications of antibodies, and processes for antibody engineering are well-known for the skilled person. In particular, a chimeric antibody may be generated by using standard DNA techniques as described in Sambrook et al., 1989,

Molecular Cloning: A laboratory Manual, New York: Cold Spring Harbor Laboratory Press, Ch. 15. Thus, the chimeric antibody may be a genetically or an enzymatically engineered recombinant antibody. It is within the knowledge of the skilled person to generate a chimeric antibody, and thus, generation of the chimeric antibody may be performed by
5 other methods than those described herein. Chimeric monoclonal antibodies for therapeutic applications in humans are developed to reduce anticipated antibody immunogenicity of non-human antibodies, e.g. rodent antibodies. They may typically contain non-human (e.g. murine or rabbit) variable regions, which are specific for the antigen of interest, and human constant antibody heavy and light chain domains. The
10 terms "variable region" or "variable domain" as used in the context of chimeric antibodies, refer to a region which comprises the CDRs and framework regions of both the heavy and light chains of an immunoglobulin, as described below.

The term "humanized antibody" as used herein, refers to a genetically engineered non-human antibody, which contains human antibody constant domains and non-human
15 variable domains modified to contain a high level of sequence homology to human variable domains. This can be achieved by grafting of the six non-human antibody CDRs, which together form the antigen-binding site, onto a homologous human acceptor framework region (FR) (see WO92/22653 and EP0629240). In order to fully reconstitute the binding affinity and specificity of the parental antibody, the substitution of
20 framework residues from the parental antibody (i.e. the non-human antibody) into the human framework regions (back-mutations) may be required. Structural homology modeling may help to identify the amino acid residues in the framework regions that are important for the binding properties of the antibody. Thus, a humanized antibody may comprise non-human CDR sequences, primarily human framework regions optionally
25 comprising one or more amino acid back-mutations to the non-human amino acid sequence, and fully human constant regions. Optionally, additional amino acid modifications, which are not necessarily back-mutations, may be applied to obtain a humanized antibody with preferred characteristics, such as affinity and biochemical properties and/or additional amino acid mutations may be introduced in the constant
30 region.

As used herein, a protein which is "derived from" another protein, e.g., a parent protein, means that one or more amino acid sequences of the protein are identical or similar to one or more amino acid sequences in the other or parent protein. For example, in an antibody, binding arm, antigen-binding region, constant region, or the
35 like which is derived from another or a parent antibody, binding arm, antigen-binding region, or constant region, one or more amino acid sequences are identical or similar to those of the other or parent antibody, binding arm, antigen-binding region, or constant region. Examples of such one or more amino acid sequences include, but are not limited

to, those of the VH and VL CDRs and/or one or more or all of the framework regions, VH, VL, CL, hinge, or CH regions. For example, a humanized antibody can be described herein as "derived from" a non-human parent antibody, meaning that at least the VL and VH CDR sequences are identical or similar to the VH and VL CDR sequences of said non-human parent antibody. A chimeric antibody can be described herein as being "derived from" a non-human parent antibody, meaning that typically the VH and VL sequences may be identical or similar to those of the non-human parent antibody. Another example is a binding arm or an antigen-binding region which may be described herein as being "derived from" a particular parent antibody, meaning that said binding arm or antigen-binding region typically comprises identical or similar VH and/or VL CDRs, or VH and/or VL sequences to the binding arm or antigen-binding region of said parent antibody. As described elsewhere herein, however, amino acid modifications such as mutations can be made in the CDRs, constant regions or elsewhere in the antibody, binding arm, antigen-binding region or the like, to introduce desired characteristics. When used in the context of one or more sequences derived from a first or parent protein, a "similar" amino acid sequence preferably has a sequence identity of at least about 50%, such as at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, or at least about 97%, 98% or 99%.

Non-human antibodies can be generated in a number of different species, such as mouse, rabbit, chicken, guinea pig, llama and goat.

Monoclonal antibodies can be produced by a variety of techniques, including conventional monoclonal antibody methodology, e.g., the standard somatic cell hybridization technique of Kohler and Milstein, *Nature* 256: 495 (1975). Other techniques for producing monoclonal antibodies can be employed, e.g., viral or oncogenic transformation of B-lymphocytes or phage display techniques using libraries of antibody genes, and such methods are well known to a person skilled in the art.

Hybridoma production in such non-human species is a very well established procedure. Immunization protocols and techniques for isolation of splenocytes of immunized animals/non-human species for fusion are known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known.

The term "human antibody" as used herein, refers to antibodies having variable and constant regions derived from human germline immunoglobulin sequences. Human antibodies may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo). However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

Human monoclonal antibodies can be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system.

The term "immunoglobulin" refers to a class of structurally related glycoproteins consisting of two pairs of polypeptide chains, one pair of light (L) low molecular weight chains and one pair of heavy (H) chains, all four inter-connected by disulfide bonds. The structure of immunoglobulins has been well characterized. See for instance Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)). Briefly, each heavy chain (abbreviated "HC") typically is comprised of a heavy chain variable region (abbreviated herein as V_H or VH) and a heavy chain constant region (abbreviated herein as C_H or CH). The heavy chain constant region typically is comprised of three domains, C_{H1} , C_{H2} , and C_{H3} . The heavy chain may typically further comprise a hinge region. Each light chain (abbreviated "LC") typically is comprised of a light chain variable region (abbreviated herein as V_L or VL) and a light chain constant region (abbreviated herein as C_L or CL). The light chain constant region typically is comprised of one domain, C_L . The V_H and V_L regions may be further subdivided into regions of hypervariability (or hypervariable regions which may be hypervariable in sequence and/or form of structurally defined loops), also termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FRs). Each V_H and V_L is typically composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4 (see also Chothia and Lesk J. Mol. Biol. 196, 901-917 (1987)). Unless otherwise stated or contradicted by context, CDR sequences herein are identified according to IMGT rules using DomainGapAlign (Program version: 4.9.1; 2013-12-19) (Lefranc MP., Nucleic Acids Research 1999;27:209-212, and Ehrenmann F., Kaas Q. and Lefranc M.-P. Nucleic Acids Research 2010;38, D301-307; see also internet [http address www.imgt.org/](http://www.imgt.org/)).

Unless otherwise stated or contradicted by context, reference to amino acid positions in the constant regions in the present invention is according to the EU-numbering (Edelman et al., Proc Natl Acad Sci U S A. 1969 May;63(1):78-85; Kabat et al., Sequences of Proteins of Immunological Interest, Fifth Edition. 1991 NIH Publication No. 91-3242).

The term "antibody" (Ab) in the context of the present invention refers to a molecule comprising at least one antibody variable domain such as an immunoglobulin heavy chain variable region, or an immunoglobulin heavy chain variable region and a light chain variable region, or a fragment thereof, or a derivative of either thereof, which has the ability to specifically bind to an antigen, such as under typical physiological conditions with a half-life of significant periods of time, such as at least about 30

minutes, at least about 45 minutes, at least about one hour, at least about two hours, at least about four hours, at least about 8 hours, at least about 12 hours, about 24 hours or more, about 48 hours or more, about 3, 4, 5, 6, 7 or more days, etc., or any other relevant functionally-defined period (such as a time sufficient to induce, promote, enhance, and/or modulate a physiological response associated with antibody binding to the antigen and/or time sufficient for the antibody to recruit an effector activity). In particular the antibody may be an immunoglobulin molecule, a fragment of an immunoglobulin molecule or a derivative thereof. The variable regions of the heavy and light chains of the immunoglobulin molecule contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (such as effector cells) and components of the complement system such as C1q, the first component in the classical pathway of complement activation. As indicated above, the term antibody herein, unless otherwise stated or clearly contradicted by context, includes fragments of an antibody that are antigen-binding fragments, *i.e.*, retain the ability to specifically bind to the antigen. It has been shown that the antigen-binding function of an antibody may be performed by fragments of a full-length antibody. Examples of antigen-binding fragments encompassed within the term "antibody" include (i) a Fab' or Fab fragment, a monovalent fragment consisting of the V_L , V_H , C_L and C_{H1} domains, or a monovalent antibody as described in WO2007059782 (Genmab); (ii) $F(ab')_2$ fragments, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) an Fd fragment consisting essentially of the V_H and C_{H1} domains; (iv) an Fv fragment consisting essentially of the V_L and V_H domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., Nature 341, 544-546 (1989)), which consists essentially of a V_H domain and also called domain antibodies (Holt et al; Trends Biotechnol. 2003 Nov;21(11):484-90); (vi) camelid or nanobodies (Revets et al; Expert Opin Biol Ther. 2005 Jan;5(1):111-24) and (vii) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, V_L and V_H , are coded for by separate genes, they may be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the V_L and V_H regions pair to form monovalent molecules (known as single chain antibodies or single chain Fv (scFv), see for instance Bird et al., Science 242, 423-426 (1988) and Huston et al., PNAS USA 85, 5879-5883 (1988)). Such single chain antibodies are encompassed within the term antibody unless otherwise noted or clearly indicated by context. Although such fragments are generally included within the meaning of antibody, they collectively and each independently are unique features of the present invention, exhibiting different biological properties and utility. These and other useful antibody fragments in the context of the present invention, as well as bispecific formats

of such fragments, are discussed further herein. It also should be understood that the term antibody, unless specified otherwise, also includes polyclonal antibodies, monoclonal antibodies (mAbs), antibody-like polypeptides, such as chimeric antibodies and humanized antibodies, and antibody fragments retaining the ability to specifically bind to the antigen (antigen-binding fragments) provided by any known technique, such as enzymatic cleavage, peptide synthesis, and recombinant techniques. An antibody as generated can possess any isotype and/or subclass.

Regular antibodies; e.g. antibodies raised in any species are normally monospecific, bivalent antibodies, which means that they comprise two antigen-binding regions which bind to the same epitope.

The term "multispecific antibody" in the context of the present invention refers to an antibody having different antigen-binding regions defined by different antibody sequences. Thus a multispecific antibody may have two, three, four, five or more different antigen-binding regions. Examples of multispecific antibodies include antibodies having two different antigen-binding regions; i.e. a bispecific antibody.

Examples of multispecific antibodies comprising three or more different antigen-binding regions include but are not limited to (i) bispecific antibodies coupled with an additional single chain variable Fragment (scFv) at their Fc part (Weidle et al., Cancer Genomics Proteomics. 2013 Jan-Feb;10(1):1-18), (ii) fusion proteins consisting of three or more scFv (triabody, tetrabody; Chames et al., FEMS Microbiol Lett. 2000 Aug 1;189(1):1-8) and (iii) fusion proteins connected to scFv (Kermer et.al. Mol Cancer Ther. 2014 Jan;13(1):112-21).

The term "bispecific antibody" in the context of the present invention refers to an antibody having two different antigen-binding regions defined by different antibody sequences.

When used herein, unless contradicted by context, the term "Fab-arm" or "arm" refers to one heavy chain-light chain pair and is used interchangeably with "half molecules" herein.

The term "binding arm comprising an antigen-binding region" means an antibody molecule or fragment that comprises an antigen-binding region. Thus, a binding arm can comprise, e.g., the six VH and VL CDR sequences, the VH and VL sequences, a Fab or Fab' fragment, or a Fab-arm.

When used herein, unless contradicted by context, the term "Fc region" refers to an antibody region comprising at least a hinge region, a CH2 domain, and a CH3 domain.

As used herein, the term "isotype" refers to the specific type of immunoglobulin encoded by the HC (for instance IgG, IgD, IgA, IgE, and IgM) or LC (kappa, κ or lambda,

λ) genes. Within each isotype, there may be several subclasses, such as IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, etc.

The term "monovalent antibody" means in the context of the present invention that an antibody molecule is capable of binding a single molecule of the antigen, and thus is not capable of antigen cross-linking.

A "CD40 antibody" or "anti-CD40 antibody" is an antibody as described above, which binds specifically to the antigen CD40.

A "CD137 antibody" or "anti-CD137 antibody" is an antibody as described above, which binds specifically to the antigen CD137.

A "CD40xCD137 antibody" or "anti-CD40xCD137 antibody" is a bispecific antibody, which comprises two different antigen-binding regions, one of which binds specifically to the antigen CD40 and one of which binds specifically to the antigen CD137.

The term "specifically binds", "specifically binding", "specific binding" or other similar wording refers to the ability of an antibody to preferentially bind to a particular antigen compared to other antigens, or to a particular part (epitope) of an antigen compared to other parts of the same antigen.

As used herein, the term "binding" in the context of the binding of an antibody to a predetermined antigen or epitope typically is a binding with an affinity corresponding to a K_D of about 10^{-7} M or less, such as about 10^{-8} M or less, such as about 10^{-9} M or less, about 10^{-10} M or less, or about 10^{-11} M or even less when determined by for instance surface plasmon resonance (SPR) technology in a BIAcore 3000 instrument using the antibody as the ligand and the antigen as the analyte (or vice versa), and binds to the predetermined antigen with an affinity corresponding to a K_D that is at least ten-fold lower, such as at least 100-fold lower, for instance at least 1,000-fold lower, such as at least 10,000-fold lower, for instance at least 100,000-fold lower than the K_D for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen. The amount with which the affinity is higher is dependent on the K_D of the antibody, so that when the K_D of the antibody is very low (that is, the antibody is highly specific), then the amount with which the affinity for the antigen is higher than the affinity for a non-specific antigen may be at least 10,000 fold.

The term " k_d " (sec^{-1}), as used herein, refers to the dissociation rate constant of a particular antibody-antigen interaction. Said value is also referred to as the k_{off} value.

The term " K_D " (M), as used herein, refers to the dissociation equilibrium constant of a particular antibody-antigen interaction.

Two antibodies have the "same specificity" if they bind to the same antigen and to the same epitope. Whether an antibody to be tested recognizes the same epitope as a

certain antigen-binding antibody, i.e., the antibodies bind to the same epitope, may be tested by different methods well known to a person skilled in the art.

The competition between the antibodies can be detected by a cross-blocking assay. For example, a competitive ELISA assay may be used as a cross-blocking assay.

5 For example, target antigen may be coated on the wells of a microtiter plate and antigen-binding antibody and candidate competing test antibody may be added. The amount of the antigen-binding antibody bound to the antigen in the well indirectly correlates with the binding ability of the candidate competing test antibody that competes therewith for binding to the same epitope. Specifically, the larger the affinity
10 of the candidate competing test antibody is for the same epitope, the smaller the amount of the antigen-binding antibody bound to the antigen-coated well. The amount of the antigen-binding antibody bound to the well can be measured by labeling the antibody with detectable or measurable labeling substances.

An antibody competing for binding to an antigen with another antibody, e.g., an
15 antibody comprising heavy and light chain variable regions as described herein, or an antibody having the specificity for an antigen of another antibody, e.g., an antibody comprising heavy and light chain variable regions as described herein, may be an antibody comprising variants of said heavy and/or light chain variable regions as described herein, e.g. modifications in the CDRs and/or a certain degree of identity as
20 described herein.

An "isolated multispecific antibody" as used herein is intended to refer to a multispecific antibody which is substantially free of other antibodies having different antigenic specificities (for instance an isolated bispecific antibody that specifically binds to CD40 and CD137 is substantially free of monospecific antibodies that specifically bind
25 to CD40 or CD137).

The term "epitope" means a protein determinant capable of specific binding to an antibody. Epitopes usually consist of surface groupings of molecules such as amino acids or sugar side chains and usually have specific three-dimensional structural characteristics, as well as specific charge characteristics. Conformational and non-
30 conformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents. The epitope may comprise amino acid residues directly involved in the binding and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked or covered by the specifically antigen-binding peptide (in other words, the amino
35 acid residue is within the footprint of the specifically antigen-binding peptide).

The term "monoclonal antibody" as used herein refers to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope.

When used herein the term "heterodimeric interaction between the first and second CH3 regions" refers to the interaction between the first CH3 region and the second CH3 region in a first-CH3/second-CH3 heterodimeric antibody.

5 When used herein the term "homodimeric interactions of the first and second CH3 regions" refers to the interaction between a first CH3 region and another first CH3 region in a first-CH3/first-CH3 homodimeric antibody and the interaction between a second CH3 region and another second CH3 region in a second-CH3/second-CH3 homodimeric antibody.

10 When used herein the term "homodimeric antibody" refers to an antibody comprising two first Fab-arms or half-molecules, wherein the amino acid sequence of said Fab-arms or half-molecules is the same.

15 When used herein the term "heterodimeric antibody" refers to an antibody comprising a first and a second Fab-arm or half-molecule, wherein the amino acid sequence of said first and second Fab-arms or half-molecules are different. In particular, the CH3 region, or the antigen-binding region, or the CH3 region and the antigen-binding region of said first and second Fab-arms/half-molecules are different.

The term "reducing conditions" or "reducing environment" refers to a condition or an environment in which a substrate, such as a cysteine residue in the hinge region of an antibody, is more likely to become reduced than oxidized.

20 The present invention also provides multispecific antibodies, such as bispecific antibodies, comprising functional variants of the V_L regions, V_H regions, or one or more CDRs of the bispecific antibodies of the examples. A functional variant of a V_L , V_H , or CDR used in the context of a bispecific antibody still allows each antigen-binding region of the bispecific antibody to retain at least a substantial proportion (at least about 50%, 25 60%, 70%, 80%, 90%, 95% or more) of the affinity and/or the specificity/selectivity of the parent bispecific antibody and in some cases such a bispecific antibody may be associated with greater affinity, selectivity and/or specificity than the parent bispecific antibody.

30 Such functional variants typically retain significant sequence identity to the parent bispecific antibody. The percent identity between two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The percent identity between two nucleotide or amino acid sequences 35 may e.g. be determined using the algorithm of E. Meyers and W. Miller, Comput. Appl. Biosci 4, 11-17 (1988) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences may be

determined using the Needleman and Wunsch, J. Mol. Biol. 48, 444-453 (1970) algorithm.

In the context of the present invention the following notations are, unless otherwise indicated, used to describe a mutation; name of amino acid which is mutated, followed by the position number which is mutated, followed by what the mutation encompass. Thus if the mutation is a substitution, the name of the amino acid which replaces the prior amino acid is included, if the amino acid is deleted it is indicated by a *, if the mutation is an addition the amino acid being added is included after the original amino acid. Amino acid names may be one or three-letter codes. Thus for example; substitution of a Lysine in position 409 with an Arginine is referred to as K409R, substitution of Lysine in position 409 with any amino acid is referred to as K409X, deletion of Lysine in position 409 is referred to as K409* and addition of P after Lysine at position K409 is referred to as K409KP.

Exemplary variants include those which differ from the VH and/or VL and/or CDRs of the parent sequences mainly by conservative substitutions; for instance 12, such as 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 of the substitutions in the variant are conservative amino acid residue replacements.

In the context of the present invention, conservative substitutions may be defined by substitutions within the classes of amino acids reflected in the following:

20

Amino acid residue classes for conservative substitutions:

Acidic Residues:	Asp (D) and Glu (E)
Basic Residues:	Lys (K), Arg (R), and His (H)
Hydrophilic Uncharged Residues:	Ser (S), Thr (T), Asn (N), and Gln (Q)
25 Aliphatic Uncharged Residues:	Gly (G), Ala (A), Val (V), Leu (L), and Ile (I)
Non-polar Uncharged Residues:	Cys (C), Met (M), and Pro (P)
Aromatic Residues:	Phe (F), Tyr (Y), and Trp (W)

The first and/or second antigen-binding region of the present invention may also be a variant of a first and/or second antigen-binding region, respectively, disclosed herein.

It is well known to a person skilled in the art how to introduce modifications and that certain amino acids of the CDR sequences may be modified; e.g. by amino acid substitutions to e.g. increase affinity of an antibody to its target antigen, reduce potential immunogenicity of non-human antibodies to be used in humans and/or to increase the yield of antibodies expressed by a host cell. Such modifications can be introduced without affecting the epitope of the target antigen to which the antibody binds.

The term "recombinant host cell" (or simply "host cell" or "cell"), as used herein, is intended to refer to a cell into which a nucleic acid, such as an expression vector has been introduced, e.g. a nucleic acid, such as an expression vector encoding a multispecific antibody of the invention. Recombinant host cells include, for example, transfectomas, such as CHO, CHO-S, HEK, HEK293, HEK-293F, Expi293F, PER.C6 or NS0 cells, and lymphocytic cells.

The term "treatment" refers to the administration of an effective amount of a therapeutically active multispecific antibody of the present invention with the purpose of easing, ameliorating, arresting or eradicating (curing) symptoms or disease states.

The term "effective amount" or "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired therapeutic result. A therapeutically effective amount of a multispecific antibody may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the multispecific antibody to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the multispecific antibody or a fragment thereof, are outweighed by the therapeutically beneficial effects.

The term "anti-idiotypic antibody" refers to an antibody which recognizes unique determinants generally associated with the antigen-binding site of an antibody.

In the context of the present invention the term "induce Fc-mediated effector function to a lesser extent" used in relation to an antibody, including a multispecific antibody, means that the antibody induces Fc-mediated effector functions, such function in particular being selected from the list of IgG Fc receptor (FcγR) binding, C1q binding, ADCC or CDC, to a lesser extent compared to a human IgG1 antibody comprising (i) the same CDR sequences, in particular comprising the same first and second antigen-binding regions, as said antibody and (ii) two heavy chains comprising human IgG1 hinge, CH2 and CH3 regions.

Fc-mediated effector function may be measured by binding to FcγRs, binding to C1q, or induction of Fc-mediated cross-linking via FcγRs.

Further aspects and embodiments of the invention

The present invention relates to a molecule comprising two different antigen-binding regions, one of which has specificity for human CD40 and one of which has specificity for human CD137.

In a particular embodiment, said molecule may be a multispecific antibody.

Thus the present invention relates to a multispecific antibody comprising (i) a first antigen-binding region binding to human CD40; and (ii) a second antigen-binding region binding to human CD137.

As shown by the inventors of the present invention a bispecific antibody according to the present invention may induce intracellular signaling when binding to CD40 expressed on one cell and binding to CD137 expressed on another cell. Thus, a multispecific antibody according to the present invention is able to trans-activate two different cells. In humans, CD40 is expressed on a number of cells including antigen-presenting cells (APCs), such as dendritic cells, whereas CD137 is expressed on T cells and other cells. Thus, multispecific antibodies, such as bispecific antibodies, according to the present invention binding to CD40 and CD137 are able to bind simultaneously to APCs and T cells expressing these receptors. Without being bound by theory, multispecific antibodies, such as bispecific antibodies, according to the present invention may thus (i) mediate cell-to-cell interaction between APCs and T cells by receptor binding and (ii) activate both CD40 and CD137 at once, which is primarily induced by cross-linking and receptor clustering upon cell-to-cell interaction and not necessarily dependent on agonistic activity of the parental monospecific bivalent antibodies. Thus, these trans-activating multispecific antibodies, such as bispecific antibodies, exert co-stimulatory activity in the context of APC:T cell interactions, and can elicit a T cell response against tumor cells. As such, this mechanism of action can reflect natural T-cell activation via antigen-presentation by activated APCs, allowing for the presentation of a variety of tumor-specific antigens by the APCs to T cells. Without being limited to theory, the costimulatory activity may provide for one or more of (i) only specific T cells being activated (i.e., those that are in contact with an APC) as opposed to any T cell and (ii) re-activation of exhausted T cells, by strong co-stimulation via activated APCs and CD137 triggering and (iii) the priming of T cells by inducing antigen presentation by activated APCs and at the same time triggering CD137.

Thus, a multispecific, such as a bispecific, antibody of the present invention may be used for treatment of a disease which would benefit from activation of T cells, such as cancer.

In one embodiment, the multispecific antibody according to the present invention comprises

(I) a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2, and CDR3 selected from the group consisting of:

- a) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:3 or a sequence wherein up to four amino acids are modified in SEQ ID NO:3, and

light chain variable region CDR3 having the sequence set forth in SEQ ID NO:5 or a sequence wherein up to four amino acids are modified in SEQ ID NO:5; and

b) heavy and light chain variable region CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR3 according to a) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR3 according to a), and

(II) a second antigen-binding region binding to human CD137.

In a further embodiment, the first antigen-binding region may further comprise heavy chain variable region CDR1 having the sequence as set forth in SEQ ID NO:1 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:1, and/or heavy chain variable region CDR2 having the sequence as set forth in SEQ ID NO:2 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:2; and/or light chain variable region CDR1 having the sequence as set forth in SEQ ID NO:4 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:4, and/or light chain variable region CDR2 having the sequence YTS or a sequence wherein up to 2 amino acids are modified in YTS.

Thus, in one embodiment, the present invention relates to a multispecific antibody comprising

(I) a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2, and CDR3 selected from the group consisting of:

- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively;
- b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations; and
- c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b), and
- (II) a second antigen-binding region binding to human CD137.

In a further embodiment, said first antigen-binding region comprises a first heavy chain variable (VH) sequence, and a first light chain variable (VL) sequence, and said second antigen-binding region comprises a second heavy chain variable (VH) sequence, and a second light chain variable (VL) sequence, wherein said variable sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively, and four framework sequences, FR1, FR2, FR3 and FR4, respectively.

In a further embodiment, said multispecific antibody comprises (I) a first binding arm comprising said first antigen-binding region, and (II) a second binding arm comprising said second antigen-binding region.

In one embodiment, the first binding arm comprises said first antigen-binding region and a first heavy chain constant sequence, and the second binding arm comprises said second-antigen-binding region and a second heavy chain constant sequence.

In a further embodiment, (i) said first binding arm comprises said first antigen-binding region, wherein the first binding arm comprises a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence, and (ii) said second binding arm comprises said second antigen-binding region, wherein the second binding arm comprises a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain comprising a second light chain variable (VL) sequence.

In a further embodiment, said first light chain further comprises a first light chain constant (CL) sequence, and said second light chain further comprises a second light chain constant (CL) sequence.

In one embodiment, the first binding arm comprises a first Fab-arm comprising said first antigen-binding region, and the second binding arm comprises a second Fab-arm comprising said second antigen-binding region.

In one embodiment, said first and second antigen-binding regions of the multispecific antibody according to the present invention are derived from a humanized antibody. In one embodiment, the first and second binding arm may be derived from a humanized antibody.

In one embodiment, the first and second binding arms of the multispecific antibody according to the present invention are derived from a full-length antibody.

In one embodiment, the first and second binding arm of the multispecific antibody according to the present invention are derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.

In one embodiment, the first and second binding arms are derived from a monoclonal antibody.

In one embodiment, the first and said second heavy chains of the multispecific antibody according to the present invention are of an IgG isotype. The subclass of the first and second heavy chains may, for example, be separately selected from the group consisting of IgG1, IgG2, IgG3, and IgG4. In one embodiment, the first and second heavy chains are of the same IgG subclass, such as IgG1.

In one embodiment, the multispecific antibody according to the present invention is an isolated antibody.

In a further embodiment, each of said first and second heavy chains comprises at least a hinge region, a CH2 and a CH3 region. In a further embodiment, the CH3 regions of said first and second heavy chains comprise asymmetrical mutations.

In one embodiment, the multispecific antibody according to the present invention is a bispecific antibody.

In one embodiment, the multispecific antibody according to the present invention may cross-link one cell expressing human CD40, e.g. a first cell, and another cell expressing human CD137, e.g. a second cell.

In one embodiment, said cross-linking is determined by an assay using a first cell line expressing human CD40 and a second cell line expressing human CD137, and wherein either the first or the second cell line comprises a reporter structure resulting in the production of a measurable reporter upon NF- κ B activation.

In one embodiment said first cell may be an antigen-presenting cell and said second cell may be a T-cell, such as a CD4⁺ or a CD8⁺ T-cell.

Different methods may be used to determine cross-linking a first cell expressing CD40 and a second cell expressing CD137, and the present invention is not limited to any particular method.

In one embodiment, said cross-linking may be determined by a reporter assay e.g. as described in Example 4. In brief, said assay comprises co-culturing a reporter cell line expressing a first target antigen and transduced with a reporter gene (luciferase for instance) driven by NF- κ B response elements with a second cell line expressing a second target antigen, adding a multispecific antibody according to the present invention in concentrations from 100 ng/mL to 10,000 ng/mL to the cell co-culture, and measuring reporter gene expression, e.g. luciferase generation, wherein said first target antigen is human CD40 and said second target antigen is human CD137 or vice versa.

A multispecific antibody capable of inducing cross-linking of the CD40 and CD137 expressed on different cells, will in this assay result in measurable activation of the first target antigen based on the reporter gene expression upon NF- κ B pathway activation.

In one embodiment, the multispecific antibody according to the present invention may be able to induce reporter gene expression produced upon NF- κ B activation only

upon addition of the second cell line expressing the second target antigen without the NF- κ B reporter gene.

In one embodiment, the multispecific antibody according to the present invention may be able to induce higher reporter gene expression produced upon NF- κ B activation
5 when the second cell line expressing the second target antigen without the NF- κ B reporter gene is added compared to addition of a second cell line not expressing the second target antigen.

In one embodiment, the multispecific antibody is a bispecific antibody, and said bispecific antibody may, in one embodiment:

10 (i) induce reporter gene expression when added to a co-culture of the reporter cell line expressing CD137 and the second cell line expressing CD40, or

(ii) induce a higher amount of the reporter gene expression when added to the co-culture of the reporter cell line expressing CD137 and the second cell line expressing CD40, compared to a reference bispecific antibody comprising the same second antigen-
15 binding region binding to human CD137, but wherein the first antigen-binding region of the reference bispecific antibody binds to an irrelevant target antigen, e.g. wherein the first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:99, 100 and 101, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ
20 ID NOs:102, GVS and 103, respectively.

In one embodiment, the multispecific antibody is a bispecific antibody, and said bispecific antibody may, in one embodiment:

(i) induce reporter gene expression when added to a co-culture of the reporter cell line expressing CD40 and the second cell line expressing CD137, or

25 (ii) induce a higher amount of the reporter gene expression when added to the co-culture of the reporter cell line expressing CD40 and the second cell line expressing CD137, compared to a reference bispecific antibody comprising the same first antigen-binding region binding to human CD40, but wherein the second antigen-binding region of the reference bispecific antibody comprises heavy chain variable region CDR1, CDR2
30 and CDR3 having the sequences set forth in SEQ ID NOs:99, 100 and 101, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:102, GVS and 103, respectively.

In one embodiment, the multispecific antibody according to the present invention induces and/or enhances proliferation of T cells, e.g. wherein said T cells are CD4⁺
35 and/or CD8⁺ T cells.

Different methods for determining or measuring proliferation of T cells may be used and the present invention is not limited to any particular method.

In one embodiment, said induction or enhancement of proliferation of T cells is determined by a non-antigen-specific T-cell proliferation assay, e.g. as described in Example 5. Thus induction and/or enhancement of proliferation of T cells may be determined by sub-optimal activation of T cells in a PBMC pool (peripheral blood mononuclear lymphocyte). The sub-optimal activation may be determined by titrating the concentration of anti-CD3 antibody added to a PBMC pool, measuring T cell proliferation and choosing the anti-CD3 antibody concentration which results in low T cell proliferation but allows for further enhancement of the T cell proliferation. This concentration is PBMC-donor-dependent and is determined for each donor before the assay is performed.

In one embodiment, said induction or enhancement of proliferation of T cells is determined by activating T cells in PBMCs with said sub-optimal concentration of an anti-CD3 antibody, contacting the PBMCs with the multispecific antibody and determining proliferation of the T cells. In a further embodiment, the PBMCs may be labelled with CFSE, contacting the PBMCs with the multispecific antibody may be performed by incubation for 4 days, and proliferation of the T cells may be measured by flow cytometry.

Inducing a certain reaction or effect such as "inducing proliferation of T cells" may mean that there was no such reaction or effect such as proliferation of T cells before induction, but it may also mean that there was a certain level of reaction or effect such as proliferation of T cells before induction and after induction said reaction or effect such as proliferation of T cells is enhanced. Thus, "inducing" also includes "enhancing".

Proliferation of T cells may also be measured by an antigen-specific T cell proliferation assay, e.g., as described in Example 6, using a test antigen of interest. Thus, induction and/or enhancement of T cell proliferation may be measured by co-culturing T cells expressing a TCR specific for a peptide of the test antigen presented in major histocompatibility complex (MHC) and DCs presenting a corresponding peptide in MHC, which is then recognized by the TCR. For example, the T cells may be CD8+ T cells and the MHC may be MHC Class I, or the T cells may be CD4+ T cells and the MHC may be MHC Class II. T cells expressing a specific TCR may be generated by transduction with mRNA encoding the TCR. DCs presenting the corresponding peptide may be generated by transduction of the DCs with mRNA encoding the antigen. Co-culture of the TCR-positive T cells with the antigen-presenting cells induces T-cell proliferation; the extent of the proliferation may depend on the antigen density presented by the DCs and/or on the strength of the costimulatory signal. In one embodiment, proliferation of T cells may be measured by such an antigen-specific T-cell assay using CFSE labeled T-cells, adding antibodies to be tested and after 4 days measuring T cell proliferation by flow cytometry.

In one embodiment, said induction or enhancement of proliferation of T cells is determined using tumor-infiltrating lymphocytes (TILs) in an *ex vivo* expansion assay, e.g., as described in Example 11. The effect of the multispecific antibody of the invention on the induction or enhancement of proliferation of the TILs may be assessed by incubating a human tumor sample with interleukin-2 (IL-2) and said antibody, and retrieving and counting viable TILs after incubation for a period of about 10 to about 14 days. An induction or enhancement of proliferation of TILs can then be determined by comparison with a suitable control, e.g., a human tumor sample incubated without any multispecific antibody or with a reference (control) multispecific antibody. For example, a sample of human tumor tissue can be isolated, e.g., via a biopsy or from a surgical specimen, washed in serum-free medium, and tumor pieces with a diameter of about 1-2 mm placed into culture dishes or wells, e.g., 1 or 2 tumor pieces in 1 mL suitable medium, and incubated at 37°C. A suitable medium can be, for example, a serum-free medium (e.g. X-VIVO 15) supplemented with 10% Human Serum Albumin, 1% Pen/Strep, 1% Fungizone and IL-2 at a concentration ranging from 10 to 100 U/mL, e.g., 10 U/mL or 100 U/mL. The multispecific antibody can then be added at a suitable concentration in TIL medium. After a total culture period of 10-14 days, optionally splitting the cell culture if needed during this period, TILs can be harvested and counted, e.g., by flow cytometry, using, e.g., anti-CD3, anti-CD4, anti-CD8, anti-CD56 and 7-AAD antibodies to permit detection of viable CD4+ and CD8+ T cells as well as NK cells.

In one embodiment, the multispecific antibody is a bispecific antibody, which induces and/or enhances more proliferation of T cells compared to a reference bispecific antibody comprising a second antigen-binding region according to any aspect or embodiment described herein, but wherein the first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:99, 100 and 101, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:102, GVS and 103, respectively.

In one embodiment, the multispecific antibody is a bispecific antibody, which induces and/or enhances more proliferation of T cells compared to a reference bispecific antibody comprising a first antigen-binding region according to any aspect or embodiment described herein, but wherein the second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:99, 100 and 101, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:102, GVS and 103, respectively.

Binding to CD40

As described above, the multispecific antibody according to the present invention comprises a first antigen-binding region binding to human CD40.

5 In one embodiment, the multispecific antibody according to the present invention comprises a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable region CDR3 selected from the group consisting of:

- 10 a) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:3 or a sequence wherein up to four amino acids are modified in SEQ ID NO:3, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:5 or a sequence wherein up to four amino acids are modified in SEQ ID NO:5,
- b) heavy and light chain variable region CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR3 according to a) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR3 according to a).

In a further embodiment, the first antigen-binding region may further comprise a heavy chain variable region CDR1 having the sequence as set forth in SEQ ID NO:1 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:1, and/or heavy chain variable region CDR2 having the sequence as set forth in SEQ ID NO:2 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:2; and/or light chain variable region CDR1 having the sequence as set forth in SEQ ID NO:4 or a sequence wherein up to 2 amino acids are modified in SEQ ID NO:4, and/or light chain variable region CDR2 having the sequence YTS or a sequence wherein up to 2 amino acids are modified in YTS.

In one embodiment, the multispecific antibody according to the present invention comprises a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:

- 30 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively,
- b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations; and
- 35 c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy

and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b).

5 In one embodiment said first antigen-binding region comprises a first heavy chain variable (VH) sequence, and a first light chain variable (VL) sequence, wherein said variable sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively.

10 In one embodiment, said first antigen-binding region comprises a first heavy chain variable (VH) sequence, and a first light chain variable (VL) sequence, and wherein said variable sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively, and four framework sequences, FR1, FR2, FR3 and FR4, respectively.

15 In one embodiment, said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively. Thus the first antigen-binding region may comprise heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences of the CD40 antibody as set forth in **Table 1**.

20 An example of an antibody comprising such a first antigen-binding region is the chimeric antibody Chi Lob 7/4 and CD40-001 disclosed herein.

25 In another embodiment, said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively having a total of one to twelve mutations, such as one to eleven mutations, one to ten mutations, one to eight mutations, one to seven mutations, one to six mutations, one to five mutations, one to four mutations, one to three mutations, or one to two mutations.

 In one embodiment, said mutation may be an amino acid substitution, such as a conservative amino acid substitution.

30 In one embodiment, said mutations may be distributed across the VH CDR1, 2 and 3 and VL CDR 1, 2 and 3 so that each of the VH and VL CDR3 comprises at the most three mutations and each of the VH and VL CDR2 and CDR1 comprises at the most two amino acid mutations.

35 In a further embodiment, the first antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively, having a total of one to twelve mutations and wherein the VH and VL CDR3 each comprises up to three amino acid mutations, and the VH and VL CDR1 and CDR2 each comprises up to two amino acid mutations.

In a further embodiment, the first antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively, having a total of one to ten mutations, such as one to eight, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises up to two amino acid mutations.

In a further embodiment, the first antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively, having a total of one to six mutations, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises at the most one amino acid mutation.

It is well known to a person skilled in the art how to introduce mutations and that certain amino acids of the CDR sequences may be mutated; e.g. by amino acid substitutions to e.g. increase affinity of the antibody to its target antigen, reduce potential immunogenicity of non-human antibodies to be used in humans and/or to increase the yield of antibodies expressed by a host cell. Such mutations can be introduced without affecting the epitope of the target to which the antibody binds.

In another embodiment, said first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b).

In a further embodiment, said first antigen-binding region comprises heavy and light chain variable regions of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b).

The term "competes" refers in this context to the competition between two antibodies for binding to a target antigen. If two antibodies do not block each other for binding to a target antigen, such antibodies are non-competing and this is an indication that said antibodies do not bind to the same part, i.e. epitope of the target antigen. It is well known to a person skilled in the art how to test for competition of antibodies for binding to a target antigen. An example of such a method is a so-called cross-competition assay, which may e.g. be performed as an ELISA or by flow-cytometry.

For example, an ELISA-based assay may be performed by coating ELISA plate wells with each of the antibodies; adding the competing antibody and His-tagged extracellular domain of the target antigen and incubate; detecting whether the added antibody inhibited binding of the His-tagged protein to the coated antibody may be

performed by adding biotinylated anti-His antibody, followed by Streptavidin-poly-HRP, and further developing the reaction with ABTS and measuring the absorbance at 405 nm. For example a flow-cytometry assay may be performed by incubating cells expressing the target antigen with an excess of unlabeled antibody, incubating the cells with a sub-optimal concentration of biotin-labelled antibody, followed by incubation with fluorescently labeled streptavidin and analyzing by flow cytometry.

In one embodiment, said VH sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to at least one of SEQ ID NO:117 and SEQ ID NO:6, such as SEQ ID NO:117.

In one embodiment, said VL sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to at least one of SEQ ID NO:121 and SEQ ID NO:7, such as SEQ ID NO:121.

In one embodiment, said VH and VL sequence of the first antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to SEQ ID NO:6 and SEQ ID NO:7, respectively.

In one embodiment, said VH and VL sequence of the first antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to SEQ ID NO:117 and SEQ ID NO:121, respectively.

In one embodiment, the VH and VL sequences only deviate in the non-CDR sequences as set forth in SEQ ID NO:6 and 7, respectively.

In one embodiment, the VH and VL sequences only deviate in the non-CDR sequences as set forth in SEQ ID NO:117 and 121, respectively.

In one embodiment, the VH and VL sequences only deviate in the framework sequences.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of said VH and VL sequences.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework

sequences of the VH sequence as set forth in SEQ ID NO:6, and VL sequence as set forth in SEQ ID NO:7.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:6, and VL sequence as set forth in SEQ ID NO:7, and the heavy and light chain variable region CDR1, CDR2 and CDR3 of the first antigen-binding region has a total of one to twelve mutations compared to the heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively. In a further embodiment, said mutations may be as described above.

In an even further embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:6, and VL sequence as set forth in SEQ ID NO:7, and the first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:117, and VL sequence as set forth in SEQ ID NO:121, and the heavy and light chain variable region CDR1, CDR2 and CDR3 of the first antigen-binding region has a total of one to twelve mutations compared to the heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively. In a further embodiment, said mutations may be as described above.

In an even further embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first antigen-binding region has at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:117, and VL sequence as set forth in SEQ ID NO:121, and the first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:1, 2, 3, 4, YTS and 5, respectively.

In one embodiment, said VH sequence of the first antigen-binding region comprises the amino acid sequence of SEQ ID NO:117.

In one embodiment, said VL sequence of the first antigen-binding region comprises the amino acid sequence of SEQ ID NO:121.

5 In a further embodiment, said VH and VL sequences of the first antigen-binding region comprise the amino acid sequences of SEQ ID NO:117 and SEQ ID NO:121, respectively.

In one embodiment, said VH sequence of the first antigen-binding region comprises the amino acid sequence of SEQ ID NO:6.

10 In one embodiment, said VL sequence of the first antigen-binding region comprises the amino acid sequence of SEQ ID NO:7.

In a further embodiment, said VH and VL sequences of the first antigen-binding region comprises the amino acid sequences of SEQ ID NO:6 and SEQ ID NO:7, respectively.

15 In one embodiment, the multispecific antibody according to the present invention may comprise a first binding arm comprising said first antigen-binding region of any aspect or embodiment herein.

In one embodiment, the multispecific antibody according to the present invention
20 comprises a first binding arm comprising said first antigen-binding region and a first heavy chain constant sequence.

In one embodiment, the multispecific antibody according to the present invention comprises a first binding arm comprising said first antigen-binding region, wherein the first binding arm comprises a first heavy chain comprising a first heavy chain variable
25 (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence.

In one embodiment, said first light chain further comprises a first light chain constant (CL) sequence.

In a further embodiment, said first heavy chain comprises at least a hinge region,
30 a CH2 and a CH3 region.

In a specific embodiment, the multispecific antibody according to the present invention comprises a first Fab-arm comprising said first antigen-binding region.

In one embodiment, the first antigen-binding region may be derived from a
35 mouse antibody.

In one embodiment, the first antigen-binding region may be derived from a chimeric antibody, such as Chi Lob 7/4.

In one embodiment, the first antigen-binding region may be derived from a humanized antibody.

In one embodiment, the first binding arm may be derived from a full-length antibody.

5 In one embodiment, the first binding arm may be derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.

In one embodiment, the first binding arm may be derived from a monoclonal antibody.

10 In one embodiment, said first heavy chain may be of an IgG isotype, optionally selected from the group consisting of IgG1, IgG2, IgG3, and IgG4.

In one embodiment, the first binding arm may be derived from an antibody comprising a HC comprising SEQ ID NO:118 and an LC comprising SEQ ID NO:122, optionally with one or more mutations in the constant region of the HC, such as 1 to 10, such as 1 to 5, such as 1, 2, 3, 4 or 5 mutations.

15 In one embodiment, the first binding arm comprises a HC comprising SEQ ID NO:118, 119 or 120 and an LC comprising SEQ ID NO:122.

Binding to CD137

The multispecific antibody according to the present invention comprises a second antigen-binding region binding to human CD137.

20 In a further embodiment, the second antigen-binding region also binds to cynomolgus CD137.

In one embodiment, said second antigen-binding region comprises heavy and light chain variable region CDR3 selected from the group consisting of:

- 25 a) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:10 or a sequence wherein up to four amino acids are modified in SEQ ID NO:10, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:12 or a sequence wherein up to four amino acids are modified in SEQ ID NO:12 (CD137 clone 001),
- 30 b) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:17 or a sequence wherein up to four amino acids are modified in SEQ ID NO:17, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:19 or a sequence wherein up to four amino acids are modified in SEQ ID NO:19 (CD137 clone 002),
- 35 c) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:24 or a sequence wherein up to four amino acids are modified in SEQ ID NO:24, and light chain variable region CDR3 having the sequence set forth in

- SEQ ID NO:26 or a sequence wherein up to four amino acids are modified in SEQ ID NO:26 (CD137 clone 003),
- 5 d) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:31 or a sequence wherein up to four amino acids are modified in SEQ ID NO:31, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:33 or a sequence wherein up to four amino acids are modified in SEQ ID NO:33 (CD137 clone 004),
- 10 e) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:38 or a sequence wherein up to four amino acids are modified in SEQ ID NO:38, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:40 or a sequence wherein up to four amino acids are modified in SEQ ID NO:40 (CD137 clone 005),
- 15 f) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:45 or a sequence wherein up to four amino acids are modified in SEQ ID NO:45, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:47 or a sequence wherein up to four amino acids are modified in SEQ ID NO:47 (CD137 clone 006),
- 20 g) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:52 or a sequence wherein up to four amino acids are modified in SEQ ID NO:52, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:54 or a sequence wherein up to four amino acids are modified in SEQ ID NO:54 (CD137 clone 007),
- 25 h) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:59 or a sequence wherein up to four amino acids are modified in SEQ ID NO:59, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:61 or a sequence wherein up to four amino acids are modified in SEQ ID NO:61 (CD137 clone 008),
- 30 i) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:66 or a sequence wherein up to four amino acids are modified in SEQ ID NO:66, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:68 or a sequence wherein up to four amino acids are modified in SEQ ID NO:68 (CD137 clone 009),
- 35 j) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:73 or a sequence wherein up to four amino acids are modified in SEQ ID NO:73, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:75 or a sequence wherein up to four amino acids are modified in SEQ ID NO:75 (CD137 clone 010),

- 5 k) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:80 or a sequence wherein up to four amino acids are modified in SEQ ID NO:80, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:82 or a sequence wherein up to four amino acids are modified in SEQ ID NO:82 (CD137 clone 011),
- 10 l) heavy chain variable region CDR3 having the sequence set forth in SEQ ID NO:87 or a sequence wherein up to four amino acids are modified in SEQ ID NO:87, and light chain variable region CDR3 having the sequence set forth in SEQ ID NO:89 or a sequence wherein up to four amino acids are modified in SEQ ID NO:89 (CD137 clone 012), and
- 15 m) heavy and light chain variable region CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR3 according to any of a) to l) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable CDR3 according to any of a) to l).

In a further embodiment, said second antigen-binding region further comprises heavy and/or light chain region CDR1 and CDR2 selected from the group consisting of:

- 20 a) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:8 or a sequence wherein up to two amino acids are modified in SEQ ID NO:8, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:9 or a sequence wherein up to two amino acids are modified in SEQ ID NO:9, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:11 or a sequence wherein up to two amino acids are modified in SEQ ID NO:11, and/or light chain variable region CDR2 having the sequence KAS or a sequence wherein up to two amino acids are modified in KAS (CD137 clone 001),
- 25 b) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:15 or a sequence wherein up to two amino acids are modified in SEQ ID NO:15, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:16 or a sequence wherein up to two amino acids are modified in SEQ ID NO:16, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:18 or a sequence wherein up to two amino acids are modified in SEQ ID NO:18, and/or light chain variable region CDR2 having the sequence KAS or a sequence wherein up to two amino acids are modified in KAS (CD137 clone 002),
- 30 c) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:22 or a sequence wherein up to two amino acids are modified in SEQ ID NO:22, and/or heavy chain variable region CDR2 having the sequence set forth
- 35

- in SEQ ID NO:23 or a sequence wherein up to two amino acids are modified in SEQ ID NO:23, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:25 or a sequence wherein up to two amino acids are modified in SEQ ID NO:25, and/or light chain variable region CDR2 having the sequence
5 RTS or a sequence wherein up to two amino acids are modified in RTS (CD137 clone 003),
- d) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:29 or a sequence wherein up to two amino acids are modified in SEQ ID NO:29, and/or heavy chain variable region CDR2 having the sequence set forth
10 in SEQ ID NO:30 or a sequence wherein up to two amino acids are modified in SEQ ID NO:30, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:32 or a sequence wherein up to two amino acids are modified in SEQ ID NO:32, and/or light chain variable region CDR2 having the sequence GAS or a sequence wherein up to two amino acids are modified in GAS (CD137
15 clone 004),
- e) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:36 or a sequence wherein up to two amino acids are modified in SEQ ID NO:36, and/or heavy chain variable region CDR2 having the sequence set forth
20 in SEQ ID NO:37 or a sequence wherein up to two amino acids are modified in SEQ ID NO:37, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:39 or a sequence wherein up to two amino acids are modified in SEQ ID NO:39, and/or light chain variable region CDR2 having the sequence SAS or a sequence wherein up to two amino acids are modified in SAS (CD137
clone 005),
- f) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:43 or a sequence wherein up to two amino acids are modified in SEQ ID NO:43, and/or heavy chain variable region CDR2 having the sequence set forth
25 in SEQ ID NO:44 or a sequence wherein up to two amino acids are modified in SEQ ID NO:44, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:46 or a sequence wherein up to two amino acids are modified
30 in SEQ ID NO:46, and/or light chain variable region CDR2 having the sequence AAS or a sequence wherein up to two amino acids are modified in AAS (CD137 clone 006),
- g) heavy chain variable region CDR1 having the sequence set forth in SEQ ID
35 NO:50 or a sequence wherein up to two amino acids are modified in SEQ ID NO:50, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:51 or a sequence wherein up to two amino acids are modified in SEQ ID NO:51, and/or light chain variable region CDR1 having the sequence set

- forth in SEQ ID NO:53 or a sequence wherein up to two amino acids are modified in SEQ ID NO:53, and/or light chain variable region CDR2 having the sequence KAS or a sequence wherein up to two amino acids are modified in KAS (CD137 clone 007),
- 5 h) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:57 or a sequence wherein up to two amino acids are modified in SEQ ID NO:57, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:58 or a sequence wherein up to two amino acids are modified in SEQ ID NO:58, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:60 or a sequence wherein up to two amino acids are modified in SEQ ID NO:60, and/or light chain variable region CDR2 having the sequence RAS or a sequence wherein up to two amino acids are modified in RAS (CD137 clone 008),
- 10 i) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:64 or a sequence wherein up to two amino acids are modified in SEQ ID NO:64, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:65 or a sequence wherein up to two amino acids are modified in SEQ ID NO:65, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:67 or a sequence wherein up to two amino acids are modified in SEQ ID NO:67, and/or light chain variable region CDR2 having the sequence GAS or a sequence wherein up to two amino acids are modified in GAS (CD137 clone 009),
- 15 j) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:71 or a sequence wherein up to two amino acids are modified in SEQ ID NO:71, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:72 or a sequence wherein up to two amino acids are modified in SEQ ID NO:72, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:74 or a sequence wherein up to two amino acids are modified in SEQ ID NO:74, and/or light chain variable region CDR2 having the sequence KAS or a sequence wherein up to two amino acids are modified in KAS (CD137 clone 010),
- 20 k) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:78 or a sequence wherein up to two amino acids are modified in SEQ ID NO:78, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:79 or a sequence wherein up to two amino acids are modified in SEQ ID NO:79, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:81 or a sequence wherein up to two amino acids are modified in SEQ ID NO:81, and/or light chain variable region CDR2 having the sequence
- 25
- 30
- 35

DTS or a sequence wherein up to two amino acids are modified in DTS (CD137 clone 011), and

- 5 l) heavy chain variable region CDR1 having the sequence set forth in SEQ ID NO:85 or a sequence wherein up to two amino acids are modified in SEQ ID NO:85, and/or heavy chain variable region CDR2 having the sequence set forth in SEQ ID NO:86 or a sequence wherein up to two amino acids are modified in SEQ ID NO:86, and/or light chain variable region CDR1 having the sequence set forth in SEQ ID NO:88 or a sequence wherein up to two amino acids are modified in SEQ ID NO:88, and/or light chain variable region CDR2 having the sequence
- 10 SAS or a sequence wherein up to two amino acids are modified in SAS (CD137 clone 012).

In one embodiment said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:

- 15 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:8, 9 and 10, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:11, KAS and 12, respectively, (CD137 clone 001),
- 20 b) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:15, 16 and 17, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:18, KAS and 19, respectively, (CD137 clone 002),
- 25 c) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:22, 23, and 24, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:25, RTS and 26, respectively, (CD137 clone 003),
- 30 d) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:29, 30 and 31, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:32, GAS and 33, respectively, (CD137 clone 004),
- e) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005),
- 35 f) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:43, 44 and 45, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:46, AAS, and 47, respectively, (CD137 clone 006),

- g) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:50, 51 and 52, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:53, KAS and 54, respectively, (CD137 clone 007),
- 5 h) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:57, 58 and 59, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:60, RAS and 61, respectively, (CD137 clone 008),
- 10 i) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009),
- j) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:71, 72 and 73, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:74, KAS and 75, respectively, (CD137 clone 010),
- 15 k) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:78, 79 and 80, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:81, DTS and 82, respectively, (CD137 clone 011),
- 20 l) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:85, 86 and 87, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:88, SAS and 89, respectively, (CD137 clone 012),
- 25 m) heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to l) having a total of one to twelve mutations, and
- n) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to m) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to m).
- 30

Thus, the second antigen-binding region may comprise the heavy and light chain variable region CDR1, CDR2 and CDR3 sequences of a CD137 antibody as set forth in

35 **Table 1**; i.e. CD137 clone 001, CD137 clone 002, CD137 clone 003, CD137 clone 004, CD137 clone 005, CD137 clone 006, CD137 clone 007, CD137 clone 008, CD137 clone 009, CD137 clone 010, CD137 clone 011 or CD137 clone 012. In particular, the second antigen-binding region may comprise the heavy and light chain variable region CDR1,

CDR2 and CDR3 sequences from the same CD137 antibody clone, optionally wherein the framework regions are primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence.

5 In a further embodiment, the second antigen-binding region comprises heavy and light chain variable regions of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to m) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to m).

10 In one embodiment, said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to a higher degree than it binds to a mutant human CD137 (SEQ ID NO:93). The mutant human CD137 of SEQ ID NO:93 is also referred to as shuffle 6 herein.

15 In another embodiment, said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to a higher degree than it binds to a mutant human CD137 (SEQ ID NO:94). The mutant human CD137 of SEQ ID NO:94 is also referred to as shuffle 5 herein.

20 In a further embodiment, said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to the same degree that it binds to a mutant human CD137 (SEQ ID NO:95). The mutant human CD137 of SEQ ID NO:95 is also referred to as shuffle 4 herein.

25 In the context of the present invention "to a higher degree" means that the affinity of the second antigen-binding region is higher for human CD137 (SEQ ID NO:92) than for a mutant human CD137 (SEQ ID NO:93 and 94, shuffle 6 and 5 respectively). If there is no binding to the mutant CD137, the affinity for human CD137 will be infinitely higher than for said mutant CD137. However, in case of binding to said mutant CD137 the affinity may be 2-fold, such as 3-fold, or 4-fold, or 5-fold, or 6-fold higher for human CD137 than for the respective mutant CD137.

30 In the context of the present invention "to the same degree" means that the affinity of the second antigen-binding region is similar for human CD137 (SEQ ID NO:92) and for a mutant human CD137 (SEQ ID NO:95, shuffle 4). In particular, "similar" in this context may mean that the affinity for human CD137 and for said mutant CD137 differs at the most by 2.5-fold, such as 2.2-fold, or 2.0-fold, or 1.8-fold, or 1.75-fold or 1.5-fold.

35 The mutant human CD137 in SEQ ID NO:93 corresponds to the amino acid sequence of human CD137 wherein amino acids 24-47 (shuffle 6) were replaced by the corresponding amino acids from wild boar CD137.

Thus, in one embodiment, the second antigen-binding region binds to an epitope of human CD137 which comprises or requires one or more of the amino acids L, Q, D, P, C, S, N, C, P, A, G, T, F, C, D, N, N, R, N, Q, I, C, S and P at positions 24-47 of SEQ ID NO:92 (corresponding to SEQ ID NO:129).

5 The mutant human CD137 in SEQ ID NO:94 corresponds to the amino acid sequence of human CD137 wherein amino acids 48-88 (shuffle 5) were replaced by the corresponding amino acids from African elephant CD137.

10 Thus, in one embodiment, the second antigen-binding region binds to an epitope of human CD137 which comprises or requires one or more of the amino acids C, P, P, N, S, F, S, S, A, G, G, Q, R, T, C, D, I, C, R, Q, C, K, G, V, F, R, T, R, K, E, C, S, S, T, S, N, A, E, C, D and C at positions 48-88 of SEQ ID NO:92 (corresponding to SEQ ID NO:130).

The mutant human CD137 in SEQ ID NO:95 corresponds to the amino acid sequence of human CD137 wherein amino acids 59-114 (shuffle 4) were replaced by the corresponding amino acids from African elephant CD137.

15 Thus, in one embodiment, the second antigen-binding region does not bind to an epitope of human CD137 which comprises or requires one or more of the amino acids T, P, G, F, H, C, L, G, A, G, C, S, M, C, E, Q, D, C, K, Q, G, Q, E, L, T and K 89-114 at positions of SEQ ID NO:92 (corresponding to SEQ ID NO:131).

20 In one embodiment, binding to the mutant and human CD137 may be performed as the shuffle assay described in Example 2. Thus, binding to human CD137 (SEQ ID NO:92) and mutant human CD137 (SEQ ID NOs:93, 94 and 95) may be determined by preparing shuffle constructs derived from human CD137 in which protein domains of the human CD137 are replaced by the corresponding domain of CD137 from different species, using human CD137 and the different species of CD137 as reference constructs;
25 transducing cells with plasmids encoding the reference construct or the shuffle constructs, respectively, and measuring binding of the antibody to each these CD137 constructs by flow cytometry, such as FACS.

30 Loss of binding to certain shuffle constructs indicates that the corresponding region is likely to be involved in the antibody epitope. Thus, protein domains of human CD137 contributing to the epitope of the anti-human CD137 antibodies may thereby be determined by the shuffle assay. The different species of CD137 used to create the shuffle constructs should be chosen so that the monoclonal anti-human CD137 antibodies do not bind to the whole CD137 protein from these different species (reference construct).

35 Determination of binding to human CD137 and mutants thereof may in particular be performed with a monoclonal antibody comprising two second antigen-binding regions according to the present invention.

In one embodiment, said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:

- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:8, 9 and 10, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:11, KAS and 12, respectively, (CD137 clone 001),
- b) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:15, 16 and 17, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:18, KAS and 19, respectively, (CD137 clone 002),
- c) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005),
- d) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:43, 44 and 45, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:46, AAS and 47, respectively, (CD137 clone 006),
- e) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009),
- f) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:71, 72 and 73, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:74, KAS and 75, respectively, (CD137 clone 010),
- g) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:85, 86 and 87, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:88, SAS and 89, respectively, (CD137 clone 012),
- h) heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to g) having a total of one to twelve mutations, and
- i) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to h) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to any of a) to h).

Hence, in one embodiment, said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:

- 5 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009),
- b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations;
- 10 c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b).

15

In another embodiment, said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:

- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005),
- 20 b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations, and
- c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b).

25

30 In a particular embodiment, said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively. An example of such an antibody includes, but is not limited to, the antibody referred to herein as
35 CD137 clone 009.

In another embodiment, said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3

having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively. An example of such an antibody includes, but is not limited to, the antibody referred to herein as CD137 clone 005.

5 In another embodiment, said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009), having a total of one to twelve mutations, such as one to ten mutations, or one to eight mutations, or one to six mutations, or one to four mutations, or to two
10 mutations.

In another embodiment, said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137
15 clone 005), having a total of one to twelve mutations, such as one to ten mutations, or one to eight mutations, or one to six mutations, or one to four mutations, or one to two mutations.

In one embodiment, said mutation may be an amino acid substitution, such as a conservative amino acid substitution.

20 In one embodiment, said mutations may be distributed across the VH CDR1, 2 and 3 and VL CDR 1, 2 and 3 so that each of the VH and VL CDR3 comprises at the most three mutations and each of the VH and VL CDR2 and CDR1 comprises at the most two amino acid modifications.

Hence, in a further embodiment, the second antigen-binding region comprises
25 heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively, (CD137 clone 009), having a total of one to twelve mutations and wherein the VH and VL CDR3 each comprises up to three amino acid modifications, and the VH and VL CDR1 and CDR2 each comprises up to two amino acid modifications.

30 In a further embodiment, the second antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively, (CD137 clone 009), having a total of one to ten mutations, such as one to eight, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises up to two amino acid modifications.

35 In a further embodiment, the second antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively, (CD137 clone 009), having a total of one to six

mutations, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises at most one amino acid modification.

In another embodiment, the second antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37, 38, 39, SAS and 40, respectively, (CD137 clone 005), having a total of one to twelve mutations and wherein the VH and VL CDR3 each comprises up to three amino acid modifications, and the VH and VL CDR1 and CDR2 each comprises up to two amino acid modifications.

In a further embodiment, the second antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37, 38, 39, SAS and 40, respectively, (CD137 clone 005), having a total of one to ten mutations, such as one to eight, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises up to two amino acid modifications.

In a further embodiment, the second antigen-binding region comprises heavy and light chain CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37, 38, 39, SAS and 40, respectively, (CD137 clone 005), having a total of one to six mutations, and wherein the VH and VL CDR1, CDR2, and CDR3 each comprises at most one amino acid modification.

In a further embodiment, there may be a total of one to twelve mutations; such as one to ten mutations, or one to eight mutations, or one to six mutations, or one to four mutations, or one to two mutations; and each CDR sequence comprises at the most two amino acid substitutions.

It is well known to a person skilled in the art how to introduce mutations and that certain amino acids of the CDR sequences may be mutated; e.g., by amino acid substitutions to, e.g., increase affinity of the antibody to its target antigen or reducing immunogenicity for non-human antibodies to be used for treatment of humans. Such mutations can be introduced without affecting the epitope of the target antigen to which the antibody binds.

In one embodiment said second antigen-binding region comprises a second heavy chain variable (VH) sequence, and a second light chain variable (VL) sequence and wherein said variable sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively.

In one embodiment, said second antigen-binding region comprises a second heavy chain variable (VH) sequence, and a second light chain variable (VL) sequence and wherein said variable sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively, and four framework sequences, FR1, FR2, FR3 and FR4, respectively.

In one embodiment, said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to an amino acid sequence selected from the group consisting of:

- 5 a) a VH sequence as set forth in SEQ ID NO:123 (humanized CD137 clone 009)
- b) a VH sequence as set forth in SEQ ID NO:13 (CD137 clone 001)
- c) a VH sequence as set forth in SEQ ID NO:20 (CD137 clone 002)
- d) a VH sequence as set forth in SEQ ID NO:27 (CD137 clone 003)
- e) a VH sequence as set forth in SEQ ID NO:34 (CD137 clone 004)
- 10 f) a VH sequence as set forth in SEQ ID NO:41 (CD137 clone 005)
- g) a VH sequence as set forth in SEQ ID NO:48 (CD137 clone 006)
- h) a VH sequence as set forth in SEQ ID NO:55 (CD137 clone 007)
- i) a VH sequence as set forth in SEQ ID NO:62 (CD137 clone 008)
- j) a VH sequence as set forth in SEQ ID NO:69 (CD137 clone 009)
- 15 k) a VH sequence as set forth in SEQ ID NO:76 (CD137 clone 010)
- l) a VH sequence as set forth in SEQ ID NO:83 (CD137 clone 011)
- m) a VH sequence as set forth in SEQ ID NO:90 (CD137 clone 012)

In one embodiment, said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to an amino acid sequence selected from the group consisting of:

- a) a VL sequence as set forth in SEQ ID NO:127 (humanized CD137 clone 009)
- b) a VL sequence as set forth in SEQ ID NO:14 (CD137 clone 001)
- 25 c) a VL sequence as set forth in SEQ ID NO:21 (CD137 clone 002)
- d) a VL sequence as set forth in SEQ ID NO:28 (CD137 clone 003)
- e) a VL sequence as set forth in SEQ ID NO:35 (CD137 clone 004)
- f) a VL sequence as set forth in SEQ ID NO:42 (CD137 clone 005)
- g) a VL sequence as set forth in SEQ ID NO:49 (CD137 clone 006)
- 30 h) a VL sequence as set forth in SEQ ID NO:56 (CD137 clone 007)
- i) a VL sequence as set forth in SEQ ID NO:63 (CD137 clone 008)
- j) a VL sequence as set forth in SEQ ID NO:70 (CD137 clone 009)
- k) a VL sequence as set forth in SEQ ID NO:77 (CD137 clone 010)
- l) a VL sequence as set forth in SEQ ID NO:84 (CD137 clone 011)
- 35 m) a VL sequence as set forth in SEQ ID NO:91 (CD137 clone 012)

In one embodiment, said VH and VL sequences of the second antigen-binding region each comprise an amino acid sequence having at least 70%, at least 75%, at

least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to an amino acid sequence selected from the group consisting of:

- a) a VH sequence as set forth in SEQ ID NO:123 and a VL sequence as set forth in SEQ ID NO:127 (humanized CD137 clone 009)
- 5 b) a VH sequence as set forth in SEQ ID NO:13 and a VL sequence as set forth in SEQ ID NO:14 (CD137 clone 001)
- c) a VH sequence as set forth in SEQ ID NO:20 and a VL sequence as set forth in SEQ ID NO:21 (CD137 clone 002)
- d) a VH sequence as set forth in SEQ ID NO:27 and a VL sequence as set forth in SEQ ID NO:28 (CD137 clone 003)
- 10 e) a VH sequence as set forth in SEQ ID NO:34 and a VL sequence as set forth in SEQ ID NO:35 (CD137 clone 004)
- f) a VH sequence as set forth in SEQ ID NO:41 and a VL sequence as set forth in SEQ ID NO:42 (CD137 clone 005)
- 15 g) a VH sequence as set forth in SEQ ID NO:48 and a VL sequence as set forth in SEQ ID NO:49 (CD137 clone 006)
- h) a VH sequence as set forth in SEQ ID NO:55 and a VL sequence as set forth in SEQ ID NO:56 (CD137 clone 007)
- i) a VH sequence as set forth in SEQ ID NO:62 and a VL sequence as set forth in SEQ ID NO:63 (CD137 clone 008)
- 20 j) a VH sequence as set forth in SEQ ID NO:69 and a VL sequence as set forth in SEQ ID NO:70 (CD137 clone 009)
- k) a VH sequence as set forth in SEQ ID NO:76 and a VL sequence as set forth in SEQ ID NO:77 (CD137 clone 010)
- 25 l) a VH sequence as set forth in SEQ ID NO:83 and a VL sequence as set forth in SEQ ID NO:84 (CD137 clone 011)
- m) a VH sequence as set forth in SEQ ID NO:90 and a VL sequence as set forth in SEQ ID NO:91 (CD137 clone 012).

30 In one embodiment, said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to an amino acid sequence selected from the group consisting of:

- a) a VH sequence as set forth in SEQ ID NO:123 (humanized CD137 clone 009)
- 35 b) a VH sequence as set forth in SEQ ID NO:41 (CD137 clone 005)
- c) a VH sequence as set forth in SEQ ID NO:69 (CD137 clone 009)

In one embodiment, said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to an amino acid sequence selected from the group consisting of:

- 5 a) a VL sequence as set forth in SEQ ID NO:127 (humanized CD137 clone 009)
- b) a VL sequence as set forth in SEQ ID NO:42 (CD137 clone 005)
- c) a VL sequence as set forth in SEQ ID NO:70 (CD137 clone 009)

10 In one embodiment, said VH and VL sequences of the second antigen-binding region each comprise an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to an amino acid sequence selected from the group consisting of:

- a) a VH sequence as set forth in SEQ ID NO:123 and a VL sequence as set forth in SEQ ID NO:127 (humanized CD137 clone 009)
- 15 b) a VH sequence as set forth in SEQ ID NO:41 and a VL sequence as set forth in SEQ ID NO:42 (CD137 clone 005)
- c) a VH sequence as set forth in SEQ ID NO:69 and a VL sequence as set forth in SEQ ID NO:70 (CD137 clone 009)

20 In one embodiment, said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:41 (CD137 clone 005).

25 In one embodiment, said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:69 (CD137 clone 009).

30 In one embodiment, said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NO:123 (humanized CD137 clone 009).

35 In one embodiment, said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:42 (CD137 clone 005).

In one embodiment, said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at

least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:70 (CD137 clone 009).

In one embodiment said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NO:127 (humanized CD137 clone 009).

In one embodiment, said VH and said VL sequence of the second antigen-binding region each comprise an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:41; and SEQ ID NO:42 (CD137 clone 005), respectively.

In one embodiment, said VH and VL sequence of the second antigen-binding region each comprise an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:69 and SEQ ID NO:70 (CD137 clone 009), respectively.

In one embodiment, said VH and VL sequence of the second antigen-binding region each comprise an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NO:123 and SEQ ID NO:127, respectively (humanized CD137 clone 009).

In one embodiment, said VH and VL sequences only deviate in the framework sequences.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first and/or second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of said VH and VL sequences.

In one embodiment, the VH and VL sequences only deviate in the non-CDR sequences as set forth in SEQ ID NO:41 and 42, respectively, (CD137 clone 005).

In one embodiment the VH and VL sequences only deviate in the non-CDR sequences as set forth in SEQ ID NO:69 and 70, respectively, (CD137 clone 009).

In one embodiment the VH and VL sequences only deviate in the non-CDR sequences as set forth in SEQ ID NO:123 and 127, respectively, (humanized CD137 clone 009).

In one embodiment, the VH and VL sequences only deviate in the framework sequences.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first and/or second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at

least 97%, at least 99%, or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of said VH and VL sequences.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:41, and VL sequence as set forth in SEQ ID NO:42.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:41, and VL sequence as set forth in SEQ ID NO:42, and the heavy and light chain variable region CDR1, CDR2 and CDR3 of the second antigen-binding region have a total of one to twelve mutations compared to the heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:36, 37, 38, 39, SAS and 40, respectively. In a further embodiment said mutations may be as described above.

In an even further embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:41, and VL sequence as set forth in SEQ ID NO:42, and the second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:36, 37, 38, 39, SAS and 40, respectively.

In one embodiment the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:69, and VL sequence as set forth in SEQ ID NO:70.

In one embodiment the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:69, and VL sequence as set forth in SEQ ID NO:70, and the heavy and light chain variable region

CDR1, CDR2 and CDR3 of the second antigen-binding region have a total of one to twelve mutations compared to the heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively. In a further embodiment said mutations may be as described above.

5 In an even further embodiment the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:69, and VL
10 sequence as set forth in SEQ ID NO:70, and the second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively.

In one embodiment, the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at
15 least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:123, and VL sequence as set forth in SEQ ID NO:127.

In one embodiment the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at least 70%, at
20 least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:123, and VL sequence as set forth in SEQ ID NO:127, and the heavy and light chain variable region
25 CDR1, CDR2 and CDR3 of the second antigen-binding region have a total of one to twelve mutations compared to the heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively. In a further embodiment said mutations may be as described above.

In an even further embodiment the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the second antigen-binding region have at
30 least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework sequences of the VH sequence as set forth in SEQ ID NO:123, and VL sequence as set forth in SEQ ID NO:127, and the second antigen-
35 binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 having the sequences as set forth in SEQ ID NOs:64, 65, 66, 67, GAS and 68, respectively.

In one embodiment, said VH sequence of the second antigen-binding region comprises SEQ ID NO:123 (humanized CD137 clone 009).

In one embodiment, said VL sequence of the second antigen-binding region comprises SEQ ID NO:127 (humanized CD137 clone 009).

5 In one embodiment, said VH and VL sequences of the second antigen-binding region comprise SEQ ID NO:123 and SEQ ID NO:127, respectively.

In one embodiment, said VH sequence of the second antigen-binding region comprises a VH sequence selected from the group consisting of:

- a) SEQ ID NO:41 (CD137 clone 005)
- 10 b) SEQ ID NO:69 (CD137 clone 009)

In one embodiment, said VL sequence of the second antigen-binding region comprises a VL sequence selected from the group consisting of:

- a) SEQ ID NO:42 (CD137 clone 005)
- b) SEQ ID NO:70 (CD137 clone 009)

15 In one embodiment, said VH and VL sequences of the second antigen-binding are selected from the group consisting of:

- a) a VH sequence as set forth in SEQ ID NO:41 and a VL sequence as set forth in SEQ ID NO:42 (CD137 clone 005),
- b) a VH sequence as set forth in SEQ ID NO:69 and a VL sequence as set forth in
- 20 SEQ ID NO:70 (CD137 clone 009).

In one embodiment, the multispecific antibody according to the present invention comprises a second binding arm comprising said second antigen-binding region.

25 In one embodiment, the multispecific antibody according to the present invention comprises a second binding arm comprising said second-antigen-binding region and a second heavy chain constant sequence.

In one embodiment, the multispecific antibody according to the present invention comprises a second binding arm comprising said second antigen-binding region, wherein the second binding arm comprises a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a

30 second light chain comprising a second light chain variable (VL) sequence.

In one embodiment said second light chain further comprises a second light chain constant sequence.

In a further embodiment, said second heavy chain comprises at least a hinge

35 region, a CH2 and a CH3 region.

In a specific embodiment, the multispecific antibody according to the present invention comprises a second Fab-arm comprising said second antigen-binding region.

In one embodiment, the second antigen-binding region is derived from a rabbit antibody, such as any of anti-CD137 clones 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12, in particular any of clones 5 and 9, disclosed herein.

5 In one embodiment, the second antigen-binding region is derived from a chimeric antibody, such as an antibody comprising a variable region from any of the anti-CD137 clones 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 and 12, in particular any of clones 5 and 9, disclosed herein.

In one embodiment, the second antigen-binding region is derived from a humanized antibody.

10 In one embodiment, the second binding arm is derived from a full-length antibody.

In one embodiment the second binding arm is derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.

15 In one embodiment, the second binding arm is derived from a monoclonal antibody.

In one embodiment, said second heavy chain is of an IgG isotype, optionally having a subclass selected from the group consisting of IgG1, IgG2, IgG3, and IgG4.

20 In one embodiment, the first binding arm may be derived from an antibody comprising a HC comprising SEQ ID NO:124 and an LC comprising SEQ ID NO:128, optionally with one or more mutations in the constant region of the HC, such as 1 to 10, such as 1 to 5, such as 1, 2, 3, 4 or 5 mutations.

In one embodiment, the first binding arm comprises a HC comprising SEQ ID NO:124, 125 or 126 and an LC comprising SEQ ID NO:128.

Binding to CD40 and CD137

25 In some embodiments, the present invention relates to a multispecific antibody comprising:

(I) a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable regions selected from the group consisting of:

- 30 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively,
- b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a)
- 35 having a total of one to twelve mutations; and

- 5 c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b), and
- (II) a second antigen-binding region binding to human CD137, wherein said second antigen-binding region comprises heavy and light chain variable regions selected from the group consisting of:
- 10 x) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, 68 and 69, respectively, (CD137 clone 009),
- y) heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) having a total of one to twelve mutations; and
- 15 z) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) or y) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) or y).
- 20

In another embodiment, the present invention relates to a multispecific antibody comprising:

- (I) a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable regions selected from the group consisting of:
- 25 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, 5 and 6, respectively,
- 30 b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations; and
- c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b), and
- 35

(II) a second antigen-binding region binding to human CD137, wherein said second antigen-binding region comprises heavy and light chain variable regions selected from the group consisting of:

- 5 x) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005),
- y) heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) having a total of one to twelve mutations; and
- 10 z) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) or y) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to x) or y).

15

Hence, in one embodiment, said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:4, YTS and 5, respectively; and said second antigen-binding region comprises

- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively (CD136 clone 009), or
- 25 b) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40 (CD137 clone 005), respectively.

30 In another embodiment, said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively; and said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009).

35

In another embodiment said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively; and said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005).

In a further embodiment, said first antigen-binding region of the multispecific antibody according to the present invention comprises a first heavy chain variable (VH) sequence, and a first light chain variable (VL) sequence, and said second antigen-binding region of the multispecific antibody according to the present invention comprises a second heavy chain variable (VH) sequence, and a second light chain variable (VL) sequence and wherein said variable sequences each comprise three CDR sequences, CDR1, CDR2 and CDR3, respectively, and four framework sequences, FR1, FR2, FR3 and FR4, respectively.

In a further embodiment said VH and VL sequence of the first antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:6 and the VL sequence as set forth in SEQ ID NO:7, respectively, and said VH and said VL sequence of the second antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:41 and the VL sequence as set forth in SEQ ID NO:42, respectively, (CD137 clone 005).

In another further embodiment, said VH and VL sequence of the first antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:6 and the VL sequence as set forth in SEQ ID NO:7, respectively, and said VH and said VL sequence of the second antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:69 and the VL sequence as set forth in SEQ ID NO:70, respectively, (CD137 clone 009).

In another further embodiment, said VH and VL sequence of the first antigen-binding region each comprises a sequence having at least 70%, at least 75%, at least

80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:117 and the VL sequence as set forth in SEQ ID NO:121, respectively, and said VH and said VL sequence of the second antigen-binding region each comprise a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:123 and the VL sequence as set forth in SEQ ID NO:127, respectively, (humanized CD137 clone 009).

10 In a particular embodiment, the present invention relates to a bispecific antibody comprising

(I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain constant (CL) sequence, and wherein said heavy first chain variable sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and said first light chain sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively, and

(II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence, and a second light chain variable (VL) sequence, wherein said second heavy chain variable sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and said second light chain sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, GAS and 68, respectively, (CD137 clone 009);

wherein the first and second heavy chain are of a human IgG1 isotype and wherein the first and second light chain is of IgG1, κ , and wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and second constant heavy chain are F, E, and A, respectively, and wherein (a) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is R; or (b) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is L.

In a further particular embodiment, the present invention relates to a bispecific antibody comprising

- (I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain constant (CL) sequence, and wherein said heavy first chain variable sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and said first light chain sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively, and
- (II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence, and a second light chain variable (VL) sequence, wherein said second heavy chain variable sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and said second light chain sequence comprises CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005); and

wherein the first and second heavy chain are of a human IgG1 isotype and wherein the first and second light chain is of IgG1, κ , and wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and second constant heavy chain are F, E, and A, respectively, and wherein (a) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is R; or (b) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is L.

In a specific embodiment, the present invention relates to a bispecific antibody comprising

- (I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain constant (CL) sequence, and wherein said first VH and VL sequences each comprises a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence

of the VH sequence as set forth in SEQ ID NO:117 and the VL sequence as set forth in SEQ ID NO:121, respectively, and

(II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence, and a second light chain variable (VL) sequence, wherein said second VH and VL sequences comprise a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:123 and the VL sequence as set forth in SEQ ID NO:127, respectively, (humanized CD137 clone 009).

In a further specific embodiment, the present invention relates to a bispecific antibody comprising

(I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain constant (CL) sequence, and wherein said first VH sequence comprises SEQ ID NO:117 and said first VL sequence comprises SEQ ID NO:121, and

(II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence, and a second light chain variable (VL) sequence, wherein said second VH sequence comprises SEQ ID NO:123 and said second VL sequence comprises SEQ ID NO:127, (humanized CD137 clone 009).

In a specific embodiment, the present invention relates to a bispecific antibody comprising

(I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain constant (CL) sequence, and wherein said first VH and VL sequences comprise a sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:117 and the VL sequence as set forth in SEQ ID NO:121, respectively, and

(II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence, and a second light chain variable (VL) sequence, wherein said second VH and VL sequences comprise a sequence having at least 70%, at least 75%, at least 80%, at

least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to the amino acid sequence of the VH sequence as set forth in SEQ ID NO:123 and the VL sequence as set forth in SEQ ID NO:127, respectively, (humanized CD137 clone 009),

wherein the first and second heavy chain are of a human IgG1 isotype and
5 wherein the first and second light chain is of IgG1, κ , and wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and second constant heavy chain are F, E, and A, respectively, and wherein (a) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is L, and
10 the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is R; or (b) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is L.

15 In a further specific embodiment, the present invention relates to a bispecific antibody comprising

(I) a first binding arm comprising a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence and a first light chain
20 constant (CL) sequence, and wherein said first VH sequence comprises SEQ ID NO:117 and said first VL sequence comprises SEQ ID NO:121, and

(II) a second binding arm comprising a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain further comprises a second light chain constant (CL) sequence,
25 and a second light chain variable (VL) sequence, wherein said second VH sequence comprises SEQ ID NO:123 and said second VL sequence comprises SEQ ID NO:127, (humanized CD137 clone 009),

wherein the first and second heavy chain are of a human IgG1 isotype and wherein the first and second light chain is of IgG1, κ , and wherein the positions
30 corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and second constant heavy chain are F, E, and A, respectively, and wherein (a) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU
35 numbering of the second constant heavy chain is R; or (b) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first constant heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second constant heavy chain is L.

In another aspect, the present invention relates to a bispecific antibody comprising a first binding arm binding to human CD40 and a second binding arm binding to human CD137, wherein

5 (i) said first binding arm comprises a heavy chain (HC) amino acid sequence comprising or consisting of SEQ ID NO:118 and a light chain (LC) amino acid sequence comprising or consisting of SEQ ID NO:122, and

(ii) said second binding arm comprises a HC amino acid sequence comprising or consisting of SEQ ID NO:124 and a LC amino acid sequence comprising or consisting of SEQ ID NO:128,

10 optionally wherein SEQ ID NOS:118, SEQ ID NO:124 or both comprise one or more mutations in the constant region of the HC, such as 1 to 10, such as 1 to 5, such as 1, 2, 3, 4 or 5 mutations.

In another aspect, the present invention relates to a bispecific antibody comprising a first binding arm binding to human CD40 and a second binding arm binding
15 to human CD137, wherein

(i) said first binding arm comprises a HC amino acid sequence comprising or consisting of SEQ ID NO:119 and a LC amino acid sequence comprising or consisting of SEQ ID NO:122, and

20 (ii) said second binding arm comprises a HC amino acid sequence comprising or consisting of SEQ ID NO:125 and a LC amino acid sequence comprising or consisting of SEQ ID NO:128.

In another aspect, the present invention relates to a bispecific antibody comprising a first binding arm binding to human CD40 and a second binding arm binding to human CD137, wherein

25 (i) said first binding arm comprises a HC amino acid sequence comprising or consisting of SEQ ID NO:120 and a LC amino acid sequence comprising or consisting of SEQ ID NO:122, and

30 (ii) said second binding arm comprises a HC amino acid sequence comprising or consisting of SEQ ID NO:126 and a LC amino acid sequence comprising or consisting of SEQ ID NO:128.

Bispecific formats

In a particular embodiment the multispecific antibody according to the present invention is a bispecific antibody.

35 The present invention provides bispecific CD40xCD137 antibodies which are able of cross-linking cells expressing CD40 and cells expressing CD137; such as antigen-presenting cells and T cells, respectively. Depending on the desired functional properties

for a particular use, particular antigen-binding regions can be selected from the set of antibodies or antigen-binding regions provided by the present invention. Many different formats and uses of bispecific antibodies are known in the art, and were reviewed by Kontermann; Drug Discov Today, 2015 Jul;20(7):838-47 and; MAbs, 2012 Mar-Apr;4(2):182-97.

A bispecific antibody according to the present invention is not limited to any particular bispecific format or method of producing it.

Examples of bispecific antibody molecules which may be used in the present invention comprise (i) a single antibody that has two binding arms comprising different antigen-binding regions; (ii) a single chain antibody that has specificity to two different epitopes, e.g., via two scFvs linked in tandem by an extra peptide linker; (iii) a dual-variable-domain antibody (DVD-Ig), where each light chain and heavy chain contains two variable domains in tandem through a short peptide linkage (Wu et al., Generation and Characterization of a Dual Variable Domain Immunoglobulin (DVD-Ig™) Molecule, In: Antibody Engineering, Springer Berlin Heidelberg (2010)); (iv) a chemically-linked bispecific (Fab')₂ fragment; (v) a Tandab, which is a fusion of two single chain diabodies resulting in a tetravalent bispecific antibody that has two binding sites for each of the target antigens; (vi) a flexibody, which is a combination of scFvs with a diabody resulting in a multivalent molecule; (vii) a so-called "dock and lock" molecule, based on the "dimerization and docking domain" in Protein Kinase A, which, when applied to Fabs, can yield a trivalent bispecific binding protein consisting of two identical Fab fragments linked to a different Fab fragment; (viii) a so-called Scorpion molecule, comprising, e.g., two scFvs fused to both termini of a human Fab-arm; and (ix) a diabody.

In one embodiment, the bispecific antibody of the present invention is a diabody, a cross-body, or a bispecific antibody obtained via a controlled Fab-arm exchange (such as described in WO2011131746 (Genmab)).

Examples of different classes of bispecific antibodies include, but are not limited to, (i) IgG-like molecules with complementary CH3 domains to force heterodimerization; (ii) recombinant IgG-like dual targeting molecules, wherein the two sides of the molecule each contain the Fab fragment or part of the Fab fragment of at least two different antibodies; (iii) IgG fusion molecules, wherein full length IgG antibodies are fused to extra Fab fragment or parts of Fab fragment; (iv) Fc fusion molecules, wherein single chain Fv molecules or stabilized diabodies are fused to heavy-chain constant-domains, Fc-regions or parts thereof; (v) Fab fusion molecules, wherein different Fab-fragments are fused together, fused to heavy-chain constant-domains, Fc-regions or parts thereof; and (vi) ScFv- and diabody-based and heavy chain antibodies (e.g., domain antibodies, nanobodies) wherein different single chain Fv molecules or different diabodies or different heavy-chain antibodies (e.g. domain antibodies, nanobodies) are fused to each

other or to another protein or carrier molecule fused to heavy-chain constant-domains, Fc-regions or parts thereof.

Examples of IgG-like molecules with complementary CH3 domain molecules include, but are not limited to, the Triomab/Quadroma molecules (Trion Pharma/Fresenius Biotech; Roche, WO2011069104), the so-called Knobs-into-Holes molecules (Genentech, WO9850431), CrossMAbs (Roche, WO2011117329) and the electrostatically-matched molecules (Amgen, EP1870459 and WO2009089004; Chugai, US201000155133; Oncomed, WO2010129304), the LUZ-Y molecules (Genentech, Wranik et al. J. Biol. Chem. 2012, 287(52): 43331-9, doi: 10.1074/jbc.M112.397869. Epub 2012 Nov 1), DIG-body and PIG-body molecules (Pharmabicine, WO2010134666, WO2014081202), the Strand Exchange Engineered Domain body (SEEDbody) molecules (EMD Serono, WO2007110205), the Biclonics molecules (Merus, WO2013157953), Fc Δ Adp molecules (Regeneron, WO201015792), bispecific IgG1 and IgG2 molecules (Pfizer/Rinat, WO11143545), Azymetric scaffold molecules (Zymeworks/Merck, WO2012058768), mAb-Fv molecules (Xencor, WO2011028952), bivalent bispecific antibodies (WO2009080254) and the DuoBody® molecules (Genmab A/S, WO2011131746).

Examples of recombinant IgG-like dual targeting molecules include, but are not limited to, Dual Targeting (DT)-Ig molecules (WO2009058383), Two-in-one Antibody (Genentech; Bostrom, et al 2009. Science 323, 1610-1614.), Cross-linked MAbs (Karmanos Cancer Center), mAb2 (F-Star, WO2008003116), Zyboby molecules (Zyngenia; LaFleur et al. MAbs. 2013 Mar-Apr;5(2):208-18), approaches with common light chain (Crucell/Merus, US7,262,028), κ LBodies (NovImmune, WO2012023053) and CovX-body (CovX/Pfizer; Doppalapudi, V.R., et al 2007. Bioorg. Med. Chem. Lett. 17,501-506.).

Examples of IgG fusion molecules include, but are not limited to, Dual Variable Domain (DVD)-Ig molecules (Abbott, US7,612,181), Dual domain double head antibodies (Unilever; Sanofi Aventis, WO20100226923), IgG-like Bispecific molecules (ImClone/Eli Lilly, Lewis et al. Nat Biotechnol. 2014 Feb;32(2):191-8), Ts2Ab (MedImmune/AZ; Dimasi et al. J Mol Biol. 2009 Oct 30;393(3):672-92) and BsAb molecules (Zymogenetics, WO2010111625), HERCULES molecules (Biogen Idec, US007951918), scFv fusion molecules (Novartis), scFv fusion molecules (Changzhou Adam Biotech Inc, CN 102250246) and TvAb molecules (Roche, WO2012025525, WO2012025530).

Examples of Fc fusion molecules include, but are not limited to, ScFv/Fc Fusions (Pearce et al., Biochem Mol Biol Int. 1997 Sep;42(6):1179-88), SCORPION molecules (Emergent BioSolutions/Trubion, Blankenship JW, et al. AACR 100th Annual meeting 2009 (Abstract # 5465); Zymogenetics/BMS, WO2010111625), Dual Affinity Retargeting

Technology (Fc-DART) molecules (MacroGenics, WO2008157379, WO2010080538) and Dual(ScFv)₂-Fab molecules (National Research Center for Antibody Medicine – China).

Examples of Fab fusion bispecific antibodies include, but are not limited to, F(ab)₂ molecules (Medarex/AMGEN; Deo et al J Immunol. 1998 Feb 15;160(4):1677-86.), Dual-Action or Bis-Fab molecules (Genentech, Bostrom, et al 2009. Science 323, 1610–1614.), Dock-and-Lock (DNL) molecules (ImmunoMedics, WO2003074569, WO2005004809), Bivalent Bispecific molecules (Biotechnol, Schoonjans, J Immunol. 2000 Dec 15;165(12):7050-7.) and Fab-Fv molecules (UCB-Celltech, WO 2009040562 A1).

Examples of ScFv-, diabody-based and domain antibodies include, but are not limited to, Bispecific T Cell Engager (BiTE) molecules (Micromet, WO2005061547), Tandem Diabody molecules (TandAb) (Affimed) Le Gall et al., Protein Eng Des Sel. 2004 Apr;17(4):357-66.), Dual Affinity Retargeting Technology (DART) molecules (MacroGenics, WO2008157379, WO2010080538), Single-chain Diabody molecules (Lawrence, FEBS Lett. 1998 Apr 3;425(3):479-84), TCR-like Antibodies (AIT, ReceptorLogics), Human Serum Albumin ScFv Fusion (Merrimack, WO2010059315) and COMBODY molecules (Epigen Biotech, Zhu et al. Immunol Cell Biol. 2010 Aug;88(6):667-75.), dual targeting nanobodies (Ablynx, Hmila et al., FASEB J. 2010) and dual targeting heavy chain only domain antibodies.

In one embodiment, each of said first and second heavy chains comprises at least a hinge region, a CH2 and a CH3 region.

In a further embodiment, the CH3 regions of the first and second heavy chains comprise asymmetrical mutations, such as asymmetrical mutations (also referred to as modifications herein) yielding a stable heterodimeric antibody.

In one embodiment, the bispecific antibody of the invention comprises a first heavy chain comprising a first CH3 region, and a second heavy chain comprising a second CH3 region, wherein the sequences of the first and second CH3 regions are different and are such that the heterodimeric interaction between said first and second CH3 regions is stronger than each of the homodimeric interactions of said first and second CH3 regions. More details on these interactions and how they can be achieved are provided in, e.g., WO 2011/131746 and WO 2013/060867 (Genmab), which are hereby incorporated by reference.

As described further herein, a stable bispecific CD40xCD137 antibody can be obtained at high yield using a particular method on the basis of one homodimeric parental CD40 antibody and one homodimeric parental CD137 antibody containing only a few, fairly conservative, asymmetrical mutations in the CH3 regions. Asymmetrical mutations mean that the sequences of said first and second CH3 regions contain amino acid substitutions at non-identical positions.

Accordingly, in one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the sequences of said first and second heavy chain CH3 regions contain asymmetrical mutations, e.g., a mutation at the position corresponding to position 405 in a human IgG1 heavy chain according to EU numbering in one of the CH3 regions, and a mutation at the position corresponding to position 409 in a human IgG1 heavy chain according to EU numbering in the other CH3 region.

In one aspect, the bispecific antibody as defined in any of the embodiments disclosed herein comprises first and second heavy chains, wherein in said first heavy chain at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370, D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and in said second heavy chain at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370, D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and wherein said first and said second heavy chains are not substituted in the same positions.

In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at a position selected from the group consisting of: 366, 368, 370, 399, 405, 407 and 409 in a human IgG1 heavy chain according to EU numbering, and the second heavy chain has an amino acid substitution at a position selected from the group consisting of: 366, 368, 370, 399, 405, 407 and 409 in a human IgG1 heavy chain according to EU numbering, and wherein the first and second heavy chains are not substituted in the same positions.

In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 366, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of: 368, 370, 399, 405, 407 and 409. In one embodiment the amino acid at position 366 is selected from Ala, Asp, Glu, His, Asn, Val, or Gln.

In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 368, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of: 366, 370, 399, 405, 407 and 409.

In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 370, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of: 366, 368, 399, 405, 407 and 409.

In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 399, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of:366, 368, 370, 405, 407 and 409.

5 In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 405, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of:366, 368, 370, 399, 407 and 409.

10 In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 407, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of:366, 368, 370, 399, 405, and 409.

15 In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid substitution at position 409, and said second heavy chain has an amino acid substitution at a position selected from the group consisting of:366, 368, 370, 399, 405, and 407.

Accordingly, in one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the sequences of said first and second CH3 regions contain asymmetrical mutations, i.e. mutations at different positions in the two CH3 regions, e.g. a mutation at position 405 in one of the CH3 regions and a mutation at position 409 in the other CH3 region.

20 In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an amino-acid substitution at a position selected from the group consisting of:366, 368, 370, 399, 405 and 407. In one such embodiment, said first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an amino acid other than Phe, e.g. Gly, Ala, Val, Ile, Ser, Thr, Lys, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, Cys, Lys, or Leu, at position 405. In a further embodiment hereof, said first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an amino acid other than Phe, Arg or Gly, e.g. Leu, Ala, Val, Ile, Ser, Thr, Met, Lys, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 405.

35 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises a Phe at position 405 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg,

His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain comprises an amino acid other than Phe, e.g. Gly, Ala, Val, Ile, Ser, Thr, Lys, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, Leu, Met, or Cys, at position 405 and a Lys at position 409. In a further embodiment hereof, said first heavy chain comprises a Phe at position 405 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain comprises an amino acid other than Phe, Arg or Gly, e.g. Leu, Ala, Val, Ile, Ser, Thr, Met, Lys, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 405 and a Lys at position 409.

10 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises a Phe at position 405 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain comprises a Leu at position 405 and a Lys at position 409. In a further
15 embodiment hereof, said first heavy chain comprises a Phe at position 405 and an Arg at position 409 and said second heavy chain comprises an amino acid other than Phe, Arg or Gly, e.g. Leu, Ala, Val, Ile, Ser, Thr, Lys, Met, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 405 and a Lys at position 409. In another embodiment, said first heavy chain comprises Phe at position 405 and an Arg at position 409 and said second
20 heavy chain comprises a Leu at position 405 and a Lys at position 409.

In a further embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain comprises a Lys at position 409, a Thr at position 370 and a Leu at position 405. In a further embodiment,
25 said first heavy chain comprises an Arg at position 409 and said second heavy chain comprises a Lys at position 409, a Thr at position 370 and a Leu at position 405.

In an even further embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises a Lys at position 370, a Phe at position 405 and an Arg at position 409 and said second heavy chain comprises a Lys at position 409, a Thr at position 370 and a Leu at position 405.
30

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain comprises a Lys at position 409 and: a) an Ile at position 350 and a Leu at position 405, or b) a Thr at position 370 and a Leu at position 405.
35

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises an Arg at position 409 and said second heavy chain comprises a Lys at position 409 and: a) an Ile at position 350 and a Leu at position 405, or b) a Thr at position 370 and a Leu at position 405.

5 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises a Thr at position 350, a Lys at position 370, a Phe at position 405 and an Arg at position 409 and said second heavy chain comprises a Lys at position 409 and: a) an Ile at position 350 and a Leu at position 405, or b) a Thr at position 370 and a Leu at position 405.

10 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain comprises a Thr at position 350, a Lys at position 370, a Phe at position 405 and an Arg at position 409 and said second heavy chain comprises an Ile at position 350, a Thr at position 370, a Leu at position 405 and a Lys at position 409.

15 In one embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has an amino acid other than Lys, Leu or Met at position 409 and said second heavy chain has an amino acid other than Phe at position 405, such as other than Phe, Arg or Gly at position 405; or said first CH3 region has an amino acid other than Lys, Leu or Met at position 409 and said second
20 CH3 region has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first heavy chain having an amino acid other than Lys, Leu or Met at position 409 and a second heavy chain having an amino acid other than Tyr,
25 Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first heavy chain having a Tyr at position 407 and an amino acid other than Lys, Leu or Met at position 409 and a second heavy chain having an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and a
30 Lys at position 409.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first heavy chain having a Tyr at position 407 and an Arg at position 409 and a second heavy chain having an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr at position 407 and a Lys at position 409.

35 In another embodiment, said first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr, e.g. Leu, Met, Gly, Ala, Val, Ile, His, Asn,

Pro, Trp, or Cys, at position 407. In another embodiment, said first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407.

5 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has a Gly, Leu, Met, Asn or Trp at position 407.

10 In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has an amino acid other than Tyr, Asp, Glu, Phe, Lys, Gln, Arg, Ser or Thr, e.g. Leu, Met,
15 Gly, Ala, Val, Ile, His, Asn, Pro, Trp, or Cys, at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain
20 has an Ala, Gly, His, Ile, Leu, Met, Asn, Val or Trp at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His,
25 Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409 and said second heavy chain has a Gly, Leu, Met, Asn or Trp at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an Arg at position 409 and said second heavy chain has an amino acid other than Tyr, Asp,
30 Glu, Phe, Lys, Gln, Arg, Ser or Thr, e.g. Leu, Met, Gly, Ala, Val, Ile, His, Asn, Pro, Trp, or Cys, at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an Arg at position 409 and said second heavy chain has an Ala, Gly, His, Ile, Leu, Met, Asn,
35 Val or Trp at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, said first heavy chain has a Tyr at position 407 and an

Arg at position 409 and said second heavy chain has a Gly, Leu, Met, Asn or Trp at position 407 and a Lys at position 409.

In another embodiment of the bispecific antibody as defined in any of the embodiments disclosed herein, the first heavy chain has an amino acid other than Lys, Leu or Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Phe, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 409, and the second heavy chain has

(i) an amino acid other than Phe, Leu and Met, e.g. Gly, Ala, Val, Ile, Ser, Thr, Lys, Arg, His, Asp, Asn, Glu, Gln, Pro, Trp, Tyr, or Cys, at position 368, or

(ii) a Trp at position 370, or

(iii) an amino acid other than Asp, Cys, Pro, Glu or Gln, e.g. Phe, Leu, Met, Gly, Ala, Val, Ile, Ser, Thr, Lys, Arg, His, Asn, Trp, Tyr, or Cys, at position 399 or

(iv) an amino acid other than Lys, Arg, Ser, Thr, or Trp, e.g. Phe, Leu, Met, Ala, Val, Gly, Ile, Asn, His, Asp, Glu, Gln, Pro, Tyr, or Cys, at position 366.

In one embodiment, the first heavy chain has an Arg, Ala, His or Gly at position 409, and the second heavy chain has

(i) a Lys, Gln, Ala, Asp, Glu, Gly, His, Ile, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or

(ii) a Trp at position 370, or

(iii) an Ala, Gly, Ile, Leu, Met, Asn, Ser, Thr, Trp, Phe, His, Lys, Arg or Tyr at position 399, or

(iv) an Ala, Asp, Glu, His, Asn, Val, Gln, Phe, Gly, Ile, Leu, Met, or Tyr at position 366.

In one embodiment, the first heavy chain has an Arg at position 409, and the second heavy chain has

(i) an Asp, Glu, Gly, Asn, Arg, Ser, Thr, Val, or Trp at position 368, or

(ii) a Trp at position 370, or

(iii) a Phe, His, Lys, Arg or Tyr at position 399, or

(iv) an Ala, Asp, Glu, His, Asn, Val, Gln at position 366.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises first and second heavy chains, wherein (i) the amino acid in the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering is L in said first heavy chain, and the amino acid in the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering is R in said second heavy chain, or (ii) the amino acid in the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering is R in said first heavy chain, and the amino acid in the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering is L in said second heavy chain.

In a further embodiment said first and second heavy chain are of a human IgG1 isotype.

In another further embodiment said first and second heavy chain are of a human IgG2 isotype.

5 In another further embodiment said first and second heavy chain are of a human IgG3 isotype.

In another embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises first and second heavy chains of the human IgG4 isotype, wherein (i) the amino acid in the position corresponding to S228 in a
10 human IgG4 heavy chain according to EU numbering is P in said first heavy chain, and the amino acid in the position corresponding to S228, F405 and R409 in a human IgG4 heavy chain according to EU numbering is P, L and K, respectively, in said second heavy chain, or (ii) the amino acid in the position corresponding to S228, F405 and R409 in a
15 human IgG4 heavy chain according to EU numbering is P, L and K, respectively, in said first heavy chain, and the amino acid in the position corresponding to S228 in a human IgG4 heavy chain according to EU numbering is P in said second heavy chain.

If reference is made herein to amino acids at certain positions of the first heavy chain and/or amino acids at certain positions of the second heavy chain, such reference is to be understood to include embodiments wherein the amino acids at certain positions
20 of the first heavy chain are present at the corresponding positions of the second heavy chain rather than the first heavy chain and/or the amino acids at certain positions of the second heavy chain are present at the corresponding positions of the first heavy chain rather than the second heavy chain.

In addition to the above-specified amino-acid substitutions, said first and second
25 heavy chains may contain further amino-acid substitutions, deletion or insertions relative to wild-type heavy chain sequences.

In a further embodiment, said first and second Fab-arms (or heavy chain constant domains) comprising the first and second heavy chains comprise, except for the specified mutations, a CH3 sequence independently selected from the following:

30 (IgG1m(a)) (SEQ ID NO:106), (IgG1m(f)) (SEQ ID NO:107), and (IgG1m(ax)) (SEQ ID NO:108).

In one embodiment, neither said first nor said second heavy chain comprises a Cys-Pro-Ser-Cys sequence in the (core) hinge region.

In a further embodiment, both said first and said second heavy chain comprise a
35 Cys-Pro-Pro-Cys sequence in the (core) hinge region.

In separate and specific embodiments, one or both Fab-arms comprise a heavy-chain constant region sequence independently selected from SEQ ID NO:109, 110, 111, 112, 113 and 116 (see **Table 1**).

Methods of preparing bispecific antibodies

Traditional methods such as the hybrid hybridoma and chemical conjugation methods (Marvin and Zhu (2005) Acta Pharmacol Sin 26:649) can be used in the preparation of the bispecific antibodies of the invention. Co-expression in a host cell of two antibodies, consisting of different heavy and light chains, leads to a mixture of possible antibody products in addition to the desired bispecific antibody, which can then be isolated by, e.g., affinity chromatography or similar methods.

Strategies favoring the formation of a functional bispecific product upon co-expression of different antibody constructs can also be used, e.g., by the method described by Lindhofer et al. (1995 J Immunol 155:219). Fusion of rat and mouse hybridomas producing different antibodies leads to a limited number of heterodimeric proteins because of preferential species-restricted heavy/light chain pairing. Another strategy to promote formation of heterodimers over homodimers is a "knob-into-hole" strategy in which a protuberance is introduced on a first heavy-chain polypeptide and a corresponding cavity in a second heavy-chain polypeptide, such that the protuberance can be positioned in the cavity at the interface of these two heavy chains so as to promote heterodimer formation and hinder homodimer formation. "Protuberances" are constructed by replacing small amino-acid side-chains from the interface of the first polypeptide with larger side chains. Compensatory "cavities" of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino-acid side-chains with smaller ones (US patent 5,731,168). EP1870459 (Chugai) and WO 2009/089004 (Amgen) describe other strategies for favoring heterodimer formation upon co-expression of different antibody domains in a host cell. In these methods, one or more residues that make up the CH3-CH3 interface in both CH3 domains are replaced with a charged amino acid such that homodimer formation is electrostatically unfavorable and heterodimerization is electrostatically favorable. WO2007110205 (Merck) describe yet another strategy, wherein differences between IgA and IgG CH3 domains are exploited to promote heterodimerization.

A preferred method for preparing the bispecific CD40xCD137 antibodies of the present invention includes the methods described in WO 2011/131746 and WO 2013/060867 (Genmab) comprising the following steps:

a) providing a first antibody comprising an Fc region, said Fc region comprising a first CH3 region;

b) providing a second antibody comprising a second Fc region, said Fc region comprising a second CH3 region,

wherein the first antibody is a CD40 antibody comprising two first antigen-binding regions as described herein and the second antibody is a CD137 antibody comprising

two second antigen-binding regions as described herein, or vice versa; and wherein the sequences of said first and second CH3 regions are different and are such that the heterodimeric interaction between said first and second CH3 regions is stronger than each of the homodimeric interactions of said first and second CH3 regions;

- 5 c) incubating said first antibody together with said second antibody under reducing conditions; and
 d) obtaining said bispecific CD40xCD137 antibody.

 In one embodiment, said first antibody is incubated together with said second antibody under reducing conditions sufficient to allow the cysteines in the hinge region
10 to undergo disulfide-bond isomerization, wherein the heterodimeric interaction between said first and second antibodies in the resulting heterodimeric antibody is such that no Fab-arm exchange occurs at 0.5 mM GSH after 24 hours at 37° C.

 Without being limited to theory, in step c), the heavy-chain disulfide bonds in the hinge regions of the parent antibodies (first and second antibody in step a) and b)) are
15 reduced and the resulting cysteines are then able to form inter heavy-chain disulfide bond with cysteine residues of another parent antibody molecule (originally with a different specificity). In one embodiment of this method, the reducing conditions in step c) comprise the addition of a reducing agent, e.g. a reducing agent selected from the group consisting of: 2-mercaptoethylamine (2-MEA), dithiothreitol (DTT), dithioerythritol (DTE), glutathione, tris(2-carboxyethyl)phosphine (TCEP), L-cysteine and beta-mercapto-ethanol, preferably a reducing agent selected from the group consisting of: 2-mercaptoethylamine, dithiothreitol and tris(2-carboxyethyl)phosphine. In a further
20 embodiment, step c) comprises restoring the conditions to become non-reducing or less reducing, for example by removal of a reducing agent, e.g. by desalting. In one particular embodiment, bispecific antibodies are generated as follows: the two parental complementary antibodies, both in the same amount, are incubated with 75 mM 2-mercaptoethylamine-HCl (2-MEA) in buffer (e.g., PBS or Tris-EDTA) at 31°C for 5 hours; the reduction reaction is stopped by removing the reducing agent 2-MEA using spin columns (e.g., Microcon centrifugal filters, 30k, Millipore) (Labrijn et al. Nature
25 Protocols, Vol 9 No 10, p2450-2463; 2014). In another particular embodiment, the method is that of Example 3.

 For this method, any of the CD40 and CD137 antibodies disclosed herein may be used. In a particular embodiment the first and second antibodies, binding to human CD40 and CD137, respectively, may be chosen so as to obtain a bispecific antibody as
35 described herein.

 In one embodiment of this method, said first and/or second antibodies are full-length antibodies.

The Fc regions of the first and second antibodies may be of any isotype, including, but not limited to, an IgG isotype having a subclass selected from the group consisting of IgG1, IgG2, IgG3 and IgG4. In one embodiment of this method, the Fc regions of both said first and said second antibodies are of the IgG1 isotype. In another embodiment, one of the Fc regions of said antibodies is of the IgG1 isotype and the other of the IgG4 isotype. In the latter embodiment, the resulting bispecific antibody comprises an Fc region of an IgG1 and an Fc region of IgG4.

In a further embodiment, one of the parental antibodies has been engineered to not bind Protein A, thus allowing separation of the heterodimeric antibody from said parental homodimeric antibodies by passing the product over a protein A column.

As described above, the sequences of the first and second CH3 regions of the homodimeric parental antibodies are different and are such that the heterodimeric interaction between said first and second CH3 regions is stronger than each of the homodimeric interactions of said first and second CH3 regions. More details on these interactions and how they can be achieved are provided in WO 2011/131746 and WO 2013/060867 (Genmab), which are hereby incorporated by reference in their entirety.

In particular, a stable bispecific CD40xCD137 antibody can be obtained at high yield using the above method of the invention on the basis of two homodimeric antibodies which bind CD40 and CD137, respectively, and contain only a few, fairly conservative, asymmetrical mutations in the CH3 regions. Asymmetrical mutations mean that the sequences of said first and second CH3 regions contain amino acid substitutions at non-identical positions.

The bispecific antibodies of the invention may also be obtained by co-expression of constructs encoding the first and second polypeptides in a single cell. Thus, in a further aspect, the invention relates to a method for producing a bispecific antibody, said method comprising the following steps:

a) providing a first nucleic-acid construct encoding a first polypeptide comprising a first Fc region and a first antigen-binding region binding to human CD40 according to any aspect or embodiment herein, said first Fc region comprising a first CH3 region,

b) providing a second nucleic-acid construct encoding a second polypeptide comprising a second Fc region and a second antigen-binding region binding to human CD137 according to any aspect or embodiment herein, said second Fc region comprising a second CH3 region,

wherein the sequences of said first and second CH3 regions are different and are such that the heterodimeric interaction between said first and second CH3 regions is stronger than each of the homodimeric interactions of said first and second CH3 regions, and wherein in said first CH3 region at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370,

D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and in said second CH3 region at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370, D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and wherein said first and said second heavy chains are not substituted in the same positions,

optionally wherein said first and second nucleic acid constructs encode light chain sequences of said first and second antibodies,

c) co-expressing said first and second nucleic-acid constructs in a host cell, and

10 d) obtaining said heterodimeric protein from the cell culture.

Thus, the present invention also relates to a recombinant eukaryotic or prokaryotic host cell which produces a bispecific antibody of the present invention.

In one embodiment of the present invention, the bispecific antibody is obtained by any of the methods according to the present invention.

15 Suitable expression vectors, including promoters, enhancers, etc., and suitable host cells for the production of antibodies are well-known in the art. Examples of host cells include yeast, bacterial and mammalian cells, such as CHO or HEK cells.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises first and second CH3 regions, except for the specified mutations, comprising the sequence of SEQ ID NO:107 (IgG1m(f)).

20 In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein neither said first nor said second Fc-region comprises a Cys-Pro-Ser-Cys sequence in the hinge region.

25 In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein both of said first and said second Fc-region comprise a Cys-Pro-Pro-Cys sequence in the hinge region.

30 In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein the first and second Fc-regions are human antibody Fc-regions.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein said first and second Fc region, except for the specified mutations, comprise a sequence independently selected from the group consisting of SEQ ID NOS:109, 110, 111, 112, 113 and 116.

In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein the first and second antigen-binding regions are from heavy-chain antibodies.

5 In one embodiment, the bispecific antibody as defined in any of the embodiments disclosed herein comprises a first Fc-region and a second Fc-region, wherein the first and second antigen-binding regions comprise a first and second light chain.

In further embodiments, the co-expression method according to the invention comprises any of the further features described under the in vitro method above.

Inert format

10 The effector functions mediated by the Fc region of an antibody allow for the destruction of foreign entities, such as the killing of pathogens and the clearance and degradation of antigens. Antibody-dependent cell-mediated cytotoxicity (ADCC) and antibody-dependent cell-mediated phagocytosis (ADCP) are initiated by binding of the Fc region to Fc receptor (FcR)-bearing cells, whereas complement-dependent cytotoxicity
15 (CDC) and complement-dependent cell-mediated cytotoxicity (CDCC) are initiated by binding of the Fc region to C1q, which initiates the classical route of complement activation.

Fc-mediated effector function, such as ADCC and complement activation, have been suggested to contribute to the therapeutic efficacy of monoclonal antibodies used
20 for the treatment of cancer (Weiner et al. Cell 2012, 148:1081-1084).

The multispecific antibody, such as a bispecific antibody, according to the present invention binds to CD137 which is expressed on T-cells, e.g. CD4⁺ and/or CD8⁺ T-cells. By concomitant binding of the antibody to CD40, which is expressed on e.g. antigen-presenting cells (APCs), provides stimulation to both APCs expressing CD40 and T-cells
25 expressing CD137 and thereby e.g. T-cell proliferation can be increased.

In general, binding of an antibody to a target antigen expressed by a cell may lead to interactions with effector molecules such as Fc receptors or complement proteins which may induce Fc-mediated effector functions, such as ADCC or complement activation, which may result in killing of the cell expressing said target antigen.

30 The use of the multispecific antibody, such as a bispecific antibody, according to the present invention is based on its ability to provide co-stimulation to APCs and T cells.

It is, in a particular embodiment, preferred that the multispecific antibody does not bind to FcRs, e.g. FcγRs, and therefore does not induce FcR-mediated cross-linking.

It is, in a further embodiment, preferred that the multispecific antibody does not
35 engage effector functions so as to avoid killing of the CD40 and/or CD137 expressing cells.

In one aspect of the present invention, the multispecific CD40xCD137 antibody according to the present invention comprises (i) a first binding arm comprising a first heavy chain and a first antigen-binding region and (ii) a second binding arm comprising a second heavy chain and a second antigen-binding region, according to any aspect or
5 embodiment described herein.

In one embodiment the multispecific antibody according to present invention comprises a first and a second heavy chain, wherein said antibody induces and/or enhances Fc-mediated effector function to a lesser extent compared to a multispecific antibody comprising the same first and second antigen-binding regions as said antibody,
10 and comprising two heavy chains comprising a human IgG1 hinge, CH2 and CH3 regions.

In one embodiment, the multispecific antibody according to present invention comprises a first and a second antigen-binding region and a first and a second heavy chain, each of the first and second heavy chains comprising a human IgG1 hinge, CH2
15 and CH3 regions, wherein at least one of the first and second heavy chain comprises a modification so as to induce and/or enhance Fc-mediated effector function to a lesser extent compared to a reference multispecific antibody comprising the same first and second antigen-binding regions as said antibody, and comprising two heavy chains comprising a human IgG1 hinge, CH2 and CH3 regions without said modification.

20 In one embodiment, said first and second heavy chains are modified so that the multispecific antibody induces and/or enhances Fc-mediated effector function to a lesser extent compared to a multispecific antibody which is identical except for comprising non-modified first and second heavy chains.

In one embodiment, said Fc-mediated effector function may be measured by
25 binding to Fcγ-receptors, binding to C1q, or induction of Fc-mediated cross-linking of FcRs.

In one embodiment, said Fc-mediated effector function is measured by binding to C1q.

In one embodiment, said first and second heavy and light chain constant
30 sequences have been modified so that binding of C1q to said multispecific antibody is reduced compared to a wild-type multispecific antibody by at least 70%, at least 80%, at least 90%, at least 95%, at least 97%, or 100%, wherein C1q binding is determined by ELISA.

Human IgG1 is known for its ability to induce Fc-mediated effector functions,
35 while other human isotypes, such as IgG4, are less able to induce Fc-mediated effector functions.

The first and second heavy chains may each be of any isotype, including, but not limited to, an IgG1 isotype selected from the groups consisting of IgG1, IgG2, IgG3 and

IgG4, and may optionally comprise one or more mutations or modifications. In one embodiment, each of the first and second heavy chains is of the IgG4 isotype or derived therefrom, optionally with one or more mutations or modifications. In one embodiment, each of the first and second heavy chains is of the IgG1 isotype or derived therefrom, optionally with one or more mutations or modifications. In another embodiment, one of the heavy chains is of the IgG1 isotype and the other of the IgG4 isotype, or is derived from such respective isotypes, optionally with one or more mutations or modifications.

In one embodiment, one or both of the first and heavy chains are such that an antibody comprising two first or two second heavy chains would be effector-function-deficient. For example, the first and second heavy chains may be of an IgG4 isotype, or a non-IgG4 type, e.g. IgG1, IgG2 or IgG3, which has been mutated such that the ability to mediate effector functions, such as ADCC, has been reduced or even eliminated compared to non-mutated heavy chains. Such mutations have e.g. been described in Dall'Acqua WF et al., J Immunol. 177(2):1129-1138 (2006) and Hezareh M, J Virol.; 75(24):12161-12168 (2001). The multispecific antibody according to the present invention may comprise modifications in the first and second heavy chains compared to a wild type human IgG1 sequence. A multispecific antibody comprising such modifications in the Fc region of the antibody may become an inert, or non-activating, multispecific antibody. The term "inertness", "inert" or "non-activating" as used herein, refers to an Fc region which is at least not able to bind any Fcγ(gamma) receptors, bind to C1q, or induce Fc-mediated cross-linking of FcRs. The inertness of an Fc region, or the first and/or second heavy chain of a multispecific antibody of the present invention may be tested with a bivalent, monospecific antibody comprising said Fc region, or two first heavy chains or two second heavy chains. It may also be tested with a multispecific antibody comprising a first and a second heavy chain.

Several variants can be constructed to make the Fc region of an antibody inactive for interactions with Fcγ receptors and C1q for therapeutic antibody development. The present invention is not limited to any specific mutation relevant for reducing Fc-mediated effector functions. Examples of such variants are described herein.

Thus, amino acids in the Fc region that play a dominant role in the interactions with C1q and the Fcγ receptors may be modified. Examples of amino acid positions that may be modified include positions L234, L235 and P331.

Hence, in one embodiment, in at least one of said first and second heavy chains the amino acid in at least one position corresponding to L234, L235 and P331 in a human IgG1 heavy chain according to EU numbering, may be A, A and S, respectively. (Xu et al., 2000, Cell Immunol. 200(1):16-26; Oganessian et al., 2008, Acta Cryst. (D64):700-4). Also, L234F and L235E amino acid substitutions can result in Fc regions with abrogated interactions with Fcγ receptors and C1q (Canfield et al., 1991, J.

Exp.Med. (173):1483-91; Duncan et al., 1988, Nature (332):738-40). Hence, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to EU numbering, may be F and E, respectively. A D265A amino acid substitution can decrease binding to all Fc gamma receptors and prevent ADCC (Shields et al., 2001, J. Biol. Chem. (276):6591-604). Hence, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to D265 in a human IgG1 heavy chain according to EU numbering, may be A. Binding to C1q can be abrogated by mutating positions D270, K322, P329, and P331. Mutating these positions to either D270A or K322A or P329A or P331A can make the antibody deficient in CDC activity Idusogie EE, et al., 2000, J Immunol. 164: 4178-84). Hence, in one embodiment, in at least one of said first and second heavy chain, the amino acids in at least one position corresponding to D270, K322, P329 and P331 in a human IgG1 heavy chain according to EU numbering, may be A, A, A, and A, respectively.

An alternative approach to minimize the interaction of the Fc region with Fcγ receptors and C1q is by removal of the glycosylation site of an antibody. Mutating position N297 to e.g. Q, A, or E removes a glycosylation site which is critical for IgG-Fcγ receptor interactions. Hence, in one embodiment, in at least one of said first and second heavy chains, the amino acid in a position corresponding to N297, may be G, Q, A or E in a human IgG1 heavy chain according to EU numbering (Leabman et al., 2013, MABs; 5(6):896-903). Another alternative approach to minimize interaction of the Fc region with Fcγ receptors may be obtained by the following mutations; P238A, A327Q, P329A or E233P/L234V/L235A/G236del (Shields et al., 2001, J. Biol. Chem. (276):6591-604).

Alternatively, human IgG2 and IgG4 subclasses are considered naturally compromised in their interactions with C1q and Fc gamma Receptors although interactions with Fcγ receptors were reported (Parren et al., 1992, J. Clin Invest. 90: 1537-1546; Bruhns et al., 2009, Blood 113: 3716-3725). Mutations abrogating these residual interactions can be made in both isotypes, resulting in reduction of unwanted side-effects associated with FcR binding. For IgG2, these include V234A and G237A, and for IgG4, L235E. Hence, in one embodiment, in at least one of said first and second heavy chains, such as in both said first and second heavy chains, the amino acid in a position corresponding to V234 and G237 in a human IgG2 heavy chain according to EU numbering, may be A and A, respectively. In one embodiment, the amino acid in a position corresponding to L235 in a human IgG4 heavy chain according to EU numbering, may be E.

Other approaches to further minimize the interaction with Fcγ receptors and C1q in IgG2 antibodies include those described in WO2011066501 and Lightle, S., et al., 2010, Protein Science (19):753-62.

The hinge region of the antibody can also be of importance with respect to interactions with Fcγ receptors and complement (Brekke et al., 2006, J Immunol 177:1129-1138; Dall'Acqua WF, et al., 2006, J Immunol 177:1129-1138). Accordingly, mutations in or deletion of the hinge region can influence effector functions of an antibody.

In one embodiment, the multispecific antibody comprises a first and a second heavy chain, wherein in at least one of said first and second immunoglobulin heavy chains one or more amino acids in the positions corresponding to positions L234, L235, D265, N297, and P331 in a human IgG1 heavy chain according to EU numbering, are not L, L, D, N, and P, respectively.

In one embodiment, in both the first and second heavy chains one or more amino acids in the position corresponding to positions L234, L235, D265, N297, and P331 in a human IgG1 heavy chain according to EU numbering, are not L, L, D, N, and P, respectively.

In another embodiment, in at least one of the first and second heavy chains one or more amino acids in the positions corresponding to positions L234, L235 and D265 in a human IgG1 heavy chain according to EU numbering, are not L, L and D, respectively, and the amino acids in the positions corresponding to N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, one or both of the heavy chains comprise a mutation removing the acceptor site for Asn-linked glycosylation or is otherwise manipulated to change the glycosylation properties. For example, in an IgG1 Fc-region, an N297Q mutation can be used to remove an Asn-linked glycosylation site. Accordingly, in a specific embodiment, one or both heavy chains comprise an IgG1 wildtype sequence with an N297Q mutation.

The term "amino acid corresponding to position" as used herein refers to an amino acid position number in a human IgG1 heavy chain. Corresponding amino acid positions in other immunoglobulins may be found by alignment with human IgG1. Unless otherwise stated or contradicted by context, the amino acids of the constant region sequences are herein numbered according to the EU-index of numbering (described in Kabat, E.A. et al., 1991, Sequences of proteins of immunological interest. 5th Edition - US Department of Health and Human Services, NIH publication No. 91-3242, pp 662, 680, 689). Thus, an amino acid or segment in one sequence that "corresponds to" an amino acid or segment in another sequence is one that aligns with the other amino acid or segment using a standard sequence alignment program such as ALIGN, ClustalW or similar, typically at default settings and has at least 50%, at least 80%, at least 90%, or at least 95% identity to a human IgG1 heavy chain. It is considered well-known in the art how to align a sequence or segment in a sequence and thereby determine the

corresponding position in a sequence to an amino acid position according to the present invention.

In the context of the present invention, the amino acid position may be defined as described above.

5 The term "the amino acid is not" or similar wording when referring to amino acids in a heavy chain is to be understood to mean that the amino acid is any other amino acid than the specific amino acid mentioned. For example, the amino acid in the position corresponding to L234 in a human IgG1 heavy chain is not L, means that the amino acid may be any of the other naturally or non-naturally occurring amino acids than L.

10 In one embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering, is not D.

15 In one embodiment, in said first and second heavy chains the amino acid in the position corresponding to D265 in a human IgG1 heavy chain according to EU numbering, is not D, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

20 In one embodiment, in said first and second heavy chains the amino acids in the positions corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is hydrophobic or polar amino acids.

The term "hydrophobic" as used herein in relation to an amino acid residue, refers to an amino acid residue selected from the group consisting of: A, C, F, G, H, I, L, M, R, T, V, W, and Y.

25 Thus, in one embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group of amino acids consisting of: A, C, F, G, H, I, L, M, R, T, V, W and Y.

30 The term "polar" as used herein in relation to amino acid residues, refers to any amino acid residue selected from the group consisting of: C, D, E, H, K, N, Q, R, S, and T. Thus, in one embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: C, E, H, K, N, Q, R, S, and T.

35 In another embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is an aliphatic uncharged, aromatic or acidic amino acid.

The term "aliphatic uncharged" as used herein in relation to amino acid residues, refers to any amino acid residue selected from the group consisting of: A, G, I, L, and V.

Thus, in one embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, G, I, L, and V.

5 The term "aromatic" as used herein in relation to amino acid residues, refers to any amino acid residue selected from the group consisting of: F, T, and W.

Thus, in one embodiment, in said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: F, T, and W.

10 The term "acidic" as used herein in relation to amino acid residues, refers to any amino acid residue chosen from the group consisting of: D and E.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: D and E.

15 Thus, in one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: D and E.

In a particular embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W.

20 In a particular embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W.

25 In one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering, is not D.

30 In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering, is not D.

In one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to D265 in a human IgG1 heavy chain according to EU numbering, is not D, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

35 In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to D265 in a human IgG1 heavy chain according to EU numbering, is not D, and the amino acids in the positions corresponding to positions

N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is hydrophobic or polar amino acid.

In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is hydrophobic or polar amino acid.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group of amino acids consisting of: A, C, F, G, H, I, L, M, R, T, V, W and Y.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human heavy chain according to EU numbering is selected from the group consisting of: C, E, H, K, N, Q, R, S, and T. In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group of amino acids consisting of: A, C, F, G, H, I, L, M, R, T, V, W and Y.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to position D265 in a human heavy chain according to EU numbering is selected from the group consisting of: C, E, H, K, N, Q, R, S, and T.

In another embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is aliphatic uncharged, aromatic or acidic amino acids.

Thus, in one embodiment, in least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, G, I, L, and V.

Thus, in one embodiment, in least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: F, T, and W.

Thus, in one embodiment, in least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering are selected from the group consisting of: D and E.

In a particular embodiment, in least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W.

In another embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is aliphatic uncharged, aromatic or acidic amino acids.

Thus, in one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, G, I, L, and V.

Thus, in one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: F, T, and W.

Thus, in one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering are selected from the group consisting of: D and E.

In a particular embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W.

In further embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position N297 in a human IgG1 heavy chain according to EU numbering, is not N.

In one embodiment, in at least one of the first and second heavy chains the amino acid in the position corresponding to N297 in a human IgG1 heavy chain according to EU numbering, is not N, and the amino acid in the position corresponding to position P331 in a human IgG1 heavy chain according to EU numbering, is P.

In one embodiment, in said first and second heavy chains the amino acid in the position corresponding to positions N297 in a human IgG1 heavy chain according to EU numbering, is not N.

In one embodiment, in both the first and second heavy chains the amino acid in the position corresponding to N297 in a human IgG1 heavy chain according to EU numbering, is not N, and the amino acid in the position corresponding to position P331 in a human IgG1 heavy chain according to EU numbering, is P.

In further embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are not L and L, respectively.

In one embodiment, in at least one of the first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to EU numbering, are not L and L, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in at least one of said first and second heavy chains the amino acids corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are selected from the group consisting of: A, C, D, E, F, G, H, I, K, M, N, P, Q, R, S, T, Y, V.

5 In one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy according to EU numbering chain are hydrophobic or polar amino acids.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human
10 IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, F, G, H, I, M, R, T, V, W, and Y.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group of amino
15 acids consisting of: C, D, E, H, K, N, Q, R, S, and T.

In a particular embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, D, E, F, G, H, I, K, M, N, Q, R, S, T, V, W, and Y.

20 In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are not L and L, respectively.

In one embodiment, in both the first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to
25 EU numbering, are not L and L, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to
30 EU numbering are hydrophobic or polar amino acids.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, F, G, H, I, M, R, T, V, W, and Y.

35 In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group of amino acids consisting of: C, D, E, H, K, N, Q, R, S, and T.

In a particular embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, D, E, F, G, H, I, K, M, N, Q, R, S, T, V, W, and Y.

5 In another embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are aliphatic uncharged, aromatic or acidic amino acids.

10 Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, G, I, and V.

15 Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: F, T, and W.

20 Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: D and E.

In a particular embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to L234 and L235 are each selected from the group consisting of: A, D, E, F, G, I, T, V, and W.

25 In one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are F and E; or A and A, respectively.

30 In one embodiment, in at least one of the first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to EU numbering, are F and E; or A and A, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are F and E; or A and A, respectively.

35 In one embodiment, in both the first and second heavy chains the amino acids in the positions corresponding to L234 and L235 in a human IgG1 heavy chain according to EU numbering, are F and E; or A and A, respectively, and the amino acids in the

positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In a particular embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human
5 IgG1 heavy chain according to EU numbering, are F and E, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are F and E, respectively.

In one embodiment, in at least one of said first and second heavy chains at least
10 the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are A and A, respectively.

In one embodiment, in both said first and second heavy chains at least the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering, are A and A, respectively.

15 In one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are not L, L, and D, respectively.

In one embodiment, in at least one of the first and second heavy chains the amino acids in the positions corresponding to L234, L235, and D265 in a human IgG1
20 heavy chain according to EU numbering, are not L, L and D, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in at least one of said first and second heavy chains the amino acids corresponding to positions L234 and L235 in a human IgG1 heavy chain
25 according to EU numbering are selected from the group consisting of: A, C, D, E, F, G, H, I, K, M, N, P, Q, R, S, T, Y, V, and W, and the amino acid corresponding to position D265 is selected from the group consisting of: A, C, E, F, G, H, I, K, L, M, N, P, Q, R, S, T, Y, V, and W.

In one embodiment, in at least one of said first and second heavy chains the
30 amino acids in the positions corresponding to positions L234, L235 and D265 in a human IgG1 heavy chain according to EU numbering are hydrophobic or polar amino acids.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group of amino acids consisting of:
35 A, C, F, G, H, I, L, M, R, T, V, W and Y, and the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, F, G, H, I, M, R, T, V, W, and Y.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group of amino acids consisting of: C, D, E, H, K, N, Q, R, S, and T, the amino acid in the position
5 corresponding to position D265 in a human heavy chain according to EU numbering is selected from the group consisting of: C, E, H, K, N, Q, R, S, and T.

In a particular embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group
10 consisting of: A, C, D, E, F, G, H, I, K, M, N, Q, R, S, T, V, W, and Y, and the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, C, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W, and Y.

In one embodiment, in both said first and second heavy chains the amino acids in
15 the positions corresponding to L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering are hydrophobic or polar amino acids.

In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group of amino acids consisting of: A, C, F, G, H, I, L,
20 M, R, T, V, W and Y, and the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, F, G, H, I, M, R, T, V, W, and Y.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain
25 according to EU numbering are each selected from the group of amino acids consisting of: C, D, E, H, K, N, Q, R, S, and T, the amino acid in the position corresponding to position D265 in a human heavy chain according to EU numbering is selected from the group consisting of: C, E, H, K, N, Q, R, S, and T.

In a particular embodiment, in both said first and second heavy chains the amino
30 acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, C, D, E, F, G, H, I, K, M, N, Q, R, S, T, V, W, and Y, and the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, C, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W,
35 and Y.

In another embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235 and D265 in a human

IgG1 heavy chain according to EU numbering are aliphatic uncharged, aromatic or acidic amino acids.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, G, I, L, and V, and the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: A, G, I, and V.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235 and D265 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: F, T, and W.

Thus, in one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of: D and E.

In a particular embodiment, in at least one of said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W, and the amino acids in the positions corresponding to L234 and L235 are each selected from the group consisting of: A, D, E, F, G, I, T, V, and W.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235 and D265 in a human IgG1 heavy chain according to EU numbering, are not L, L, and D, respectively.

In one embodiment, in both the first and second heavy chains the amino acids in the positions corresponding to L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are not L, L, and D, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering are aliphatic uncharged, aromatic or acidic amino acids.

In one embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of; A, G, I, L, and V, and the amino acids in the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of; A, G, I, and V.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering are each selected from the group consisting of; D and E.

5 In a particular embodiment, in both said first and second heavy chains the amino acid in the position corresponding to position D265 in a human IgG1 heavy chain according to EU numbering is selected from the group consisting of: A, E, F, G, I, L, T, V, and W, and the amino acids in the positions corresponding to L234 and L235 are each selected from the group consisting of: A, D, E, F, G, I, T, V, and W.

10 In one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A; or A, A, and A, respectively.

In one embodiment, in at least one of the first and second heavy chains the
15 amino acids in the positions corresponding to L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A; or A, A, and A, respectively, and the amino acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in
20 the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A; or A, A, and A, respectively.

In one embodiment, in both the first and second heavy chains the amino acids in the positions corresponding to L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A; or A, A, and A, respectively, and the amino
25 acids in the positions corresponding to positions N297 and P331 in a human IgG1 heavy chain according to EU numbering, are N and P, respectively.

In a particular embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A, respectively.

30 In a particularly preferred embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are F, E, and A, respectively.

In one embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a
35 human IgG1 heavy chain according to EU numbering, are A, A, and A, respectively.

In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering, are A, A, and A, respectively.

In another embodiment, in at least one of said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, D265, N297, and P331 in a human IgG1 heavy chain according to EU numbering, are F, E, A, Q, and S, respectively.

5 In one embodiment, in both said first and second heavy chains the amino acids in the positions corresponding to positions L234, L235, D265, N297, and P331 in a human IgG1 heavy chain according to EU numbering, are F, E, A, Q, and S, respectively.

In a particular embodiment said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID
10 NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively; and said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ
15 ID NOs:67, GAS and 68, respectively, (CD137 clone 009), and in at least one of the first and second heavy chains, such as both said first and second heavy chain, the amino acids in positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain, are F, E, and A, respectively.

In another embodiment said first antigen-binding region comprises heavy chain
20 variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively; and said second antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable
25 region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, SAS and 40, respectively, (CD137 clone 005), and in at least one of the first and second heavy chains, such as both said first and second heavy chain, the amino acids in positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain, are F, E, and A, respectively.

30 A non-activating Fc region prevents the antibody from interacting with Fc-receptors present on blood cells, such as monocytes, or with C1q to activate the classical complement pathway. Reduction of the Fc activity was tested in antibody variants that contain different combinations of amino acid substitutions in the Fc region. Three amino acid substitutions were introduced in the parental antibodies of the present invention,
35 which include the mutations L234F, L235E, and D265A. Substitutions in these three amino acid positions were introduced in the K409R and/or F405L IgG1 backbone. The resulting non-activating antibody variant is termed with the suffix "FEAR" or "FEAL",

respectively. Said parental antibodies were used in to generate bispecific antibodies of the present invention as described in the examples.

In one aspect, the multispecific antibodies according to the invention may be modified in the light chain and/or heavy chain to increase the expression level and/or production yield. In one embodiment, the antibodies according to the invention may be modified in the light chain. Such modifications are known in the art and may be performed according to the methods described in e.g. Zheng, L., Goddard, J.-P., Baumann, U., & Reymond, J.-L. (2004). Expression improvement and mechanistic study of the retro-Diels-Alderase catalytic antibody 10F11 by site-directed mutagenesis. *Journal of Molecular Biology*, 341(3), 807–14.

In a further embodiment of the invention, one or both of the antibodies forming part of the multispecific antibody of the invention have been engineered to reduce or increase the binding to the neonatal Fc receptor (FcRn) in order to manipulate the serum half-life of the multispecific antibody. Techniques for increasing or reducing the serum half-life are well-known in the art. See for example Dall'Acqua et al. 2006, *J. Biol. Chem.*, 281:23514-24; Hinton et al. 2006, *J. Immunol.*, 176:346-56; and Zalevsky et al. 2010 *Nat. Biotechnol.*, 28:157-9.

In one aspect, the multispecific antibody as defined in any of the embodiments disclosed herein comprises a first constant heavy chain (HC) and a first constant light chain (LC), wherein the positions corresponding to positions L234, L235, and D265 in the human IgG1 heavy chain of SEQ ID NO:109 of both the first heavy chain and the second heavy chain are F, E, and A, respectively.

In one embodiment, the multispecific antibody as defined in any of the embodiments disclosed herein comprises a first and second constant heavy chain (HC) and a first and second constant light chain (LC), wherein the positions corresponding to positions L234 and L235 in the human IgG1 heavy chain of SEQ ID NO:109 of both the first heavy chain and the second heavy chain are F and E, respectively.

In one embodiment, the multispecific antibody comprises a first and a second heavy chain, wherein the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering of both the first heavy chain and the second heavy chain are F and E, respectively, and wherein (i) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is R, or (ii) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is L.

In one embodiment, the multispecific antibody comprises a first and a second heavy chain, wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and the second heavy chain are F, E, and A, respectively, and wherein the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is R. Thus in a further embodiment, said first heavy chain comprises the constant heavy chain sequence as set forth in SEQ ID NO:113; and the second heavy chain comprises the constant heavy chain sequence as set forth in SEQ ID NO:112.

In one embodiment, the multispecific antibody comprises a first and a second heavy chain, wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first and second heavy chain are F, E, and A, respectively, and wherein the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is L. Thus in a further embodiment, said first heavy chain comprises the constant heavy chain sequence as set forth in SEQ ID NO:112; and the second heavy chain comprises the constant heavy chain sequence as set forth in SEQ ID NO:113.

Nucleic acids

The present invention also relates to a nucleic acid encoding one or more amino acid sequences according to any aspect or embodiment disclosed herein.

The present invention also relates to a nucleic acid encoding a multispecific antibody as defined in any aspect or embodiment disclosed herein.

The present invention also relates to an expression vector comprising a nucleic acid of the present invention.

The present invention also relates to a host cell comprising a nucleic acid or an expression vector according to the present invention.

In one embodiment said host cell is a recombinant eukaryotic, recombinant prokaryotic, or recombinant microbial host cell.

In a further embodiment, the expression vector further comprises a nucleotide sequence encoding the constant region of a light chain, a heavy chain or both light and heavy chains of an antibody, e.g. a human antibody.

An expression vector in the context of the present invention may be any suitable vector, including chromosomal, non-chromosomal, and synthetic nucleic acid vectors (a

nucleic acid sequence comprising a suitable set of expression control elements). Examples of such vectors include derivatives of SV40, bacterial plasmids, phage DNA, baculovirus, yeast plasmids, vectors obtained from combinations of plasmids and phage DNA, and viral nucleic acid (RNA or DNA) vectors. In one embodiment, a nucleic acid is comprised in a naked DNA or RNA vector, including, for example, a linear expression element (as described in for instance Sykes and Johnston, *Nat Biotech* 17, 355 59 (1997)), a compacted nucleic acid vector (as described in for instance US 6,077, 835 and/or WO 00/70087), a plasmid vector such as pBR322, pUC 19/18, or pUC 118/119, a "midge" minimally-sized nucleic acid vector (as described in for instance Schakowski et al., *Mol Ther* 3, 793 800 (2001)), or as a precipitated nucleic acid vector construct, such as a CaP04-precipitated construct (as described in for instance WO200046147, Benvenisty and Reshef, *PNAS USA* 83, 9551 55 (1986), Wigler et al., *Cell* 14, 725 (1978), and Coraro and Pearson, *Somatic Cell Genetics* 7, 603 (1981)). Such nucleic acid vectors and the usage thereof are well known in the art (see for instance US 5,589,466 and US 5,973,972).

In one embodiment, the vector is suitable for expression of the CD40 antibody and/or the CD137 antibody in a bacterial cell. Examples of such vectors include expression vectors such as BlueScript (Stratagene), pIN vectors (Van Heeke & Schuster, *J Biol Chem* 264, 5503 5509 (1989), pET vectors (Novagen, Madison WI) and the like).

An expression vector may also or alternatively be a vector suitable for expression in a yeast system. Any vector suitable for expression in a yeast system may be employed. Suitable vectors include, for example, vectors comprising constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH (reviewed in: F. Ausubel et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley InterScience New York (1987), and Grant et al., *Methods in Enzymol* 153, 516 544 (1987)).

An expression vector may also or alternatively be a vector suitable for expression in mammalian cells, e.g. a vector comprising glutamine synthetase as a selectable marker, such as the vectors described in Bebbington (1992) *Biotechnology (NY)* 10:169-175.

A nucleic acid and/or vector may also comprises a nucleic acid sequence encoding a secretion/localization sequence, which can target a polypeptide, such as a nascent polypeptide chain, to the periplasmic space or into cell culture media. Such sequences are known in the art, and include secretion leader or signal peptides.

The expression vector may comprise or be associated with any suitable promoter, enhancer, and other expression-facilitating elements. Examples of such elements include strong expression promoters (e. g., human CMV IE promoter/enhancer as well as RSV, SV40, SL3 3, MMTV, and HIV LTR promoters), effective poly (A) termination sequences,

an origin of replication for plasmid product in *E. coli*, an antibiotic resistance gene as selectable marker, and/or a convenient cloning site (e.g., a polylinker). Nucleic acids may also comprise an inducible promoter as opposed to a constitutive promoter such as CMV IE.

5 In one embodiment, the CD40 and/or CD137 antibody-encoding expression vector may be positioned in and/or delivered to the host cell or host animal via a viral vector.

In an even further aspect, the invention relates to a host cell comprising the first and second nucleic-acid constructs specified herein above.

10 Thus the present invention also relates to a recombinant eukaryotic or prokaryotic host cell which produces a multispecific antibody of the present invention, such as a transfectoma.

The first, CD40-specific, antibody may be expressed in a recombinant eukaryotic or prokaryotic host cell, such as a transfectoma, which produces an antibody as defined
15 herein. The second, CD137-specific, antibody may likewise be expressed in a recombinant eukaryotic or prokaryotic host cell, such as a transfectoma, which produces an antibody. Such antibodies may be used to prepare a multispecific antibody according to the present invention. A multispecific antibody according to the present invention may also be expressed in a recombinant eukaryotic or prokaryotic host cell, such as a
20 transfectoma.

Examples of host cells include yeast, bacterial, plant and mammalian cells, such as CHO, CHO-S, HEK, HEK293, HEK-293F, Expi293F, PER.C6 or NS0 cells or lymphocytic cells. For example, in one embodiment, the host cell may comprise a first and second nucleic acid construct stably integrated into the cellular genome. In another
25 embodiment, the present invention provides a cell comprising a non-integrated nucleic acid, such as a plasmid, cosmid, phagemid, or linear expression element, which comprises a first and second nucleic acid construct as specified above.

In an even further aspect, the invention relates to a transgenic non-human animal or plant comprising nucleic acids encoding one or two sets of a human heavy
30 chain and a human light chain, wherein the animal or plant produces a multispecific antibody of the invention.

The first, CD40-specific, antibody and/or second, CD137-specific, antibody may also be produced by a hybridoma, a transgenic non-human animal or plant comprising nucleic acids encoding one or two sets of a human heavy chain and a human light chain,
35 wherein the animal or plant produces an antibody for use in a multispecific antibody or a multispecific antibody of the invention.

In one aspect, the invention relates to a nucleic acid encoding one or more amino acid sequences set out in **Table 1**.

In one aspect, the invention relates to an expression vector comprising

(i) a nucleic acid sequence encoding a heavy chain sequence of a first binding arm according to any one of the embodiments disclosed herein;

5 (ii) a nucleic acid sequence encoding a light chain sequence of a first binding arm according to any one of the embodiments disclosed herein;

(iii) a nucleic acid sequence encoding a heavy chain sequence of a second binding arm according to any one of the embodiments disclosed herein;

(iv) a nucleic acid sequence encoding a light chain sequence of a second binding arm according to any one of the of the embodiments disclosed herein;

10 (v) the nucleic acid set forth in (i) and the nucleic acid set forth in (ii);

(vi) the nucleic acid set forth in (iii) and the nucleic acid set forth in (iv).

(vii) the nucleic acid set forth in (i), (ii), (iii) and (iv).

In a particular embodiment, the nucleic acid may encode a heavy chain variable
15 region comprising the VH CDR1, CDR2 and CDR3 of the CD40 antibody listed in **Table 1** and encoding a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116.

In another embodiment, the nucleic acid may encode a heavy chain variable
20 region comprising the VH CDR1, CDR2 and CDR3 of one the CD137 antibodies listed in **Table 1**, i.e. any one of clones 001-012, and encoding a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116.

In separate and specific embodiments, a nucleic acid, nucleic acid construct, a
25 combination of a first and a second nucleic acid construct, an expression vector, or a combination of a first and a second expression vector according to the present invention may encode

(a) a HC comprising (i) a VH comprising the VH CDR1, CDR2 and CDR3 of the CD40 antibody in **Table 1**, and primarily human framework regions, optionally
30 comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

(b) a HC comprising (i) a VH comprising the VH CDR1, CDR2 and CDR3 of one the CD137 antibodies listed in **Table 1**, i.e. any one of clones 001-012, and primarily
35 human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

(c) an LC comprising (i) a VL comprising the VL CDR1, CDR2 and CDR3 of the CD40 antibody in **Table 1**, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a light chain constant region having the sequence of SEQ ID NO:114;

5 (d) an LC comprising (i) a VL comprising the VL CDR1, CDR2 and CDR3 of one the CD137 antibodies listed in **Table 1**, i.e. any of clones 001-012, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a light chain constant region having the sequence of SEQ ID NO:114;

10 (e) both (a) and (b);
(f) both (a) and (c);
(g) both (b) and (d);
(h) both (c) and (d); or
(i) both (a), (b), (c) and (d).

15

In other separate and specific embodiments, a nucleic acid, nucleic acid construct, a combination of a first and a second nucleic acid construct, an expression vector, or a combination of a first and a second expression vector according to the present invention may encode

20 (a) a HC comprising (i) a VH comprising the VH CDR1, CDR2 and CDR3 of SEQ ID NOS:1, 2 and 3, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

25 (b) a HC comprising (i) a VH comprising the VH CDR1, CDR2 and CDR3 of SEQ ID NO: 64, 65 and 66, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

30 (c) an LC comprising (i) a VL comprising the VL CDR1, CDR2 and CDR3 of SEQ ID NO:4, YTS and SEQ ID NO:5, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a light chain constant region having the sequence of SEQ ID NO:114;

(d) an LC comprising (i) a VL comprising the VL CDR1, CDR2 and CDR3 of SEQ ID
35 NO: 67, GAS and SEQ ID NO:68, and primarily human framework regions, optionally comprising one or more amino acid back-mutations to the non-human amino acid sequence, and (ii) a light chain constant region having the sequence of SEQ ID NO:114;

(e) both (a) and (b);

- (f) both (a) and (c);
- (g) both (b) and (d);
- (h) both (c) and (d); or
- (i) both (a), (b), (c) and (d).

5

In other separate and specific embodiments, a nucleic acid, nucleic acid construct, a combination of a first and a second nucleic acid construct, an expression vector, or a combination of a first and a second expression vector according to the present invention may encode

- 10 (a) a HC comprising a VH comprising SEQ ID NO:117 and a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

- (b) a HC comprising a VH comprising SEQ ID NO:123 and a human IgG1 heavy chain having a sequence selected from the group consisting of SEQ ID NO:110, 111, 112, 113 and 116;

- 15 (c) an LC comprising a VL comprising SEQ ID NO:121 and a light chain constant region having the sequence of SEQ ID NO:114;

- (d) an LC comprising a VL comprising SEQ ID NO:127 and a light chain constant region having the sequence of SEQ ID NO:114;

- 20 (e) both (a) and (b);
- (f) both (a) and (c);
- (g) both (b) and (d);
- (h) both (c) and (d); or
- (i) both (a), (b), (c) and (d).

25

In other separate and specific embodiments, a nucleic acid, a nucleic acid construct, a combination of a first and a second nucleic acid construct, an expression vector, or a combination of a first and a second expression vector according to the present invention may encode

- 30 (a) a HC comprising SEQ ID NO:118 (CD40-001-HC6, IgG1);
- (b) a HC comprising SEQ ID NO:119 (CD40-001-HC6-FEAL);
- (c) a HC comprising SEQ ID NO:120 (CD40-001-HC6-FEAR);
- (d) a HC comprising SEQ ID NO:124 (CD137-009-HC7);
- (e) a HC comprising SEQ ID NO:125 (CD137-009-HC7-FEAR);
- 35 (f) a HC comprising SEQ ID NO:126 (CD137-009-HC7-FEAL);
- (g) an LC comprising SEQ ID NO:122 (CD40-001-LC1);
- (h) an LC comprising SEQ ID NO:128 (CD137-009-LC2);
- (i) both (a) and (g);

- (j) both (b) and (g);
(k) both (c) and (g);
(l) both (d) and (h);
(m) both (e) and (h);
5 (n) both (f) and (h);
(o) both (b) and (e);
(p) both (c) and (f);
(q) both (g) and (h);
(r) both (b), (e), (g) and (h);
10 (s) both (c), (f), (g) and (h).

In one aspect, the invention relates to a method for producing a bispecific antibody according to any one of the embodiments as disclosed herein, comprising the steps of

- 15 a) culturing a host cell as disclosed herein comprising an expression vector as disclosed herein expressing the first antibody as disclosed herein and purifying said antibody from the culture media;
b) culturing a host cell as disclosed herein comprising an expression vector as disclosed herein expressing the second antibody as disclosed herein and purifying said
20 antibody from the culture media;
c) incubating said first antibody together with said second antibody under reducing conditions sufficient to allow the cysteines in the hinge region to undergo disulfide-bond isomerization, and
d) obtaining said bispecific antibody.

- 25 In one aspect, the invention relates to a host cell comprising an expression vector as defined above. In one embodiment, the host cell is a recombinant eukaryotic, recombinant prokaryotic, or recombinant microbial host cell.

Compositions

- The present invention also relates to a composition comprising a multispecific
30 antibody according to the present invention, a nucleic acid according to the present invention, an expression vector to the present invention or a host cell according to the present invention.

In a further embodiment the composition according to the present invention is a pharmaceutical composition.

- 35 In an even further embodiment, the pharmaceutical composition according to the present invention further comprises a pharmaceutically acceptable carrier.

In a further aspect, the invention relates to a pharmaceutical composition comprising:

- a multispecific CD40xCD137 antibody as defined in any of the embodiments disclosed herein, and

5 - a pharmaceutically acceptable carrier.

The pharmaceutical composition of the present invention may contain one multispecific antibody of the present invention or a combination of different multispecific antibodies of the present invention.

The pharmaceutical compositions may be formulated in accordance with
10 conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995. A pharmaceutical composition of the present invention may e.g. include diluents, fillers, salts, buffers, detergents (e.g., a nonionic detergent, such as Tween-20 or Tween-80), stabilizers (e. g., sugars or protein-free amino acids), preservatives, tissue fixatives,
15 solubilizers, and/or other materials suitable for inclusion in a pharmaceutical composition.

Pharmaceutically acceptable carriers include any and all suitable solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonicity agents, antioxidants and absorption delaying agents, and the like that are physiologically
20 compatible with a multispecific antibody of the present invention. Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the present invention include water, saline, phosphate buffered saline, ethanol, dextrose, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, carboxymethyl cellulose colloidal
25 solutions, tragacanth gum and injectable organic esters, such as ethyl oleate, and/or various buffers. Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. Proper fluidity may be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in
30 the case of dispersions, and by the use of surfactants.

Pharmaceutical compositions of the present invention may also comprise pharmaceutically acceptable antioxidants for instance (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate,
35 butylated hydroxyanisole, butylated hydroxytoluene, lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Pharmaceutical compositions of the present invention may also comprise isotonicity agents, such as sugars, polyalcohols, such as mannitol, sorbitol, glycerol or sodium chloride in the compositions.

5 The pharmaceutical compositions of the present invention may also contain one or more adjuvants appropriate for the chosen route of administration such as preservatives, wetting agents, emulsifying agents, dispersing agents, preservatives or buffers, which may enhance the shelf life or effectiveness of the pharmaceutical composition. The pharmaceutical composition of the present invention may be prepared with carriers that will protect the multispecific antibody against rapid release, such as a
10 controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Such carriers may include gelatin, glyceryl monostearate, glyceryl distearate, biodegradable, biocompatible polymers such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid alone or with a wax, or other materials well known in the art. Methods for
15 the preparation of such formulations are generally known to those skilled in the art.

Sterile injectable solutions may be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients e.g. as enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active
20 compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients e.g. from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, examples of methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered
25 solution thereof.

The actual dosage levels of the active ingredients in the pharmaceutical compositions may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The
30 selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the
35 age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The pharmaceutical composition may be administered by any suitable route and mode. In one embodiment, a pharmaceutical composition of the present invention is

administered parenterally. "Administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and include epidermal, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, intratendinous, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, intracranial, intrathoracic, epidural and intrasternal injection and infusion.

In one embodiment the pharmaceutical composition is administered by intravenous or subcutaneous injection or infusion.

Uses

10 The present invention also relates to the multispecific antibody according to the present invention, the nucleic acid according to the present invention, the expression vector according to the present invention, the host cell according to the present invention, the composition according to the present invention, or the pharmaceutical composition according to the present invention for use as a medicament.

15 The present invention also relates to the multispecific antibody according to the present invention, the nucleic acid according to the present invention, the expression vector according to the present invention, the host cell according to the present invention, the composition according to the present invention, or the pharmaceutical composition according to the present invention for use in the treatment of a disease, such as cancer or an infectious disease.

20 According to the invention, the term "disease" refers to any pathological state, in particular cancer, infectious diseases, inflammatory diseases, metabolic diseases, autoimmune disorders, degenerative diseases, apoptosis-associated diseases and transplant rejections.

25 As used herein, the term "cancer" includes a disease characterized by aberrantly regulated cellular growth, proliferation, differentiation, adhesion, and/or migration. By "cancer cell" is meant an abnormal cell that grows by a rapid, uncontrolled cellular proliferation and continues to grow after the stimuli that initiated the new growth cease.

30 The term "cancer" according to the invention comprises leukemias, seminomas, melanomas, teratomas, lymphomas, neuroblastomas, gliomas, rectal cancer, endometrial cancer, kidney cancer, adrenal cancer, thyroid cancer, blood cancer, skin cancer, cancer of the brain, cervical cancer, intestinal cancer, liver cancer, colon cancer, stomach cancer, intestine cancer, head and neck cancer, gastrointestinal cancer, lymph node cancer, esophagus cancer, colorectal cancer, pancreas cancer, ear, nose and throat (ENT) cancer, breast cancer, prostate cancer, cancer of the uterus, ovarian cancer and lung cancer and the metastases thereof. Examples thereof are lung carcinomas, mamma

carcinomas, prostate carcinomas, colon carcinomas, renal cell carcinomas, cervical carcinomas, or metastases of the cancer types or tumors described above.

The term "cancer" according to the invention also comprises cancer metastases. By "metastasis" is meant the spread of cancer cells from its original site to another part of the body. The formation of metastasis is a very complex process and depends on detachment of malignant cells from the primary tumor, invasion of the extracellular matrix, penetration of the endothelial basement membranes to enter the body cavity and vessels, and then, after being transported by the blood, infiltration of target organs. Finally, the growth of a new tumor, i.e. a secondary tumor or metastatic tumor, at the target site depends on angiogenesis. Tumor metastasis often occurs even after the removal of the primary tumor because tumor cells or components may remain and develop metastatic potential. In one embodiment, the term "metastasis" according to the invention relates to "distant metastasis" which relates to a metastasis which is remote from the primary tumor and the regional lymph node system.

The term "infectious disease" refers to any disease which can be transmitted from individual to individual or from organism to organism, and is caused by a microbial agent (e.g. common cold).

Examples of infectious diseases include viral infectious diseases, such as AIDS (HIV), hepatitis A, B or C, herpes, herpes zoster (chicken-pox), German measles (rubella virus), yellow fever, dengue etc. flaviviruses, influenza viruses, hemorrhagic infectious diseases (Marburg or Ebola viruses), and severe acute respiratory syndrome (SARS), bacterial infectious diseases, such as Legionnaire's disease (*Legionella*), sexually transmitted diseases (e.g. chlamydia or gonorrhea), gastric ulcer (*Helicobacter*), cholera (*Vibrio*), tuberculosis, diphtheria, infections by *E. coli*, *Staphylococci*, *Salmonella* or *Streptococci* (tetanus); infections by protozoan pathogens such as malaria, sleeping sickness, leishmaniasis; toxoplasmosis, i.e. infections by *Plasmodium*, *Trypanosoma*, *Leishmania* and *Toxoplasma*; or fungal infections, which are caused, e.g., by *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis* or *Candida albicans*.

The term "inflammatory disease" refers to any disease, which is characterized by or associated with high levels of inflammation in tissues, in particular connective tissues, or degeneration of these tissues. A chronic inflammatory disease is a medical condition which is characterized by persistent inflammation. Examples of (chronic) inflammatory diseases include celiac disease, vasculitis, lupus, chronic obstructive pulmonary disease (COPD), irritable bowel disease, atherosclerosis, arthritis, ankylosing spondylitis, Crohn's disease, colitis, chronic active hepatitis, dermatitis and psoriasis.

The term "metabolic disease" refers to any disease or disorder that disrupts normal metabolism. Examples include cystinosis, diabetes, dyslipidemia,

hyperthyroidism, hypothyroidism, hyperlipidemia, hypolipidemia, galactosemia, Gaucher's disease, obesity and phenylketonuria.

The term "autoimmune disorder" refers to any disease/disorder in which the body produces an immunogenic (i.e. immune system) response to some constituent of its own tissue. In other words, the immune system loses its ability to recognize some tissue or system within the body as self and targets and attacks it as if it were foreign. Autoimmune diseases can be classified into those in which predominantly one organ is affected (e.g. hemolytic anemia and anti-immune thyroiditis), and those in which the autoimmune disease process is diffused through many tissues (e.g. systemic lupus erythematosus). For example, multiple sclerosis is thought to be caused by T cells attacking the sheaths that surround the nerve fibers of the brain and spinal cord. This results in loss of coordination, weakness, and blurred vision. Autoimmune diseases are known in the art and include, for instance, Hashimoto's thyroiditis, Grave's disease, lupus, multiple sclerosis, rheumatic arthritis, hemolytic anemia, anti-immune thyroiditis, systemic lupus erythematosus, celiac disease, Crohn's disease, colitis, diabetes, scleroderma, psoriasis, and the like.

The term "degenerative disease" refers to any disease in which the function or structure of the affected tissues or organs will increasingly deteriorate over time. Examples include Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), Huntington's disease, macular degeneration, multiple sclerosis, muscular dystrophy, Niemann Pick disease, osteoporosis and rheumatoid arthritis.

The term "apoptosis-associated diseases" refers to any disease in which alterations of apoptosis are involved. Examples include cancer, neurological disorders, such as Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis (ALS) and stroke, heart diseases, such as ischemia reperfusion and chronic heart failure, infectious diseases and autoimmune diseases.

The term "transplant rejection" refers to the rejection of a transplanted tissue or organ by the recipient's immune system, which may, ultimately, destroy the transplanted tissue or organ.

In one embodiment, the use of the multispecific antibody, nucleic acid, expression vector, host cell or composition for use according to the present invention may be for treating cancer.

In one embodiment, the use of the multispecific antibody, nucleic acid, expression vector, host cell or composition for use according to the present invention may be for treating an infectious disease.

The present invention also relates to a method of treatment of a disease, such as cancer or an infectious disease, comprising administering the multispecific antibody according to the present invention, the nucleic acid according to the present invention,

the expression vector according to the present invention, the host cell according to claim the present invention, the composition according to the present invention, or the pharmaceutical composition according to the present invention to a subject in need thereof.

5 The present invention also relates to use of a multispecific antibody according to the present invention, the nucleic acid according to the present invention, the expression vector according to the present invention, the host cell according to the present invention, the composition according to the present invention, or the pharmaceutical composition according to the present invention for the manufacture of a medicament.

10 In one embodiment the method or use according to the present invention is for use in combination with one or more further therapeutic agent, such as a chemotherapeutic agent.

 In one aspect, the invention relates to the multispecific antibody, such as a bispecific antibody, according to any one of the embodiments disclosed herein, the
15 composition as disclosed herein, or the pharmaceutical composition as disclosed herein for use as a medicament.

 In another aspect, the present invention relates to the use of a multispecific antibody according to the present invention in the manufacture of a medicament for the treatment of a disease, such as cancer or an infectious disease.

20 In one aspect, the invention relates to the multispecific antibody according to any one of the embodiments disclosed herein, the composition as disclosed herein, or the pharmaceutical composition as disclosed herein for use in the treatment of a disease, such as cancer or an infectious disease.

 In one aspect, the invention relates to a method of treatment of a disease
25 comprising administering the multispecific antibody according to any one of the embodiments disclosed herein, the composition as disclosed herein, or the pharmaceutical composition as disclosed herein to a subject in need thereof.

 The multispecific antibodies of the invention may be used for a number of
30 purposes. In particular, the multispecific antibodies of the invention may be used for the treatment of various forms of cancer, including metastatic cancer and refractory cancer.

 In one embodiment the use according to the present invention is in combination with one or more further therapeutic agent, such as a chemotherapeutic agent.

 In particular, the multispecific antibodies according to the invention may be
35 useful in therapeutic settings in which increasing proliferation of T cells is relevant. An example of such a therapeutic setting includes but is not limited to cancer or tumors, such as hematological and solid tumors, e.g., melanoma, lung cancer, breast cancer, non-small cell lung cancer (NSCLC), colon cancer, renal cancer, cervical cancer and

prostate cancer, such as melanoma or NSCLC. Examples thereof are lung carcinomas, mamma carcinomas, prostate carcinomas, colon carcinomas, renal cell carcinomas, cervical carcinomas, or metastases of such cancer types or tumors.

- 5 The present invention also relates to a method for treating cancer, comprising
- a) selecting a subject suffering from a cancer, and
 - b) administering to the subject the multispecific antibody of the present invention or a pharmaceutical composition of the present invention.

Also, the invention relates to the use of a multispecific antibody that binds to
10 human CD40 and human CD137 for the preparation of a medicament for the treatment of cancer, such as one of the specific cancer indications mentioned herein.

The invention further relates to a multispecific antibody for use in the treatment of cancer, such as one of the cancer indications mentioned above.

In one embodiment the method or use according to the present invention is for
15 use in combination with one or more further therapeutic agent, such as a chemotherapeutic agent.

For the above mentioned uses any multispecific antibody, such as a bispecific antibody, of the present invention may be used.

20 In one aspect, the invention relates to a diagnostic composition comprising a multispecific antibody according to any one of the embodiments as disclosed herein.

In one embodiment, the diagnostic composition is a companion diagnostic which is used to screen and select those patients who will benefit from treatment with the multispecific antibody.

25 In a further embodiment of the methods of treatment of the present invention, the efficacy of the treatment is being monitored during the therapy, e.g. at predefined points in time, by determining tumor burden.

Dosage regimens in the above methods of treatment and uses are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a
30 single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage.

The efficient dosages and the dosage regimens for the multispecific antibodies
35 depend on the disease or condition to be treated and may be determined by the persons skilled in the art. An exemplary, non-limiting range for a therapeutically effective amount of a multispecific antibody of the present invention is about 0.001-30 mg/kg.

A physician or veterinarian having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the multispecific antibody employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a multispecific antibody of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Administration may e.g. be parenteral, such as intravenous, intramuscular or subcutaneous. In one embodiment, the multispecific antibodies may be administered by infusion in a weekly dosage of calculated by mg/m². Such dosages can, for example, be based on the mg/kg dosages provided above according to the following: dose (mg/kg) x 70: 1.8. Such administration may be repeated, e.g., 1 to 8 times, such as 3 to 5 times. The administration may be performed by continuous infusion over a period of from 2 to 24 hours, such as from 2 to 12 hours. In one embodiment, the multispecific antibodies may be administered by slow continuous infusion over a long period, such as more than 24 hours, in order to reduce toxic side effects.

In one embodiment the multispecific antibodies may be administered in a weekly dosage of calculated as a fixed dose for up to 8 times, such as from 4 to 6 times when given once a week. Such regimen may be repeated one or more times as necessary, for example, after 6 months or 12 months. Such fixed dosages can, for example, be based on the mg/kg dosages provided above, with a body weight estimate of 70 kg. The dosage may be determined or adjusted by measuring the amount of multispecific antibody of the present invention in the blood upon administration by for instance taking out a biological sample and using anti-idiotypic antibodies which target the CD137 antigen-antigen-binding region of the multispecific antibodies of the present invention.

In one embodiment, the multispecific antibodies may be administered as maintenance therapy, such as, e.g., once a week for a period of 6 months or more.

A multispecific antibody may also be administered prophylactically in order to reduce the risk of developing cancer, delay the onset of the occurrence of an event in cancer progression, and/or reduce the risk of recurrence when a cancer is in remission.

The multispecific antibodies of the invention may also be administered in combination therapy, i.e., combined with other therapeutic agents relevant for the disease or condition to be treated. Accordingly, in one embodiment, the multispecific antibody-containing medicament is for combination with one or more further therapeutic agents, such as a cytotoxic, chemotherapeutic or anti-angiogenic agent.

Such combined administration may be simultaneous, separate or sequential. For simultaneous administration the agents may be administered as one composition or as

separate compositions, as appropriate. The present invention thus also provides methods for treating a disorder, which methods comprise administration of a multispecific antibody of the present invention combined with one or more additional therapeutic agents as described below.

5 In one embodiment, the present invention provides a method for treating a disorder, which method comprises administration of a therapeutically effective amount of a multispecific antibody of the present invention, and optionally at least one additional therapeutic agent, to a subject in need thereof.

10 In one embodiment, the present invention provides a method for treating or preventing cancer, which method comprises administration of a therapeutically effective amount of a multispecific antibody of the present invention and at least one additional therapeutic agent to a subject in need thereof.

Pharmaceutical compositions of the invention can also be administered in combination therapy, i.e., combined with other agents, or combined with other treatment regimen. For example the multispecific antibodies may be combined with
15 radiotherapy and/or surgery and/or autologous or allogeneic peripheral stem cell or bone marrow transplantation.

Biomarkers

Thus, in one aspect, the present invention also relates to use of the multispecific
20 antibody as a biomarker.

In another aspect, the invention relates to a kit for detecting cross-linking between CD40- and CD137-expressing cells, in a sample obtained from a patient, such as a blood sample, lymph node sample or bone marrow sample, comprising

- 25 i) a multispecific antibody according to any one of the embodiments as disclosed herein; and
ii) instructions for use of said kit.

In a further aspect, the invention relates to a method for detecting whether cross-linking between CD40- and CD137-expressing cells occurs in a sample obtained from a patient, such as a blood sample, lymph node sample or bone marrow sample,
30 upon administration of a bispecific antibody according to any one of the embodiments as disclosed herein, comprising the steps of:

- (i) contacting the sample with a multispecific antibody according to any one of the embodiments as disclosed herein under conditions that allow for formation of a complex between said multispecific antibody and the CD40- and CD137-expressing cells;
35 and
(ii) analyzing whether a complex has been formed.

Detection of the complex can be done by methods known in the art, such as performed in Example 4, 5, 6, 10, 11 or 12.

Anti-idiotypic antibodies

5 In another aspect, the invention relates to an anti-idiotypic antibody which binds to the first and/or second antigen-binding region as defined in any one of the embodiments disclosed herein.

In a particular embodiment, the anti-idiotypic antibody binds to the first and/or second antigen-binding region of a multispecific antibody, wherein

- 10 - the first antigen-binding region comprises a VH sequence as set forth in SEQ ID NO:117 and a VL sequence as set forth in SEQ ID NO:121, and
- the second antigen-binding region comprises a VH sequence comprising SEQ ID NO:123 and a VL sequence comprising SEQ ID NO:127.

In one embodiment, the anti-idiotypic antibody binds to the first antigen-binding region defined in any one of the embodiments disclosed herein. In a specific
15 embodiment, the first antigen-binding region comprises a VH sequence comprising SEQ ID NO:117 and a VL sequence comprising SEQ ID NO:121.

In another embodiment, the anti-idiotypic antibody binds to the second antigen-binding region defined in any one of the embodiments disclosed herein. In a specific embodiment, the second antigen-binding region comprises a VH sequence comprising
20 SEQ ID NO:123 and a VL sequence comprising SEQ ID NO:127.

The present invention is further illustrated by the following examples, which should not be construed as limiting the scope of the invention.

EXAMPLES

25 *Example 1: Generation of antibodies*

The CD40 and each of the CD137 antibodies (i.e. clones 1-12) mentioned in **Table 1** above were produced with the VH and VL sequences described in **Table 1** and with a human κ light chain, and with a human IgG1 heavy chain. The CD40 antibody was produced with two different human IgG1 heavy chains; 1) a human IgG1 heavy chain
30 containing the following amino acid mutations: L234F, L235E, D265A and F405L (FEAL) wherein the amino acid position number is according to EU numbering (corresponding to SEQ ID NO:113); and 2) a human IgG1 heavy chain containing the following amino acid

mutations: L234F, L235E, D265A and K409R (FEAR) wherein the amino acid position number is according to EU numbering (corresponding to SEQ ID NO:112).

The CD137 antibodies were all produced with a human IgG1 heavy chain containing the following amino acid mutations: L234F, L235E, D265A and K409R (FEAR) wherein the amino acid position number is according to EU numbering (correspond to SEQ ID NO:112).

Similarly, a b12 antibody was produced comprising the VH and VL sequences mentioned in **Table 1**, and with a human IgG1 light chain and a human IgG heavy chain containing the following amino acid mutations: L234F, L235E, D265A and F405L (FEAL) wherein the amino acid position number is according to EU numbering (correspond to SEQ ID NO:113).

Antibodies were produced under serum-free conditions by co-transfecting relevant heavy and light chain expression vectors in Expi293F™ cells (ThermoFisher catalogue number A14527), using ExpiFectamine™ 293 (ThermoFisher catalogue number A14525), according to the manufacturer's instructions. Antibodies were purified by protein A affinity chromatography and buffer exchanged into 12.6 mM NaH₂PO₄, 140 mM NaCl, pH 7.4 buffer (B.Braun or Thermo Fisher). After buffer exchange, samples were sterile filtered over 0.2 µm dead-end filters. Purified proteins were analyzed by CE-SDS and HP-SEC. Concentration was measured by absorbance at 280 nm. Purified antibodies were stored at 2-8°C.

Example 2: DNA shuffling between wild boar or elephant and human CD137 to determine domains important for binding of anti CD137 antibodies

To determine domains important for binding of anti CD137 antibodies to human CD137, DNA shuffling was performed between human and wild boar CD137 (*sus scrofa*; XP_005665023) or between human and African elephant CD137 (*loxodonta africana*; XP_003413533). Shuffle constructs were prepared from DNA encoding human CD137, by replacing human domains with wild boar (shuffle construct 1-4, 6) or elephant (shuffle construct 5) domains. If a domain in human CD137 is important for binding of an anti CD137 antibody, binding will be lost upon replacement of that domain with the wild boar or African elephant domain. Requirement is that the antibody does not bind to the whole CD137 elephant or wild boar sequence Homology between human and wild boar and between human and African elephant CD137 is 70.2 and 74.5%, respectively. **Figure 1** shows sequence alignments of human, wild boar and African elephant CD137. **Figure 2** shows the constructs for human CD137 containing wild boar CD137 or African elephant domains, as indicated.

3 x 10⁶ HEK293T-17 cells were seeded in T75 culture flasks (Greiner Bio-One, cat. no. 658175) in 20 mL RPMI 1640 GlutaMAX medium containing 10% FCS (Biochrom, cat. no. S 0115). After O/N incubation, cells were transiently transduced with expression vectors encoding the shuffle constructs or the wild boar, African elephant or human CD137 downstream of a constitutively active human elongation factor-1 alpha (EF-1 alpha) promoter using TransIT®-LT1 Transfection Reagent, Mirus Bio (VWR International, cat. no. 731-0029), according to the manufacturer's instructions. The next day, cells were harvested using 1.5 mL Accutase (Sigma Aldrich, cat. no. A6964) (incubation at 37°C for 5 min.) and flow cytometry was performed, essentially as described in Example 4, to measure surface expression of the shuffle constructs and the human, African elephant and wild boar CD137 and to measure binding of the antibody clones to the different shuffle constructs. To measure cell surface expression of the constructs, transduced cells were incubated with 1 µg/mL goat polyclonal anti-human CD137 (R&D Systems, cat. no. AF838) in FACS buffer (D-PBS supplemented with 5 mM EDTA [Sigma Aldrich, cat. no. 03690] and 5% (v/v) fetal bovine serum [FBS, Biochrom, cat. no. S 0115]) (4°C, 20 min.), followed by incubation with APC-labeled anti-goat IgG (H+L) (R&D Systems, cat. no. F0108) (4°C, 20 min.). Binding of the different CD137 antibody clones to cells expressing the shuffle constructs was measured by incubation of the transduced cells with 1 µg/mL of the antibody clones, followed by APC-labeled AffiniPure F(ab')₂ Fragment (1:50 final dilution; Jackson, cat. no. 109-136-127).

All CD137 shuffle constructs, as well as human, African elephant and wild boar CD137, were expressed on the cell surface with similar expression levels (**Figure 3**).

Table 2 and **Figure 4** show that all clones, except for clone 1, showed loss of binding to at least African elephant or wild boar CD137. Clones 2, 3, 4, 5, 6, 7, 8, 9, 10 and 11 showed loss of binding to at least one of the shuffle constructs. Clone 1 showed reduced binding to African elephant and shuffle construct 5, as compared to binding to human CD137. Clone 12 did not show loss of binding to any of the shuffle constructs, but showed reduced binding to shuffle construct 5. None of the clones showed loss of binding or reduced binding to shuffle constructs 1 and 2.

Table 2: Summary of binding of the CD137 antibodies to the shuffle constructs

	Binding similar to human CD137 binding	Binding decreased compared to human CD137 binding	No binding
Wild boar CD137	Clone 1, 2, 6, 10	None	Clone 3, 4, 5, 7, 8, 9, 11, 12
African elephant CD137	None	Clone 1,3	Clone 2, 4, 5, 6, 7-12
Shuffle 1 (aa 162-	Clone 1-12	None	None

196)			
Shuffle 2(139-161)	Clone 1-12	None	None
Shuffle 3 (115-138)	Clone 1, 2, 5, 6, 9, 10, 12	Clone 3, 7, 8	Clone 4, 11
Shuffle 4 (89-114)	Clone 1, 2, 5, 6, 9, 10, 12	None	Clone 3, 4, 7, 8, 11
Shuffle 5 (48-88)	Clone 3, 4, 5, 7, 8, 11	Clone 1, 12	Clone 2, 6, 9, 10
Shuffle 6 (aa 24-47)	Clone 1, 2, 3, 4, 6, 7-12	None	Clone 5

Example 3: Generation of bispecific antibodies by 2-MEA-induced Fab-arm exchange

A method to produce stable IgG1-based bispecific antibodies is described in WO2011131746 (Genmab). The bispecific antibody product generated by this method described below will no longer participate in Fab-arm exchange. The basis for this method was the use of complimentary CH3 domains, which promote the formation of heterodimers under specific assay conditions. To enable the production of bispecific antibodies by this method, IgG1 molecules carrying certain mutations in the CH3 domain were generated: in one parental IgG1 antibody T350I, K370T and F405L mutations (or minimally F405L), in the other parental IgG1 antibody a K409R mutation.

The concentrations of parental IgG1 antibodies that minimally contained either an F405L or a K409R point mutation were measured using their absorption at 280 nm. Specific extinction coefficients based upon the amino acid sequence were used to infer the protein concentration.

The Cube system is Genmab's flexible robotic work cell. The system was designed and built in collaboration with Peak Analysis and Automation (PAA), Farnborough UK.

Bispecific antibodies were generated by combining the following antibodies from Example 1:

- CD40-FEAL antibody combined with each of the CD137-FEAR antibodies,
- CD40-FEAR antibody combined with the b12-FEAL antibody, and
- Each of the CD137-FEAR antibodies combined with the b12-FEAL antibody

The bispecific antibody discovery process is performed in an automated fashion on the Cube system, as shown in **Figure 5** and described below.

To generate bispecific antibodies, the following (automated) steps are performed:

- Depending on the volume required, deep well source plates (96 well clear V-bottom 2 mL polypropylene deep well plate, Corning, cat. no. 3960; 48 well Riplate® SW 5mL, Ritter, cat. no. 43001-1062 ; 24 well Riplate SW 10 mL, Ritter, cat. no. 43001-1066) are filled with parental antibodies (F405L- and K409R-containing antibodies in

different plates) at a concentration of 1.0 mg/mL (in 1x PBS, B.Braun) (**Figure 5**, left plates).

- From these source plates, the pre-grid plates (96 well V-bottom, Corning) are prepared by the Cube, according to **Figure 5**, middle plates. For each combination with a parental antibody in the exchange grid, 67.5 μ L parental antibody is added to the appropriate pre-grid plate.

- After the pre-grid, the exchange is performed. Here, two parental antibodies (67.5 μ L, 1.0 mg/mL each), each from a different pre-grid plate are added to an exchange plate (96 well round-bottom polypropylene plate, Greiner, cat. no. 650293), each antibody at a final concentration of 0.5 mg/mL (equimolar concentration) (**Figure 5**, right plates).

- The exchange reaction is started by adding 15 μ L 75 mM 2-mercaptoethylamine-HCl (2-MEA) (in 1x PBS, B.Braun) to the exchange plate. The total volume in the exchange plate is now 150 μ L (final concentration 2-MEA 7.5 mM)

- The exchange plates are incubated at 31°C for 5 hours in the Cytomat 6000 automated incubator (Thermo Scientific).

- The reducing agent 2-MEA is removed by using desalting columns (PhyTip desalting columns, 600 μ L resin, PhyNexus, cat. no. PDR 91-60-06), for which flow is based on gravity.

o The columns are conditioned by placing an adapter with 96 columns on a waste position, adding two times 450 μ L 1x PBS (B.Braun) and allowing the solutions to flow through the columns into the waste.

o After conditioning, 100 μ L sample from the exchange plate is added, thereby pushing the remaining PBS out of the columns.

o After allowing the solutions to flow through the columns into the waste, the adapter with columns is placed on a desalting (or destination) plate (96 well round-bottom, Greiner).

o The remaining sample from the exchange plate is added to the columns.

o After allowing the samples to flow through the columns into the desalting plate, 225 μ L 1x PBS (B.Braun) is added to the columns and the sample is eluted into the desalting plate.

o The 2-MEA remains inside the columns. Where appropriate, the columns can be regenerated by washing with 1x PBS (B.Braun).

- The desalting plates are stored in the Cytomat 6001 automated incubator (Thermo Scientific) at 8°C. These plates now contain the bispecific antibodies.

The final bispecific antibody samples were filtered over 0.2 μ m dead-end filters and the absorbance at 280 nm (A280) of bispecific products was measured to determine

the final concentration. Samples were stored at 2-8 °C for at least 24 hours before further use.

The bispecific antibody exchange efficiency was quantified using High Pressure Liquid Chromatography (HPLC) – hydrophobic interaction chromatography (HIC) using a Butyl-NPR, 2.5 µm, 4.6 x 35 mm HIC-HPLC column (Tosoh Bioscience) with a flow rate of 1 mL/min. Parental antibodies and analysis samples were normalized in concentration and diluted two-fold with HIC eluent A (15.4 mM K₂HPO₄, 9.6 mM KH₂PO₄, 1.5 M (NH₄)₂SO₄; pH 7.0). 50 µL of sample was injected and elution was performed with a 12-min gradient of HIC eluent A to HIC eluent B (15.4 mM K₂HPO₄, 9.6 mM KH₂PO₄; pH 7.0) with detection at 280 nm. Alternatively, HPLC - analytical cation exchange chromatography (CIEX) was used to quantify the bispecific antibody exchange efficiency. Parental antibodies and analysis samples at 1 mg/mL in mobile Phase A (10 mM NaPO₄, pH 7.0) were injected onto the HPLC. The differently charged IgG molecules were separated by using a ProPac WCX-10, 4 mm x 250 mm, analytical column with a flow rate of 1 mL/min. 50 µL of sample was injected and elution was performed with a gradient of Mobile Phase A (10 mM NaPO₄, pH 7.0; prepared from a 0.1 M stock of sodium phosphate buffer, that was obtained by adding 10.3 g Na₂HPO₄·2H₂O and 5.07 g NaH₂PO₄ per liter Milli-Q) to Mobile Phase B (10 mM NaPO₄, pH 7.0, 0.25 M NaCl) with detection at 280 nm. Empower 3 software (Waters) was used to assign peaks as parental antibodies or bispecific reaction products, and to integrate peak areas to quantify extent of the bispecific antibody exchange reaction. Bispecific antibody reaction products were further analyzed using analytical size exclusion chromatography, using a TSK HP-SEC column (G3000SWxl; Tosoh Biosciences, via Omnilabo, Breda, The Netherlands) and Capillary Electrophoresis – Sodium Dodecyl Sulfate (CE-SDS) using a LabChip GXII (Caliper Life Sciences, MA) on a HT Protein Express LabChip (Caliper Life Sciences, MA) under reducing and non-reducing conditions according to manufacturer's instructions.

Example 4: Reporter assay measuring trans-activation by bispecific antibodies binding to CD40 and CD137

CD40 is predominantly expressed on antigen-presenting cells (APCs), such as dendritic cells, whereas CD137 is predominantly expressed on activated T cells. Thus, bispecific antibodies binding to CD40 and CD137 can bind simultaneously to APCs and T cells expressing these receptors. Thereby, these bispecific antibodies can mediate cell-cell contact between APCs and T cells by receptor binding and activate both receptors. This receptor activation can be induced by cross-linking and receptor clustering upon cell-cell interaction and is not necessarily dependent on agonistic activity of the parental

monospecific bivalent antibodies. Thus, these trans-activating bispecific antibodies can exert co-stimulatory activity in the context of interactions between APCs and T cells.

A reporter assay system was established to measure activation of each receptor by the bispecific antibodies. NF- κ B/293/GFP-LucTM Transcriptional Reporter Cell Line (System Biosciences; cat. no. TR860A-1) is a reporter cell line designed for monitoring the NF- κ B signal transduction pathway in vitro. Activation of the NF- κ B pathway can be monitored by the detection of green fluorescent protein (GFP) fluorescence as well as luciferase activity for quantitative transcription activation reporter assays. NF- κ B/293/GFP-Luc cells were stably transduced with expression vectors encoding full length human CD40 or CD137 downstream of a constitutively active human elongation factor-1 alpha (EF-1 alpha) promoter, using TransIT®-LT1 Transfection Reagent, Mirus Bio (VWR International, cat. no. 731-0029), according to the manufacturer's instructions. Stable clones were selected using 10 mg/mL blasticidin (Invivogen, cat. no. ant-bl-1). In addition, K562 cells were stably transduced, as described supra, with human CD40 and CD137 to generate cell lines that can provide the corresponding target antigen for the other arm of the bispecific antibody. Cell surface expression of the receptors was measured by flow cytometry. 0.3×10^6 cells were spun down (460 x g, 5 min.) and washed in FACS buffer (D-PBS supplemented with 5 mM EDTA [Sigma Aldrich, cat. no. 03690] and 5% (v/v) fetal bovine serum [FBS, Biochrom, cat. no. S 0115]) (460 x g, 5 min.). 50 μ L of 1:50 diluted allophycocyanin (APC)-labeled anti-human CD40 (BD Biosciences, clone 5C3, cat. no. 555591) or phycoerythrin (PE)-labeled anti-human CD137 (BD Biosciences, clone 4B4-1, cat. no. 555956) was added to the cell pellet and incubated at 4°C in the dark for 20 minutes. After washing three times with FACS buffer, cells were resuspended in 100 μ L FACS buffer and binding of the antibodies was detected by flow cytometry on a FACSCanto II (BD Biosciences). Cell surface expression of CD40 and CD137 on transduced NF- κ B/293/GFP-Luc cells (**Figure 6A**) and K562 cells (**Figure 6B**) was nicely shown.

The reporter assay measuring trans-activation was set up as follows: NF- κ B/293/GFP-Luc cells expressing one of the two indicated TNF receptors were seeded at 10,000 cells/well in 30 μ L RPMI 1640 medium with GlutaMAX supplement (Life Technologies, cat. no. 61870) in white opaque 384-well cell culture plates (PerkinElmer, cat. no. 6007680). Bispecific antibodies binding with one arm to CD40 and with the other arm to CD137 and the corresponding monospecific, monovalent (containing one irrelevant control arm [b12]) control antibodies were added in 10 μ L/well to the reporter cells in serial dilutions (in medium), ranging from 0.078 μ g/mL to 10 μ g/mL final concentration, including a buffer control. 17,000 K562 cells expressing the second TNF receptor or wildtype K562 (K562_wt) cells were added in 10 μ L medium to each well and incubated at 37°C and 5% CO₂ for 18 hours. Thus, the bispecific antibodies are able to

bind to the first TNF receptor on the NF- κ B/293/GFP-Luc cell line and, at the same time, to the second TNF receptor on the K562 cell line. Only receptor activation of the first TNF receptor on NF- κ B/293/GFP-Luc cells is measured by luciferase activity induced upon NF- κ B signaling. Thus, bispecific antibodies targeting CD40 and CD137 were analyzed by two reporter assays: the first assay measuring CD137 activation induced by simultaneous binding of CD137 on the reporter cell line and CD40 on the K562 cells (HEK293_NFK_CD137_gfp_luc + K562_CD40) and the second assay measuring CD40 activation induced by simultaneous binding of CD40 on the reporter cell line and CD137 on the K562 cells (HEK293_NFK_CD40_gfp_luc + K562_CD137). Luciferase activity was measured as relative luminescence units on an Envision plate reader (PerkinElmer) after addition of 50 μ L/well Steady-Glo® reagent (Promega; cat. no. E2520) reconstituted in Glo Lysis Buffer (Promega; cat. no. E266A) and incubation at room temperature for 30 min.

Only the bispecific CD40xCD137 antibodies (**Figure 7 A-L**, lower panels, first and third graph) induced luciferase activity (at concentrations of about 100 ng/mL and higher) in NF- κ B/293/GFP-Luc cells transduced either with CD137 or with CD40, under trans-activation conditions (incubation with K562-CD40 or K562-CD137, respectively). None of the monospecific, monovalent (containing one irrelevant control arm [b12]) control antibodies induced luciferase activity in the transduced NF- κ B/293/GFP-Luc cells (upper panels). Furthermore, in the absence of trans-activation conditions (using wildtype K562 cells) no luciferase activity was induced by the bispecific CD40xCD137 antibodies (lower panels, second and fourth panel).

Example 5: Non-antigen-specific T-cell proliferation assay to measure trans-activation by bispecific antibodies binding to CD40 and CD137

To measure non-antigen-specific proliferation, T cells in a peripheral blood mononuclear cell (PBMC) population were incubated with a sub-optimal concentration of anti-CD3 antibody (clone UCHT1), combined with CD40xCD137 bispecific or control antibodies. Within this PBMC population, antigen-presenting cells expressing CD40 can be bound by the CD40-specific arm of the bispecific antibody, whereas the T cells in the population can be bound by the CD137-specific arm. Trans-activation of the T cells induced by cross-linking to the antigen-presenting cells via the bispecific antibody is measured as T-cell proliferation.

PBMCs were obtained from buffy coats of healthy donors (Transfusionszentrale, University Hospital, Mainz, Germany) using a Ficoll gradient (VWR, cat. no. 17-5446-02). PBMCs were labeled using 1.6 μ M carboxyfluorescein succinimidyl ester (CFSE) (Thermo Fisher, cat. no. C34564) in PBS, according to the manufacturer's instructions. 75,000

CFSE-labeled PBMCs were seeded per well in a 96-well round-bottom plate (Sigma Aldrich, CLS3799-50EA) and incubated with a sub-optimal concentration of anti-CD3 antibody (R&D Systems, clone UCHT1, cat. no. MAB100; 0,01-0,1 µg/mL final concentration) that was pre-determined for each donor, and bispecific or control antibodies, in 150 µL IMDM GlutaMAX supplemented with 5% human AB serum, at 37°C, 5% CO₂, for four days. Proliferation of CD4⁺ and CD8⁺ T cells was analyzed by flow cytometry, essentially as described supra. 30 µL containing PE-labeled CD4 antibody (BD Biosciences, cat. no. 555347; 1:80 final dilution), PE-Cy7-labeled CD8α antibody (clone RPA-T8, eBioscience, cat. no. 25-0088-41; 1:80 final dilution) APC-labeled CD56 antibody (eBiosciences, cat. no. 17-0567; 1:80 final dilution) and 7-AAD (Beckman Coulter, cat. no. A07704; 1:80 final dilution) in FACS buffer was used to stain the cells and exclude natural killer (NK) cells (CD56) and dead cells (7-AAD) from the analysis. Samples were measured on a FACSCanto II (BD Biosciences). Detailed analyses of T-cell proliferation based on CFSE-peaks indicating cell divisions were made by FlowJo 7.6.5 software. Mean percentages of T cells that went into division (% divided cells) and the average number of divisions of cells that went into division (proliferation index) were calculated.

Figure 8A shows that only the CD40xCD137 bispecific antibodies efficiently enhanced proliferation of CD8⁺ T cells. The control monospecific, monovalent antibodies (b12xCD40; b12xCD137) and the combination of monospecific, monovalent CD40 with monospecific, monovalent CD137 antibodies (b12xCD40 + b12xCD137) did not induce more proliferation than observed in the control (only the weakly activated PBMCs, ctrl, w/o). The flow cytometry histograms, for the different antibodies at different concentrations, were quantified to indicate percentage of divided cells (**Figure 8B**) and proliferation index (**Figure 8C**), as described supra. These figures show that only the bispecific antibodies were capable of inducing proliferation of CD8⁺ cells, with an optimum between 0.04 and 0.2 µg/mL.

Example 6: Antigen-specific CD8⁺ T cell proliferation assay to measure trans-activation by bispecific antibodies binding to CD40 and CD137

To measure induction of proliferation by the bispecific antibodies in an antigen-specific assay, dendritic cells (DCs) were transfected with claudin 6 in vitro-transcribed RNA (IVT-RNA,) to express the claudin 6 antigen. T cells were transfected with the claudin-6-specific, HLA-A2-restricted T cell receptor (TCR). This TCR can recognize the claudin-6-derived epitope presented in HLA-A2 on the DC. The CD40xCD137 bispecific antibody can cross-link CD40 on the dendritic cell and CD137 on the T cell, leading to

activation of the DC and a co-stimulatory signal to the T cell, resulting in T-cell proliferation.

HLA-A2⁺ PBMCs were obtained from healthy donors (Transfusionszentrale, University Hospital, Mainz, Germany). Monocytes were isolated from PBMCs by
5 magnetic-activated cell sorting (MACS) technology using anti-CD14 MicroBeads (Miltenyi; cat. no. 130-050-201), according to the manufacturer's instructions. The peripheral blood lymphocytes (PBLs, CD14-negative fraction) were frozen for future T-cell isolation. For differentiation into immature DCs (iDCs), 1×10^6 monocytes/mL were cultured in RPMI GlutaMAX (Life technologies GmbH, cat. no. 61870-044) containing 5%
10 human AB serum (Sigma-Aldrich Chemie GmbH, cat. no. H4522-100ML), sodium pyruvate (Life technologies GmbH, cat. no. 11360-039), non-essential amino acids (Life technologies GmbH, cat. no. 11140-035), 100 IU/mL penicillin-streptomycin (Life technologies GmbH, cat. no. 15140-122), 1000 IU/mL granulocyte-macrophage colony-stimulating factor (GM-CSF; Miltenyi, cat. no. 130-093-868) and 1000 IU/mL interleukin
15 4 (IL-4; Miltenyi, cat. no. 130-093-924), for five days. Once during these five days, half of the medium was replaced with fresh medium. iDCs were harvested by collecting non-adherent cells; adherent cells were detached by incubation with PBS containing 2mM EDTA for 10 min at 37°. After washing, iDCs were frozen in RPMI GlutaMAX containing 10% v/v DMSO (AppliChem GmbH, cat. no. A3672,0050) and 50% v/v human AB serum
20 for future antigen-specific T cell assays.

One day before T-cell assays were started, frozen PBLs and iDCs, from the same donor, were thawed. PBLs were used for isolation of CD8⁺ T cells by MACS technology using anti-CD8 MicroBeads (Miltenyi, cat. no. 130-045-201), according to the manufacturer's instructions. About $10\text{--}15 \times 10^6$ CD8⁺ T cells were electroporated with 10
25 µg IVT-RNA encoding the alpha-chain plus 10 µg of IVT-RNA encoding the beta-chain of a claudin-6-specific murine TCR (HLA-A2-restricted; described in WO 2015150327 A1) in 250 µL X-Vivo15 (Biozym Scientific GmbH, cat. no. 881026) in a 4-mm electroporation cuvette (VWR International GmbH, cat. no. 732-0023) using the BTX ECM® 830 Electroporation System device (BTX; 500 V, 1 x 3 ms pulse). Immediately after
30 electroporation, cells were transferred into fresh IMDM medium (Life Technologies GmbH, cat. no. 12440-061) supplemented with 5% human AB serum and rested at 37°C, 5% CO₂ for at least 1 hour. T cells were labeled using 1,6 µM carboxyfluorescein succinimidyl ester (CFSE; Invitrogen, cat. no. C34564) in PBS, according to the manufacturer's instructions, and incubated in IMDM medium supplemented with 5%
35 human AB serum, O/N.

Up to 5×10^6 thawed iDCs were electroporated with 0,4–5 µg IVT-RNA encoding full length claudin-6 (Uniprot P56747), in 250 µL X-Vivo15 medium, using the

electroporation system as described above (300 V, 1x12 ms pulse) and incubated in IMDM medium supplemented with 5% human AB serum, O/N.

The next day, cells were harvested. Cell surface expression of claudin-6 on DCs and TCR on T cells were checked by flow cytometry. Therefore, DCs were stained with an Alexa647-conjugated CLDN6-specific antibody (not commercially available; in-house production) and T cells were stained with an anti-mouse TCR β Chain antibody (Becton Dickinson GmbH, cat. no. 553174). 5,000 electroporated DCs were incubated with 50,000 electroporated, CFSE-labeled T cells in the presence of bispecific or control antibodies in IMDM GlutaMAX (Life Technologies, cat. no. 12440-061) supplemented with 5% human AB serum in a 96-well round-bottom plate. T-cell proliferation was measured after 5 days by flow cytometry. Detailed analyses of T-cell proliferation based on CFSE-peaks indicating cell divisions were made by FlowJo 7.6.5 software. Mean percentages of T cells that went into division (% divided cells) and the average number of divisions of cells that went into division (proliferation index) were calculated.

Figure 9A shows that only the CD40xCD137 bispecific antibodies efficiently enhanced proliferation of CD8⁺ T cells. The control monospecific, monovalent antibodies (b12xCD40; b12xCD137) and the combination of monospecific, monovalent CD40 with monospecific, monovalent CD137 antibodies (b12xCD40 + b12xCD137) did not induce more proliferation than observed in the control (only the weakly activated PBMCs, ctrl, w/o). The same is also reflected in the percentage of divided cells (**Figure 9B**) and is very clear from the proliferation index (**Figure 9C**). **Figure 9D** shows that the induction of antigen-specific proliferation by the CD40xCD137 bispecific antibodies was concentration dependent, with an optimum around 0.1 μ g/mL in this assay.

Example 7: Humanization of murine and rabbit antibodies

Humanized antibody sequences from the antibodies mouse anti-CD40-001 and rabbit anti-CD137-009 were generated at Antitope (Cambridge, UK). Humanized antibody sequences were generated using germline humanization (CDR-grafting) technology. Humanized V region genes were designed based upon human germline sequences with closest homology to the VH and Vk amino acid sequences of the murine and rabbit antibodies. A series of four to six VH and four or five Vk (VL) germline humanized V-region genes were designed for each of the non-human parental antibodies. Structural models of the non-human parental antibody V regions were produced using Swiss PDB and analyzed in order to identify amino acids in the V region frameworks that may be important for the binding properties of the antibody. These amino acids were noted for incorporation into one or more variant CDR-grafted

antibodies. The closest matching germline sequences used as the basis for the humanized designs are shown in **Table 3**.

Table 3: Closest matching human germline V segment and J segment sequences.

Antibody	Heavy chain		Light chain (κ)	
	Human V region germline segment	Human J region germline segment	Human V region germline segment	Human J region germline segment
Mouse anti-CD40-001	hIGHV1-46*01	hIGHJ4	hIGKV1-33*01	IGKJ4
Rabbit anti-CD137-009	hIGHV3-49*04	hIGHJ4	hIGKV1-33*01	IGKJ4

- 5 Variant sequences with the lowest incidence of potential T cell epitopes were then selected using Antitope's proprietary *in silico* technologies, iTope™ and TCED™ (T Cell Epitope Database) (Perry, L.C.A, Jones, T.D. and Baker, M.P. New Approaches to Prediction of Immune Responses to Therapeutic Proteins during Preclinical Development (2008). *Drugs in R&D* 9 (6): 385-396; 20; Bryson, C.J., Jones, T.D. and Baker, M.P. Prediction of Immunogenicity of Therapeutic Proteins (2010). *Biodrugs* 24 (1):1-8).
 10 Finally, the nucleotide sequences of the designed variants were codon-optimized for expression in human cells.

The variable region sequences of the humanized CD40 and CD137 antibodies are shown in the Sequence Listing and in Table 1 above.

15 *Example 8: Expression constructs for antibodies, transient expression and purification*

- For antibody expression the VH and VL sequences were cloned in expression vectors (pcDNA3.3) containing, in case of the VH, the relevant constant heavy chain (HC), in certain cases containing a F405L or K409R mutation, and/or L234F, L235E and D265A, and, in case of the VL, light chain (LC) regions. Antibodies were expressed as
 20 IgG1, κ . Plasmid DNA mixtures encoding both heavy and light chains of antibodies were transiently transfected in Expi293F cells (Life technologies, USA) using 293fectin (Life technologies) essentially as described by Vink et al. (Vink et al., *Methods*, 65 (1), 5-10 2014). Next, antibodies were purified by immobilized protein G chromatography.

Example 9: Non-specific T-cell proliferation assay to test the functionality of a humanized bispecific antibody binding to CD40 and CD137

To measure the functionality of a humanized bispecific antibody binding to CD40 and CD137, a non-antigen-specific T-cell proliferation assay was performed as described
5 supra. In short, PBMCs of one donor were CFSE-labeled and incubated with a sub-optimal concentration of anti-CD3 antibody (clone UCHT1; 0.01 µg/mL as determined for this donor) and 0.008, 0.04, 0.2 or 1 µg/mL humanized CD40xCD137 bispecific antibody, the parental bispecific antibody or IgG1 control antibody.

Proliferation of CD8⁺ T cells was analyzed by flow cytometry, essentially as
10 described supra. Detailed analyses of T-cell proliferation based on CFSE-peaks indicating cell divisions were made by FlowJo 7.6.5 software. Mean percentages of T cells that went into division (% divided cells) and the average number of divisions of cells that went into division (proliferation index) were calculated.

Figure 10 shows that the humanized CD40xCD137 bispecific antibody (BisG1-
15 CD40-001-HC6LC1-FEALxCD137-009-HC7LC2-FEAR) efficiently enhanced proliferation of CD8⁺ T cells. The humanized bispecific antibody enhanced both the percentage of divided cells and the average number of divisions of CD8⁺ cells. Efficacy of the humanized bispecific antibody was comparable to that of the parental bispecific antibody (CD40-001xCD137-009).

20 *Example 10: Antigen-specific CD8⁺ T-cell proliferation assay to test the functionality of the humanized bispecific antibody binding to CD40 and CD137*

To measure the functionality of the humanized bispecific antibody binding to CD40 and CD137, an antigen-specific CD8⁺ T-cell proliferation assay was performed as described *supra*. In short, CFSE-labeled, CLDN6-TCR transfected CD8⁺ T-cells were
25 incubated with CLDN6 RNA-electroporated DCs in the presence of humanized CD40xCD137 bispecific antibody, the parental antibody or IgG1 control antibody. T-cell proliferation was measured by flow cytometry after 4 days. Detailed analyses of T-cell proliferation based on CFSE-peaks indicating cell divisions were made by FlowJo 7.6.5 software. Mean percentages of T cells that went into division (% divided cells) and the
30 average number of divisions of cells that went into division (proliferation index) were calculated.

Figure 11 shows that the humanized CD40xCD137 bispecific antibody (BisG1-
CD40-001-H6LC1-FEALxCD137-009-HC7LC2-FEAR) efficiently enhanced proliferation of CD8⁺ T cells. Efficacy of the humanized bispecific antibody was comparable to that of the
35 parental bispecific antibody (CD40-001xCD137-009). Both the humanized and the

parental bispecific antibody enhanced the percentage of divided cells as well as the proliferation index of the CD8+ cells in this assay.

Example 11: Ex vivo TIL expansion assay to evaluate the effects of CD40xCD137 bispecific antibodies on tumor infiltrating lymphocytes

5 To evaluate the effects of CD40xCD137 bispecific antibody (BisG1-CD40-001-FEAL/CD137-009-FEAR) on tumor infiltrating lymphocytes (TIL), *ex vivo* culture of human tumor tissue was performed as follows. Freshly human tumor tissue resections were washed three times by transferring the isolated tumor chunks from one wash medium-containing well of a six-well plate (Fisher Scientific cat. no. 10110151) to the
10 next using a spatula or serological pipette. Wash medium was composed of X-VIVO 15 (Biozym, cat. no. 881024) supplemented with 1% Pen/Strep (Thermo Fisher, cat. no. 15140-122) and 1% Fungizone (Thermo Fisher, cat. no. 15290-026). Next, the tumor was dissected with a surgical knife (Braun/Roth, cat. no. 5518091 BA223) and cut into tumor pieces with a diameter of about 1-2 mm. Two pieces each were put into one well
15 of a 24-well plate (VWR international, cat. no. 701605) containing 1 mL TIL medium (X-VIVO 15, 10% Human Serum Albumin (HSA, CSL Behring, cat. no. PZN-6446518), 1% Pen/Strep, 1% Fungizone and IL-2 (Proleukin®S, Novartis Pharma, cat. no. 02238131) at the indicated concentration. Bispecific antibody binding to CD40 and CD137 was added at the indicated final concentrations. Culture plates were incubated at 37°C and
20 5% CO₂ for 72 hours and 1 mL fresh TIL medium containing the indicated IL-2 concentration and the indicated concentration of bispecific antibody was added to each well. Wells were monitored for the occurrence of TIL clusters using a Leica DMI1 microscope equipped with a 5.0 megapixel camera, every other day. Wells were split on an individual basis, when more than 25 TIL microclusters were detected. To split TIL
25 cultures, cells were re-suspended and transferred to a well of a 6-well plate and supplemented with another 2 mL of TIL medium.

After a total culture period of 10-14 days, TILs were harvested and analyzed by flow cytometry. To allow for quantitative comparison of the different treatment groups, cell pellets were re-suspended after the last washing step with FACS-buffer
30 supplemented with BD™ CompBeads (BD biosciences, cat. no. 51-90-9001291). Flow cytometric analysis was performed on a BD FACSCanto™ II flow cytometer (Becton Dickinson) and acquired data was analyzed using FlowJo 7.6.5 software. The relative viable TIL count (7-AAD-negative cells) per 1,000 beads was calculated for each well.

35 **Figure 12** shows the analysis of a TIL expansion from a human melanoma tissue. Here, 100 U/mL IL-2 was used as supplement for the TIL medium. Moreover, the

following concentrations of the bispecific antibody binding to CD40 and CD137 (BisG1-CD40-001-FEAL/CD137-009-FEAR) were added: 0.016, 0.08, 0.4, 2.0 and 10.0 µg/mL; wells without antibody addition served as negative control. After 14 days of culture, TILs were harvested and analyzed by flow cytometry. Five samples for each antibody concentration, derived from different wells of the 24-well plate, were measured. In all samples cultured with the bispecific antibody binding to CD40 and CD137 the viable count of TIL was substantially increased in comparison to the control samples without antibody. Overall, about a 100-fold increase of the mean relative viable TIL count was observed (**Figure 12**).

10

Figure 13 shows the analysis of a TIL expansion from a non-small cell lung cancer (NSCLC) tissue. Here, 10 U/mL IL-2 was used as supplement for the TIL medium. In addition, the following final concentrations of the bispecific antibody binding to CD40 and CD137 (BisG1-CD40-001-FEAL/CD137-009-FEAR) were administered: 0.01, 0.1, and 1.0 µg/mL; wells without antibody addition served as negative control. After 10 days of culture, TILs were harvested and analyzed by flow cytometry. Five samples for each antibody concentration, derived from different wells of the 24-well plate, were measured. In all samples cultured with the bispecific antibody binding to CD40 and CD137, the viable count of TIL was substantially increased in comparison to the control samples without antibody. Overall, an up to 10-fold increase of the mean relative viable TIL count was observed at 0.1 or 1 µg/mL (**Figure 13**).

15
20

CLAIMS

1. A multispecific antibody comprising
 - (I) a first antigen-binding region binding to human CD40, wherein said first antigen-binding region comprises heavy and light chain variable region CDR1, CDR2, and CDR3 selected from the group consisting of:
 - a) heavy chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:4, YTS and 5, respectively;
 - b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations; and
 - c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD40 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD40 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b), and
 - (II) a second antigen-binding region binding to human CD137.
2. The multispecific antibody according to claim 1, wherein said first antigen-binding region comprises a first heavy chain variable (VH) sequence and a first light chain variable (VL) sequence, said second antigen-binding region comprises a second VH sequence and a second VL sequence, and said first and second VH and VL sequences each comprises three CDR sequences, CDR1, CDR2 and CDR3, respectively, and four framework sequences, FR1, FR2, FR3 and FR4, respectively.
3. The multispecific antibody according to any one of the preceding claims, wherein said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:4, YTS and 5, respectively.
4. The multispecific antibody according to any one of claims 2-3, wherein said VH sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to at least one of SEQ ID NOS:117 and 6.

5. The multispecific antibody according to any one of claims 2-4, wherein said VL sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to at least one of SEQ ID NOS:121 and 7.
6. The multispecific antibody according to any one of claims 2-5, wherein said VH sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NO:117; and wherein said VL sequence of the first antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NOS:121.
7. The multispecific antibody according to any one of the preceding claims, wherein the second antigen-binding region binds to cynomolgus CD137.
8. The multispecific antibody according to any one of the preceding claims, wherein said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to a higher degree than it binds to a mutant human CD137 (SEQ ID NO:93).
9. The multispecific antibody according to any one of claims 1-7, wherein said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to a higher degree than it binds to a mutant human CD137 (SEQ ID NO:94).
10. The multispecific antibody according to any one of preceding claims, wherein said second antigen-binding region binds to human CD137 (SEQ ID NO:92) to the same degree that it binds to a mutant human CD137 (SEQ ID NO:95).
11. The multispecific antibody according to any one of claims 8-10, wherein binding to said human CD137 (SEQ ID NO:92) and said mutant human CD137 (SEQ ID NO:93, 94 and 95, respectively) is determined by preparing shuffle constructs derived from human CD137 in which protein domains of the human CD137 are replaced by the corresponding domain of CD137 from different species, using human CD137 and the different species of CD137 as reference constructs; transducing cells with plasmids encoding the reference construct or the shuffle constructs, respectively, and

measuring binding of the antibody to each these CD137 constructs by flow cytometry.

- 5 12. The multispecific antibody according to any one of claims 1-8 and 10-11, wherein said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:
- 10 a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOS:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOS:67, 68 and 68, respectively, and
- 15 b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations.
- c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) or b) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a), or b).
- 20 13. The multispecific antibody according to any one of claims 1-8 and 10-12, wherein said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99%, or 100% identity to at least one of SEQ ID NOS:123 and 69.
- 25 14. The multispecific antibody according to any one of claims 1-8 and 10-13, wherein said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to at least one of SEQ ID NO:127 and 70.
- 30 15. The multispecific antibody according to any one of claims 1-8 and 10-14, wherein said VH and said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 99% or 100% identity to SEQ ID NO:123 and SEQ ID NO:127, respectively.
- 35

16. The multispecific antibody according to any one of claims 1-7 and 9-11, wherein said second antigen-binding region comprises heavy and light chain variable region CDR1, CDR2 and CDR3 selected from the group consisting of:
- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, 40 and 41, respectively, and
 - b) heavy and light chain variable region CDR1, CDR2 and CDR3 according to a) having a total of one to twelve mutations;
 - c) heavy and light chain variable region CDR1, CDR2 and CDR3 of an antibody which (i) competes for human CD137 binding with an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a), or b) and/or (ii) has the specificity for CD137 of an antibody comprising heavy and light chain variable region CDR1, CDR2 and CDR3 according to a), or b).
17. The multispecific antibody according to any one of claims 1-7, 9-11, and 16 wherein said VH sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:41.
18. The multispecific antibody according to any one of claims 1-7, 9-11, and 16-17, wherein said VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:42.
19. The multispecific antibody according to any one of claims 1-7, 9-11, and 16-18, wherein said VH and VL sequence of the second antigen-binding region comprises an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97% or at least 99% identity to SEQ ID NO:41; and SEQ ID NO:42, respectively.
20. The multispecific antibody according to any one of claims 4-19, wherein said first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:1, 2 and 3, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the amino acid sequences set forth in SEQ ID NOs:4, 5 and 6, respectively; and said second antigen-binding region comprises

- a) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:64, 65 and 66, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:67, 68 and 69, respectively, or
- 5 b) heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:36, 37 and 38, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:39, 40 and 41, respectively.
- 10 21. The multispecific antibody according to any one of claims 4-20, wherein the respective FR1, FR2, FR3 and FR4 framework sequences of the VH and VL sequences of the first and/or second antigen-binding region have at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, or at least 99% amino acid sequence identity to the respective FR1, FR2, FR3 and FR4 framework
- 15 sequences of said VH and VL sequences.
22. The multispecific antibody according to any one of the preceding claims, wherein said antibody comprises (I) a first binding arm comprising said first antigen-binding region, and (II) a second binding arm comprising said second antigen-binding region.
- 20 23. The multispecific antibody according to claim 22, wherein said first binding arm comprises a first heavy chain constant sequence, and said second binding arm comprises a second heavy chain constant sequence.
- 25 24. The multispecific antibody according to any one of claims 22-23, wherein (I) said first binding arm comprises a first heavy chain comprising a first heavy chain variable (VH) sequence and a first heavy chain constant (CH) sequence, and a first light chain comprising a first light chain variable (VL) sequence, and (II) said second binding arm comprises a second heavy chain comprising a second heavy chain variable (VH) sequence and a second heavy chain constant (CH) sequence, and a second light chain comprising a second light chain variable (VL) sequence.
- 30 25. The multispecific antibody according to claim 24, wherein said first light chain further comprises a first light chain constant (CL) sequence, and said second light chain further comprises a second light chain constant (CL) sequence.
- 35 26. The multispecific antibody according to any one of the preceding claims, wherein the first antigen-binding region is derived from a mouse antibody.

27. The multispecific antibody according to any one of the preceding claims, wherein the first antigen-binding region is derived from a humanized antibody.
- 5 28. The multispecific antibody according to any one of claims 22-27, wherein the first binding arm is derived from a full-length antibody.
29. The multispecific antibody according to any one of claims 22-28, wherein the first binding arm is derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.
- 10 30. The multispecific antibody according to any one of the preceding claims, wherein the second antigen-binding region is derived from a rabbit antibody.
- 15 31. The multispecific antibody according to any one of the preceding claims, wherein the second antigen-binding region is derived from a humanized antibody.
32. The multispecific antibody according to any one of claims 22-31, wherein the second binding arm is derived from a full-length antibody.
- 20 33. The multispecific antibody according to any one of claims 22-32, wherein the second binding arm is derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.
- 25 34. The multispecific antibody according to any one of the preceding claims, wherein each of the first and second antigen-binding region is derived from a humanized antibody.
- 30 35. The multispecific antibody according to any one of claims 22-34, wherein each of the first and second binding arm is derived from a full-length antibody.
36. The multispecific antibody according to any one of claims 22-35, wherein each of the first and second binding arm is derived from a full-length IgG1, λ (lambda) or IgG1, κ (kappa) antibody.
- 35 37. The multispecific antibody according to any one of claims 22-36, wherein said first and said second heavy chains are of an IgG isotype, having a subclass selected from the group consisting of IgG1, IgG2, IgG3, and IgG4.

38. The multispecific antibody according to any of claims 24-37, wherein each of said first and second heavy chains comprise at least a hinge region, a CH2 and a CH3 region.
- 5
39. The multispecific antibody according to claim 38, wherein the CH3 regions of the first and second heavy chains comprise asymmetrical mutations.
40. The multispecific antibody according to any one of claims 24-39, wherein in said first heavy chain at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370, D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and in said second heavy chain at least one of the amino acids in the positions corresponding to a position selected from the group consisting of T366, L368, K370, D399, F405, Y407, and K409 in a human IgG1 heavy chain according to EU numbering has been substituted, and wherein said first and said second heavy chains are not substituted in the same positions.
- 10
- 15
41. The multispecific antibody according to claim 40, wherein (i) the amino acid in the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering is L in said first heavy chain, and the amino acid in the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering is R in said second heavy chain, or (ii) the amino acid in the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering is R in said first heavy chain, and the amino acid in the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering is L in said second heavy chain.
- 20
- 25
42. The multispecific antibody according to any one of the preceding claims, wherein said antibody comprises a first and a second heavy chain, and wherein said antibody induces Fc-mediated effector function to a lesser extent compared to a multispecific antibody comprising the same first and second antigen-binding regions and two heavy chains comprising human IgG1 hinge, CH2 and CH3 regions.
- 30
43. The multispecific antibody according to claim 42, wherein said first and second heavy chains are modified so that the multispecific antibody induces Fc-mediated effector function to a lesser extent compared to a multispecific antibody which is identical except for comprising non-modified first and second heavy chains.
- 35

44. The multispecific antibody according to any one of claims 42-43, wherein said Fc-mediated effector function is measured by binding to Fcγ receptors, binding to C1q, or induction of Fc-mediated cross-linking of FcRs.
- 5 45. The multispecific antibody according to claim 44, wherein said Fc-mediated effector function is measured by binding to C1q.
- 10 46. The multispecific antibody according to claim 45, and wherein said first and second heavy and light chain constant sequences have been modified so that binding of C1q to said multispecific antibody is reduced compared to a wild-type bispecific antibody by at least 70%, at least 80%, at least 90%, at least 95%, at least 97%, or 100%, wherein C1q binding is determined by ELISA.
- 15 47. The multispecific antibody according to any one of the preceding claims, wherein said antibody comprises a first and a second heavy chain, wherein in at least one of said first and second heavy chains one or more amino acids in the positions corresponding to positions L234, L235, D265, N297, and P331 in a human IgG1 heavy chain according to EU numbering, are not L, L, D, N, and P, respectively.
- 20 48. The multispecific antibody according to claim 47, wherein the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering are F and E, respectively, in said first and second heavy chains.
- 25 49. The multispecific antibody according to claim 48, wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering are F, E, and A, respectively, in said first and second heavy chains.
- 30 50. The multispecific antibody according to claim 49, wherein the positions corresponding to positions L234, L235, and D265 in a human IgG1 heavy chain according to EU numbering of both the first heavy chain and the second heavy chain are F, E, and A, respectively, and wherein (i) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is R, or (ii) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is L.
- 35

51. The multispecific antibody according to claim 48, wherein the positions corresponding to positions L234 and L235 in a human IgG1 heavy chain according to EU numbering of both the first heavy chain and the second heavy chain are F and E, respectively, and wherein (i) the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is L, and the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is R, or (ii) the position corresponding to K409 in a human IgG1 heavy chain according to EU numbering of the first heavy chain is R, and the position corresponding to F405 in a human IgG1 heavy chain according to EU numbering of the second heavy chain is L.
52. The multispecific antibody according to any of the preceding claims, wherein said antibody is able to cross-link a first cell expressing human CD40 and a second cell expressing human CD137.
53. The multispecific antibody according to claim 52, wherein said cross-linking is determined by an assay using a first cell line expressing human CD40 and a second cell line expressing human CD137, and wherein either the first or the second cell line comprises a reporter structure resulting in the production of a measurable reporter upon NF- κ B activation.
54. The multispecific antibody according to any one of the preceding claims, wherein said antibody induces and/or enhances proliferation of T cells.
55. The multispecific antibody according to claim 54, wherein said T cells are CD4⁺ T cells and/or CD8⁺ T cells.
56. The multispecific antibody according to any one of claims 54-55, wherein said induction or enhancement of proliferation of T cells is determined by sub-optimal activating of T cells in a PBMC pool.
57. The multispecific antibody according to claim 56, wherein sub-optimal activation is determined by titrating the concentration of anti-CD3 antibody added to a PBMC pool, measuring T cell proliferation and choosing the anti-CD3 antibody concentration which results in low T cell proliferation but allows for further enhancement of the T cell proliferation.

58. The multispecific antibody according to any one of claims 54-55, wherein proliferation of T cells is measured by co-culturing T-cells expressing a specific T-cell receptor (TCR) with dendritic cells (DCs) presenting the corresponding antigen on the major histocompatibility complex, which is recognized by the TCR.
- 5
59. The multispecific antibody according to any one of claims 54-55, wherein said T cells are tumor-infiltrating lymphocytes (TILs), and said induction or enhancement of proliferation of the TILs is determined by incubating a human tumor sample with interleukin-2 (IL-2) and said antibody, and retrieving and counting TILs after
- 10 incubation for a period of about 10 to about 14 days.
60. The multispecific antibody of claim 59, wherein said human tumor is a melanoma or a non-small-cell lung cancer (NSCLC) tumor.
- 15 61. The multispecific antibody according to any one of claims 54-60, wherein said antibody induces or enhances more proliferation of T cells compared to a bispecific antibody comprising a second antigen-binding region according to any one of claims 1-2, but wherein the first antigen-binding region comprises heavy chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:99, 100
- 20 and 101, respectively, and light chain variable region CDR1, CDR2 and CDR3 having the sequences set forth in SEQ ID NOs:102, GVS and 103, respectively.
62. The antibody according to any one of the preceding claims, wherein said antibody is a bispecific antibody.
- 25
63. A nucleic acid encoding one or more amino acid sequences of claims 1-62.
64. A nucleic acid encoding a multispecific antibody as defined in any one of claims 1-62.
- 30 65. An expression vector comprising a nucleic acid according to any one of claims 63-64.
66. A host cell comprising a nucleic acid according to any one of claims 63 or 64, or an expression vector of claim 65.
- 35 67. The host cell according to claim 66, wherein said host cell is a recombinant eukaryotic, recombinant prokaryotic, or recombinant microbial host cell.

68. A composition comprising a multispecific antibody according to any one of claims 1-62, a nucleic acid according to any one of claims 63 and 64, an expression vector according to claim 65 or a host cell according to any one of claims 66 and 67.
- 5 69. The composition according to claim 68, which is a pharmaceutical composition.
70. The pharmaceutical composition according to claim 69, which further comprises a pharmaceutically acceptable carrier.
- 10 71. The multispecific antibody according to any one of claims 1-62, the nucleic acid according to claim 63 or 64, the expression vector according to claim 65, the host cell according to claim 66 or 67, the composition according to claim 68, or the pharmaceutical composition according to claim 69 or 70 for use as a medicament.
- 15 72. The multispecific antibody according to any one of claims 1-62, the nucleic acid according to claim 63 or 64, the expression vector according to claim 65, the host cell according to claim 66 or 67, the composition according to claim 68, or the pharmaceutical composition according to claim 69 or 70 for use in the treatment of a disease.
- 20 73. The multispecific antibody, nucleic acid, expression vector, host cell or composition for use according to claim 72 wherein the disease is an infectious disease.
74. The multispecific antibody, nucleic acid, expression vector, host cell or composition
25 for use according to claim 72, wherein the disease is cancer.
75. The multispecific antibody, nucleic acid, expression vector, host cell or composition for use according to claim 74, wherein the cancer is selected from the group consisting of melanoma, lung cancer, breast cancer, colon cancer, renal cancer,
30 cervical cancer and prostate cancer, such as non-small cell lung cancer (NSCLC) or melanoma.
76. A method of treatment of a disease comprising administering the multispecific antibody according to any one of claims 1-62, the nucleic acid according to claim 63
35 or 64, the expression vector according to claim 65, the host cell according to claim 66 or 67, the composition according to claim 68, or the pharmaceutical composition according to claim 69 or 70 to a subject in need thereof.

- 5 77. Use of a multispecific antibody according to any one of claims 1-62, the nucleic acid according to claim 63 or 64, the expression vector according to claim 65, the host cell according to claim 66 or 67, the composition according to claim 68, or the pharmaceutical composition according to claim 69 or 70 for the manufacture of a medicament.
- 10 78. The method or use according to any one of claims 71-77, wherein the method or use is for use in combination with one or more further therapeutic agent, such as a chemotherapeutic agent.
- 15 79. A method for producing a bispecific antibody according to claim 62, comprising the steps of
- a) providing a first antibody comprising an Fc region, said Fc region comprising a first CH3 region;
 - 20 b) providing a second antibody comprising a second Fc region, said Fc region comprising a second CH3 region,
wherein the first antibody is a CD40 antibody comprising two first antigen-binding regions according to any of claims 1-62, and the second antibody is a CD137 antibody comprising two second antigen-binding regions according to any of claims 1-62, or vice versa; and wherein the sequences of said first and second CH3 regions are different and are such that the heterodimeric interaction between said first and second CH3 regions is stronger than each of the homodimeric interactions of said first and second CH3 regions;
 - 25 c) incubating said first antibody together with said second antibody under reducing conditions; and
 - d) obtaining said bispecific CD40xCD137 antibody.
- 30 80. A method for detecting whether cross-linking between CD40- and CD137-expressing cells occurs in a sample obtained from a patient, such as a blood sample, lymph node sample or bone marrow sample, upon administration of a multispecific antibody according to any one of claims 1-62, comprising the steps of:
- (i) contacting the sample with a multispecific antibody according to any one of claims 1-62, under conditions that allow for formation of a complex between said multispecific antibody and the CD40- and CD137-expressing cells; and
 - 35 (ii) analyzing whether a complex has been formed.

81. A kit for detecting cross-linking between CD40- and CD137-expressing cells, in a sample obtained from a patient such as a blood sample, lymph node sample or bone marrow sample, comprising

- 5 i) a multispecific antibody according to any one of claims 1-62; and
- ii) instructions for use of said kit.

82. An anti-idiotypic antibody which binds to the second or first and second antigen-binding region as defined in any of claims 1-62.