

Pharmaceutical Combination Comprising Ponesimod

5 The present invention relates to pharmaceutical combinations comprising a first active ingredient which is ponesimod and a second active ingredient which is selected from the group consisting of teriflunomide and leflunomide. The pharmaceutical combinations of the present invention are suitable for the prevention and/or treatment of diseases or disorders in mammals, such as especially humans, that are associated with a lymphocyte-mediated
10 activated immune system.

Description of the Figures:

Fig. 1: Average clinical score in acute monophasic EAE rats treated once daily with ponesimod, teriflunomide or the combination thereof from day 0. Data are presented as
15 mean values + SEM; n = 10-20/group. EAE = experimental autoimmune encephalomyelitis; SEM = standard error of the mean.

Fig. 2: Effect of ponesimod, teriflunomide or the combination thereof on severity of the acute monophasic rat EAE disease, represented as the maximal clinical score reached over the 21 days study. Data are presented as mean values + SEM; n = 10-20/group.

20 ** p<0.01, *** p< 0.001 using Kruskal-wallis test, followed by Dunn's multiple comparisons test.

Fig. 3: Average body weight in acute monophasic EAE rats treated once daily with ponesimod, teriflunomide or the combination thereof from day 0. Data are presented as mean values + SEM; n = 10-20/group.

25 Fig. 4: Effect of ponesimod, teriflunomide or the combination thereof on the average of body weight loss in percentage over the 21 days study in the acute monophasic rat EAE model. Data are presented as mean values + SEM; n = 10-20/group.

** p<0.01 using Kruskal-wallis test, followed by Dunn's multiple comparisons test.

30 Description of the invention:

1) In a first embodiment the present invention relates to a pharmaceutical combination comprising a first active ingredient which is ponesimod and a second active ingredient which is selected from the group consisting of teriflunomide and leflunomide.

Ponesimod (chemical name: (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-
([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one) is a selective S1P₁ receptor agonist and oral
administration thereof results in a consistent, sustained, and dose-dependent reduction in
the number of peripheral blood lymphocytes. Ponesimod has been described to be useful
5 in the treatment and/or prevention of diseases or disorders associated with an activated
immune system (see e.g. WO 2005/054215 and WO 2009/115954). In particular
ponesimod has shown clinical benefit in phase II trials in patients with moderate to severe
chronic plaque psoriasis and in patients with relapsing-remitting multiple sclerosis.
Currently ponesimod is in phase III clinical development for relapsing multiple sclerosis.
10 Ponesimod may be prepared according to any procedure as disclosed in WO
2005/054215, WO 2008/062376 and WO 2014/027330.

Teriflunomide (chemical name: (Z)-2-cyano-3-hydroxy-but-2-enoic acid-(4-
trifluoromethylphenyl)-amide) has been described in the US patents US 4,965,276, US
15 5,459,163 and US 5,679,709 to have immunomodulating properties and to be suitable for
the treatment of chronic graft-versus-host diseases and autoimmune diseases such as
systemic lupus erythematosus. WO 02/080897 discloses that teriflunomide is useful for the
treatment of multiple sclerosis. In particular, teriflunomide (Aubagio®) has been approved
for the treatment of relapsing forms of multiple sclerosis. Teriflunomide can be prepared
20 according to procedures known in the art, for example as described in US 6,894,184. 2-
Cyano-3-hydroxy-but-2-enoic acid-(4-trifluoromethylphenyl)-amide can interconvert
between the Z- and E-stereoisomers with the Z-enol being the most stable and therefore
the most predominant form.

25 Leflunomide (chemical name: 5-methyl-N-(4-(trifluoromethyl)phenyl)isoxazole-4-
carboxamide) was disclosed generically in US patent US 4,087,535 and specifically in US
4,284,786 wherein it was disclosed that the compound could be used for the treatment of
multiple sclerosis. *In vivo* leflunomide is metabolized to the active metabolite teriflunomide
which is responsible for leflunomide's activity *in vivo*. Leflunomide (Arava®) has been
30 approved for the treatment of adults with active rheumatoid arthritis and active psoriatic
arthritis. Leflunomide can be prepared according to procedures known in the art, for
example as described in US 4,284,786.

2) A further embodiment of the invention relates to the pharmaceutical combination
35 according to embodiment 1), wherein the first active ingredient is ponesimod and the
second active ingredient is teriflunomide.

3) A further embodiment of the invention relates to the pharmaceutical combination according to embodiment 1), wherein the first active ingredient is ponesimod and the second active ingredient is leflunomide.

5

4) A further embodiment of the invention relates to the pharmaceutical combination according to any one of embodiments 1) to 3), wherein the first and the second active ingredient are comprised in a single pharmaceutical composition.

10 In a special case of embodiment 4) where e.g. the first active ingredient is administered once daily and the second ingredient is administered twice daily, only one of the two pharmaceutical compositions needed per day will contain both the first and the second active ingredient whereas the other will only contain the second active ingredient. Preferably the first and the second active ingredients are administered once daily.

15

Moreover, in case of a pharmaceutical combination according to embodiment 4) where the first and/or the second active ingredient is initially administered according to a specific dosing regimen (see for example the up-titration regimens disclosed in WO 2009/115954 and WO 2016/091996 for ponesimod) the pharmaceutical compositions needed for the initial dosing
20 regimen will contain the amounts of active ingredient required for the different steps of the dosing regimen.

5) A further embodiment of the invention relates to the pharmaceutical combination according to any one of embodiments 1) to 3), wherein the first and the second active
25 ingredient are comprised in separated pharmaceutical compositions.

In case the first and the second active ingredient are comprised in separated pharmaceutical compositions, they can be administered simultaneously, sequentially or separately; preferably the separated pharmaceutical compositions are administered
30 simultaneously or sequentially, especially sequentially. In case the first active ingredient is for example administered once daily and the second active ingredient twice daily, then the separated pharmaceutical compositions are preferably administered one time per day simultaneously or sequentially, especially sequentially. If administered sequentially or separately, the separated pharmaceutical compositions may be administered in one or the
35 other order. The number of administrations per day may be the same or different for the separated pharmaceutical compositions. For instance, one pharmaceutical composition

may be administered once daily and the other pharmaceutical composition may be administered once or twice daily. Preferably the separated pharmaceutical compositions are administered once daily. Further, the separated pharmaceutical compositions may be administered by the same or different routes of administration, preferably by the same
5 route of administration. Most preferably the separated pharmaceutical compositions are administered orally.

6) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use as a medicament.

10

7) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use in the prevention or treatment, preferably treatment, of a disease or disorder associated with a lymphocyte-mediated activated immune system.

15

8) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use in the prevention or treatment, preferably treatment, of a disease or disorder selected from the group consisting of rejection of transplanted organs such as kidney, liver, heart, lung, pancreas, cornea, and skin; graft-versus-host disease; autoimmune syndromes including rheumatoid arthritis, multiple
20 sclerosis, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, psoriasis, psoriatic arthritis, thyroiditis such as Hashimoto's thyroiditis, and uveo-retinitis; atopic diseases such as rhinitis, conjunctivitis, and dermatitis; asthma; type I diabetes; post-infectious autoimmune diseases including rheumatic fever and post-infectious
25 glomerulonephritis; solid cancers; and tumor metastasis.

9) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use in the prevention or treatment, preferably treatment, of a disease or disorder selected from the group consisting of rejection of
30 transplanted organs selected from kidney, liver, heart and lung; graft-versus-host disease; autoimmune syndromes selected from rheumatoid arthritis, multiple sclerosis, psoriasis, psoriatic arthritis, Crohn's disease, and Hashimoto's thyroiditis; and atopic dermatitis.

10) A preferred embodiment of the invention relates to a pharmaceutical combination
35 according to any one of embodiments 1) to 5) for use in the prevention or treatment,

preferably treatment, of graft-versus-host disease, especially chronic graft-versus-host disease.

11) A most preferred embodiment of the invention relates to a pharmaceutical combination
5 according to any one of embodiments 1) to 5) for use in the prevention or treatment, preferably treatment, of multiple sclerosis.

12) Another very preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use in the prevention or
10 treatment, preferably treatment, of relapsing multiple sclerosis.

13) Another very preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5) for use in the prevention or treatment, preferably treatment, of relapsing-remitting multiple sclerosis.

15 The present invention also relates to a method for the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13) comprising administering to a subject (preferably a human subject) in need thereof a pharmaceutically active amount of a pharmaceutical combination of any one of embodiments 1) to 5).

20 14) A further embodiment of the invention relates to a pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient, for use as a medicament, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing teriflunomide or leflunomide and at
25 least one therapeutically inert excipient.

15) A further embodiment of the invention relates to a pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient, for use as a medicament, wherein the pharmaceutical composition is to be administered in combination
30 with a second pharmaceutical composition containing teriflunomide and at least one therapeutically inert excipient.

16) A further embodiment of the invention relates to a pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient, for use as a
35 medicament, wherein the pharmaceutical composition is to be administered in combination

with a second pharmaceutical composition containing leflunomide and at least one therapeutically inert excipient.

17) A further embodiment of the invention relates to a pharmaceutical composition
5 containing ponesimod and at least one therapeutically inert excipient, for use in the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13), wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing teriflunomide or leflunomide and at least one therapeutically inert excipient.

10

18) A further embodiment of the invention relates to a pharmaceutical composition containing teriflunomide or leflunomide and at least one therapeutically inert excipient, for use as a medicament, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing ponesimod and at least
15 one therapeutically inert excipient.

19) A further embodiment of the invention relates to a pharmaceutical composition containing teriflunomide and at least one therapeutically inert excipient, for use as a medicament, wherein the pharmaceutical composition is to be administered in combination
20 with a second pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient.

20) A further embodiment of the invention relates to a pharmaceutical composition containing leflunomide and at least one therapeutically inert excipient, for use as a
25 medicament, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient.

21) A further embodiment of the invention relates to a pharmaceutical composition
30 containing teriflunomide or leflunomide and at least one therapeutically inert excipient, for use in the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13), wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient.

35

22) A further embodiment of the invention relates to a kit of parts comprising a first pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient; and a second pharmaceutical composition containing teriflunomide or leflunomide and at least one therapeutically inert excipient.

5

23) A further embodiment of the invention relates to the kit of parts according to embodiment 22), wherein the second pharmaceutical composition contains teriflunomide and at least one therapeutically inert excipient.

10

24) A further embodiment of the invention relates to the kit of parts according to embodiment 22), wherein the second pharmaceutical composition contains leflunomide and at least one therapeutically inert excipient.

15

25) A further embodiment of the invention relates to the kit of parts according to any one of embodiments 22) to 24) further comprising instructions for the simultaneous, sequential or separate administration of the pharmaceutical compositions.

20

26) A further embodiment of the invention relates to the kit of parts according to any one of embodiments 22) to 25) for use as a medicament.

27) A further embodiment of the invention relates to the kit of parts according to any one of embodiments 22) to 25) for use in the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13).

25

28) A further embodiment of the invention relates to the use of ponesimod and a second active ingredient which is selected from the group consisting of teriflunomide or leflunomide, for the manufacture of a medicament for use in the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13).

30

29) A further embodiment of the invention relates to the use according to embodiment 28), wherein the second active ingredient is teriflunomide.

30) A further embodiment of the invention relates to the use according to embodiment 28), wherein the second active ingredient is leflunomide.

35

31) A further embodiment of the invention relates to the use of ponesimod for the manufacture of a medicament for use, in combination with a second medicament comprising teriflunomide or leflunomide, in the prevention or treatment, preferably treatment, of a disease or disorder listed in any one of embodiments 7) to 13).

5

32) A further embodiment of the invention relates to the use according to embodiment 31), wherein the second medicament comprises teriflunomide.

33) A further embodiment of the invention relates to the use according to embodiment 31),
10 wherein the second medicament comprises leflunomide.

34) A further embodiment of the invention relates to the use of teriflunomide or leflunomide for the manufacture of a medicament for use, in combination with a second medicament comprising ponesimod, in the prevention or treatment, preferably treatment, of a disease or
15 disorder listed in any one of embodiments 7) to 13).

35) A further embodiment of the invention relates to the use of teriflunomide for the manufacture of a medicament for use, in combination with a second medicament comprising ponesimod, in the prevention or treatment, preferably treatment, of a disease or
20 disorder listed in any one of embodiments 7) to 13).

36) A further embodiment of the invention relates to the use of leflunomide for the manufacture of a medicament for use, in combination with a second medicament comprising ponesimod, in the prevention or treatment, preferably treatment, of a disease or
25 disorder listed in any one of embodiments 7) to 13).

Based on the dependencies of the different embodiments 1) to 36) as disclosed hereinabove, the following dependent embodiments are thus possible and intended and herewith specifically disclosed in individualised form: 2+1, 3+1, 4+1, 4+2+1, 4+3+1, 5+1,
30 5+2+1, 5+3+1, 6+1, 6+2+1, 6+3+1, 6+4+1, 6+4+2+1, 6+4+3+1, 6+5+1, 6+5+2+1, 6+5+3+1, 7+1, 7+2+1, 7+3+1, 7+4+1, 7+4+2+1, 7+4+3+1, 7+5+1, 7+5+2+1, 7+5+3+1, 8+1, 8+2+1, 8+3+1, 8+4+1, 8+4+2+1, 8+4+3+1, 8+5+1, 8+5+2+1, 8+5+3+1, 9+1, 9+2+1, 9+3+1, 9+4+1, 9+4+2+1, 9+4+3+1, 9+5+1, 9+5+2+1, 9+5+3+1, 10+1, 10+2+1, 10+3+1, 10+4+1, 10+4+2+1, 10+4+3+1, 10+5+1, 10+5+2+1, 10+5+3+1, 11+1, 11+2+1, 11+3+1, 11+4+1,
35 11+4+2+1, 11+4+3+1, 11+5+1, 11+5+2+1, 11+5+3+1, 12+1, 12+2+1, 12+3+1, 12+4+1, 12+4+2+1, 12+4+3+1, 12+5+1, 12+5+2+1, 12+5+3+1, 13+1, 13+2+1, 13+3+1, 13+4+1,

13+4+2+1, 13+4+3+1, 13+5+1, 13+5+2+1, 13+5+3+1, 23+22, 24+22, 25+22, 25+23+22, 25+24+22, 26+22, 26+23+22, 26+24+22, 26+25+22, 26+25+23+22, 26+25+24+22, 27+22, 27+23+22, 27+24+22, 27+25+22, 27+25+23+22, 27+25+24+22, 29+28, 30+28, 32+31, and 33+31.

- 5 In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas "+" indicates the dependency from another embodiment. The different individualised embodiments are separated by commas. In other words, "6+4+1" for example refers to embodiment 6) depending on embodiment 4), depending on embodiment 1), i.e. embodiment "6+4+1" corresponds to the pharmaceutical combination
10 of embodiment 1) further limited by the features of the embodiments 4) and 6).

- Definitions provided herein are intended to apply uniformly to the subject matter as defined in any one of embodiments 1) to 36), and, mutatis mutandis, throughout the description and the claims unless an otherwise expressly set out definition provides a broader or narrower
15 definition. It is well understood that a definition or preferred definition of a term or expression defines and may replace the respective term or expression independently of (and in combination with) any definition or preferred definition of any or all other terms or expressions as defined herein.

- 20 Any reference herein to the active ingredient ponesimod, teriflunomide or leflunomide is to be understood as referring to such active ingredient in any form (in the case of teriflunomide also to the *E*-enolic form and mixtures of the *Z*- and *E*-enolic forms, preferably to the *Z*-enolic form) that is suitable to be used for pharmaceutical compositions, including the active ingredient in free or pharmaceutically acceptable salt form, as well as e.g. a
25 hydrate or solvate thereof, or mixtures of any suitable forms. Such forms also encompass all types and polymorphic variations of crystalline forms of the active ingredient (or mixtures of any such crystalline forms) including crystalline forms of the active ingredient in non-ionic or ionic form, optionally together with one or more organic and/or inorganic component in non-ionic or ionic form, such as for example crystalline forms of the mere compound, as
30 well as hydrates, solvates, and salts (including hydrates and solvates of salts) of the compound, and co-crystals (when the active ingredient, or a salt thereof, can be co-crystallized with one or more co-crystal formers optionally together with water and/or solvent) including ionic co-crystals, provided the crystalline forms are suitable to be used for pharmaceutical compositions.

In a preferred embodiment any reference to the active ingredient ponesimod, teriflunomide or leflunomide is meant to refer to such active ingredient in free form, i.e. not in salt form.

5 In another preferred embodiment any reference to ponesimod is meant to refer to ponesimod in crystalline form A or C, most preferably in crystalline form C. The preparation of the crystalline forms A and C of ponesimod is disclosed in WO 2010/046835.

10 The term "pharmaceutically acceptable salt(s)" refers to salts that retain the desired biological activity of the subject compound and exhibit minimal undesired toxicological effects. Such salts include inorganic or organic acid and/or base addition salts depending on the presence of basic and/or acidic groups in the subject compound. For reference see for example 'Handbook of Pharmaceutical Salts. Properties, Selection and Use.', P. Heinrich Stahl, Camille G. Wermuth (Eds.), Wiley-VCH, 2008 and 'Pharmaceutical Salts and Co-crystals', Johan Wouters and Luc Quéré (Eds.), RSC Publishing, 2012.

15

The term "pharmaceutical combination", as used herein, refers to a combination of two or more, preferably two, active ingredients, wherein the active ingredients are comprised in a single pharmaceutical composition or in separated pharmaceutical compositions.

20 The term "active ingredient", as used herein, refers to a pharmaceutically active component of a pharmaceutical composition. Examples of active ingredients, as used herein, are ponesimod, teriflunomide and leflunomide.

25 The term "simultaneous" or "simultaneously", when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) is still ongoing when the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started. Especially, the term "simultaneous" or "simultaneously" means that two active ingredients (or two pharmaceutical compositions, respectively) are administered at the same time, i.e. with the same starting and end time, as is for instance the case for the administration of two active ingredients comprised in a single pharmaceutical composition.

30

35 The term "sequential" or "sequentially", when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started less

than one hour after the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) has been finalized.

5 The term "separate" or "separately", when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started one hour or more (and up to about twelve hours, up to about 24 hours, or up to about 7 days) after the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) has been finalized.

10

The expressions "to be administered in combination" or "for use, in combination" mean simultaneous, sequential or separate, preferably sequential, administration of the pharmaceutical compositions/medicaments.

15 The term "route of administration", as used herein, refers to the path by which an active ingredient (e.g. in form of a pharmaceutical composition in a particular dosage form) enters the body. The active ingredients may be administered by enteral (especially oral) or parenteral administration (including topical application or inhalation). Examples of dosage forms which may be used for the administration of the active ingredients are tablets, capsules, pills, powders, solutions, suspensions, emulsions, injectable aqueous or oily
20 solutions or suspensions, suppositories, creams, gels, ear or eye drops, nasal spray, skin patches, or aerosols. Dosage forms for oral administration, such as tablets, capsules, pills, solutions or suspensions are preferred. In case the two active ingredients are comprised in separated pharmaceutical compositions, said separated pharmaceutical compositions may
25 be administered by the same or different routes of administration using the same or different dosage forms. Most preferably ponesimod, teriflunomide and leflunomide are administered orally, especially in the form of tablets.

For the sake of clarity, relapsing multiple sclerosis means relapsing forms of multiple
30 sclerosis which includes forms of multiple sclerosis with relapses. Examples of relapsing multiple sclerosis are relapsing-remitting multiple sclerosis, secondary progressive multiple sclerosis with relapses, and progressive relapsing multiple sclerosis.

The production of pharmaceutical compositions can be effected in a manner which will be
35 familiar to any person skilled in the art (see for example Remington, The Science and Practice of Pharmacy, 21st Edition (2005), Part 5, "Pharmaceutical Manufacturing"

[published by Lippincott Williams & Wilkins]) by bringing ponesimod and/or a second active ingredient selected from the group consisting of teriflunomide and leflunomide, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants. Particular formulations for teriflunomide are described in e.g. WO 2011/032929.

A preferred maintenance dose of ponesimod, when administered alone, especially for the treatment of multiple sclerosis, is 20 mg orally once daily. Teriflunomide is approved for the treatment of patients with relapsing forms of multiple sclerosis at dosages of 7 mg or 14 mg orally once daily. Leflunomide is approved for the treatment of adults with active rheumatoid arthritis at a recommended maintenance dose of 10 mg to 20 mg orally once daily and for the treatment of adults with active psoriatic arthritis at a recommended maintenance dose of 20 mg orally once daily.

The optimal dosing (i.e., the magnitude of the dose and the dosing frequency) for each of the two active ingredients of the pharmaceutical combinations of the present invention may vary depending upon the route of administration, the dosage form, the disease or disorder to be treated, and the particular second active ingredient applied. In particular, pharmaceutical combinations, like those of the present invention, that exhibit additional beneficial effects, such as synergism, have the advantage that depending on the selected dosing of the active ingredients of the pharmaceutical combination they may achieve better efficacy while keeping an acceptable/comparable or even improved safety profile, or achieve non-inferior efficacy while improving the safety profile compared to when administering the individual active ingredients of the pharmaceutical combination alone.

Further, the dose and/or the dosing frequency may be different during the initial phase and the later phase of the treatment for the first and/or the second active ingredient of the pharmaceutical combination. Preferred dosing regimens for ponesimod have been disclosed in WO 2009/115954 and WO 2016/091996. In psoriatic arthritis for example leflunomide therapy is started with a loading dose of 100 mg once daily for 3 days.

The present invention also includes isotopically labelled, especially ^2H (deuterium) labelled active ingredients, which active ingredients are identical to the active ingredients of the present invention except that one or more atoms have each been replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually found in nature. Isotopically labelled, especially ^2H (deuterium) labelled active ingredients

- and pharmaceutically acceptable salts thereof are within the scope of the present invention. Substitution of hydrogen with the heavier isotope ^2H (deuterium) may lead to greater metabolic stability, resulting e.g. in increased in-vivo half-life and/or reduced dosage requirements, and/or may reduce the formation of undesirable metabolites resulting e.g. in an improved safety profile. In one embodiment only one of the two active ingredients of the pharmaceutical combination is isotopically labelled. In a preferred embodiment of the invention, the active ingredients are not isotopically labelled, or one active ingredient is not isotopically labelled and the other active ingredient is labelled only with one or more deuterium atoms, or both active ingredients are each labelled only with one or more deuterium atoms. In a most preferred embodiment, the active ingredients are not isotopically labelled at all. Isotopically labelled active ingredients may be prepared in analogy to the methods described for the not isotopically labelled active ingredients, but using the appropriate isotopic variation of suitable reagents or starting materials.
- 15 The term "about" placed before a numerical value "X" refers in the current application to an interval extending from X minus 10% of X to X plus 10% of X.

Biological assay:

- 20 The efficacy of teriflunomide and ponesimod in acute monophasic models of experimental autoimmune encephalomyelitis (EAE) can be determined in a pilot experiment.

Female Lewis rats are immunized with an emulsion of myelin basic protein (MBP) from Guinea pig in complete Freund's adjuvant. A total of 200 micrograms of MBP is injected per rat at two sites subcutaneously (right paw and base of tail). Within 10 days the rats develop signs of paralysis which will be graded on a scale from 0 to 15 assessing the tail and each limb individually according to the method originally described by Weaver A. et al., *FASEB J.*, 2005, 19(12): 1668-1670. The disease takes an acute monophasic course and is self-remitting. Usually by day 21 clinical scores reach values below 3.0.

- 30 Groups of 10 rats are dosed orally starting at the day of disease induction (day 0) with different doses of teriflunomide in the range between 1 and 10 mg/kg once daily (q.d.) or with 30 mg/kg ponesimod q.d.. Clinical scores are assessed on a daily basis and disease development is compared between vehicle-treated rats and rats receiving teriflunomide or ponesimod (vehicle: 0.5% Methylcellulose/0.5% Tween® 80/water). In parallel, body weights of the rats are monitored on a daily basis to follow general health. Hematology
- 35

parameters, including lymphocyte counts, are measured at 3 different time points: 6 and 24 hours after the oral gavage at day 20 and at the termination of the experiment at day 21, 3 hours post dosing. Plasma samples can be taken for compound concentration determination and for the measurement of protein markers.

5

Based on the first pilot experiment described above one dose of teriflunomide is selected for the combination efficacy experiment. The dose selected exerts partial efficacy on EAE scores: a strong delay of disease onset without affecting the severity of the disease (dose: 1 mg/kg). Ponesimod at the dose of 30 mg/kg showed partial reduction of lymphocytes over 24 hours.

10

The combination efficacy experiment is performed in the same rat EAE model as described for the first pilot experiment. It consists of four treatment groups:

1. Vehicle (0.5% Methylcellulose/0.5% Tween® 80/water) q.d., from day 0
- 15 2. Teriflunomide (1 mg/kg) q.d., from day 0
3. Ponesimod (30 mg/kg) q.d., from day 0
4. Teriflunomide (1 mg/kg) q.d., from day 0 + ponesimod (30 mg/kg) q.d., from day 0

In life and termination endpoints are the same as described for the first pilot experiment.

- 20 The goal of this experiment is to show that the combination of a dose of ponesimod, that is partially efficacious on lymphocyte count, shows synergism or added benefit to a partially efficacious dose of teriflunomide in a model of EAE.

- 25 The results from the combination efficacy experiment are shown in Fig. 1-4. Data collected in the pilot experiment are similar to those obtained in the combination study and therefore are used to increase the monotherapy groups size in order to increase sensitivity as described by Bate S. et al., *PLoS One.*, 2014, 9(12): e114872. As can be seen from Fig. 1, both compounds alone, ponesimod and teriflunomide exhibit moderate efficacy on the clinical course of EAE in the rat monophasic EAE model. Ponesimod (30 mg/kg) showed a moderate effect during the ascending phase (Fig. 1) and a reduction in body weight loss (Fig. 3 and Fig. 4). Teriflunomide alone exerts partial efficacy on EAE scores: a strong delay of disease onset without affecting the severity of the disease (Fig. 1 and Fig. 2). When combined, the two compounds benefit from each other. As shown in Fig. 2, when combined the two compound showed synergistic efficacy on the severity of the disease, 35 where ponesimod or teriflunomide alone had no effect.

Claims:

1. Use of a pharmaceutical combination comprising a first active ingredient which is ponesimod and a second active ingredient which is selected from the group consisting of teriflunomide and leflunomide in the manufacture of a medicament for the treatment of multiple sclerosis.
5
2. Use of a pharmaceutical combination according to claim 1, wherein the first active ingredient is ponesimod and the second active ingredient is teriflunomide.
10
3. Use of a pharmaceutical combination according to claim 1, wherein the first active ingredient is ponesimod and the second active ingredient is leflunomide.
4. Use of a pharmaceutical combination according to any one of claims 1 to 3, wherein the first and the second active ingredient are comprised in a single pharmaceutical composition.
15
5. Use of a pharmaceutical combination according to any one of claims 1 to 3, wherein the first and the second active ingredient are comprised in separated pharmaceutical compositions.
20
6. Use of a pharmaceutical composition containing ponesimod and at least one therapeutically inert excipient in the manufacture of a medicament, for the prevention or treatment of multiple sclerosis, wherein said medicament is to be administered in combination with a second medicament containing teriflunomide or leflunomide and at least one therapeutically inert excipient.
25
7. Use of a pharmaceutical composition containing teriflunomide or leflunomide and at least one therapeutically inert excipient, in the manufacture of a medicament for the prevention or treatment of multiple sclerosis, wherein said medicament is to be administered in combination with a second medicament containing ponesimod and at least one therapeutically inert excipient.
30