

MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING MYOTONIC DYSTROPHY

RELATED APPLICATIONS

[0001] This application claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Application Serial No. 63/143827, entitled “MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING MYOTONIC DYSTROPHY”, filed on January 30, 2021, to U.S. Provisional Application Serial No. 63/069075, entitled “MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING MYOTONIC DYSTROPHY”, filed on August 23, 2020, and to U.S. Provisional Application Serial No. 63/055749, entitled “MUSCLE TARGETING COMPLEXES AND USES THEREOF FOR TREATING MYOTONIC DYSTROPHY”, filed on July 23, 2020; the contents of each of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] The present application relates to targeting complexes for delivering molecular payloads (e.g., oligonucleotides) to cells and uses thereof, particularly uses relating to treatment of disease.

REFERENCE TO SEQUENCE LISTING SUBMITTED AS A TEXT FILE VIA EFS-WEB

[0003] The instant application contains a sequence listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on July 8, 2021, is named D082470038WO00-SEQ-DWY and is 268,942 bytes in size.

BACKGROUND OF INVENTION

[0004] Myotonic dystrophy (DM) is a dominantly inherited genetic disease that is characterized by myotonia, muscle loss or degeneration, diminished muscle function, insulin resistance, cardiac arrhythmia, smooth muscle dysfunction, and neurological abnormalities. DM is the most common form of adult-onset muscular dystrophy, with a worldwide incidence of about 1 in 8000 people worldwide. Two types of the disease, myotonic dystrophy type 1 (DM1) and myotonic dystrophy type 2 (DM2), have been described. DM1, the more common form of the disease, results from a repeat expansion of a CTG trinucleotide repeat in the 3' non-coding region of DMPK on chromosome 19; DM2 results from a repeat expansion of a CCTG tetranucleotide repeat in the first intron of ZNF9 on chromosome 3. In DM1 patients, the repeat

expansion of a CTG trinucleotide repeat, which may comprise greater than ~50 to ~3,000+ total repeats, leads to generation of toxic RNA repeats capable of forming hairpin structures that bind essential intracellular proteins, e.g. muscleblind-like proteins, with high affinity resulting in protein sequestration and the loss-of-function phenotypes that are characteristic of the disease. Apart from supportive care and treatments to address the symptoms of the disease, no effective therapeutic for DM1 is currently available.

SUMMARY OF INVENTION

[0005] In some aspects, the disclosure provides complexes that target muscle cells for purposes of delivering molecular payloads to those cells. In some embodiments, complexes provided herein are particularly useful for delivering molecular payloads that inhibit the expression or activity of a DMPK allele comprising an expanded disease-associated-repeat, e.g., in a subject having or suspected of having myotonic dystrophy. Accordingly, in some embodiments, complexes provided herein comprise muscle-targeting agents (e.g., muscle targeting antibodies) that specifically bind to receptors on the surface of muscle cells for purposes of delivering molecular payloads to the muscle cells. In some embodiments, the complexes are taken up into the cells via a receptor mediated internalization, following which the molecular payload may be released to perform a function inside the cells. For example, complexes engineered to deliver oligonucleotides may release the oligonucleotides such that the oligonucleotides can inhibit mutant DMPK expression in the muscle cells. In some embodiments, the oligonucleotides are released by endosomal cleavage of covalent linkers connecting oligonucleotides and muscle-targeting agents of the complexes.

[0006] One aspect of the present disclosure relates to a complex comprising an anti-transferrin receptor (TfR) antibody covalently linked to a molecular payload configured for reducing expression or activity of DMPK, wherein the anti-TfR antibody comprises:

(i) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 75;

(ii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 69; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;

(iii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 71; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;

(iv) a heavy chain variable region (VH) comprising an amino acid sequence at

least 95% identical to SEQ ID NO: 72; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;

(v) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 74;

(vi) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 75;

(vii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 74;

(viii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 78;

(ix) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 79; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 80; or

(x) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 80.

[0007] In some embodiments, the antibody comprises:

(i) a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of SEQ ID NO: 75;

(ii) a VH comprising an amino acid sequence of SEQ ID NO: 69 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(iii) a VH comprising an amino acid sequence of SEQ ID NO: 71 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(iv) a VH comprising an amino acid sequence of SEQ ID NO: 72 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(v) a VH comprising an amino acid sequence of SEQ ID NO: 73 and a VL comprising an amino acid sequence of SEQ ID NO: 74;

(vi) a VH comprising an amino acid sequence of SEQ ID NO: 73 and a VL comprising an amino acid sequence of SEQ ID NO: 75;

(vii) a VH comprising an amino acid sequence of SEQ ID NO: 76 and a VL comprising an amino acid sequence of SEQ ID NO: 74;

(viii) a VH comprising an amino acid sequence of SEQ ID NO: 77 and a VL comprising an amino acid sequence of SEQ ID NO: 78;

(ix) a VH comprising an amino acid sequence of SEQ ID NO: 79 and a VL comprising an amino acid sequence of SEQ ID NO: 80; or

(x) a VH comprising an amino acid sequence of SEQ ID NO: 77 and a VL comprising an amino acid sequence of SEQ ID NO: 80.

[0008] In some embodiments, the antibody is selected from the group consisting of a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, an scFv, an Fv, and a full-length IgG. In some embodiments, the antibody is a Fab fragment.

[0009] In some embodiments, the antibody comprises:

(i) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(ii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 97; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 98; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iv) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 99; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(v) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(vi) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(vii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(viii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 93;

(ix) a heavy chain comprising an amino acid sequence at least 85% identical to

SEQ ID NO: 103; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95; or

(x) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95.

[00010] In some embodiments, the antibody comprises:

(i) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(ii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 97 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 98 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iv) a heavy chain comprising the amino acid sequence of SEQ ID NO: 99 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(v) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(vi) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(vii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(viii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 93;

(ix) a heavy chain comprising the amino acid sequence of SEQ ID NO: 103 and a light chain comprising the amino acid sequence of SEQ ID NO: 95; or

(x) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

[00011] In some embodiments, the antibody does not specifically bind to the transferrin binding site of the transferrin receptor and/or the antibody does not inhibit binding of transferrin to the transferrin receptor. In some embodiments, the antibody is cross-reactive with extracellular epitopes of two or more of a human, non-human primate and rodent transferrin receptor. In some embodiments, the complex is configured to promote transferrin receptor mediated internalization of the molecular payload into a muscle cell.

[00012] In some embodiments, the molecular payload is an oligonucleotide. In some embodiments, the oligonucleotide comprises at least 15 consecutive nucleotides of SEQ ID NOs: 148-383 and 621-638, wherein any one or more of the thymidine bases (T's) in the

oligonucleotide may optionally be a uridine base (U) and/or any one or more of the U's may optionally be a T. In some embodiments, the oligonucleotide comprises a sequence comprising any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264, wherein any one or more of the U's in the oligonucleotide may optionally be a T. In some embodiments, the oligonucleotide comprises a region of complementarity to at least 15 consecutive nucleotides of any one of SEQ ID NO: 384-619.

[00013] In some embodiments, the oligonucleotide mediates RNase H-mediated cleavage of a DMPK mRNA transcript.

[00014] In some embodiments, the oligonucleotide comprises a 5'-X-Y-Z-3' formula, wherein X and Z are flanking regions comprising one or more 2'-modified nucleosides selected from the group consisting of: 2'-O-methyl, 2'-fluoro, 2'-O-methoxyethyl, and 2',4'- bridged nucleosides, and wherein Y is a gap region and each nucleoside in Y is a 2'-deoxyribonucleoside.

[00015] In some embodiments, the oligonucleotide comprises one or more phosphorothioate internucleoside linkages.

[00016] In some embodiments, the antibody is covalently linked to the molecular payload via a cleavable linker. In some embodiments, the cleavable linker comprises a valine-citrulline sequence.

[00017] In some embodiments, the antibody is covalently linked to the molecular payload via conjugation to a lysine residue or a cysteine residue of the antibody.

[00018] In some embodiments, reducing expression comprises reducing RNA levels of DMPK, optionally wherein the reduced RNA levels are in the nucleus of a cell, optionally wherein the cell is a muscle cell. In some embodiments, the DMPK is encoded from an allele comprising a disease-associated repeat.

[00019] Another aspect of the present disclosure relates to a method of reducing DMPK expression in a cell, the method comprising contacting the cell with a complex disclosed herein in an effective amount for promoting internalization of the molecular payload in the cell, optionally wherein the cell is a muscle cell.

[00020] Another aspect of the present disclosure relates to a method of treating a subject having an expansion of a disease-associated-repeat of a DMPK allele that is associated with myotonic dystrophy, the method comprising administering to the subject an effective amount of a complex disclosed herein. In some embodiments, the disease-associated-repeat comprises repeating units of a CTG trinucleotide sequence. In some embodiments, the complex is intravenously administered to the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

[00021] **FIG. 1** depicts a non-limiting schematic showing the effect of transfecting Hepa 1-6 cells with an antisense oligonucleotide that targets DMPK (ASO300) on expression levels of DMPK relative to a vehicle transfection.

[00022] **FIG. 2A** depicts a non-limiting schematic showing an HIL-HPLC trace obtained during purification of a muscle targeting complex comprising an anti-transferrin receptor antibody covalently linked to a DMPK antisense oligonucleotide.

[00023] **FIG. 2B** depicts a non-limiting image of an SDS-PAGE analysis of a muscle targeting complex.

[00024] **FIG. 3** depicts a non-limiting schematic showing the ability of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to reduce expression levels of DMPK.

[00025] **FIGs. 4A-4E** depict non-limiting schematics showing the ability of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to reduce expression levels of DMPK in mouse muscle tissues *in vivo*, relative to a vehicle treatment, treatment with naked ASO300, or treatment with a control non-targeting complex (DTX-C-007). (N=3 C57Bl/6 WT mice)

[00026] **FIGs. 5A-5B** depict non-limiting schematics showing the tissue selectivity of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300. The muscle targeting complex (DTX-C-008) comprising ASO300 does not reduce expression levels of DMPK in mouse brain or spleen tissues *in vivo*, relative to a vehicle treatment, treatment with naked ASO300, or treatment with a control non-targeting complex (DTX-C-007). (N=3 C57Bl/6 WT mice).

[00027] **FIGs. 6A-6F** depict non-limiting schematics showing the ability of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to reduce expression levels of DMPK in mouse muscle tissues *in vivo*, relative to a vehicle treatment, treatment with naked ASO300, or treatment with a control non-targeting complex (DTX-C-007). (N=5 C57Bl/6 WT mice)

[00028] **FIGs. 7A-7L** depict non-limiting schematics showing the ability of a muscle targeting antibody-oligonucleotide complex (DTX-C-012) comprising ASO300 covalently linked to an anti-hTfR antibody to reduce expression levels of DMPK in cynomolgus monkey muscle tissues *in vivo*, relative to a vehicle treatment (saline) and compared to naked ASO300. (N=3 male cynomolgus monkeys)

[00029] **FIGs. 8A-8B** depict non-limiting schematics showing the ability of a muscle targeting antibody-oligonucleotide complex (DTX-C-012) comprising ASO300 covalently linked to an anti-hTfR antibody to reduce expression levels of DMPK in cynomolgus monkey smooth muscle tissues *in vivo*, relative to a vehicle treatment (saline) and compared to naked ASO300. (N=3 male cynomolgus monkeys)

[00030] **FIGs. 9A-9D** depict non-limiting schematics showing the tissue selectivity of a muscle targeting antibody-oligonucleotide complex (DTX-C-012) comprising ASO300 covalently linked to an anti-hTfR antibody. The muscle targeting complex comprising DMPK-ASO does not reduce expression levels of DMPK in cynomolgus monkey kidney, brain, or spleen tissues *in vivo*, relative to a vehicle treatment. (N=3 male cynomolgus monkeys)

[00031] **FIG. 10** shows normalized DMPK mRNA tissue expression levels across several tissue types in cynomolgus monkeys. (N=3 male cynomolgus monkeys)

[00032] **FIGs. 11A-11B** depict non-limiting schematics showing the ability of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to reduce expression levels of DMPK in mouse muscle tissues *in vivo* for up to 28 days after dosing with DTX-C-008, relative to a vehicle treatment (saline) and compared to naked ASO300.

[00033] **FIG. 12** shows that a single dose of a muscle targeting complex (DTX-C-012) comprising ASO300 covalently linked to an anti-hTfR antibody is safe and tolerated in cynomolgus monkeys. (N=3 male cynomolgus monkeys)

[00034] **FIGs. 13A-13B** depict non-limiting schematics showing the ability of a muscle targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to reduce expression levels of DMPK in mouse muscle tissues *in vivo* for up to twelve weeks after dosing with DTX-C-008, relative to a vehicle treatment (PBS); and compared to a control IgG2a Fab antibody-oligonucleotide complex (DTX-C-007) and naked DMPK ASO (ASO300). (N=5 C57Bl/6 WT mice)

[00035] **FIGs. 14A-14B** depict non-limiting schematics showing the ability of a muscle-targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 to target nuclear mutant DMPK RNA in a mouse model. (N=6 mice)

[00036] **FIGs. 15A-15B** depict non-limiting schematics showing the ability of a muscle-targeting RI7 217 Fab antibody-ASO complex (DTX-Actin) comprising an oligonucleotide that targets actin to dose-dependently reduce expression levels of actin and functional grades of myotonia in muscle tissues. (N=2 HSA^{LR} mice)

[00037] **FIGs. 16A-16C** depict non-limiting schematics showing that a muscle-targeting RI7 217 Fab antibody-oligonucleotide complex (DTX-C-008) comprising ASO300 is capable of

significantly reducing the prolonged QTc interval in a mouse model for validation of the functional correction of arrhythmia in a DM1 cardiac model. (N=10 mice)

[00038] **FIGs. 17A-17B** depict non-limiting schematics showing that a muscle-targeting antibody-oligonucleotide complex (DTX-C-012) comprising ASO300 antisense oligonucleotide covalently linked to an anti-hTfR antibody is capable of reducing expression levels of DMPK and correcting splicing of a DMPK-specific target gene (Bin1) in human cells from a DM1 patient. (N=3)

[00039] **FIGs. 18A-18C** depict non-limiting schematics showing the dose response of selected antisense oligonucleotides in DMPK knockdown in human DM1 myotubes. ASO300 was used as control. All tested oligonucleotides showed activity in DMPK knockdown. Statistical analysis: One-way ANOVA with Tukey's HSD post-hoc test vs. naked ASO300 treatment; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

[00040] **FIGs. 19A-19B** depict non-limiting schematics showing the dose response of selected antisense oligonucleotides in DMPK knockdown in non-human primate (NHP) DM1 myotubes. ASO300 was used as control. All tested oligonucleotides showed activity in DMPK knockdown.

[00041] **FIG. 20** shows the serum stability of the linker used for linking an anti-TfR antibody and a molecular payload (e.g., an oligonucleotide) in various species over time after intravenous administration.

[00042] **FIGs. 21A-21F** show binding of humanized anti-TfR Fabs to human TfR1 (hTfR1) or cynomolgus monkey TfR1 (cTfR1), as measured by ELISA. FIG. 21A shows binding of humanized 3M12 variants to hTfR1. FIG. 21B shows binding of humanized 3M12 variants to cTfR1. FIG. 21C shows binding of humanized 3A4 variants to hTfR1. FIG. 21D shows binding of humanized 3A4 variants to cTfR1. FIG. 21E shows binding of humanized 5H12 variants to hTfR1. FIG. 21F shows binding of humanized 5H12 variants to hTfR1.

[00043] **FIGs. 22** shows the quantified cellular uptake of anti-TfR Fab conjugates into rhabdomyosarcoma (RD) cells. The molecular payload in the tested conjugates are DMPK-targeting oligonucleotides and the uptake of the conjugates were facilitated by indicated anti-TfR Fabs. Conjugates having a negative control Fab (anti-mouse TfR) or a positive control Fab (anti-human TfR1) are also included this assay. Cells were incubated with indicated conjugate at a concentration of 100 nM for 4 hours. Cellular uptake was measured by mean Cypher5e fluorescence.

[00044] **FIGs. 23A-23F** show binding of oligonucleotide-conjugated or unconjugated humanized anti-TfR Fabs to human TfR1 (hTfR1) and cynomolgus monkey TfR1 (cTfR1), as measured by ELISA. FIG. 23A shows the binding of humanized 3M12 variants alone or in

conjugates with a DMPK targeting oligo to hTfR1. FIG. 23B shows the binding of humanized 3M12 variants alone or in conjugates with a DMPK targeting oligo to cTfR1. FIG. 23C shows the binding of humanized 3A4 variants alone or in conjugates with a DMPK targeting oligo to hTfR1. FIG. 23D shows the binding of humanized 3A4 variants alone or in conjugates with a DMPK targeting oligo to cTfR1. FIG. 23E shows the binding of humanized 5H12 variants alone or in conjugates with a DMPK targeting oligo to hTfR1. FIG. 23F shows the binding of humanized 5H12 variants alone or in conjugates with a DMPK targeting oligo to cTfR1. The respective EC₅₀ values are also shown.

[00045] FIG. 24 shows DMPK expression in RD cells treated with DMPK-targeting oligonucleotides relative to cells treated with PBS. The duration treatment was 3 days. DMPK-targeting oligonucleotides were delivered to the cells as free oligonucleotides (gymnotic uptake, “free”) or with transfection reagent (“trans”).

[00046] FIG. 25 shows DMPK expression in RD cells treated with various concentrations of conjugates containing the indicated humanized anti-TfR antibodies conjugated to a DMPK-targeting antisense oligonucleotide (ASO300). The duration of treatment was 3 days. ASO300 delivered using transfection agents (labeled “Trans”) was used as control.

[00047] FIG. 26 shows results of splicing correction in Atp2a1 by an anti-TfR1 antibody-oligonucleotide conjugate (Ab-ASO) in the HSA-LR mouse model of DM1, measured in the gastrocnemius muscle. The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide is targeting skeletal actin.

[00048] FIG. 27 shows splicing correction in more than 30 different RNAs related to DM1, measured in the gastrocnemius muscle of HSA-LR mice treated anti-TfR1 antibody-oligonucleotide (Ab-ASO) conjugate or saline. The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide is targeting human skeletal actin.

[00049] FIG. 28 shows splicing derangement in quadriceps, gastrocnemius, or tibialis anterior muscles of HSA-LR mice treated with anti-TfR1 antibody-oligonucleotide conjugate (Ab-ASO) or saline. The data represent composite splicing derangement measured in the more than 30 RNAs shown in FIG. 27.

[00050] FIG. 29 shows myotonia grade measured in quadriceps, gastrocnemius, and tibialis anterior muscles of HSA-LR mice treated with saline, unconjugated oligonucleotide (ASO), or anti-TfR1 antibody-oligonucleotide conjugate (Ab-ASO). Myotonia was measured by electromyography (EMG), and graded 0, 1, 2, or 3 based on the frequency of myotonic discharge.

[00051] FIGs. 30A-30E show *in vivo* activity of conjugates containing designated anti-TfR Fabs (control, 3M12 VH3/VK2, 3M12 VH4/VK3, and 3A4 VH3 N54S/VK4) conjugated to

DMPK-targeting oligonucleotide in reducing DMPK mRNA expression in mice expressing human TfR1 (hTfR1 knock-in mice). FIG. 30A shows the experimental design (e.g., IV dosage, dosing frequency). DMPK mRNA levels were measured 14 days post first dose in the tibialis anterior (FIG. 30B), gastrocnemius (FIG. 30C), heart (FIG. 30D), and diaphragm (FIG. 30E), of the mice.

[00052] FIGs. 31A-31C show that conjugates containing anti-TfR antibody conjugated to DMPK-targeting oligonucleotide corrected splicing and reduced foci in CM-DM1-32F primary cells expressing a DMPK mutant mRNA containing 380 CUG repeats. FIG. 31A shows that the conjugates reduced mutant DMPK mRNA expression. FIG. 31B shows that the conjugates corrected BIN1 Exon 11 splicing. FIG. 31C shows images of a fluorescence in situ hybridization (FISH) analysis and quantification of the images, demonstrating that the conjugated reduced nuclear foci formed by the mutant DMPK mRNA. In the microscopy images shown in the top panels of FIG. 31C, the light rounded shapes show cell nuclei, and the bright puncta within the nuclei of the DM1 cells (right three microscopy panels) show CUG foci.

[00053] FIG. 32 shows ELISA measurements of binding of anti-TfR Fab 3M12 VH4/Vk3 to recombinant human (circles), cynomolgus monkey (squares), mouse (upward triangles), or rat (downward triangles) TfR1 protein, at a range of concentrations from 230 pM to 500 nM of the Fab. Measurement results show that the anti-TfR Fab is reactive with human and cynomolgus monkey TfR1. Binding was not observed to mouse or rat recombinant TfR1. Data is shown as relative fluorescent units normalized to baseline.

[00054] FIG. 33 shows results of an ELISA testing the affinity of anti-TfR Fab 3M12 VH4/Vk3 to recombinant human TfR1 or TfR2 over a range of concentrations from 230 pM to 500 nM of Fab. The data are presented as relative fluorescence units normalized to baseline. The results demonstrate that the Fab does not bind recombinant human TfR2.

[00055] FIG. 34 shows the serum stability of the linker used for linking anti-TfR Fab 3M12 VH4/Vk3 to a control antisense oligonucleotide over 72 hours incubation in PBS or in rat, mouse, cynomolgus monkey or human serum.

[00056] FIGs. 35A-35B show splicing correction in more than 30 different RNAs known to be mis-spliced in DM1 patients, measured in the tibialis anterior (FIG. 35A) or the quadriceps (FIG. 35B) of HSA-LR mice treated with a single dose of anti-TfR antibody-oligonucleotide (Ab-ASO) conjugate or saline. The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide targets skeletal actin (ACTA1).

[00057] FIGs. 36A-36C show EMG myotonia grade in quadriceps (FIG. 36A), gastrocnemius (FIG. 36B), and tibialis anterior (FIG. 36C) of HSA-LR mice treated with vehicle, a single dose of unconjugated ASO, or a single dose of anti-TfR antibody-ASO

conjugate (Ab-ASO). The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide targets human skeletal actin (ACTA1).

[00058] FIG. 37 shows human ACTA1 expression measured by qPCR in HSA^{LR} DM1 mice after a single dose of naked ASO or dose equivalent of anti-TfR antibody-ASO conjugate (Ab-ASO), relative to vehicle-treated mice. The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide targets human skeletal actin (ACTA1).

[00059] FIGs. 38A-38C show ACTA1 expression in quadriceps (FIG. 38A), gastrocnemius (FIG. 38B), and tibialis anterior (FIG. 38C) in HSA^{LR} DM1 mice after a single dose of 10 mg/kg naked ASO, 20 mg/kg naked ASO, or dose equivalents of anti-TfR antibody-ASO conjugate (Ab-ASO), relative to vehicle-treated mice. The anti-TfR antibody used is RI7 217 Fab and the oligonucleotide targets human skeletal actin (ACTA1). (* $p < 0.05$; *** $p < 0.001$)

DETAILED DESCRIPTION OF INVENTION

[00060] Aspects of the disclosure relate to a recognition that while certain molecular payloads (*e.g.*, oligonucleotides, peptides, small molecules) can have beneficial effects in muscle cells, it has proven challenging to effectively target such cells. As described herein, the present disclosure provides complexes comprising muscle-targeting agents covalently linked to molecular payloads in order to overcome such challenges. In some embodiments, the complexes are particularly useful for delivering molecular payloads that inhibit the expression or activity of target genes in muscle cells, *e.g.*, in a subject having or suspected of having a rare muscle disease. For example, in some embodiments, complexes are provided for targeting a DMPK allele that comprises an expanded disease-associated-repeat to treat subjects having DM1. In some embodiments, complexes provided herein may comprise oligonucleotides that inhibit expression of a DMPK allele comprising an expanded disease-associated-repeat. As another example, complexes may comprise oligonucleotides that interfere with the binding of a disease-associated DMPK mRNA to a muscleblind-like protein (*e.g.*, MBNL1, 2, and/or (*e.g.*, and) 3), thereby reducing a toxic effect of a disease-associated DMPK allele. In some embodiments, synthetic nucleic acid payloads (*e.g.*, DNA or RNA payloads) may be used that express one or more proteins that reduce a toxic effect of a disease-associated DMPK allele. In some embodiments, complexes may comprise molecular payloads of synthetic cDNAs and/or (*e.g.*, and) synthetic mRNAs, *e.g.*, that express one or more muscleblind-like-proteins (*e.g.*, MBNL1, 2, and/or (*e.g.*, and) 3) or fragments thereof. In some embodiments, complexes may comprise molecular payloads such as guide molecules (*e.g.*, guide RNAs) that are capable of targeting nucleic acid programmable nucleases (*e.g.*, Cas9) to a sequence at or near a disease-associated

repeat sequence of DMPK. In some embodiments, such nucleic programmable nucleases could be used to cleave part or all of a disease-associated repeat sequence from a DMPK gene.

[00061] Further aspects of the disclosure, including a description of defined terms, are provided below.

I. Definitions

[00062] **Administering:** As used herein, the terms “administering” or “administration” means to provide a complex to a subject in a manner that is physiologically and/or (e.g., and) pharmacologically useful (e.g., to treat a condition in the subject).

[00063] **Approximately:** As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[00064] **Antibody:** As used herein, the term “antibody” refers to a polypeptide that includes at least one immunoglobulin variable domain or at least one antigenic determinant, e.g., paratope that specifically binds to an antigen. In some embodiments, an antibody is a full-length antibody. In some embodiments, an antibody is a chimeric antibody. In some embodiments, an antibody is a humanized antibody. However, in some embodiments, an antibody is a Fab fragment, a Fab', a F(ab')₂ fragment, a Fv fragment or a scFv fragment. In some embodiments, an antibody is a nanobody derived from a camelid antibody or a nanobody derived from shark antibody. In some embodiments, an antibody is a diabody. In some embodiments, an antibody comprises a framework having a human germline sequence. In another embodiment, an antibody comprises a heavy chain constant domain selected from the group consisting of IgG, IgG1, IgG2, IgG2A, IgG2B, IgG2C, IgG3, IgG4, IgA1, IgA2, IgD, IgM, and IgE constant domains. In some embodiments, an antibody comprises a heavy (H) chain variable region (abbreviated herein as VH), and/or (e.g., and) a light (L) chain variable region (abbreviated herein as VL). In some embodiments, an antibody comprises a constant domain, e.g., an Fc region. An immunoglobulin constant domain refers to a heavy or light chain constant domain. Human IgG heavy chain and light chain constant domain amino acid sequences and their functional variations are known. With respect to the heavy chain, in some embodiments, the heavy chain of an antibody described herein can be an alpha (α), delta (Δ), epsilon (ϵ), gamma (γ) or mu (μ) heavy chain. In some embodiments, the heavy chain of an antibody described

herein can comprise a human alpha (α), delta (Δ), epsilon (ϵ), gamma (γ) or mu (μ) heavy chain. In a particular embodiment, an antibody described herein comprises a human gamma 1 CH1, CH2, and/or (e.g., and) CH3 domain. In some embodiments, the amino acid sequence of the VH domain comprises the amino acid sequence of a human gamma (γ) heavy chain constant region, such as any known in the art. Non-limiting examples of human constant region sequences have been described in the art, e.g., see U.S. Pat. No. 5,693,780 and Kabat E A et al., (1991) *supra*. In some embodiments, the VH domain comprises an amino acid sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or at least 99% identical to any of the variable chain constant regions provided herein. In some embodiments, an antibody is modified, e.g., modified via glycosylation, phosphorylation, sumoylation, and/or (e.g., and) methylation. In some embodiments, an antibody is a glycosylated antibody, which is conjugated to one or more sugar or carbohydrate molecules. In some embodiments, the one or more sugar or carbohydrate molecule are conjugated to the antibody via N-glycosylation, O-glycosylation, C-glycosylation, glypiation (GPI anchor attachment), and/or (e.g., and) phosphoglycosylation. In some embodiments, the one or more sugar or carbohydrate molecule are monosaccharides, disaccharides, oligosaccharides, or glycans. In some embodiments, the one or more sugar or carbohydrate molecule is a branched oligosaccharide or a branched glycan. In some embodiments, the one or more sugar or carbohydrate molecule includes a mannose unit, a glucose unit, an N-acetylglucosamine unit, an N-acetylgalactosamine unit, a galactose unit, a fucose unit, or a phospholipid unit. In some embodiments, an antibody is a construct that comprises a polypeptide comprising one or more antigen binding fragments of the disclosure linked to a linker polypeptide or an immunoglobulin constant domain. Linker polypeptides comprise two or more amino acid residues joined by peptide bonds and are used to link one or more antigen binding portions. Examples of linker polypeptides have been reported (see e.g., Holliger, P., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R. J., et al. (1994) *Structure* 2:1121-1123). Still further, an antibody may be part of a larger immunoadhesion molecule, formed by covalent or noncovalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov, S. M., et al. (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S. M., et al. (1994) *Mol. Immunol.* 31:1047-1058).

[00065] CDR: As used herein, the term "CDR" refers to the complementarity determining region within antibody variable sequences. A typical antibody molecule comprises a heavy chain variable region (VH) and a light chain variable region (VL), which are usually involved in

antigen binding. The VH and VL regions can be further subdivided into regions of hypervariability, also known as “complementarity determining regions” (“CDR”), interspersed with regions that are more conserved, which are known as “framework regions” (“FR”). Each VH and VL is typically composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The extent of the framework region and CDRs can be precisely identified using methodology known in the art, for example, by the Kabat definition, the IMGT definition, the Chothia definition, the AbM definition, and/or (e.g., and) the contact definition, all of which are well known in the art. See, e.g., Kabat, E.A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; IMGT®, the international ImMunoGeneTics information system® <http://www.imgt.org>, Lefranc, M.-P. et al., Nucleic Acids Res., 27:209-212 (1999); Ruiz, M. et al., Nucleic Acids Res., 28:219-221 (2000); Lefranc, M.-P., Nucleic Acids Res., 29:207-209 (2001); Lefranc, M.-P., Nucleic Acids Res., 31:307-310 (2003); Lefranc, M.-P. et al., In Silico Biol., 5, 0006 (2004) [Epub], 5:45-60 (2005); Lefranc, M.-P. et al., Nucleic Acids Res., 33:D593-597 (2005); Lefranc, M.-P. et al., Nucleic Acids Res., 37:D1006-1012 (2009); Lefranc, M.-P. et al., Nucleic Acids Res., 43:D413-422 (2015); Chothia et al., (1989) Nature 342:877; Chothia, C. et al. (1987) J. Mol. Biol. 196:901-917, Al-lazikani et al (1997) J. Molec. Biol. 273:927-948; and Almagro, J. Mol. Recognit. 17:132-143 (2004). See also hgmp.mrc.ac.uk and bioinf.org.uk/abs. As used herein, a CDR may refer to the CDR defined by any method known in the art. Two antibodies having the same CDR means that the two antibodies have the same amino acid sequence of that CDR as determined by the same method, for example, the IMGT definition.

[00066] There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term “CDR set” as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the antigen. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Kabat *et al.*, Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs. These CDRs may be referred to as Kabat CDRs. Sub-portions of CDRs may be designated as L1, L2 and L3 or H1, H2 and H3 where the “L” and the “H” designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (FASEB J. 9:133-139 (1995))

and MacCallum (J Mol Biol 262(5):732-45 (1996)). Still other CDR boundary definitions may not strictly follow one of the above systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems. Examples of CDR definition systems are provided in Table 1.

Table 1. CDR Definitions

	IMGT ¹	Kabat ²	Chothia ³
CDR-H1	27-38	31-35	26-32
CDR-H2	56-65	50-65	53-55
CDR-H3	105-116/117	95-102	96-101
CDR-L1	27-38	24-34	26-32
CDR-L2	56-65	50-56	50-52
CDR-L3	105-116/117	89-97	91-96

¹ IMGT®, the international ImMunoGeneTics information system®, imgt.org, Lefranc, M.-P. et al., Nucleic Acids Res., 27:209-212 (1999)

² Kabat et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242

³ Chothia et al., J. Mol. Biol. 196:901-917 (1987))

[00067] CDR-grafted antibody: The term "CDR-grafted antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species but in which the sequences of one or more of the CDR regions of VH and/or (e.g., and) VL are replaced with CDR sequences of another species, such as antibodies having murine heavy and light chain variable regions in which one or more of the murine CDRs (e.g., CDR3) has been replaced with human CDR sequences.

[00068] Chimeric antibody: The term "chimeric antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

[00069] Complementary: As used herein, the term "complementary" refers to the capacity for precise pairing between two nucleotides or two sets of nucleotides. In particular, complementary is a term that characterizes an extent of hydrogen bond pairing that brings about binding between two nucleotides or two sets of nucleotides. For example, if a base at one position of an oligonucleotide is capable of hydrogen bonding with a base at the corresponding position of a target nucleic acid (e.g., an mRNA), then the bases are considered to be complementary to each other at that position. Base pairings may include both canonical Watson-Crick base pairing and non-Watson-Crick base pairing (e.g., Wobble base pairing and Hoogsteen base pairing). For example, in some embodiments, for complementary base pairings, adenosine-type bases (A) are complementary to thymidine-type bases (T) or uracil-type bases

(U), that cytosine-type bases (C) are complementary to guanosine-type bases (G), and that universal bases such as 3-nitropyrrole or 5-nitroindole can hybridize to and are considered complementary to any A, C, U, or T. Inosine (I) has also been considered in the art to be a universal base and is considered complementary to any A, C, U or T.

[00070] Conservative amino acid substitution: As used herein, a “conservative amino acid substitution” refers to an amino acid substitution that does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Fourth Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2012, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

[00071] Covalently linked: As used herein, the term “covalently linked” refers to a characteristic of two or more molecules being linked together via at least one covalent bond. In some embodiments, two molecules can be covalently linked together by a single bond, e.g., a disulfide bond or disulfide bridge, that serves as a linker between the molecules. However, in some embodiments, two or more molecules can be covalently linked together via a molecule that serves as a linker that joins the two or more molecules together through multiple covalent bonds. In some embodiments, a linker may be a cleavable linker. However, in some embodiments, a linker may be a non-cleavable linker.

[00072] Cross-reactive: As used herein and in the context of a targeting agent (e.g., antibody), the term “cross-reactive,” refers to a property of the agent being capable of specifically binding to more than one antigen of a similar type or class (e.g., antigens of multiple homologs, paralogs, or orthologs) with similar affinity or avidity. For example, in some embodiments, an antibody that is cross-reactive against human and non-human primate antigens of a similar type or class (e.g., a human transferrin receptor and non-human primate transferrin receptor) is capable of binding to the human antigen and non-human primate antigens with a similar affinity or avidity. In some embodiments, an antibody is cross-reactive against a human antigen and a rodent antigen of a similar type or class. In some embodiments, an antibody is cross-reactive against a rodent antigen and a non-human primate antigen of a similar type or class. In some embodiments, an antibody is cross-reactive against a human antigen, a non-human primate antigen, and a rodent antigen of a similar type or class.

[00073] Disease-associated-repeat: As used herein, the term “disease-associated-repeat” refers to a repeated nucleotide sequence at a genomic location for which the number of units of the repeated nucleotide sequence is correlated with and/or (e.g., and) directly or indirectly contributes to, or causes, genetic disease. Each repeating unit of a disease associated repeat may be 2, 3, 4, 5 or more nucleotides in length. For example, in some embodiments, a disease associated repeat is a dinucleotide repeat. In some embodiments, a disease associated repeat is a trinucleotide repeat. In some embodiments, a disease associated repeat is a tetranucleotide repeat. In some embodiments, a disease associated repeat is a pentanucleotide repeat. In some embodiments, the disease-associated-repeat comprises CAG repeats, CTG repeats, CUG repeats, CGG repeats, CCTG repeats, or a nucleotide complement of any thereof. In some embodiments, a disease-associated-repeat is in a non-coding portion of a gene. However, in some embodiments, a disease-associated-repeat is in a coding region of a gene. In some embodiments, a disease-associated-repeat is expanded from a normal state to a length that directly or indirectly contributes to, or causes, genetic disease. In some embodiments, a disease-associated-repeat is in RNA (e.g., an RNA transcript). In some embodiments, a disease-associated-repeat is in DNA (e.g., a chromosome, a plasmid). In some embodiments, a disease-associated-repeat is expanded in a chromosome of a germline cell. In some embodiments, a disease-associated-repeat is expanded in a chromosome of a somatic cell. In some embodiments, a disease-associated-repeat is expanded to a number of repeating units that is associated with congenital onset of disease. In some embodiments, a disease-associated-repeat is expanded to a number of repeating units that is associated with childhood onset of disease. In some embodiments, a disease-associated-repeat is expanded to a number of repeating units that is associated with adult onset of disease.

[00074] DMPK: As used herein, the term “DMPK” refers to a gene that encodes myotonin-protein kinase (also known as myotonic dystrophy protein kinase or dystrophia myotonica protein kinase), a serine/threonine protein kinase. Substrates for this enzyme may include myogenin, the beta-subunit of the L-type calcium channels, and phospholemman. In some embodiments, DMPK may be a human (Gene ID: 1760), non-human primate (e.g., Gene ID: 456139, Gene ID: 715328), or rodent gene (e.g., Gene ID: 13400). In humans, a CTG repeat expansion in the 3' non-coding, untranslated region of DMPK is associated with myotonic dystrophy type I (DM1). In addition, multiple human transcript variants (e.g., as annotated under GenBank RefSeq Accession Numbers: NM_001081563.2, NM_004409.4, NM_001081560.2, NM_001081562.2, NM_001288764.1, NM_001288765.1, and NM_001288766.1) have been characterized that encode different protein isoforms.

[00075] DMPK allele: As used herein, the term "DMPK allele" refers to any one of alternative forms (e.g., wild-type or mutant forms) of a DMPK gene. In some embodiments, a DMPK allele may encode for wild-type myotonin-protein kinase that retains its normal and typical functions. In some embodiments, a DMPK allele may comprise one or more disease-associated-repeat expansions. In some embodiments, normal subjects have two DMPK alleles comprising in the range of 5 to 37 repeat units. In some embodiments, the number of CTG repeat units in subjects having DM1 is in the range of ~50 to ~3,000+ with higher numbers of repeats leading to an increased severity of disease. In some embodiments, mildly affected DM1 subjects have at least one DMPK allele having in the range of 50 to 150 repeat units. In some embodiments, subjects with classic DM1 have at least one DMPK allele having in the range of 100 to 1,000 or more repeat units. In some embodiments, subjects having DM1 with congenital onset may have at least one DMPK allele comprising more than 2,000 repeat units.

[00076] Framework: As used herein, the term "framework" or "framework sequence" refers to the remaining sequences of a variable region minus the CDRs. Because the exact definition of a CDR sequence can be determined by different systems, the meaning of a framework sequence is subject to correspondingly different interpretations. The six CDRs (CDR-L1, CDR-L2, and CDR-L3 of light chain and CDR-H1, CDR-H2, and CDR-H3 of heavy chain) also divide the framework regions on the light chain and the heavy chain into four sub-regions (FR1, FR2, FR3 and FR4) on each chain, in which CDR1 is positioned between FR1 and FR2, CDR2 between FR2 and FR3, and CDR3 between FR3 and FR4. Without specifying the particular sub-regions as FR1, FR2, FR3 or FR4, a framework region, as referred by others, represents the combined FRs within the variable region of a single, naturally occurring immunoglobulin chain. As used herein, a FR represents one of the four sub-regions, and FRs represents two or more of the four sub-regions constituting a framework region. Human heavy chain and light chain acceptor sequences are known in the art. In one embodiment, the acceptor sequences known in the art may be used in the antibodies disclosed herein.

[00077] Human antibody: The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the disclosure may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

[00078] Humanized antibody: The term "humanized antibody" refers to antibodies which comprise heavy and light chain variable region sequences from a non-human species (*e.g.*, a mouse) but in which at least a portion of the VH and/or (*e.g.*, and) VL sequence has been altered to be more "human-like", *i.e.*, more similar to human germline variable sequences. One type of humanized antibody is a CDR-grafted antibody, in which human CDR sequences are introduced into non-human VH and VL sequences to replace the corresponding nonhuman CDR sequences. In one embodiment, humanized anti-transferrin receptor antibodies and antigen binding portions are provided. Such antibodies may be generated by obtaining murine anti-transferrin receptor monoclonal antibodies using traditional hybridoma technology followed by humanization using *in vitro* genetic engineering, such as those disclosed in Kasaian et al PCT publication No. WO 2005/123126 A2.

[00079] Internalizing cell surface receptor: As used herein, the term, "internalizing cell surface receptor" refers to a cell surface receptor that is internalized by cells, *e.g.*, upon external stimulation, *e.g.*, ligand binding to the receptor. In some embodiments, an internalizing cell surface receptor is internalized by endocytosis. In some embodiments, an internalizing cell surface receptor is internalized by clathrin-mediated endocytosis. However, in some embodiments, an internalizing cell surface receptor is internalized by a clathrin-independent pathway, such as, for example, phagocytosis, macropinocytosis, caveolae- and raft-mediated uptake or constitutive clathrin-independent endocytosis. In some embodiments, the internalizing cell surface receptor comprises an intracellular domain, a transmembrane domain, and/or (*e.g.*, and) an extracellular domain, which may optionally further comprise a ligand-binding domain. In some embodiments, a cell surface receptor becomes internalized by a cell after ligand binding. In some embodiments, a ligand may be a muscle-targeting agent or a muscle-targeting antibody. In some embodiments, an internalizing cell surface receptor is a transferrin receptor.

[00080] Isolated antibody: An "isolated antibody", as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (*e.g.*, an isolated antibody that specifically binds transferrin receptor is substantially free of antibodies that specifically bind antigens other than transferrin receptor). An isolated antibody that specifically binds transferrin receptor complex may, however, have cross-reactivity to other antigens, such as transferrin receptor molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or (*e.g.*, and) chemicals.

[00081] Kabat numbering: The terms "Kabat numbering", "Kabat definitions and "Kabat labeling" are used interchangeably herein. These terms, which are recognized in the art, refer to a system of numbering amino acid residues which are more variable (*i.e.* hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an

antigen binding portion thereof (Kabat et al. (1971) Ann. NY Acad. Sci. 190:382-391 and, Kabat, E. A., *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

[00082] Molecular payload: As used herein, the term “molecular payload” refers to a molecule or species that functions to modulate a biological outcome. In some embodiments, a molecular payload is linked to, or otherwise associated with a muscle-targeting agent. In some embodiments, the molecular payload is a small molecule, a protein, a peptide, a nucleic acid, or an oligonucleotide. In some embodiments, the molecular payload functions to modulate the transcription of a DNA sequence, to modulate the expression of a protein, or to modulate the activity of a protein. In some embodiments, the molecular payload is an oligonucleotide that comprises a strand having a region of complementarity to a target gene.

[00083] Muscle-targeting agent: As used herein, the term, “muscle-targeting agent,” refers to a molecule that specifically binds to an antigen expressed on muscle cells. The antigen in or on muscle cells may be a membrane protein, for example an integral membrane protein or a peripheral membrane protein. Typically, a muscle-targeting agent specifically binds to an antigen on muscle cells that facilitates internalization of the muscle-targeting agent (and any associated molecular payload) into the muscle cells. In some embodiments, a muscle-targeting agent specifically binds to an internalizing, cell surface receptor on muscles and is capable of being internalized into muscle cells through receptor mediated internalization. In some embodiments, the muscle-targeting agent is a small molecule, a protein, a peptide, a nucleic acid (*e.g.*, an aptamer), or an antibody. In some embodiments, the muscle-targeting agent is linked to a molecular payload.

[00084] Muscle-targeting antibody: As used herein, the term, “muscle-targeting antibody,” refers to a muscle-targeting agent that is an antibody that specifically binds to an antigen found in or on muscle cells. In some embodiments, a muscle-targeting antibody specifically binds to an antigen on muscle cells that facilitates internalization of the muscle-targeting antibody (and any associated molecular payment) into the muscle cells. In some embodiments, the muscle-targeting antibody specifically binds to an internalizing, cell surface receptor present on muscle cells. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds to a transferrin receptor.

[00085] Myotonic dystrophy (DM): As used herein, the term “Myotonic dystrophy (DM)” refers to a genetic disease caused by mutations in the DMPK gene or CNBP (ZNF9) gene that is characterized by muscle loss, muscle weakening, and muscle function. Two types of the disease, myotonic dystrophy type 1 (DM1) and myotonic dystrophy type 2 (DM2), have been described. DM1 is associated with an expansion of a CTG trinucleotide repeat in the 3' non-coding region of DMPK. DM2 is associated with an expansion of a CCTG tetranucleotide repeat in the first intron of ZNF9. In both DM1 and DM2, the nucleotide expansions lead to toxic RNA repeats capable of forming hairpin structures that bind critical intracellular proteins, e.g., muscleblind-like proteins, with high affinity. Myotonic dystrophy, the genetic basis for the disease, and related symptoms are described in the art (see, e.g. Thornton, C.A., “Myotonic Dystrophy” *Neurol Clin.* (2014), 32(3): 705–719.; and Konieczny et al. “Myotonic dystrophy: candidate small molecule therapeutics” *Drug Discovery Today* (2017), 22:11.) In some embodiments, subjects are born with a variation of DM1 called congenital myotonic dystrophy. Symptoms of congenital myotonic dystrophy are present from birth and include weakness of all muscles, breathing problems, clubfeet, developmental delays and intellectual disabilities. DM1 is associated with Online Mendelian Inheritance in Man (OMIM) Entry # 160900. DM2 is associated with OMIM Entry # 602668.

[00086] Oligonucleotide: As used herein, the term “oligonucleotide” refers to an oligomeric nucleic acid compound of up to 200 nucleotides in length. Examples of oligonucleotides include, but are not limited to, RNAi oligonucleotides (e.g., siRNAs, shRNAs), microRNAs, gapmers, mixmers, phosphorodiamidite morpholinos, peptide nucleic acids, aptamers, guide nucleic acids (e.g., Cas9 guide RNAs), etc. Oligonucleotides may be single-stranded or double-stranded. In some embodiments, an oligonucleotide may comprise one or more modified nucleotides (e.g. 2'-O-methyl sugar modifications, purine or pyrimidine modifications). In some embodiments, an oligonucleotide may comprise one or more modified internucleotide linkage. In some embodiments, an oligonucleotide may comprise one or more phosphorothioate linkages, which may be in the Rp or Sp stereochemical conformation.

[00087] Recombinant antibody: The term “recombinant human antibody”, as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described in more details in this disclosure), antibodies isolated from a recombinant, combinatorial human antibody library (Hoogenboom H. R., (1997) *TIB Tech.* 15:62-70; Azzazy H., and Highsmith W. E., (2002) *Clin. Biochem.* 35:425-445; Gavidondo J. V., and Larrick J. W. (2002) *BioTechniques* 29:128-145; Hoogenboom H., and Chames P. (2000) *Immunology Today* 21:371-378), antibodies isolated from an animal (e.g., a

mouse) that is transgenic for human immunoglobulin genes (see e.g., Taylor, L. D., et al. (1992) Nucl. Acids Res. 20:6287-6295; Kellermann S-A., and Green L. L. (2002) Current Opinion in Biotechnology 13:593-597; Little M. et al (2000) Immunology Today 21:364-370) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*. One embodiment of the disclosure provides fully human antibodies capable of binding human transferrin receptor which can be generated using techniques well known in the art, such as, but not limited to, using human Ig phage libraries such as those disclosed in Jeremius et al., PCT publication No. WO 2005/007699 A2.

[00088] Region of complementarity: As used herein, the term “region of complementarity” refers to a nucleotide sequence, e.g., of an oligonucleotide, that is sufficiently complementary to a cognate nucleotide sequence, e.g., of a target nucleic acid, such that the two nucleotide sequences are capable of annealing to one another under physiological conditions (e.g., in a cell). In some embodiments, a region of complementarity is fully complementary to a cognate nucleotide sequence of target nucleic acid. However, in some embodiments, a region of complementarity is partially complementary to a cognate nucleotide sequence of target nucleic acid (e.g., at least 80%, 90%, 95% or 99% complementarity). In some embodiments, a region of complementarity contains 1, 2, 3, or 4 mismatches compared with a cognate nucleotide sequence of a target nucleic acid.


[00089] Specifically binds: As used herein, the term “specifically binds” refers to the ability of a molecule to bind to a binding partner with a degree of affinity or avidity that enables the molecule to be used to distinguish the binding partner from an appropriate control in a binding assay or other binding context. With respect to an antibody, the term, “specifically binds”, refers to the ability of the antibody to bind to a specific antigen with a degree of affinity or avidity, compared with an appropriate reference antigen or antigens, that enables the antibody to be used to distinguish the specific antigen from others, e.g., to an extent that permits preferential targeting to certain cells, e.g., muscle cells, through binding to the antigen, as described herein. In some embodiments, an antibody specifically binds to a target if the antibody has a K_D for binding the target of at least about 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, 10^{-8} M,

10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, 10^{-13} M, or less. In some embodiments, an antibody specifically binds to the transferrin receptor, e.g., an epitope of the apical domain of transferrin receptor.

[00090] Subject: As used herein, the term “subject” refers to a mammal. In some embodiments, a subject is non-human primate, or rodent. In some embodiments, a subject is a human. In some embodiments, a subject is a patient, e.g., a human patient that has or is suspected of having a disease. In some embodiments, the subject is a human patient who has or is suspected of having a disease resulting from a disease-associated-repeat expansion, e.g., in a DMPK allele.

[00091] Transferrin receptor: As used herein, the term, “transferrin receptor” (also known as TFRC, CD71, p90, TFR, or TFR1) refers to an internalizing cell surface receptor that binds transferrin to facilitate iron uptake by endocytosis. In some embodiments, a transferrin receptor may be of human (NCBI Gene ID 7037), non-human primate (e.g., NCBI Gene ID 711568 or NCBI Gene ID 102136007), or rodent (e.g., NCBI Gene ID 22042) origin. In addition, multiple human transcript variants have been characterized that encoded different isoforms of the receptor (e.g., as annotated under GenBank RefSeq Accession Numbers: NP_001121620.1, NP_003225.2, NP_001300894.1, and NP_001300895.1).

[00092] 2'-modified nucleoside: As used herein, the terms “2'-modified nucleoside” and “2'-modified ribonucleoside” are used interchangeably and refer to a nucleoside having a sugar moiety modified at the 2' position. In some embodiments, the 2'-modified nucleoside is a 2'-4' bicyclic nucleoside, where the 2' and 4' positions of the sugar are bridged (e.g., via a methylene, an ethylene, or a (S)-constrained ethyl bridge). In some embodiments, the 2'-modified nucleoside is a non-bicyclic 2'-modified nucleoside, e.g., where the 2' position of the sugar moiety is substituted. Non-limiting examples of 2'-modified nucleosides include: 2'-deoxy, 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), 2'-O-N-methylacetamido (2'-O-NMA), locked nucleic acid (LNA, methylene-bridged nucleic acid), ethylene-bridged nucleic acid (ENA), and (S)-constrained ethyl-bridged nucleic acid (cEt). In some embodiments, the 2'-modified nucleosides described herein are high-affinity modified nucleotides and oligonucleotides comprising the 2'-modified nucleotides have increased affinity to a target sequences, relative to an unmodified oligonucleotide. Examples of structures of 2'-modified nucleosides are provided below:



[00093]

II. Complexes

[00094] Provided herein are complexes that comprise a targeting agent, e.g. an antibody, covalently linked to a molecular payload. In some embodiments, a complex comprises a muscle-targeting antibody covalently linked to an oligonucleotide. A complex may comprise an antibody that specifically binds a single antigenic site or that binds to at least two antigenic sites that may exist on the same or different antigens.

[00095] A complex may be used to modulate the activity or function of at least one gene, protein, and/or (e.g., and) nucleic acid. In some embodiments, the molecular payload present with a complex is responsible for the modulation of a gene, protein, and/or (e.g., and) nucleic acids. A molecular payload may be a small molecule, protein, nucleic acid, oligonucleotide, or any molecular entity capable of modulating the activity or function of a gene, protein, and/or (e.g., and) nucleic acid in a cell. In some embodiments, a molecular payload is an oligonucleotide that targets a disease-associated repeat in muscle cells.

[00096] In some embodiments, a complex comprises a muscle-targeting agent, e.g. an anti-transferrin receptor antibody, covalently linked to a molecular payload, e.g. an antisense oligonucleotide that targets a disease-associated repeat, e.g. DMPK allele.

A. Muscle-Targeting Agents

[00097] Some aspects of the disclosure provide muscle-targeting agents, e.g., for delivering a molecular payload to a muscle cell. In some embodiments, such muscle-targeting agents are capable of binding to a muscle cell, e.g., via specifically binding to an antigen on the

muscle cell, and delivering an associated molecular payload to the muscle cell. In some embodiments, the molecular payload is bound (*e.g.*, covalently bound) to the muscle targeting agent and is internalized into the muscle cell upon binding of the muscle targeting agent to an antigen on the muscle cell, *e.g.*, via endocytosis. It should be appreciated that various types of muscle-targeting agents may be used in accordance with the disclosure. For example, the muscle-targeting agent may comprise, or consist of, a nucleic acid (*e.g.*, DNA or RNA), a peptide (*e.g.*, an antibody), a lipid (*e.g.*, a microvesicle), or a sugar moiety (*e.g.*, a polysaccharide). Exemplary muscle-targeting agents are described in further detail herein, however, it should be appreciated that the exemplary muscle-targeting agents provided herein are not meant to be limiting.

[00098] Some aspects of the disclosure provide muscle-targeting agents that specifically bind to an antigen on muscle, such as skeletal muscle, smooth muscle, or cardiac muscle. In some embodiments, any of the muscle-targeting agents provided herein bind to (*e.g.*, specifically bind to) an antigen on a skeletal muscle cell, a smooth muscle cell, and/or (*e.g.*, and) a cardiac muscle cell.

[00099] By interacting with muscle-specific cell surface recognition elements (*e.g.*, cell membrane proteins), both tissue localization and selective uptake into muscle cells can be achieved. In some embodiments, molecules that are substrates for muscle uptake transporters are useful for delivering a molecular payload into muscle tissue. Binding to muscle surface recognition elements followed by endocytosis can allow even large molecules such as antibodies to enter muscle cells. As another example molecular payloads conjugated to transferrin or anti-transferrin receptor antibodies can be taken up by muscle cells via binding to transferrin receptor, which may then be endocytosed, *e.g.*, via clathrin-mediated endocytosis.

[000100] The use of muscle-targeting agents may be useful for concentrating a molecular payload (*e.g.*, oligonucleotide) in muscle while reducing toxicity associated with effects in other tissues. In some embodiments, the muscle-targeting agent concentrates a bound molecular payload in muscle cells as compared to another cell type within a subject. In some embodiments, the muscle-targeting agent concentrates a bound molecular payload in muscle cells (*e.g.*, skeletal, smooth, or cardiac muscle cells) in an amount that is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times greater than an amount in non-muscle cells (*e.g.*, liver, neuronal, blood, or fat cells). In some embodiments, a toxicity of the molecular payload in a subject is reduced by at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, or 95% when it is delivered to the subject when bound to the muscle-targeting agent.

[000101] In some embodiments, to achieve muscle selectivity, a muscle recognition element (*e.g.*, a muscle cell antigen) may be required. As one example, a muscle-targeting agent may be a small molecule that is a substrate for a muscle-specific uptake transporter. As another example, a muscle-targeting agent may be an antibody that enters a muscle cell via transporter-mediated endocytosis. As another example, a muscle targeting agent may be a ligand that binds to cell surface receptor on a muscle cell. It should be appreciated that while transporter-based approaches provide a direct path for cellular entry, receptor-based targeting may involve stimulated endocytosis to reach the desired site of action.

i. Muscle-Targeting Antibodies

[000102] In some embodiments, the muscle-targeting agent is an antibody. Generally, the high specificity of antibodies for their target antigen provides the potential for selectively targeting muscle cells (*e.g.*, skeletal, smooth, and/or (*e.g.*, and) cardiac muscle cells). This specificity may also limit off-target toxicity. Examples of antibodies that are capable of targeting a surface antigen of muscle cells have been reported and are within the scope of the disclosure. For example, antibodies that target the surface of muscle cells are described in Arahata K., et al. “Immunostaining of skeletal and cardiac muscle surface membrane with antibody against Duchenne muscular dystrophy peptide” *Nature* 1988; 333: 861-3; Song K.S., et al. “Expression of caveolin-3 in skeletal, cardiac, and smooth muscle cells. Caveolin-3 is a component of the sarcolemma and co-fractionates with dystrophin and dystrophin-associated glycoproteins” *J Biol Chem* 1996; 271: 15160-5; and Weisbart R.H. et al., “Cell type specific targeted intracellular delivery into muscle of a monoclonal antibody that binds myosin IIB” *Mol Immunol.* 2003 Mar, 39(13):78309; the entire contents of each of which are incorporated herein by reference.

a. Anti-Transferrin Receptor Antibodies

[000103] Some aspects of the disclosure are based on the recognition that agents binding to transferrin receptor, *e.g.*, anti-transferrin-receptor antibodies, are capable of targeting muscle cell. Transferrin receptors are internalizing cell surface receptors that transport transferrin across the cellular membrane and participate in the regulation and homeostasis of intracellular iron levels. Some aspects of the disclosure provide transferrin receptor binding proteins, which are capable of binding to transferrin receptor. Accordingly, aspects of the disclosure provide binding proteins (*e.g.*, antibodies) that bind to transferrin receptor. In some embodiments, binding proteins that bind to transferrin receptor are internalized, along with any bound molecular payload, into a muscle cell. As used herein, an antibody that binds to a transferrin receptor may be referred to interchangeably as an, transferrin receptor antibody, an anti-transferrin receptor antibody, or an anti-TfR antibody. Antibodies that bind, *e.g.* specifically

bind, to a transferrin receptor may be internalized into the cell, e.g. through receptor-mediated endocytosis, upon binding to a transferrin receptor.

[000104] It should be appreciated that anti-transferrin receptor antibodies may be produced, synthesized, and/or (e.g., and) derivatized using several known methodologies, e.g. library design using phage display. Exemplary methodologies have been characterized in the art and are incorporated by reference (Díez, P. et al. “High-throughput phage-display screening in array format”, *Enzyme and microbial technology*, 2015, 79, 34-41.; Christoph M. H. and Stanley, J.R. “Antibody Phage Display: Technique and Applications” *J Invest Dermatol.* 2014, 134:2.; Engleman, Edgar (Ed.) “Human Hybridomas and Monoclonal Antibodies.” 1985, Springer.). In other embodiments, an anti-transferrin receptor antibody has been previously characterized or disclosed. Antibodies that specifically bind to transferrin receptor are known in the art (see, e.g. US Patent. No. 4,364,934, filed 12/4/1979, “Monoclonal antibody to a human early thymocyte antigen and methods for preparing same”; US Patent No. 8,409,573, filed 6/14/2006, “Anti-CD71 monoclonal antibodies and uses thereof for treating malignant tumor cells”; US Patent No. 9,708,406, filed 5/20/2014, “Anti-transferrin receptor antibodies and methods of use”; US 9,611,323, filed 12/19/2014, “Low affinity blood brain barrier receptor antibodies and uses therefor”; WO 2015/098989, filed 12/24/2014, “Novel anti-Transferrin receptor antibody that passes through blood-brain barrier”; Schneider C. et al. “Structural features of the cell surface receptor for transferrin that is recognized by the monoclonal antibody OKT9.” *J Biol Chem.* 1982, 257:14, 8516-8522.; Lee et al. “Targeting Rat Anti-Mouse Transferrin Receptor Monoclonal Antibodies through Blood-Brain Barrier in Mouse” 2000, *J Pharmacol. Exp. Ther.*, 292: 1048-1052.).

[000105] Provided herein, in some aspects, are new anti-TfR antibodies for use as the muscle targeting agents (e.g., in muscle targeting complexes). In some embodiments, the anti-TfR antibody described herein binds to transferrin receptor with high specificity and affinity. In some embodiments, the anti-TfR antibody described herein specifically binds to any extracellular epitope of a transferrin receptor or an epitope that becomes exposed to an antibody. In some embodiments, anti-TfR antibodies provided herein bind specifically to transferrin receptor from human, non-human primates, mouse, rat, *etc.* In some embodiments, anti-TfR antibodies provided herein bind to human transferrin receptor. In some embodiments, the anti-TfR antibody described herein binds to an amino acid segment of a human or non-human primate transferrin receptor, as provided in SEQ ID NOs: 105-108. In some embodiments, the anti-TfR antibody described herein binds to an amino acid segment corresponding to amino acids 90-96 of a human transferrin receptor as set forth in SEQ ID NO: 105, which is not in the apical domain of the transferrin receptor.

[000106] An example human transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_003225.2 (transferrin receptor protein 1 isoform 1, homo sapiens) is as follows:

MMDQARSASFNLFGGEPLSYTRFSLARQVDGDN SHVEMKLAVDEEENADNNTKANVT
KPKRCSGSICYGTIAVIVFFLIGFMIGYLG YCKGVEPKTECERLAGTESPVREEPGEDFPA
ARRLYWDDLKRKLSEKLDSTDFGTIKLLNENSYVPREAGSQKDENLALYVENQFREF
KLSKVWRDQHFVKIQVKDSAQNSVIIVDKNGRLVYLVENPGGYVAYS KAATVTGKLV
HANFGTKKDFEDLYTPVNGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNA
ELSFFGHAHLGTGDPYTPGFPSFNHTQFPSSRSSGLPNIPVQTISR AAAEKLFGNMEGDCP
SDWKTDSTCRMVTSESKNVKLTVSNVLKEIKILNIFGVIKGFVEPDHYVVVGAQRDAW
GPGA AKSGVGTALLLKLAQMFS DMVLKDG FQPSRSIIFASWSAGDFG SVGATEWLEGY
LSSLHLKAFTYINLDKAVLGTSNFKVSASPLLYTLIEKTMQNVKHPVTGQFLYQDSNWA
SKVEKLTLDNAAFPFLAYSGIPAVSFCFCEDTDYPYLGTTMDTYKELIERIPELNKVARA
AAEVAGQFVIKLTHDVELNLDYERYNSQLLSFVRDLNQYRADIKEMGLSLQWLYSARG
DFFRATSRLTTDFGNAEKTDRFVMKKLNDRVMRVEYHFLSPYVSPKESPRHVFWSG
SHTLPALLENLKL RKQNNGAFNETLFRNQLALATWTIQGAANALSGDVWDIDNEF
(SEQ ID NO: 105).

[000107] An example non-human primate transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_001244232.1(transferrin receptor protein 1, Macaca mulatta) is as follows:

MMDQARSASFNLFGGEPLSYTRFSLARQVDGDN SHVEMKLG VDEEENTDNNTKPNGT
KPKRCGGNICYGTIAV IIFLIGFMIGYLG YCKGVEPKTECERLAGTESPAREEPEEDFPA
APRLYWDDLKRKLSEKLDTTDFSTIKLLNENLYVPREAGSQKDENLALYIENQFREFK
LSKVWRDQHFVKIQVKDSAQNSVIIVDKNGGLVYLVENPGGYVAYS KAATVTGKLVH
ANFGTKKDFEDLDSPVNGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVKAD
LSFFGHAHLGTGDPYTPGFPSFNHTQFPSSQS SGLPNIPVQTISR AAAEKLFGNMEGDCPS
DWKTDSTCKMVTSENKSVKLTVSNVLKETKILNIFGVIKGFVEPDHYVVVGAQRDAW
GPGA AKSSVGTALLLKLAQMFS DMVLKDG FQPSRSIIFASWSAGDFG SVGATEWLEGY
LSSLHLKAFTYINLDKAVLGTSNFKVSASPLLYTLIEKTMQDVKHPVTGRSLYQDSNWA
SKVEKLTLDNAAFPFLAYSGIPAVSFCFCEDTDYPYLGTTMDTYKELVERIPELNKVAR
AAAEVAGQFVIKLTHDTELNLDYERYNSQLLLFLRDLNQYRADVKEMGLSLQWLYSA
RGDFFRATSRLTTDFRNAEKRDKFVMKKLNDRVMRVEYYFLSPYVSPKESPRHVFWSG
SGSHTLSALLESKLRRQNNSAFNETLFRNQLALATWTIQGAANALSGDVWDIDNEF
(SEQ ID NO: 106)

[000108] An example non-human primate transferrin receptor amino acid sequence, corresponding to NCBI sequence XP_005545315.1 (transferrin receptor protein 1, *Macaca fascicularis*) is as follows:

MMDQARSASFNLFGGEPLSYTRFSLARQVDGDN SHVEMKLGVD E EENTDNNTKANGT
KPKRCGGNICYGTIAVIIFFLIGFMIGYLG YCKGVEPKTECERLAGTESPAREEPEEDFPA
APRLYWDDLKRKLSEKLDTTDFTSTIKLLNENLYVPREAGSQKDENLALYIENQFREFK
LSKVWRDQHFVKIQVKDSAQNSV IVDKNGGLVYLVENPGGYVAYS KAATVTGKLVH
ANFGTKKDFEDLDSPVNGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVKAD
LSFFGHAHLGTGDPYTPGFPSFNHTQFPPSQSSGLPNIPVQTISR AAEKLFGNMEGDCPS
DWKTDSTCKMVTSENKSVKLT VSNVLKETKILNIFGVIKGFVEPDHYVVVGAQRDAW
GPGA AKSSVGTALLLKL AQMFSDMVLKDG FQPSRSIIFASWSAGDFG SVGATEWLEGY
LSSLHLKAFTYINLDKAVLGTSNFKVSASPLLYTLIEKTMQDV KHPVTGRSLYQDSNWA
SKVEKLTLDNAAFPFLAYSGIPAVSFCFCEDTDYPYLGTTMDTYKELVERIPELNKVAR
AAAEVAGQFVIKLT HDTELNDYERYNSQLLLFLRDLNQYRADVKEMGLSLQWL YSA
RGDFFRATSRLTTDFRNAEK RDKFVMKKLNDRVMRVEYYFLSPYVSPKESPF RHVFWG
SGSHTLSALLESLKLRRQNNSAFNETLFRNQLALATWTIQGAANALSGDVWDIDNEF
(SEQ ID NO: 107).

[000109] An example mouse transferrin receptor amino acid sequence, corresponding to NCBI sequence NP_001344227.1 (transferrin receptor protein 1, *mus musculus*) is as follows:

MMDQARSASFNLFGGEPLSYTRFSLARQVDGDN SHVEMKLAAD E EENADNNMKASV
RKPKRFNGRLCFAAIALVIFFLIGFMSGYLG YCKRVEQKEECVKLAETEETDKSETMETE
DVPTSSRLYWADLKTLLSEKLSNIEFADTIKLSQNTYTPREAGSQKDESLAYYIENQFH
EFKFSKVWRDEHYVKIQVKSSIGQNMVTIVQSNGNLDPVESPEGYVAFSKPTEVSGKLV
HANFGTKKDFEELSYSVNGSLVIVRAGEITFAEKVANAQSFNAIGVLIYMDKNKFPVVE
ADLALFGHAHLGTGDPYTPGFPSFNHTQFPPSQSSGLPNIPVQTISR AAEKLF GKMEGS
CPARWNIDSSCKLELSQNQN VKLIVKNVLKERRILNIFGVIKGYEEPDRYVVVGAQRDA
LGAGVAAKSSVGTGLLLKL AQVFSDMISKDGRPSRSIIFASWTAGDFGAVGATEWLEG
YLSSLHLKAFTYINLDKVVLGTSNFKVSASPLLYTLMGKIMQDV KHPVDGKSLYRDSN
WISKVEKLSFDNAAYPFLAYSGIPAVSFCFCEDADYPYLGTRLDTYEALTQKVPQLNQM
VRTAAEVAGQLIKLTHDVELNDYEMYN SKLLSFMKDLNQFKTDIRDMGLSLQWL YS
ARGDYFRATSRLTTDFHNAEKTNR FVMREINDRIMKVEYHFLSPYVSPRESPR RHIFWG
SGSHTLSALVENLKL RQKNITAFNETLFRNQLALATWTIQGVANALSGDIWNIDNEF
(SEQ ID NO: 108)

[000110] In some embodiments, an anti-transferrin receptor antibody binds to an amino acid segment of the receptor as follows:

FKVIQVKDSAQNSVIIVDKNGRLVYLVENPGGYVAYSKAATVTGKLVHANFGTKKDFE
DLYTPVNGSIVIVRAGKITFAEKVANAESLNAIGVLIYMDQTKFPIVNAELSSFFGHAHLG
TGDPYTPGFPSFNHTQFPPSRSSGLPNIPVQTISRAAAEKLFNGMEGDCPSDWKTDSTCR
MVTSESKNVKLTVSNVLKE (SEQ ID NO: 109) and does not inhibit the binding interactions between transferrin receptors and transferrin and/or (e.g., and) human hemochromatosis protein (also known as HFE). In some embodiments, the anti-transferrin receptor antibody described herein does not bind an epitope in SEQ ID NO: 109.

[000111] Appropriate methodologies may be used to obtain and/or (e.g., and) produce antibodies, antibody fragments, or antigen-binding agents, e.g., through the use of recombinant DNA protocols. In some embodiments, an antibody may also be produced through the generation of hybridomas (see, e.g., Kohler, G and Milstein, C. "Continuous cultures of fused cells secreting antibody of predefined specificity" *Nature*, 1975, 256: 495-497). The antigen-of-interest may be used as the immunogen in any form or entity, e.g., recombinant or a naturally occurring form or entity. Hybridomas are screened using standard methods, e.g. ELISA screening, to find at least one hybridoma that produces an antibody that targets a particular antigen. Antibodies may also be produced through screening of protein expression libraries that express antibodies, e.g., phage display libraries. Phage display library design may also be used, in some embodiments, (see, e.g. U.S. Patent No 5,223,409, filed 3/1/1991, "Directed evolution of novel binding proteins"; WO 1992/18619, filed 4/10/1992, "Heterodimeric receptor libraries using phagemids"; WO 1991/17271, filed 5/1/1991, "Recombinant library screening methods"; WO 1992/20791, filed 5/15/1992, "Methods for producing members of specific binding pairs"; WO 1992/15679, filed 2/28/1992, and "Improved epitope displaying phage"). In some embodiments, an antigen-of-interest may be used to immunize a non-human animal, e.g., a rodent or a goat. In some embodiments, an antibody is then obtained from the non-human animal, and may be optionally modified using a number of methodologies, e.g., using recombinant DNA techniques. Additional examples of antibody production and methodologies are known in the art (see, e.g. Harlow et al. "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory, 1988.).

[000112] In some embodiments, an antibody is modified, e.g., modified via glycosylation, phosphorylation, sumoylation, and/or (e.g., and) methylation. In some embodiments, an antibody is a glycosylated antibody, which is conjugated to one or more sugar or carbohydrate molecules. In some embodiments, the one or more sugar or carbohydrate molecule are conjugated to the antibody via N-glycosylation, O-glycosylation, C-glycosylation, glypiation (GPI anchor attachment), and/or (e.g., and) phosphoglycosylation. In some embodiments, the one or more sugar or carbohydrate molecules are monosaccharides, disaccharides,

oligosaccharides, or glycans. In some embodiments, the one or more sugar or carbohydrate molecule is a branched oligosaccharide or a branched glycan. In some embodiments, the one or more sugar or carbohydrate molecule includes a mannose unit, a glucose unit, an N-acetylglucosamine unit, an N-acetylgalactosamine unit, a galactose unit, a fucose unit, or a phospholipid unit. In some embodiments, there are about 1-10, about 1-5, about 5-10, about 1-4, about 1-3, or about 2 sugar molecules. In some embodiments, a glycosylated antibody is fully or partially glycosylated. In some embodiments, an antibody is glycosylated by chemical reactions or by enzymatic means. In some embodiments, an antibody is glycosylated in vitro or inside a cell, which may optionally be deficient in an enzyme in the N- or O- glycosylation pathway, e.g. a glycosyltransferase. In some embodiments, an antibody is functionalized with sugar or carbohydrate molecules as described in International Patent Application Publication WO2014065661, published on May 1, 2014, entitled, "*Modified antibody, antibody-conjugate and process for the preparation thereof*".

[000113] In some embodiments, the anti-TfR antibody of the present disclosure comprises a VL domain and/or (e.g., and) VH domain of any one of the anti-TfR antibodies selected from Table 2, and comprises a constant region comprising the amino acid sequences of the constant regions of an IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule, any class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2), or any subclass (e.g., IgG2a and IgG2b) of immunoglobulin molecule. Non-limiting examples of human constant regions are described in the art, e.g., see Kabat E A et al., (1991) supra.

[000114] In some embodiments, agents binding to transferrin receptor, e.g., anti-TfR antibodies, are capable of targeting muscle cell and/or (e.g., and) mediate the transportation of an agent across the blood brain barrier. Transferrin receptors are internalizing cell surface receptors that transport transferrin across the cellular membrane and participate in the regulation and homeostasis of intracellular iron levels. Some aspects of the disclosure provide transferrin receptor binding proteins, which are capable of binding to transferrin receptor. Antibodies that bind, e.g. specifically bind, to a transferrin receptor may be internalized into the cell, e.g. through receptor-mediated endocytosis, upon binding to a transferrin receptor.

[000115] Provided herein, in some aspects, are humanized antibodies that bind to transferrin receptor with high specificity and affinity. In some embodiments, the humanized anti-TfR antibody described herein specifically binds to any extracellular epitope of a transferrin receptor or an epitope that becomes exposed to an antibody. In some embodiments, the humanized anti-TfR antibodies provided herein bind specifically to transferrin receptor from human, non-human primates, mouse, rat, *etc.* In some embodiments, the humanized anti-TfR antibodies provided herein bind to human transferrin receptor. In some embodiments, the

humanized anti-TfR antibody described herein binds to an amino acid segment of a human or non-human primate transferrin receptor, as provided in SEQ ID NOs: 105-108. In some embodiments, the humanized anti-TfR antibody described herein binds to an amino acid segment corresponding to amino acids 90-96 of a human transferrin receptor as set forth in SEQ ID NO: 105, which is not in the apical domain of the transferrin receptor. In some embodiments, the humanized anti-TfR antibodies described herein binds to TfR1 but does not bind to TfR2.

[000116] In some embodiments, an anti-TfR antibody specifically binds a TfR1 (e.g., a human or non-human primate TfR1) with binding affinity (e.g., as indicated by K_d) of at least about 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, 10^{-13} M, or less. In some embodiments, the anti-TfR antibodies described herein binds to TfR1 with a K_D of sub-nanomolar range. In some embodiments, the anti-TfR antibodies described herein selectively binds to transferrin receptor 1 (TfR1) but do not bind to transferrin receptor 2 (TfR2). In some embodiments, the anti-TfR antibodies described herein binds to human TfR1 and cyno TfR1 (e.g., with a K_d of 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M, 10^{-12} M, 10^{-13} M, or less), but does not bind to a mouse TfR1. The affinity and binding kinetics of the anti-TfR antibody can be tested using any suitable method including but not limited to biosensor technology (e.g., OCTET or BIACORE). In some embodiments, binding of any one of the anti-TfR antibody described herein does not complete with or inhibit transferrin binding to the TfR1. In some embodiments, binding of any one of the anti-TfR antibody described herein does not complete with or inhibit HFE-beta-2-microglobulin binding to the TfR1.

[000117] The anti-TfR antibodies described herein are humanized antibodies. The CDR and variable region amino acid sequences of the mouse monoclonal anti-TfR antibody from which the humanized anti-TfR antibodies described herein are derived are provided in Table 2.

Table 2. Mouse Monoclonal Anti-TfR Antibodies

Ab		No. System		
		IMGT	Kabat	Chothia
3-A4	CDR-H1	GFNIKDDY (SEQ ID NO: 1)	DDYMY (SEQ ID NO: 7)	GFNIKDD (SEQ ID NO: 12)
	CDR-H2	IDPENGDT (SEQ ID NO: 2)	WIDPENGDT EYASKFQD (SEQ ID NO: 8)	ENG (SEQ ID NO: 13)
	CDR-H3	TLWLRRGLDY (SEQ ID NO: 3)	WLRRGLDY (SEQ ID NO: 9)	LRRGLD (SEQ ID NO: 14)
	CDR-L1	KSLLHSNGYTY (SEQ ID NO: 4)	RSSKSLLHSNGYTYLF (SEQ ID NO: 10)	SKSLLHSNGYTY (SEQ ID NO: 15)
	CDR-L2	RMS (SEQ ID NO: 5)	RMSNLAS (SEQ ID NO: 11)	RMS (SEQ ID NO: 5)
	CDR-L3	MQHLEYPFT (SEQ ID NO: 6)	MQHLEYPFT (SEQ ID NO: 6)	HLEYPF (SEQ ID NO: 16)
	VH	EVQLQQSGAELVRPGASVKLSCTASGFNIKDDYMYWVKQRPEQGLEWIGWIDPENGDT EYASKFQDKATVTADTSSNTAYLQSSLTSED TAVYYCTLWLRRGLDYWGQGTSVTVS S (SEQ ID NO: 17)		

	VL	DIVMTQAAPSVPVTPGESVSISCRSSKSLHNSGYTYLFWFLQRPQGSPQLLIYRMSNLA SGVPDRFSGSGSGTAFTLRISRVEAEDVGYYCMQHLEYPFTFGGGTKLEIK (SEQ ID NO: 18)		
3-A4 N54T*	CDR- H1	GFNIKDDY (SEQ ID NO: 1)	DDYMY (SEQ ID NO: 7)	GFNIKDD (SEQ ID NO: 12)
	CDR- H2	IDPETGDT (SEQ ID NO: 19)	WIDPETGDTEYASKFQD (SEQ ID NO: 20)	ETG (SEQ ID NO: 21)
	CDR- H3	TLWLRRGLDY (SEQ ID NO: 3)	WLRRGLDY (SEQ ID NO: 9)	LRRGLD (SEQ ID NO: 14)
	CDR- L1	KSLHNSGYTY (SEQ ID NO: 4)	RSSKSLHNSGYTYLF (SEQ ID NO: 10)	SKSLHNSGYTY (SEQ ID NO: 15)
	CDR- L2	RMS (SEQ ID NO: 5)	RMSNLAS (SEQ ID NO: 11)	RMS (SEQ ID NO: 5)
	CDR- L3	MQHLEYPFT (SEQ ID NO: 6)	MQHLEYPFT (SEQ ID NO: 6)	HLEYPF (SEQ ID NO: 16)
	VH	EVQLQQSGAELVRPGASVKLSCTASGFNIKDDYMYWVKQRPEQGLEWIGWIDPETGDT EYASKFQDKATVTADTSSNTAYLQLSSLTSEDYAVYYCTLWLRRGLDYWGQGTSVTVS S (SEQ ID NO: 22)		
	VL	DIVMTQAAPSVPVTPGESVSISCRSSKSLHNSGYTYLFWFLQRPQGSPQLLIYRMSNLA SGVPDRFSGSGSGTAFTLRISRVEAEDVGYYCMQHLEYPFTFGGGTKLEIK (SEQ ID NO: 18)		
3-A4 N54S*	CDR- H1	GFNIKDDY (SEQ ID NO: 1)	DDYMY (SEQ ID NO: 7)	GFNIKDD (SEQ ID NO: 12)
	CDR- H2	IDPESGDT (SEQ ID NO: 23)	WIDPESGDTEYASKFQD (SEQ ID NO: 24)	ESG (SEQ ID NO: 25)
	CDR- H3	TLWLRRGLDY (SEQ ID NO: 3)	WLRRGLDY (SEQ ID NO: 9)	LRRGLD (SEQ ID NO: 14)
	CDR- L1	KSLHNSGYTY (SEQ ID NO: 4)	RSSKSLHNSGYTYLF (SEQ ID NO: 10)	SKSLHNSGYTY (SEQ ID NO: 15)
	CDR- L2	RMS (SEQ ID NO: 5)	RMSNLAS (SEQ ID NO: 11)	RMS (SEQ ID NO: 5)
	CDR- L3	MQHLEYPFT (SEQ ID NO: 6)	MQHLEYPFT (SEQ ID NO: 6)	HLEYPF (SEQ ID NO: 16)
	VH	EVQLQQSGAELVRPGASVKLSCTASGFNIKDDYMYWVKQRPEQGLEWIGWIDPESGDT EYASKFQDKATVTADTSSNTAYLQLSSLTSEDYAVYYCTLWLRRGLDYWGQGTSVTVS S (SEQ ID NO: 26)		
	VL	DIVMTQAAPSVPVTPGESVSISCRSSKSLHNSGYTYLFWFLQRPQGSPQLLIYRMSNLA SGVPDRFSGSGSGTAFTLRISRVEAEDVGYYCMQHLEYPFTFGGGTKLEIK (SEQ ID NO: 18)		
3-M12	CDR- H1	GYSITSGYY (SEQ ID NO: 27)	SGYYWN (SEQ ID NO: 33)	GYSITSGY (SEQ ID NO: 38)
	CDR- H2	ITFDGAN (SEQ ID NO: 28)	YITFDGANNYNPSLKN (SEQ ID NO: 34)	FDG (SEQ ID NO: 39)
	CDR- H3	TRSSYDYDVL DY (SEQ ID NO: 29)	SSYDYDVL DY (SEQ ID NO: 35)	SYDYDVL DY (SEQ ID NO: 40)
	CDR- L1	QDISNF (SEQ ID NO: 30)	RASQDISNFLN (SEQ ID NO: 36)	SQDISNF (SEQ ID NO: 41)
	CDR- L2	YTS (SEQ ID NO: 31)	YTSRLHS (SEQ ID NO: 37)	YTS (SEQ ID NO: 31)
	CDR- L3	QQGHTLPYT (SEQ ID NO: 32)	QQGHTLPYT (SEQ ID NO: 32)	GHTLPY (SEQ ID NO: 42)
	VH	DVQLQESGPGLVKPSQSLTCSVTGYSITSGYYWNWIRQFPNGKLEWMGYITFDGAN NYNPSLKNRISITRDTSKNQFFLKLTSVTEDTATYYCTRSSYDYDVL DYWGQGTTTLTV SS (SEQ ID NO: 43)		
	VL	DIQMTQTSSLSASLGDRVTISCRASQDISNFLNWWYQRPDGTVKLLIYYTSRLHSGVPS RFGSGSGTDFSLTVSNLEQEDIATYFCQQGHTLPYTFGGGTKLEIK (SEQ ID NO: 44)		
5-H12	CDR- H1	GYSFTDYC (SEQ ID NO: 45)	DYCIN (SEQ ID NO: 51)	GYSFTDY (SEQ ID NO: 56)
	CDR- H2	IYPGSGNT (SEQ ID NO: 46)	WIYPGSGNTRYSERFKG (SEQ ID NO: 52)	GSG (SEQ ID NO: 57)

	CDR-H3	AREDYYPYHGMDY (SEQ ID NO: 47)	EDYYPYHGMDY (SEQ ID NO: 53)	DYYPYHGMD (SEQ ID NO: 58)
	CDR-L1	ESVDGYDNSF (SEQ ID NO: 48)	RASESVDGYDNSFMH (SEQ ID NO: 54)	SESVDGYDNSF (SEQ ID NO: 59)
	CDR-L2	RAS (SEQ ID NO: 49)	RASNLES (SEQ ID NO: 55)	RAS (SEQ ID NO: 49)
	CDR-L3	QQSSEDPWT (SEQ ID NO: 50)	QQSSEDPWT (SEQ ID NO: 50)	SSDPW (SEQ ID NO: 60)
	VH	QIQLQQSGPELVPRGASVKISCKASGYSFTDYCNWVNQRPGGLEWIGWIYPGSGNTRYSERFKGKATLTVDTSSNTAYMQLSSLTSEDSAVYFCAREDYYPYHGMDYWGQGTSTVTVSS (SEQ ID NO: 61)		
	VL	DIVLTQSPTSLAVSLGQRATISCRASESVDGYDNSFMHWYQQKPGQPPKLLIFRASNLES GIPARFSGSGSRDTFTLTINPVEAADVATYYCQQSSEDPWTFGGGTKLEIK (SEQ ID NO: 62)		
5-H12 C33Y*	CDR-H1	GYSFTDYY (SEQ ID NO: 63)	DYYIN (SEQ ID NO: 64)	GYSFTDY (SEQ ID NO: 56)
	CDR-H2	IYPGSGNT (SEQ ID NO: 46)	WIYPGSGNTRYSERFKG (SEQ ID NO: 52)	GSG (SEQ ID NO: 57)
	CDR-H3	AREDYYPYHGMDY (SEQ ID NO: 47)	EDYYPYHGMDY (SEQ ID NO: 53)	DYYPYHGMD (SEQ ID NO: 58)
	CDR-L1	ESVDGYDNSF (SEQ ID NO: 48)	RASESVDGYDNSFMH (SEQ ID NO: 54)	SESVDGYDNSF (SEQ ID NO: 59)
	CDR-L2	RAS (SEQ ID NO: 49)	RASNLES (SEQ ID NO: 55)	RAS (SEQ ID NO: 49)
	CDR-L3	QQSSEDPWT (SEQ ID NO: 50)	QQSSEDPWT (SEQ ID NO: 50)	SSDPW (SEQ ID NO: 60)
	VH	QIQLQQSGPELVPRGASVKISCKASGYSFTDYYINWVNQRPGGLEWIGWIYPGSGNTRYSERFKGKATLTVDTSSNTAYMQLSSLTSEDSAVYFCAREDYYPYHGMDYWGQGTSTVTVSS (SEQ ID NO: 65)		
	VL	DIVLTQSPTSLAVSLGQRATISCRASESVDGYDNSFMHWYQQKPGQPPKLLIFRASNLES GIPARFSGSGSRDTFTLTINPVEAADVATYYCQQSSEDPWTFGGGTKLEIK (SEQ ID NO: 62)		
5-H12 C33D*	CDR-H1	GYSFTDYD (SEQ ID NO: 66)	DYDIN (SEQ ID NO: 67)	GYSFTDY (SEQ ID NO: 56)
	CDR-H2	IYPGSGNT (SEQ ID NO: 46)	WIYPGSGNTRYSERFKG (SEQ ID NO: 52)	GSG (SEQ ID NO: 57)
	CDR-H3	AREDYYPYHGMDY (SEQ ID NO: 47)	EDYYPYHGMDY (SEQ ID NO: 53)	DYYPYHGMD (SEQ ID NO: 58)
	CDR-L1	ESVDGYDNSF (SEQ ID NO: 48)	RASESVDGYDNSFMH (SEQ ID NO: 54)	SESVDGYDNSF (SEQ ID NO: 59)
	CDR-L2	RAS (SEQ ID NO: 49)	RASNLES (SEQ ID NO: 55)	RAS (SEQ ID NO: 49)
	CDR-L3	QQSSEDPWT (SEQ ID NO: 50)	QQSSEDPWT (SEQ ID NO: 50)	SSDPW (SEQ ID NO: 60)
	VH	QIQLQQSGPELVPRGASVKISCKASGYSFTDYDINWVNQRPGGLEWIGWIYPGSGNTRYSERFKGKATLTVDTSSNTAYMQLSSLTSEDSAVYFCAREDYYPYHGMDYWGQGTSTVTVSS (SEQ ID NO: 68)		
	VL	DIVLTQSPTSLAVSLGQRATISCRASESVDGYDNSFMHWYQQKPGQPPKLLIFRASNLES GIPARFSGSGSRDTFTLTINPVEAADVATYYCQQSSEDPWTFGGGTKLEIK (SEQ ID NO: 62)		

* mutation positions are according to Kabat numbering of the respective VH sequences containing the mutations

[000118] In some embodiments, the anti-TfR antibody of the present disclosure is a humanized variant of any one of the anti-TfR antibodies provided in Table 2. In some embodiments, the anti-TfR antibody of the present disclosure comprises a CDR-H1, a CDR-H2, a CDR-H3, a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 in any one of the anti-TfR antibodies provided in Table 2, and comprises a

humanized heavy chain variable region and/or (e.g., and) a humanized light chain variable region.

[000119] Humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementarity determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some embodiments, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences, but are included to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Antibodies may have Fc regions modified as described in WO 99/58572. Other forms of humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs derived from one or more CDRs from the original antibody. Humanized antibodies may also involve affinity maturation.

[000120] Humanized antibodies and methods of making them are known, e.g., as described in Almagro et al., *Front. Biosci.* 13:1619-1633 (2008); Riechmann et al., *Nature* 332:323-329 (1988); Queen et al., *Proc. Nat'l Acad. Sci. USA* 86:10029-10033 (1989); U.S. Pat. Nos. 5,821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri et al., *Methods* 36:25-34 (2005); Padlan et al., *Mol. Immunol.* 28:489-498 (1991); Dall'Acqua et al., *Methods* 36:43-60 (2005); Osbourn et al., *Methods* 36:61-68 (2005); and Klimka et al., *Br. J. Cancer*, 83:252-260 (2000), the contents of all of which are incorporated herein by reference. Human framework regions that may be used for humanization are described in e.g., Sims et al. *J. Immunol.* 151:2296 (1993); Carter et al., *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta et al., *J. Immunol.*, 151:2623 (1993); Almagro et al., *Front. Biosci.* 13:1619-1633 (2008)); Baca et al., *J. Biol. Chem.* 272:10678-10684 (1997); and Rosok et al., *J Biol. Chem.* 271:22611-22618 (1996), the contents of all of which are incorporated herein by reference.

[000121] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising one or more amino acid variations (e.g., in the VH framework region) as compared with any one of the VHs listed in Table 2, and/or (e.g., and) a

humanized VL comprising one or more amino acid variations (e.g., in the VL framework region) as compared with any one of the VLs listed in Table 2.

[000122] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH of any of the anti-TfR antibodies listed in Table 2 (e.g., any one of SEQ ID NOs: 17, 22, 26, 43, 61, 65, and 68). Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL of any one of the anti-TfR antibodies listed in Table 2 (e.g., any one of SEQ ID NOs: 18, 44, and 62).

[000123] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH of any of the anti-TfR antibodies listed in Table 2 (e.g., any one of SEQ ID NOs: 17, 22, 26, 43, 61, 65, and 68). Alternatively or in addition (e.g., in addition), In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL of any of the anti-TfR antibodies listed in Table 2 (e.g., any one of SEQ ID NOs: 18, 44, and 62).

[000124] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 1 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 19, or SEQ ID NO: 23 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 3 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 17, SEQ ID NO: 22, or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 4 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 5 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 6 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23,

22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 18.

[000125] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 1 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 19, or SEQ ID NO: 23 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 3 (according to the IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 17, SEQ ID NO: 22, or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 4 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 5 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 6 (according to the IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in any one of SEQ ID NO: 18.

[000126] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 7 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 8, SEQ ID NO: 20, or SEQ ID NO: 24 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 9 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 17, SEQ ID NO: 22, or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 10 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 11 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 6 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 18.

[000127] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 7 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of

SEQ ID NO: 8, SEQ ID NO: 20, or SEQ ID NO: 24 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 9 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 17, SEQ ID NO: 22, or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 10 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 11 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 6 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in any one of SEQ ID NO: 18.

[000128] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 12 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 21, or SEQ ID NO: 25 (according to the Chothia definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 14 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 17, SEQ ID NO: 22 or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 15 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 5 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 16 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 18.

[000129] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 12 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 21, or SEQ ID NO: 25 (according to the Chothia definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 14 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: SEQ ID NO: 17, SEQ

ID NO: 22 or SEQ ID NO: 26. Alternatively or in addition (e.g., in addition), the anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 15 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 5 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 16 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in any one of SEQ ID NO: 18.

[000130] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 27 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 28 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 29 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 30 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 31 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 32 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 44.

[000131] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 27 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 28 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 29 (according to the IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 30 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 31 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 32 (according to the

IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 44.

[000132] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 33 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 34 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 35 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 36 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 37 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 32 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 44.

[000133] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 33 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 34 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 35 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 36 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 37 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 32 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 44.

[000134] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 38 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 39 (according to the Chothia definition system), a CDR-H3 having the amino

acid sequence of SEQ ID NO: 40 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 41 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 31 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 42 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 44.

[000135] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 38 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 39 (according to the Chothia definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 40 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 43. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 41 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 31 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 42 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 44.

[000136] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 45, SEQ ID NO: 63, or SEQ ID NO: 66 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 46 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 47 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, or SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a

CDR-L1 having the amino acid sequence of SEQ ID NO: 48 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 49 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 50 (according to the IMGT definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 62.

[000137] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 45, SEQ ID NO: 63, or SEQ ID NO: 66 (according to the IMGT definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 46 (according to the IMGT definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 47 (according to the IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 48 (according to the IMGT definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 49 (according to the IMGT definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 50 (according to the IMGT definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 62.

[000138] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 51, SEQ ID NO: 64, or SEQ ID NO: 67 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 52 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 53 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 54 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 55 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 50 (according to the Kabat definition system), and containing no more than 25 amino acid variations (e.g., no more

than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 62.

[000139] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 51, SEQ ID NO: 64, or SEQ ID NO: 67 (according to the Kabat definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 52 (according to the Kabat definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 53 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 54 (according to the Kabat definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 55 (according to the Kabat definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 50 (according to the Kabat definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 62.

[000140] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID NO: 56 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 57 (according to the Chothia definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 58 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 59 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 49 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 60 (according to the Chothia definition system), and containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) in the framework regions as compared with the VL as set forth in SEQ ID NO: 62.

[000141] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising a CDR-H1 having the amino acid sequence of SEQ ID

NO: 56 (according to the Chothia definition system), a CDR-H2 having the amino acid sequence of SEQ ID NO: 57 (according to the Chothia definition system), a CDR-H3 having the amino acid sequence of SEQ ID NO: 58 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VH as set forth in SEQ ID NO: 61, SEQ ID NO: 65, SEQ ID NO: 68. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising a CDR-L1 having the amino acid sequence of SEQ ID NO: 59 (according to the Chothia definition system), a CDR-L2 having the amino acid sequence of SEQ ID NO: 49 (according to the Chothia definition system), and a CDR-L3 having the amino acid sequence of SEQ ID NO: 60 (according to the Chothia definition system), and is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical in the framework regions to the VL as set forth in SEQ ID NO: 62.

[000142] Examples of amino acid sequences of the humanized anti-TfR antibodies described herein are provided in Table 3.

Table 3. Variable Regions of Humanized Anti-TfR Antibodies

Antibody	Variable Region Amino Acid Sequence**
3A4 VH3 (N54T*)/Vκ4	V _H : EVQLVQSGSELKKPGASVKVSVCTASGFNIK DDYMYWVRQPPGKGLEWIGWIDP ETGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRGLD YWGQGTLVTVSS (SEQ ID NO: 69)
	V _L : DIVMTQSPLSLPVTGPGEPAISCR SSKSLLHSNGYTYLFWFQQRPGQSPRLLIYR MSNLASGV PDRFSGSGSGTDFTLKISRVEAEDVGVYYC MQHLEY PFTFGGGTK VEIK (SEQ ID NO: 70)
3A4 VH3 (N54S*)/Vκ4	V _H : EVQLVQSGSELKKPGASVKVSVCTASGFNIK DDYMYWVRQPPGKGLEWIGWIDP ESGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRGLD YWGQGTLVTVSS (SEQ ID NO: 71)
	V _L : DIVMTQSPLSLPVTGPGEPAISCR SSKSLLHSNGYTYLFWFQQRPGQSPRLLIYR MSNLASGV PDRFSGSGSGTDFTLKISRVEAEDVGVYYC MQHLEY PFTFGGGTK VEIK (SEQ ID NO: 70)
3A4 VH3 /Vκ4	V _H : EVQLVQSGSELKKPGASVKVSVCTASGFNIK DDYMYWVRQPPGKGLEWIGWIDP ENGDTTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRGLD YWGQGTLVTVSS (SEQ ID NO: 72)
	V _L : DIVMTQSPLSLPVTGPGEPAISCR SSKSLLHSNGYTYLFWFQQRPGQSPRLLIYR MSNLASGV PDRFSGSGSGTDFTLKISRVEAEDVGVYYC MQHLEY PFTFGGGTK VEIK (SEQ ID NO: 70)
3M12 VH3/Vκ2	V _H : QVQLQESGPGLVKPSQTLSTLCSVTGYSIT SGYYWNWIRQPPGKGLEWMGYITF DGANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTR SSYD YDVL DY WGQGTTVTVSS (SEQ ID NO: 73)

Antibody	Variable Region Amino Acid Sequence**
	VL: DIQMTQSPSSLSASVGDRVITTCRASQDISNFLNWFYQQKPGQPVKLLIYYTSRLH SGVPSRFSGSGSGTDFTLTISLQPEDFATYFCQQGHTLPYTFGQGTKLEIK (SEQ ID NO: 74)
3M12 VH3/Vκ3	VH: QVQLQESGPGLVKPSQTLSTCSVTGYSTSGYYWNWIRQPPGKGLEWMGYITF DGANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVL DY WGQGTITVTVSS (SEQ ID NO: 73)
	VL: DIQMTQSPSSLSASVGDRVITTCRASQDISNFLNWFYQQKPGQPVKLLIYYTSRLH SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQGHTLPYTFGQGTKLEIK (SEQ ID NO: 75)
3M12 VH4/Vκ2	VH: QVQLQESGPGLVKPSQTLSTCTVTGYSTSGYYWNWIRQPPGKGLEWIGYITFD GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVL DY WGQGTITVTVSS (SEQ ID NO: 76)
	VL: DIQMTQSPSSLSASVGDRVITTCRASQDISNFLNWFYQQKPGQPVKLLIYYTSRLH SGVPSRFSGSGSGTDFTLTISLQPEDFATYFCQQGHTLPYTFGQGTKLEIK (SEQ ID NO: 74)
3M12 VH4/Vκ3	VH: QVQLQESGPGLVKPSQTLSTCTVTGYSTSGYYWNWIRQPPGKGLEWIGYITFD GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVL DY WGQGTITVTVSS (SEQ ID NO: 76)
	VL: DIQMTQSPSSLSASVGDRVITTCRASQDISNFLNWFYQQKPGQPVKLLIYYTSRLH SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQGHTLPYTFGQGTKLEIK (SEQ ID NO: 75)
5H12 VH5 (C33Y*)/Vκ3	VH: QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIIY PGSGNTRYSERFKGRVTITRDTSASTAYMELSSLRSED TAVYYCARE DYY PYH GMDY WGQGTITVTVSS (SEQ ID NO: 77)
	VL: DIVLTQSPDSLAVSLGERATINCRASESDGYDNSFMHWYQQKPGQPPKLLIFR ASNLESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQ QSS EDPWT FGQGTKL EIK (SEQ ID NO: 78)
5H12 VH5 (C33D*)/Vκ4	VH: QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYDINWVRQAPGQGLEWMGWIIY PGSGNTRYSERFKGRVTITRDTSASTAYMELSSLRSED TAVYYCARE DYY PYH GMDY WGQGTITVTVSS (SEQ ID NO: 79)
	VL: DIVMTQSPDSLAVSLGERATINCRASESDGYDNSFMHWYQQKPGQPPKLLIFR ASNLESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQ QSS EDPWT FGQGTKL EIK (SEQ ID NO: 80)
5H12 VH5 (C33Y*)/Vκ4	VH: QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIIY PGSGNTRYSERFKGRVTITRDTSASTAYMELSSLRSED TAVYYCARE DYY PYH GMDY WGQGTITVTVSS (SEQ ID NO: 77)
	VL: DIVMTQSPDSLAVSLGERATINCRASESDGYDNSFMHWYQQKPGQPPKLLIFR ASNLESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQ QSS EDPWT FGQGTKL EIK (SEQ ID NO: 80)

* mutation positions are according to Kabat numbering of the respective VH sequences containing the mutations

** CDRs according to the Kabat numbering system are bolded

[000143] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the CDR-H1, CDR-H2, and CDR-H3 of any one of the

anti-TfR antibodies provided in Table 2 and comprises one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) amino acid variations in the framework regions as compared with the respective humanized VH provided in Table 3. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a humanized VL comprising the CDR-L1, CDR-L2, and CDR-L3 of any one of the anti-TfR antibodies provided in Table 2 and comprises one or more (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) amino acid variations in the framework regions as compared with the respective humanized VL provided in Table 3.

[000144] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 69, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 70. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 69 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 70.

[000145] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 71, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 70. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 71 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 70.

[000146] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 72, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 70. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 72 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 70.

[000147] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 73, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 74. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 73 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 74.

[000148] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 73, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 75. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 73 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 75.

[000149] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 76, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 74. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 76 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 74.

[000150] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 76, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 75. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 76 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 75.

[000151] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 77, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 78. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 77 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 78.

[000152] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 79, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 80. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 79 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 80.

[000153] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 77, and/or (e.g., and) a humanized VL comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 80. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a humanized VH comprising the amino acid sequence of SEQ ID NO: 77 and a humanized VL comprising the amino acid sequence of SEQ ID NO: 80.

[000154] In some embodiments, the humanized anti-TfR antibody described herein is a full-length IgG, which can include a heavy constant region and a light constant region from a human antibody. In some embodiments, the heavy chain of any of the anti-TfR antibodies as described herein may comprises a heavy chain constant region (CH) or a portion thereof (e.g., CH1, CH2, CH3, or a combination thereof). The heavy chain constant region can of any suitable origin, e.g., human, mouse, rat, or rabbit. In one specific example, the heavy chain constant region is from a human IgG (a gamma heavy chain), e.g., IgG1, IgG2, or IgG4. An example of a human IgG1 constant region is given below:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 81)

[000155] In some embodiments, the heavy chain of any of the anti-TfR antibodies described herein comprises a mutant human IgG1 constant region. For example, the introduction of LALA mutations (a mutant derived from mAb b12 that has been mutated to replace the lower hinge residues Leu234 Leu235 with Ala234 and Ala235) in the CH2 domain of human IgG1 is known to reduce Fcγ receptor binding (Bruhns, P., et al . (2009) and Xu, D. et al. (2000)). The mutant human IgG1 constant region is provided below (mutations bonded and underlined):

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 82)

[000156] In some embodiments, the light chain of any of the anti-TfR antibodies described herein may further comprise a light chain constant region (CL), which can be any CL known in the art. In some examples, the CL is a kappa light chain. In other examples, the CL is a lambda light chain. In some embodiments, the CL is a kappa light chain, the sequence of which is provided below:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQ
DSKDSTYLSSTLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 83)

[000157] Other antibody heavy and light chain constant regions are well known in the art, e.g., those provided in the IMGT database (www.imgt.org) or at www.vbase2.org/vbstat.php, both of which are incorporated by reference herein.

[000158] In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region that is at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to SEQ ID NO: 81 or SEQ ID NO: 82. In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region that contains no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with SEQ ID NO: 81 or SEQ ID NO: 82. In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region as set forth in SEQ ID NO: 81. In some embodiments, the humanized anti-TfR antibody described herein comprises heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region as set forth in SEQ ID NO: 82.

[000159] In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region that is at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to SEQ ID NO: 83. In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region contains no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with SEQ ID NO: 83. In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region set forth in SEQ ID NO: 83.

[000160] Examples of IgG heavy chain and light chain amino acid sequences of the anti-TfR antibodies described are provided in Table 4 below.

Table 4. Heavy chain and light chain sequences of examples of humanized anti-TfR IgGs

Antibody	IgG Heavy Chain/Light Chain Sequences**
3A4 VH3 (N54T*)/Vκ4	<p>Heavy Chain (with wild type human IgG1 constant region) <u>EVQLVQSGSELKKGASVKVSVCTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>TGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRLDYW</u> <u>GQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWSNSGAL</u> TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPGK (SEQ ID NO: 84)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPLSLPVTGPGEASISCRSSKSLHNSNGYTYLFWFQORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMOHLEYPFTFGGGTKVEIK</u> RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES VTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 85)</p>
3A4 VH3 (N54S*)/Vκ4	<p>Heavy Chain (with wild type human IgG1 constant region) <u>EVQLVQSGSELKKGASVKVSVCTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>SGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRLDYW</u> <u>GQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWSNSGAL</u> TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPGK (SEQ ID NO: 86)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPLSLPVTGPGEASISCRSSKSLHNSNGYTYLFWFQORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMOHLEYPFTFGGGTKVEIK</u> RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES VTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 85)</p>
3A4 VH3 /Vκ4	<p>Heavy Chain (with wild type human IgG1 constant region) <u>EVQLVQSGSELKKGASVKVSVCTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>NGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSEDTAVYYCTLWLRRLDYW</u> <u>GQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWSNSGAL</u> TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS CDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPGK (SEQ ID NO: 87)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPLSLPVTGPGEASISCRSSKSLHNSNGYTYLFWFQORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMOHLEYPFTFGGGTKVEIK</u> RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES VTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 85)</p>

Antibody	IgG Heavy Chain/Light Chain Sequences**
3M12 VH3/V κ 2	<p>Heavy Chain (with wild type human IgG1 constant region) <u>QVQLQESGPGLVKPSQTLSTCSVTGYSTSGYYWNWIRPPGKGLEWMGYITFD</u> <u>GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVLDYWG</u> <u>QGTTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>SGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC</u> <u>DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN</u> <u>WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP</u> <u>APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ</u> <u>PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL</u> <u>SLSPGK (SEQ ID NO: 88)</u></p> <p>Light Chain (with kappa light chain constant region) <u>DIOMTQSPSSLSASVGDRVTTTCRASQDISNFLNWYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>SVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 89)</u></p>
3M12 VH3/V κ 3	<p>Heavy Chain (with wild type human IgG1 constant region) <u>QVQLQESGPGLVKPSQTLSTCSVTGYSTSGYYWNWIRPPGKGLEWMGYITFD</u> <u>GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVLDYWG</u> <u>QGTTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>SGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC</u> <u>DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFN</u> <u>WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP</u> <u>APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ</u> <u>PENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL</u> <u>SLSPGK (SEQ ID NO: 88)</u></p> <p>Light Chain (with kappa light chain constant region) <u>DIOMTQSPSSLSASVGDRVTTTCRASQDISNFLNWYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>PSVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS</u> <u>KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 90)</u></p>
3M12 VH4/V κ 2	<p>Heavy Chain (with wild type human IgG1 constant region) <u>QVQLQESGPGLVKPSQTLSTCTVTGYSTSGYYWNWIRPPGKGLEWIGYITFDG</u> <u>ANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVLDYWGQ</u> <u>GTTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCD</u> <u>KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW</u> <u>YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA</u> <u>PIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP</u> <u>ENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL</u> <u>LSPGK (SEQ ID NO: 91)</u></p> <p>Light Chain (with kappa light chain constant region) <u>DIOMTQSPSSLSASVGDRVTTTCRASQDISNFLNWYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>SVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 89)</u></p>
3M12 VH4/V κ 3	<p>Heavy Chain (with wild type human IgG1 constant region) <u>QVQLQESGPGLVKPSQTLSTCTVTGYSTSGYYWNWIRPPGKGLEWIGYITFDG</u> <u>ANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDYDVLDYWGQ</u> <u>GTTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCD</u> <u>KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW</u> <u>YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA</u> <u>PIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP</u> <u>ENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL</u> <u>LSPGK (SEQ ID NO: 91)</u></p>

Antibody	IgG Heavy Chain/Light Chain Sequences**
	<p>Light Chain (with kappa light chain constant region) <u>DIQMTQSPSSLSASVGDRTTTCRASQDISNFLNWYQOKPGQPVKLLIYYTSRLHS</u> <u>GVPSRFSGSGSGTDFTLTISLQPEDFATYYCOQGHTLPYTFGQGTKEIKRTVAA</u> PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS KDYSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 90)</p>
5H12 VH5 (C33Y*)/Vκ3	<p>Heavy Chain (with wild type human IgG1 constant region) QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIYP <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSED</u><u>AVYYCARE</u><u>DYYPYHGM</u> <u>DYWGQGLTVTVSS</u>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPE VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGK (SEQ ID NO: 92)</p> <p>Light Chain (with kappa light chain constant region) DIVLTQSPDSLAVSLGERATINCRASESVDGYNDFMHYQOKPGQPPKLLIFRAS <u>NLESGVPDRFSGSGSRTDFTLTISLQAE</u><u>DVAVYYCQOSSEDPWTFGQGTKEIKR</u> TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVT EQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 93)</p>
5H12 VH5 (C33D*)/Vκ4	<p>Heavy Chain (with wild type human IgG1 constant region) QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYDINWVRQAPGQGLEWMGWIYP <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSED</u><u>AVYYCARE</u><u>DYYPYHGM</u> <u>DYWGQGLTVTVSS</u>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPE VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGK (SEQ ID NO: 94)</p> <p>Light Chain (with kappa light chain constant region) DIVMTQSPDSLAVSLGERATINCRASESVDGYNDFMHYQOKPGQPPKLLIFRA <u>SNLESGVPDRFSGSGSGTDFTLTISLQAE</u><u>DVAVYYCQOSSEDPWTFGQGTKEIK</u> RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES VTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 95)</p>
5H12 VH5 (C33Y*)/Vκ4	<p>Heavy Chain (with wild type human IgG1 constant region) QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIYP <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSED</u><u>AVYYCARE</u><u>DYYPYHGM</u> <u>DYWGQGLTVTVSS</u>ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKKV EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPE VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGK (SEQ ID NO: 92)</p> <p>Light Chain (with kappa light chain constant region) DIVMTQSPDSLAVSLGERATINCRASESVDGYNDFMHYQOKPGQPPKLLIFRA <u>SNLESGVPDRFSGSGSGTDFTLTISLQAE</u><u>DVAVYYCQOSSEDPWTFGQGTKEIK</u> RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES VTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 95)</p>

* mutation positions are according to Kabat numbering of the respective VH sequences containing the mutations

** CDRs according to the Kabat numbering system are bolded; VH/VL sequences underlined

[000161] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain containing no more than 25 amino acid variations (e.g., no more than

25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain as set forth in any one of SEQ ID NOs: 84, 86, 87, 88, 91, 92, and 94. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a light chain containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain as set forth in any one of SEQ ID NOs: 85, 89, 90, 93, and 95.

[000162] In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical to any one of SEQ ID NOs: 84, 86, 87, 88, 91, 92, and 94. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody described herein comprises a light chain comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical to any one of SEQ ID NOs: 85, 89, 90, 93, and 95. In some embodiments, the anti-TfR antibody described herein comprises a heavy chain comprising the amino acid sequence of any one of SEQ ID NOs: 84, 86, 87, 88, 91, 92, and 94. Alternatively or in addition (e.g., in addition), the anti-TfR antibody described herein comprises a light chain comprising the amino acid sequence of any one of SEQ ID NOs: 85, 89, 90, 93, and 95.

[000163] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 84, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 84 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000164] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 86, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 86 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000165] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 87, and/or (e.g., and) a light chain

comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 87 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000166] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 88, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 89. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 88 and a light chain comprising the amino acid sequence of SEQ ID NO: 89.

[000167] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 88, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 90. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 88 and a light chain comprising the amino acid sequence of SEQ ID NO: 90.

[000168] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 91, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 89. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 89.

[000169] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 91, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 90. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 90.

[000170] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 92, and/or (e.g., and) a light chain

comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 93. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92 and a light chain comprising the amino acid sequence of SEQ ID NO: 93.

[000171] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 94, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 95. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 94 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

[000172] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 92, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 95. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

[000173] In some embodiments, the anti-TfR antibody is a Fab fragment, Fab' fragment, or F(ab')₂ fragment of an intact antibody (full-length antibody). Antigen binding fragment of an intact antibody (full-length antibody) can be prepared via routine methods (e.g., recombinantly or by digesting the heavy chain constant region of a full length IgG using an enzyme such as papain). For example, F(ab')₂ fragments can be produced by pepsin or papain digestion of an antibody molecule, and Fab' fragments that can be generated by reducing the disulfide bridges of F(ab')₂ fragments. In some embodiments, a heavy chain constant region in a Fab fragment of the anti-TfR1 antibody described herein comprises the amino acid sequence of:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP
SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHT (SEQ ID NO: 96)

[000174] In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region that is at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to SEQ ID NO: 96. In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region that contains no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12,

11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with SEQ ID NO: 96. In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising any one of the VH as listed in Table 3 or any variants thereof and a heavy chain constant region as set forth in SEQ ID NO: 96.

[000175] In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region that is at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical to SEQ ID NO: 83. In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region contains no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with SEQ ID NO: 83. In some embodiments, the humanized anti-TfR antibody described herein comprises a light chain comprising any one of the VL as listed in Table 3 or any variants thereof and a light chain constant region set forth in SEQ ID NO: 83.

[000176] Examples of Fab heavy chain and light chain amino acid sequences of the anti-TfR antibodies described are provided in Table 5 below.

Table 5. Heavy chain and light chain sequences of examples of humanized anti-TfR Fabs

Antibody	Fab Heavy Chain/Light Chain Sequences**
3A4 VH3 (N54T*)/Vκ4	Heavy Chain (with partial human IgG1 constant region) <u>EVQLVQSGSELKKPGASVKVSTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>TGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSED</u> <u>AVYYCTLWLRRLDYW</u> <u>GQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL</u> <u>TSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS</u> CDKTHT (SEQ ID NO: 97)
	Light Chain (with kappa light chain constant region) <u>DIVMTQSPLSLPVTPGEPASISCRSSKSLLSNGYTYLFWFQORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQHLEYPFTFGGGTKVEIK</u> <u>RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES</u> <u>VTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 85)
3A4 VH3 (N54S*)/Vκ4	Heavy Chain (with partial human IgG1 constant region) <u>EVQLVQSGSELKKPGASVKVSTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>SGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSED</u> <u>AVYYCTLWLRRLDYW</u> <u>GQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL</u> <u>TSGVHTFPAVLQSSGLYSLSSVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS</u> CDKTHT (SEQ ID NO: 98)
	Light Chain (with kappa light chain constant region) <u>DIVMTQSPLSLPVTPGEPASISCRSSKSLLSNGYTYLFWFQORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQHLEYPFTFGGGTKVEIK</u> <u>RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES</u> <u>VTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 85)

Antibody	Fab Heavy Chain/Light Chain Sequences**
3A4 VH3/Vκ4	<p>Heavy Chain (with partial human IgG1 constant region) <u>EVQLVQSGSELKKPGASVKVSTASGFNIKDDYMYWVRQPPGKGLEWIGWIDPE</u> <u>NGDTEYASKFQDRVTVTADTSTNTAYMELSSLRSED</u><u>AVYYCTLWLRRLDYW</u> <u>GQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL</u> <u>TSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKS</u> <u>CDKHTT</u> (SEQ ID NO: 99)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPSLSPVTPGEPASISCRSSKSLLSNGYTYLFWFOORPGQSPRLLIYRMS</u> <u>NLASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMOHLEYPTFGGGTKVEIK</u> <u>RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES</u> <u>VTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 85)</p>
3M12 VH3/Vκ2	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLQESGPGLVKPSOTLSLTCSVTGYSITSGYYWNWIRQPPGKGLEWMGYITFD</u> <u>GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDIDVLDYWG</u> <u>QGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>SGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC</u> <u>DKTHT</u> (SEQ ID NO: 100)</p> <p>Light Chain (with kappa light chain constant region) <u>DIQMTQSPSSLSASVGDRVTITCRASQDISNFLNYYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>SVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 89)</p>
3M12 VH3/Vκ3	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLQESGPGLVKPSOTLSLTCSVTGYSITSGYYWNWIRQPPGKGLEWMGYITFD</u> <u>GANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDIDVLDYWG</u> <u>QGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>SGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC</u> <u>DKTHT</u> (SEQ ID NO: 100)</p> <p>Light Chain (with kappa light chain constant region) <u>DIQMTQSPSSLSASVGDRVTITCRASQDISNFLNYYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 90)</p>
3M12 VH4/Vκ2	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLQESGPGLVKPSOTLSLTCTVTGYSITSGYYWNWIRQPPGKGLEWIGYITFDG</u> <u>ANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDIDVLDYWGQ</u> <u>GTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCD</u> <u>KTHT</u> (SEQ ID NO: 101)</p> <p>Light Chain (with kappa light chain constant region) <u>DIQMTQSPSSLSASVGDRVTITCRASQDISNFLNYYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>SVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 89)</p>
3M12 VH4/Vκ3	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLQESGPGLVKPSOTLSLTCTVTGYSITSGYYWNWIRQPPGKGLEWIGYITFDG</u> <u>ANNYNPSLKNRVSISRDTSKNQFSLKLSSVTAEDTATYYCTRSSYDIDVLDYWGQ</u> <u>GTTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT</u> <u>GVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCD</u> <u>KTHT</u> (SEQ ID NO: 101)</p> <p>Light Chain (with kappa light chain constant region) <u>DIQMTQSPSSLSASVGDRVTITCRASQDISNFLNYYQOKPGOPVKLLIYYTSRLHS</u> <u>GVPSRFGSGSGTDFTLTISLQPEDFATYFCQOGHTLPYTFGQGTKEIKRTVAAP</u> <u>PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK</u> <u>KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC</u> (SEQ ID NO: 90)</p>

Antibody	Fab Heavy Chain/Light Chain Sequences**
5H12 VH5 (C33Y*)/Vκ3	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIYP</u> <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSEDTAVYYCAREDDYYPYHGM</u> <u>DYWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN</u> <u>SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKV</u> <u>EPKSCDKTHT</u> (SEQ ID NO: 102)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVLTQSPDSLAVSLGERATINCRASESDGYDNSEFMHWYQOKPGQPPKLLIFRAS</u> <u>NLESGVPDRFSGSGSRTDFTLTISLQAEDVAVYYCQOSSEDPWTFGQGTKLEIKR</u> <u>TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVT</u> <u>EQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC</u> (SEQ ID NO: 93)</p>
5H12 VH5 (C33D*)/Vκ4	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIYP</u> <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSEDTAVYYCAREDDYYPYHGM</u> <u>DYWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN</u> <u>SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKV</u> <u>EPKSCDKTHT</u> (SEQ ID NO: 103)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPDSLAVSLGERATINCRASESDGYDNSEFMHWYQOKPGQPPKLLIFRA</u> <u>SNLESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQOSSEDPWTFGQGTKLEIK</u> <u>RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES</u> <u>VTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC</u> (SEQ ID NO: 95)</p>
5H12 VH5 (C33Y*)/Vκ4	<p>Heavy Chain (with partial human IgG1 constant region) <u>QVQLVQSGAEVKKPGASVKVSCKASGYSTFDYYINWVRQAPGQGLEWMGWIYP</u> <u>GSGNTRYSERFKGRVTITRDTASTAYMELSSLRSEDTAVYYCAREDDYYPYHGM</u> <u>DYWGQGTLLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN</u> <u>SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKV</u> <u>EPKSCDKTHT</u> (SEQ ID NO: 102)</p> <p>Light Chain (with kappa light chain constant region) <u>DIVMTQSPDSLAVSLGERATINCRASESDGYDNSEFMHWYQOKPGQPPKLLIFRA</u> <u>SNLESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQOSSEDPWTFGQGTKLEIK</u> <u>RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES</u> <u>VTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC</u> (SEQ ID NO: 95)</p>

* mutation positions are according to Kabat numbering of the respective VH sequences containing the mutations

** CDRs according to the Kabat numbering system are bolded; VH/VL sequences underlined

[000177] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain as set forth in any one of SEQ ID NOs: 97-103. Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody of the present disclosure comprises a light chain containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain as set forth in any one of SEQ ID NOs: 85, 89, 90, 93, and 95.

[000178] In some embodiments, the humanized anti-TfR antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical to any one of SEQ ID NOs: 97-103.

Alternatively or in addition (e.g., in addition), the humanized anti-TfR antibody described herein comprises a light chain comprising an amino acid sequence that is at least 75% (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) identical to any one of SEQ ID NOs: 85, 89, 90, 93, and 95. In some embodiments, the anti-TfR antibody described herein comprises a heavy chain comprising the amino acid sequence of any one of SEQ ID NOs: 97-103. Alternatively or in addition (e.g., in addition), the anti-TfR antibody described herein comprises a light chain comprising the amino acid sequence of any one of SEQ ID NOs: 85, 89, 90, 93, and 95.

[000179] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 97, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 97 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000180] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 98, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 98 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000181] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 99, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 85. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 99 and a light chain comprising the amino acid sequence of SEQ ID NO: 85.

[000182] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 100, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 89. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 89.

[000183] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 100, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 90. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 90.

[000184] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 101, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 89. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 89.

[000185] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 101, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 90. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 90.

[000186] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 102, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 93. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 93.

[000187] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 103, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 95. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 103 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

[000188] In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to SEQ ID NO: 102, and/or (e.g., and) a light chain comprising an amino acid sequence that is at least 80% identical (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) to SEQ ID NO: 95. In some embodiments, the humanized anti-TfR antibody of the present disclosure comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

[000189] In some embodiments, the humanized anti-TfR receptor antibodies described herein can be in any antibody form, including, but not limited to, intact (i.e., full-length) antibodies, antigen-binding fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain antibodies, bi-specific antibodies, or nanobodies. In some embodiments, humanized the anti-TfR antibody described herein is a scFv. In some embodiments, the humanized anti-TfR antibody described herein is a scFv-Fab (e.g., scFv fused to a portion of a constant region). In some embodiments, the anti-TfR receptor antibody described herein is a scFv fused to a constant region (e.g., human IgG1 constant region as set forth in SEQ ID NO: 81 or SEQ ID NO: 82, or a portion thereof such as the Fc portion) at either the N-terminus or C-terminus.

[000190] In some embodiments, conservative mutations can be introduced into antibody sequences (e.g., CDRs or framework sequences) at positions where the residues are not likely to be involved in interacting with a target antigen (e.g., transferrin receptor), for example, as determined based on a crystal structure. In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc region of an anti-TfR antibody described herein (e.g., in a CH₂ domain (residues 231-340 of human IgG1) and/or (e.g., and) CH₃ domain (residues 341-447 of human IgG1) and/or (e.g., and) the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding and/or (e.g., and) antigen-dependent cellular cytotoxicity.

[000191] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the hinge region of the Fc region (CH₁ domain) such that the number of cysteine residues in the hinge region are altered (e.g., increased or decreased) as described in, e.g., U.S. Pat. No. 5,677,425. The number of cysteine residues in the hinge region of the CH₁ domain can be altered to, e.g., facilitate assembly of the light and heavy chains, or to alter (e.g., increase or decrease) the stability of the antibody or to facilitate linker conjugation.

[000192] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc region of a muscle-targeting antibody described herein (e.g., in a CH₂ domain (residues 231-340 of human IgG1) and/or (e.g., and) CH₃ domain

(residues 341-447 of human IgG1) and/or (e.g., and) the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to increase or decrease the affinity of the antibody for an Fc receptor (e.g., an activated Fc receptor) on the surface of an effector cell. Mutations in the Fc region of an antibody that decrease or increase the affinity of an antibody for an Fc receptor and techniques for introducing such mutations into the Fc receptor or fragment thereof are known to one of skill in the art. Examples of mutations in the Fc receptor of an antibody that can be made to alter the affinity of the antibody for an Fc receptor are described in, e.g., Smith P et al., (2012) PNAS 109: 6181-6186, U.S. Pat. No. 6,737,056, and International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631, which are incorporated herein by reference.

[000193] In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to alter (e.g., decrease or increase) half-life of the antibody *in vivo*. See, e.g., International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631; and U.S. Pat. Nos. 5,869,046, 6,121,022, 6,277,375 and 6,165,745 for examples of mutations that will alter (e.g., decrease or increase) the half-life of an antibody *in vivo*.

[000194] In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to decrease the half-life of the anti-TfR antibody *in vivo*. In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to increase the half-life of the antibody *in vivo*. In some embodiments, the antibodies can have one or more amino acid mutations (e.g., substitutions) in the second constant (CH2) domain (residues 231-340 of human IgG1) and/or (e.g., and) the third constant (CH3) domain (residues 341-447 of human IgG1), with numbering according to the EU index in Kabat (Kabat E A et al., (1991) *supra*). In some embodiments, the constant region of the IgG1 of an antibody described herein comprises a methionine (M) to tyrosine (Y) substitution in position 252, a serine (S) to threonine (T) substitution in position 254, and a threonine (T) to glutamic acid (E) substitution in position 256, numbered according to the EU index as in Kabat. See U.S. Pat. No. 7,658,921, which is incorporated herein by reference. This type of mutant IgG, referred to as "YTE mutant" has been shown to display fourfold increased half-life as compared to wild-type versions of the same antibody (see Dall'Acqua W F et al., (2006) J Biol Chem 281: 23514-24). In some embodiments, an antibody comprises an IgG constant domain comprising one, two, three or more amino acid

substitutions of amino acid residues at positions 251-257, 285-290, 308-314, 385-389, and 428-436, numbered according to the EU index as in Kabat.

[000195] In some embodiments, one, two or more amino acid substitutions are introduced into an IgG constant domain Fc region to alter the effector function(s) of the anti-anti-TfR antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Pat. Nos. 5,624,821 and 5,648,260. In some embodiments, the deletion or inactivation (through point mutations or other means) of a constant region domain can reduce Fc receptor binding of the circulating antibody thereby increasing tumor localization. See, e.g., U.S. Pat. Nos. 5,585,097 and 8,591,886 for a description of mutations that delete or inactivate the constant domain and thereby increase tumor localization. In some embodiments, one or more amino acid substitutions may be introduced into the Fc region of an antibody described herein to remove potential glycosylation sites on Fc region, which may reduce Fc receptor binding (see, *e.g.*, Shields R L et al., (2001) J Biol Chem 276: 6591-604).

[000196] In some embodiments, one or more amino in the constant region of an anti-TfR antibody described herein can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or (e.g., and) reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Pat. No. 6,194,551 (Idusogie et al). In some embodiments, one or more amino acid residues in the N-terminal region of the CH2 domain of an antibody described herein are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in International Publication No. WO 94/29351. In some embodiments, the Fc region of an antibody described herein is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or (e.g., and) to increase the affinity of the antibody for an Fcγ receptor. This approach is described further in International Publication No. WO 00/42072.

[000197] In some embodiments, the heavy and/or (e.g., and) light chain variable domain(s) sequence(s) of the antibodies provided herein can be used to generate, for example, CDR-grafted, chimeric, humanized, or composite human antibodies or antigen-binding fragments, as described elsewhere herein. As understood by one of ordinary skill in the art, any variant, CDR-grafted, chimeric, humanized, or composite antibodies derived from any of the antibodies provided herein may be useful in the compositions and methods described herein and will maintain the ability to specifically bind transferrin receptor, such that the variant, CDR-grafted, chimeric, humanized, or composite antibody has at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or more binding to transferrin receptor relative to the original antibody from which it is derived.

[000198] In some embodiments, the antibodies provided herein comprise mutations that confer desirable properties to the antibodies. For example, to avoid potential complications due to Fab-arm exchange, which is known to occur with native IgG4 mAbs, the antibodies provided herein may comprise a stabilizing ‘Adair’ mutation (Angal S., et al., “A single amino acid substitution abolishes the heterogeneity of chimeric mouse/human (IgG4) antibody,” *Mol Immunol* 30, 105-108; 1993), where serine 228 (EU numbering; residue 241 Kabat numbering) is converted to proline resulting in an IgG1-like hinge sequence. Accordingly, any of the antibodies may include a stabilizing ‘Adair’ mutation.

[000199] In some embodiments, an antibody is modified, e.g., modified via glycosylation, phosphorylation, sumoylation, and/or (e.g., and) methylation. In some embodiments, an antibody is a glycosylated antibody, which is conjugated to one or more sugar or carbohydrate molecules. In some embodiments, the one or more sugar or carbohydrate molecule are conjugated to the antibody via N-glycosylation, O-glycosylation, C-glycosylation, glypiation (GPI anchor attachment), and/or (e.g., and) phosphoglycosylation. In some embodiments, the one or more sugar or carbohydrate molecules are monosaccharides, disaccharides, oligosaccharides, or glycans. In some embodiments, the one or more sugar or carbohydrate molecule is a branched oligosaccharide or a branched glycan. In some embodiments, the one or more sugar or carbohydrate molecule includes a mannose unit, a glucose unit, an N-acetylglucosamine unit, an N-acetylgalactosamine unit, a galactose unit, a fucose unit, or a phospholipid unit. In some embodiments, there are about 1-10, about 1-5, about 5-10, about 1-4, about 1-3, or about 2 sugar molecules. In some embodiments, a glycosylated antibody is fully or partially glycosylated. In some embodiments, an antibody is glycosylated by chemical reactions or by enzymatic means. In some embodiments, an antibody is glycosylated in vitro or inside a cell, which may optionally be deficient in an enzyme in the N- or O- glycosylation pathway, e.g. a glycosyltransferase. In some embodiments, an antibody is functionalized with sugar or carbohydrate molecules as described in International Patent Application Publication WO2014065661, published on May 1, 2014, entitled, “*Modified antibody, antibody-conjugate and process for the preparation thereof*”.

[000200] In some embodiments, any one of the anti-TfR1 antibodies described herein may comprise a signal peptide in the heavy and/or (e.g., and) light chain sequence (e.g., a N-terminal signal peptide). In some embodiments, the anti-TfR1 antibody described herein comprises any one of the VH and VL sequences, any one of the IgG heavy chain and light chain sequences, or any one of the Fab heavy chain and light chain sequences described herein, and further comprises a signal peptide (e.g., a N-terminal signal peptide). In some embodiments, the signal

peptide comprises the amino acid sequence of MGWSCILFLVATATGVHS (SEQ ID NO: 104).

Other known anti-transferrin receptor antibodies

[000201] Any other appropriate anti-transferrin receptor antibodies known in the art may be used as the muscle-targeting agent in the complexes disclosed herein. Examples of known anti-transferrin receptor antibodies, including associated references and binding epitopes, are listed in Table 6. In some embodiments, the anti-transferrin receptor antibody comprises the complementarity determining regions (CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and CDR-L3) of any of the anti-transferrin receptor antibodies provided herein, e.g., anti-transferrin receptor antibodies listed in Table 6.

[000202] Table 6 – List of anti-transferrin receptor antibody clones, including associated references and binding epitope information.

Antibody Clone Name	Reference(s)	Epitope / Notes
OKT9	US Patent. No. 4,364,934, filed 12/4/1979, entitled “MONOCLONAL ANTIBODY TO A HUMAN EARLY THYMOCYTE ANTIGEN AND METHODS FOR PREPARING SAME” Schneider C. et al. “Structural features of the cell surface receptor for transferrin that is recognized by the monoclonal antibody OKT9.” J Biol Chem. 1982, 257:14, 8516-8522.	Apical domain of TfR (residues 305-366 of human TfR sequence XM_052730.3, available in GenBank)
(From JCR) Clone M11 Clone M23 Clone M27 Clone B84	<ul style="list-style-type: none"> WO 2015/098989, filed 12/24/2014, “Novel anti-Transferrin receptor antibody that passes through blood-brain barrier” US Patent No. 9,994,641, filed 12/24/2014, “Novel anti-Transferrin receptor antibody that passes through blood-brain barrier” 	Apical domain (residues 230-244 and 326-347 of TfR) and protease-like domain (residues 461-473)
(From Genentech) 7A4, 8A2, 15D2, 10D11, 7B10, 15G11, 16G5, 13C3, 16G4, 16F6, 7G7, 4C2, 1B12, and 13D4	<ul style="list-style-type: none"> WO 2016/081643, filed 5/26/2016, entitled “ANTI-TRANSFERRIN RECEPTOR ANTIBODIES AND METHODS OF USE” US Patent No. 9,708,406, filed 5/20/2014, “Anti-transferrin receptor antibodies and methods of use” 	Apical domain and non-apical regions

(From Armagen) 8D3	<ul style="list-style-type: none"> • Lee et al. "Targeting Rat Anti-Mouse Transferrin Receptor Monoclonal Antibodies through Blood-Brain Barrier in Mouse" 2000, J Pharmacol. Exp. Ther., 292: 1048-1052. • US Patent App. 2010/077498, filed 9/11/2008, entitled "COMPOSITIONS AND METHODS FOR BLOOD-BRAIN BARRIER DELIVERY IN THE MOUSE" 	
OX26	<ul style="list-style-type: none"> • Haobam, B. et al. 2014. Rab17-mediated recycling endosomes contribute to autophagosome formation in response to Group A Streptococcus invasion. Cellular microbiology. 16: 1806-21. 	
DF1513	<ul style="list-style-type: none"> • Ortiz-Zapater E et al. Trafficking of the human transferrin receptor in plant cells: effects of tyrphostin A23 and brefeldin A. Plant J 48:757-70 (2006). 	
1A1B2, 66IG10, MEM-189, JF0956, 29806, 1A1B2, TFRC/1818, 1E6, 66Ig10, TFRC/1059, Q1/71, 23D10, 13E4, TFRC/1149, ER-MP21, YTA74.4, BU54, 2B6, RI7 217	<ul style="list-style-type: none"> • Commercially available anti-transferrin receptor antibodies. 	Novus Biologicals 8100 Southpark Way, A-8 Littleton CO 80120
(From INSERM) BA120g	<ul style="list-style-type: none"> • US Patent App. 2011/0311544A1, filed 6/15/2005, entitled "ANTI-CD71 MONOCLONAL ANTIBODIES AND USES THEREOF FOR TREATING MALIGNANT TUMOR CELLS" 	Does not compete with OKT9
LUCA31	<ul style="list-style-type: none"> • US Patent No. 7,572,895, filed 6/7/2004, entitled "TRANSFERRIN RECEPTOR ANTIBODIES" 	"LUCA31 epitope"
(Salk Institute) B3/25 T58/30	<ul style="list-style-type: none"> • Trowbridge, I.S. et al. "Anti-transferrin receptor monoclonal antibody and toxin-antibody conjugates affect growth of human tumour cells." Nature, 1981, volume 294, pages 171-173 	
R17 217.1.3, 5E9C11,	<ul style="list-style-type: none"> • Commercially available anti-transferrin receptor antibodies. 	BioXcell 10 Technology Dr., Suite 2B

OKT9 (BE0023 clone)		West Lebanon, NH 03784-1671 USA
BK19.9, B3/25, T56/14 and T58/1	<ul style="list-style-type: none"> Gatter, K.C. et al. "Transferrin receptors in human tissues: their distribution and possible clinical relevance." J Clin Pathol. 1983 May;36(5):539-45. 	

[000203] In some embodiments, transferrin receptor antibodies of the present disclosure include one or more of the CDR-H (*e.g.*, CDR-H1, CDR-H2, and CDR-H3) amino acid sequences from any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, transferrin receptor antibodies include the CDR-H1, CDR-H2, and CDR-H3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, anti-transferrin receptor antibodies include the CDR-L1, CDR-L2, and CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, anti-transferrin antibodies include the CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 6. The disclosure also includes any nucleic acid sequence that encodes a molecule comprising a CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, or CDR-L3 as provided for any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, antibody heavy and light chain CDR3 domains may play a particularly important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, anti-transferrin receptor antibodies of the disclosure may include at least the heavy and/or (*e.g.*, and) light chain CDR3s of any one of the anti-transferrin receptor antibodies selected from Table 6.

[000204] In some examples, any of the anti- transferrin receptor antibodies of the disclosure have one or more CDR (*e.g.*, CDR-H or CDR-L) sequences substantially similar to any of the CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and/or (*e.g.*, and) CDR-L3 sequences from one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the position of one or more CDRs along the VH (*e.g.*, CDR-H1, CDR-H2, or CDR-H3) and/or (*e.g.*, and) VL (*e.g.*, CDR-L1, CDR-L2, or CDR-L3) region of an antibody described herein can vary by one, two, three, four, five, or six amino acid positions so long as immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained (*e.g.*, substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived). For example, in some embodiments, the position defining a CDR of any antibody described

herein can vary by shifting the N-terminal and/or (e.g., and) C-terminal boundary of the CDR by one, two, three, four, five, or six amino acids, relative to the CDR position of any one of the antibodies described herein, so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived). In another embodiment, the length of one or more CDRs along the VH (e.g., CDR-H1, CDR-H2, or CDR-H3) and/or (e.g., and) VL (e.g., CDR-L1, CDR-L2, or CDR-L3) region of an antibody described herein can vary (e.g., be shorter or longer) by one, two, three, four, five, or more amino acids, so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% of the binding of the original antibody from which it is derived).

[000205] Accordingly, in some embodiments, a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (e.g., and) CDR-H3 described herein may be one, two, three, four, five or more amino acids shorter than one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (e.g., and) CDR-H3 described herein may be one, two, three, four, five or more amino acids longer than one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the amino portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (e.g., and) CDR-H3 described herein can be extended by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (e.g., CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (e.g., human transferrin receptor) is maintained (e.g., substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the carboxy portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (e.g., and) CDR-H3 described herein can be extended by one, two, three, four, five or more amino acids compared to one or more of the CDRs described

herein (*e.g.*, CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained (*e.g.*, substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the amino portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (*e.g.*, and) CDR-H3 described herein can be shortened by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (*e.g.*, CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained (*e.g.*, substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, the carboxy portion of a CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and/or (*e.g.*, and) CDR-H3 described herein can be shortened by one, two, three, four, five or more amino acids compared to one or more of the CDRs described herein (*e.g.*, CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained (*e.g.*, substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). Any method can be used to ascertain whether immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained, for example, using binding assays and conditions described in the art.

[000206] In some examples, any of the anti-transferrin receptor antibodies of the disclosure have one or more CDR (*e.g.*, CDR-H or CDR-L) sequences substantially similar to any one of the anti-transferrin receptor antibodies selected from Table 6. For example, the antibodies may include one or more CDR sequence(s) from any of the anti-transferrin receptor antibodies selected from Table 6 containing up to 5, 4, 3, 2, or 1 amino acid residue variations as compared to the corresponding CDR region in any one of the CDRs provided herein (*e.g.*, CDRs from any of the anti-transferrin receptor antibodies selected from Table 6) so long as immunospecific binding to transferrin receptor (*e.g.*, human transferrin receptor) is maintained (*e.g.*, substantially maintained, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% relative to the binding of the original antibody from which it is derived). In some embodiments, any of the amino acid variations in any of the CDRs provided herein may be conservative variations. Conservative variations can be introduced into the CDRs at positions where the residues are not likely to be involved in interacting with a transferrin receptor protein (*e.g.*, a human transferrin receptor protein), for example, as determined based on a crystal

structure. Some aspects of the disclosure provide transferrin receptor antibodies that comprise one or more of the heavy chain variable (VH) and/or (e.g., and) light chain variable (VL) domains provided herein. In some embodiments, any of the VH domains provided herein include one or more of the CDR-H sequences (e.g., CDR-H1, CDR-H2, and CDR-H3) provided herein, for example, any of the CDR-H sequences provided in any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, any of the VL domains provided herein include one or more of the CDR-L sequences (e.g., CDR-L1, CDR-L2, and CDR-L3) provided herein, for example, any of the CDR-L sequences provided in any one of the anti-transferrin receptor antibodies selected from Table 6.

[000207] In some embodiments, anti-transferrin receptor antibodies of the disclosure include any antibody that includes a heavy chain variable domain and/or (e.g., and) a light chain variable domain of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, anti-transferrin receptor antibodies of the disclosure include any antibody that includes the heavy chain variable and light chain variable pairs of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6.

[000208] Aspects of the disclosure provide anti-transferrin receptor antibodies having a heavy chain variable (VH) and/or (e.g., and) a light chain variable (VL) domain amino acid sequence homologous to any of those described herein. In some embodiments, the anti-transferrin receptor antibody comprises a heavy chain variable sequence or a light chain variable sequence that is at least 75% (e.g., 80%, 85%, 90%, 95%, 98%, or 99%) identical to the heavy chain variable sequence and/ or any light chain variable sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the homologous heavy chain variable and/or (e.g., and) a light chain variable amino acid sequences do not vary within any of the CDR sequences provided herein. For example, in some embodiments, the degree of sequence variation (e.g., 75%, 80%, 85%, 90%, 95%, 98%, or 99%) may occur within a heavy chain variable and/or (e.g., and) a light chain variable sequence excluding any of the CDR sequences provided herein. In some embodiments, any of the anti-transferrin receptor antibodies provided herein comprise a heavy chain variable sequence and a light chain variable sequence that comprises a framework sequence that is at least 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to the framework sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6.

[000209] In some embodiments, an anti-transferrin receptor antibody, which specifically binds to transferrin receptor (e.g., human transferrin receptor), comprises a light chain variable

VL domain comprising any of the CDR-L domains (CDR-L1, CDR-L2, and CDR-L3), or CDR-L domain variants provided herein, of any of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, an anti-transferrin receptor antibody, which specifically binds to transferrin receptor (*e.g.*, human transferrin receptor), comprises a light chain variable VL domain comprising the CDR-L1, the CDR-L2, and the CDR-L3 of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the anti-transferrin receptor antibody comprises a light chain variable (VL) region sequence comprising one, two, three or four of the framework regions of the light chain variable region sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the anti-transferrin receptor antibody comprises one, two, three or four of the framework regions of a light chain variable region sequence which is at least 75%, 80%, 85%, 90%, 95%, or 100% identical to one, two, three or four of the framework regions of the light chain variable region sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the light chain variable framework region that is derived from said amino acid sequence consists of said amino acid sequence but for the presence of up to 10 amino acid substitutions, deletions, and/or (*e.g.*, and) insertions, preferably up to 10 amino acid substitutions. In some embodiments, the light chain variable framework region that is derived from said amino acid sequence consists of said amino acid sequence with 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 amino acid residues being substituted for an amino acid found in an analogous position in a corresponding non-human, primate, or human light chain variable framework region.

[000210] In some embodiments, an anti-transferrin receptor antibody that specifically binds to transferrin receptor comprises the CDR-L1, the CDR-L2, and the CDR-L3 of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the antibody further comprises one, two, three or all four VL framework regions derived from the VL of a human or primate antibody. The primate or human light chain framework region of the antibody selected for use with the light chain CDR sequences described herein, can have, for example, at least 70% (*e.g.*, at least 75%, 80%, 85%, 90%, 95%, 98%, or at least 99%) identity with a light chain framework region of a non-human parent antibody. The primate or human antibody selected can have the same or substantially the same number of amino acids in its light chain complementarity determining regions to that of the light chain complementarity determining regions of any of the antibodies provided herein, *e.g.*, any of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, the primate or human light chain framework region amino acid residues are from

a natural primate or human antibody light chain framework region having at least 75% identity, at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, at least 98% identity, at least 99% (or more) identity with the light chain framework regions of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6. In some embodiments, an anti-transferrin receptor antibody further comprises one, two, three or all four VL framework regions derived from a human light chain variable kappa subfamily. In some embodiments, an anti-transferrin receptor antibody further comprises one, two, three or all four VL framework regions derived from a human light chain variable lambda subfamily.

[000211] In some embodiments, any of the anti-transferrin receptor antibodies provided herein comprise a light chain variable domain that further comprises a light chain constant region. In some embodiments, the light chain constant region is a kappa, or a lambda light chain constant region. In some embodiments, the kappa or lambda light chain constant region is from a mammal, *e.g.*, from a human, monkey, rat, or mouse. In some embodiments, the light chain constant region is a human kappa light chain constant region. In some embodiments, the light chain constant region is a human lambda light chain constant region. It should be appreciated that any of the light chain constant regions provided herein may be variants of any of the light chain constant regions provided herein. In some embodiments, the light chain constant region comprises an amino acid sequence that is at least 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to any of the light chain constant regions of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6.

[000212] In some embodiments, the anti-transferrin receptor antibody is any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6.

[000213] In some embodiments, an anti-transferrin receptor antibody comprises a VL domain comprising the amino acid sequence of any anti-transferrin receptor antibody, such as any one of the anti-transferrin receptor antibodies selected from Table 6, and wherein the constant regions comprise the amino acid sequences of the constant regions of an IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule, or a human IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule. In some embodiments, an anti-transferrin receptor antibody comprises any of the VL domains, or VL domain variants, and any of the VH domains, or VH domain variants, wherein the VL and VH domains, or variants thereof, are from the same antibody clone, and wherein the constant regions comprise the amino acid sequences of the constant regions of an IgG, IgE, IgM, IgD, IgA or IgY immunoglobulin molecule, any class (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2), or any subclass (*e.g.*, IgG2a and IgG2b) of

immunoglobulin molecule. Non-limiting examples of human constant regions are described in the art, e.g., see Kabat E A et al., (1991) supra.

[000214] In some embodiments, the muscle-targeting agent is a transferrin receptor antibody (e.g., the antibody and variants thereof as described in International Application Publication WO 2016/081643, incorporated herein by reference).

[000215] The heavy chain and light chain CDRs of the antibody according to different definition systems are provided in Table 7. The different definition systems, e.g., the Kabat definition, the Chothia definition, and/or (e.g., and) the contact definition have been described. See, e.g., (e.g., Kabat, E.A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, Chothia et al., (1989) Nature 342:877; Chothia, C. et al. (1987) J. Mol. Biol. 196:901-917, Al-lazikani et al (1997) J. Molec. Biol. 273:927-948; and Almagro, J. Mol. Recognit. 17:132-143 (2004). See also hgmp.mrc.ac.uk and bioinf.org.uk/abs).

Table 7 Heavy chain and light chain CDRs of a mouse transferrin receptor antibody

CDRs	Kabat	Chothia	Contact
CDR-H1	SYWMH (SEQ ID NO: 110)	GYTFTSY (SEQ ID NO: 116)	TSYWMH (SEQ ID NO: 118)
CDR-H2	EINPTNGRTNYIEFKS (SEQ ID NO: 111)	NPTNGR (SEQ ID NO: 117)	WIGEINPTNGRTN (SEQ ID NO: 119)
CDR-H3	GTRAYHY (SEQ ID NO: 112)	GTRAYHY (SEQ ID NO: 112)	ARGTRA (SEQ ID NO: 120)
CDR-L1	RASDNLYSNLA (SEQ ID NO: 113)	RASDNLYSNLA (SEQ ID NO: 113)	YSNLAWY (SEQ ID NO: 121)
CDR-L2	DATNLAD (SEQ ID NO: 114)	DATNLAD (SEQ ID NO: 114)	LLVYDATNLA (SEQ ID NO: 122)
CDR-L3	QHFWDGTPLT (SEQ ID NO: 115)	QHFWDGTPLT (SEQ ID NO: 115)	QHFWDGTP (SEQ ID NO: 123)

[000216] The heavy chain variable domain (VH) and light chain variable domain sequences are also provided:

[000217] VH

QVQLQQPGAELVKPGASVKLSCKASGYTFTSYWMHWVKQRPQGQLEWIGEINPTNGRTNYIEFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGTRAYHYWGQGTSTVTS (SEQ ID NO: 124)

[000218] VL

DIQMTQSPASLSVSVGETVTITCRASDNLYSNLAWYQQKQKGKSPQLLVYDATNLADGVPFRFSGSGSGTQYSLKINSLSQSEDFGTYTCQHFWDGTPLTFGAGTKLELK (SEQ ID NO: 125)

[000219] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 7. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 7.

[000220] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3, which collectively contains no more than 5 amino acid variations (e.g., no more than 5, 4, 3, 2, or 1 amino acid variation) as compared with the CDR-H1, CDR-H2, and CDR-H3 as shown in Table 7. “Collectively” means that the total number of amino acid variations in all of the three heavy chain CDRs is within the defined range. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure may comprise a CDR-L1, a CDR-L2, and a CDR-L3, which collectively contains no more than 5 amino acid variations (e.g., no more than 5, 4, 3, 2 or 1 amino acid variation) as compared with the CDR-L1, CDR-L2, and CDR-L3 as shown in Table 7.

[000221] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, and a CDR-H3, at least one of which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the counterpart heavy chain CDR as shown in Table 7. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure may comprise CDR-L1, a CDR-L2, and a CDR-L3, at least one of which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the counterpart light chain CDR as shown in Table 7.

[000222] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3, which contains no more than 3 amino acid variations (e.g., no more than 3, 2, or 1 amino acid variation) as compared with the CDR-L3 as shown in Table 7. In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3 containing one amino acid variation as compared with the CDR-L3 as shown in Table 7. In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-L3 of QHFAGTPLT (SEQ ID NO: 126) according to the Kabat and Chothia definition system) or QHFAGTPL (SEQ ID NO: 127) according to the Contact definition system). In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, a CDR-H3, a CDR-L1 and a CDR-L2 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 7, and comprises a CDR-L3 of QHFAGTPLT (SEQ ID NO: 126) according to the Kabat and Chothia definition system) or QHFAGTPL (SEQ ID NO: 127) according to the Contact definition system).

[000223] In some embodiments, the transferrin receptor antibody of the present disclosure comprises heavy chain CDRs that collectively are at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the heavy chain CDRs as shown in Table 7. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises light chain CDRs that collectively are at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the light chain CDRs as shown in Table 7.

[000224] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising the amino acid sequence of SEQ ID NO: 124. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL comprising the amino acid sequence of SEQ ID NO: 125.

[000225] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VH as set forth in SEQ ID NO: 124. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VL as set forth in SEQ ID NO: 125.

[000226] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VH as set forth in SEQ ID NO: 124. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VL as set forth in SEQ ID NO: 125.

[000227] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized antibody (e.g., a humanized variant of an antibody). In some embodiments, the transferrin receptor antibody of the present disclosure comprises a CDR-H1, a CDR-H2, a CDR-H3, a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 7, and comprises a humanized heavy chain variable region and/or (e.g., and) a humanized light chain variable region.

[000228] Humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some embodiments, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human

residues. Furthermore, the humanized antibody may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences, but are included to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Antibodies may have Fc regions modified as described in WO 99/58572. Other forms of humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs derived from one or more CDRs from the original antibody. Humanized antibodies may also involve affinity maturation.

[000229] In some embodiments, humanization is achieved by grafting the CDRs (e.g., as shown in Table 7) into the IGKV1-NL1*01 and IGHV1-3*01 human variable domains. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising one or more amino acid substitutions at positions 9, 13, 17, 18, 40, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 125, and/or (e.g., and) one or more amino acid substitutions at positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 66, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 124. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at all of positions 9, 13, 17, 18, 40, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 125, and/or (e.g., and) amino acid substitutions at all of positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 66, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 124.

[000230] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized antibody and contains the residues at positions 43 and 48 of the VL as set forth in SEQ ID NO: 125. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure is a humanized antibody and contains the residues at positions 48, 67, 69, 71, and 73 of the VH as set forth in SEQ ID NO: 124.

[000231] The VH and VL amino acid sequences of an example humanized antibody that may be used in accordance with the present disclosure are provided:

[000232] Humanized VH

EVQLVQSGAEVKKPGASVKVSCKASGYTFTSYWMHWVRQAPGQRLEWIGEINPTNGR
TNYIEKFKSRATLTVDKSASTAYMELSSLRSEDVAVYYCARGTRAYHYWGQGTMTVTV
SS (SEQ ID NO: 128)

[000233] Humanized VL

DIQMTQSPSSLSASVGDRVTITCRASDNLYSNLAWYQQKPGKSPKLLVYDATNLADGV
PSRFGSGSGTDYTLTISSLQPEDFATYYCQHFHWGTPLTFGQGTKVEIK
(SEQ ID NO: 129)

[000234] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising the amino acid sequence of SEQ ID NO: 128. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL comprising the amino acid sequence of SEQ ID NO: 129.

[000235] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VH as set forth in SEQ ID NO: 128. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the VL as set forth in SEQ ID NO: 129.

[000236] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a VH comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VH as set forth in SEQ ID NO: 128. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a VL comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to the VL as set forth in SEQ ID NO: 129.

[000237] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at one or more of positions 43 and 48 as compared with the VL as set forth in SEQ ID NO: 125, and/or (e.g., and) amino acid substitutions at one or more of positions 48, 67, 69, 71, and 73 as compared with the VH as set forth in SEQ ID NO: 124. In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising a S43A and/or (e.g., and) a V48L mutation as compared with the VL as set forth in SEQ ID NO: 125, and/or (e.g., and) one or more of A67V, L69I, V71R, and K73T mutations as compared with the VH as set forth in SEQ ID NO: 124.

[000238] In some embodiments, the transferrin receptor antibody of the present disclosure is a humanized variant comprising amino acid substitutions at one or more of positions 9, 13, 17, 18, 40, 43, 48, 45, and 70 as compared with the VL as set forth in SEQ ID NO: 125, and/or (e.g., and) amino acid substitutions at one or more of positions 1, 5, 7, 11, 12, 20, 38, 40, 44, 48, 66, 67, 69, 71, 73, 75, 81, 83, 87, and 108 as compared with the VH as set forth in SEQ ID NO: 124.

[000239] In some embodiments, the transferrin receptor antibody of the present disclosure is a chimeric antibody, which can include a heavy constant region and a light constant region from a human antibody. Chimeric antibodies refer to antibodies having a variable region or part of variable region from a first species and a constant region from a second species. Typically, in these chimeric antibodies, the variable region of both light and heavy chains mimics the variable regions of antibodies derived from one species of mammals (e.g., a non-human mammal such as mouse, rabbit, and rat), while the constant portions are homologous to the sequences in antibodies derived from another mammal such as human. In some embodiments, amino acid modifications can be made in the variable region and/or (e.g., and) the constant region.

[000240] In some embodiments, the transferrin receptor antibody described herein is a chimeric antibody, which can include a heavy constant region and a light constant region from a human antibody. Chimeric antibodies refer to antibodies having a variable region or part of variable region from a first species and a constant region from a second species. Typically, in these chimeric antibodies, the variable region of both light and heavy chains mimics the variable regions of antibodies derived from one species of mammals (e.g., a non-human mammal such as mouse, rabbit, and rat), while the constant portions are homologous to the sequences in antibodies derived from another mammal such as human. In some embodiments, amino acid modifications can be made in the variable region and/or (e.g., and) the constant region.

[000241] In some embodiments, the heavy chain of any of the transferrin receptor antibodies as described herein may comprises a heavy chain constant region (CH) or a portion thereof (e.g., CH1, CH2, CH3, or a combination thereof). The heavy chain constant region can of any suitable origin, e.g., human, mouse, rat, or rabbit. In one specific example, the heavy chain constant region is from a human IgG (a gamma heavy chain), e.g., IgG1, IgG2, or IgG4. An example of human IgG1 constant region is given below:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS
SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLG
GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLT
PSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 130)

[000242] In some embodiments, the light chain of any of the transferrin receptor antibodies described herein may further comprise a light chain constant region (CL), which can be any CL known in the art. In some examples, the CL is a kappa light chain. In other examples, the CL is a lambda light chain. In some embodiments, the CL is a kappa light chain, the sequence of which is provided below:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQ
DSKDSTYLSSTLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 83)

[000243] Other antibody heavy and light chain constant regions are well known in the art, e.g., those provided in the IMGT database (www.imgt.org) or at www.vbase2.org/vbstat.php, both of which are incorporated by reference herein.

[000244] Examples of heavy chain and light chain amino acid sequences of the transferrin receptor antibodies described are provided below:

[000245] Heavy Chain (VH + human IgG1 constant region)

QVQLQPGAELVKPGASVKLSCKASGYTFTSYWMHWVKQRPGGLEWIGEINPTNGR
TNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGTRAYHYWGQGTSTVTS
SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEL
GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPRE
EQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL
PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLT
VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 132)

[000246] Light Chain (VL + kappa light chain)

DIQMTQSPASLSVSVGETVTITCRASDNLYSNLAWYQQKQKGKSPQLLVYDATNLADGV
PSRFGSGSGGTQYSLKINSLSQSEDFGTYYCQHFHWGTPLTFGAGTKLELKRTVAAPSVFIF
PPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLS
STLTLSKADYEEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 133)

[000247] Heavy Chain (humanized VH + human IgG1 constant region)

EVQLVQSGAEVKKPGASVKVSCASGYTFTSYWMHWVRQAPGQRLEWIGEINPTNGR
TNYIEKFKSRATLTVDKSASTAYMELSSLRSEDVAVYYCARGTRAYHYWGQGTMTVTV
SSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEL
LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPR
EEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT
LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKL
TVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 134)

[000248] Light Chain (humanized VL + kappa light chain)
DIQMTQSPSSLSASVGDRVITICRASDNLYSNLAWYQQKPGKSPKLLVYDATNLADGV
PSRFGSGSGTDYTLTISSLQPEDFATYYCQHFHWGTPLEFGGQTKVEIKRTVAAPSVFIFP
PSDEQLKSGTASVCLLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKDESTYSLSS
TLTLKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 135)

[000249] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 132. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 133. In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 132. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 133.

[000250] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a heavy chain containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain as set forth in SEQ ID NO: 132. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a light chain containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain as set forth in SEQ ID NO: 133.

[000251] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 134. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising an amino acid sequence that is at least 80% (e.g., 80%, 85%, 90%, 95%, or 98%) identical to SEQ ID NO: 135. In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 134. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 135.

[000252] In some embodiments, the transferrin receptor antibody of the present disclosure comprises a heavy chain containing no more than 25 amino acid variations (e.g., no more than 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the heavy chain of humanized antibody as set forth in SEQ ID NO:

134. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody of the present disclosure comprises a light chain containing no more than 15 amino acid variations (e.g., no more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 9, 8, 7, 6, 5, 4, 3, 2, or 1 amino acid variation) as compared with the light chain of humanized antibody as set forth in SEQ ID NO: 135.

[000253] In some embodiments, the transferrin receptor antibody is an antigen binding fragment (Fab) of an intact antibody (full-length antibody). Antigen binding fragment of an intact antibody (full-length antibody) can be prepared via routine methods. For example, F(ab')₂ fragments can be produced by pepsin digestion of an antibody molecule, and Fab' fragments that can be generated by reducing the disulfide bridges of F(ab')₂ fragments. Examples of Fab amino acid sequences of the transferrin receptor antibodies described herein are provided below:

[000254] Heavy Chain Fab (VH + a portion of human IgG1 constant region)
QVQLQQPGAELVKPGASVKLSCKASGYTFTSYWMHWVKQRPQGQLEWIGEINPTNGR
TNYIEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGTRAYHYWGQGTSTVTVS
SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ
SSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP (SEQ ID
NO: 136)

[000255] Heavy Chain Fab (humanized VH + a portion of human IgG1 constant region)
EVQLVQSGAEVKKPGASVKVSKASGYTFTSYWMHWVRQAPGQRLEWIGEINPTNGR
TNYIEKFKSRATLTVDKSASTAYMELSSLRSEDTAVYYCARGTRAYHYWGQGTMTVTVS
SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL
QSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP (SEQ ID
NO: 137)

[000256] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 136. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 133.

[000257] In some embodiments, the transferrin receptor antibody described herein comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 137. Alternatively or in addition (e.g., in addition), the transferrin receptor antibody described herein comprises a light chain comprising the amino acid sequence of SEQ ID NO: 135.

[000258] The transferrin receptor antibodies described herein can be in any antibody form, including, but not limited to, intact (i.e., full-length) antibodies, antigen-binding fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain antibodies, bi-specific antibodies, or nanobodies. In some embodiments, the transferrin receptor antibody described herein is a scFv.

In some embodiments, the transferrin receptor antibody described herein is a scFv-Fab (e.g., scFv fused to a portion of a constant region). In some embodiments, the transferrin receptor antibody described herein is a scFv fused to a constant region (e.g., human IgG1 constant region as set forth in SEQ ID NO: 130).

[000259] In some embodiments, any one of the anti-TfR antibodies described herein is produced by recombinant DNA technology in Chinese hamster ovary (CHO) cell suspension culture, optionally in CHO-K1 cell (e.g., CHO-K1 cells derived from European Collection of Animal Cell Culture, Cat. No. 85051005) suspension culture.

[000260] In some embodiments, an antibody provided herein may have one or more post-translational modifications. In some embodiments, N-terminal cyclization, also called pyroglutamate formation (pyro-Glu), may occur in the antibody at N-terminal Glutamate (Glu) and/or Glutamine (Gln) residues during production. As such, it should be appreciated that an antibody specified as having a sequence comprising an N-terminal glutamate or glutamine residue encompasses antibodies that have undergone pyroglutamate formation resulting from a post-translational modification. In some embodiments, pyroglutamate formation occurs in a heavy chain sequence. In some embodiments, pyroglutamate formation occurs in a light chain sequence.

b. Other Muscle-Targeting Antibodies

[000261] In some embodiments, the muscle-targeting antibody is an antibody that specifically binds hemojuvelin, caveolin-3, Duchenne muscular dystrophy peptide, myosin IIb, or CD63. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a myogenic precursor protein. Exemplary myogenic precursor proteins include, without limitation, ABCG2, M-Cadherin/Cadherin-15, Caveolin-1, CD34, FoxK1, Integrin alpha 7, Integrin alpha 7 beta 1, MYF-5, MyoD, Myogenin, NCAM-1/CD56, Pax3, Pax7, and Pax9. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a skeletal muscle protein. Exemplary skeletal muscle proteins include, without limitation, alpha-Sarcoglycan, beta-Sarcoglycan, Calpain Inhibitors, Creatine Kinase MM/CKMM, eIF5A, Enolase 2/Neuron-specific Enolase, epsilon-Sarcoglycan, FABP3/H-FABP, GDF-8/Myostatin, GDF-11/GDF-8, Integrin alpha 7, Integrin alpha 7 beta 1, Integrin beta 1/CD29, MCAM/CD146, MyoD, Myogenin, Myosin Light Chain Kinase Inhibitors, NCAM-1/CD56, and Troponin I. In some embodiments, the muscle-targeting antibody is an antibody that specifically binds a smooth muscle protein. Exemplary smooth muscle proteins include, without limitation, alpha-Smooth Muscle Actin, VE-Cadherin, Caldesmon/CALD1, Calponin 1, Desmin, Histamine H2 R, Motilin R/GPR38, Transgelin/TAGLN, and Vimentin. However, it should be appreciated

that antibodies to additional targets are within the scope of this disclosure and the exemplary lists of targets provided herein are not meant to be limiting.

c. Antibody Features/Alterations

[000262] In some embodiments, conservative mutations can be introduced into antibody sequences (e.g., CDRs or framework sequences) at positions where the residues are not likely to be involved in interacting with a target antigen (e.g., transferrin receptor), for example, as determined based on a crystal structure. In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc region of a muscle-targeting antibody described herein (e.g., in a CH2 domain (residues 231-340 of human IgG1) and/or (e.g., and) CH3 domain (residues 341-447 of human IgG1) and/or (e.g., and) the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding and/or (e.g., and) antigen-dependent cellular cytotoxicity.

[000263] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the hinge region of the Fc region (CH1 domain) such that the number of cysteine residues in the hinge region are altered (e.g., increased or decreased) as described in, e.g., U.S. Pat. No. 5,677,425. The number of cysteine residues in the hinge region of the CH1 domain can be altered to, e.g., facilitate assembly of the light and heavy chains, or to alter (e.g., increase or decrease) the stability of the antibody or to facilitate linker conjugation.

[000264] In some embodiments, one, two or more mutations (e.g., amino acid substitutions) are introduced into the Fc region of a muscle-targeting antibody described herein (e.g., in a CH2 domain (residues 231-340 of human IgG1) and/or (e.g., and) CH3 domain (residues 341-447 of human IgG1) and/or (e.g., and) the hinge region, with numbering according to the Kabat numbering system (e.g., the EU index in Kabat)) to increase or decrease the affinity of the antibody for an Fc receptor (e.g., an activated Fc receptor) on the surface of an effector cell. Mutations in the Fc region of an antibody that decrease or increase the affinity of an antibody for an Fc receptor and techniques for introducing such mutations into the Fc receptor or fragment thereof are known to one of skill in the art. Examples of mutations in the Fc receptor of an antibody that can be made to alter the affinity of the antibody for an Fc receptor are described in, e.g., Smith P et al., (2012) PNAS 109: 6181-6186, U.S. Pat. No. 6,737,056, and International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631, which are incorporated herein by reference.

[000265] In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to alter (e.g., decrease or increase) half-

life of the antibody *in vivo*. See, e.g., International Publication Nos. WO 02/060919; WO 98/23289; and WO 97/34631; and U.S. Pat. Nos. 5,869,046, 6,121,022, 6,277,375 and 6,165,745 for examples of mutations that will alter (*e.g.*, decrease or increase) the half-life of an antibody *in vivo*.

[000266] In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to decrease the half-life of the anti-transferrin receptor antibody *in vivo*. In some embodiments, one, two or more amino acid mutations (*i.e.*, substitutions, insertions or deletions) are introduced into an IgG constant domain, or FcRn-binding fragment thereof (preferably an Fc or hinge-Fc domain fragment) to increase the half-life of the antibody *in vivo*. In some embodiments, the antibodies can have one or more amino acid mutations (*e.g.*, substitutions) in the second constant (CH2) domain (residues 231-340 of human IgG1) and/or (*e.g.*, and) the third constant (CH3) domain (residues 341-447 of human IgG1), with numbering according to the EU index in Kabat (Kabat E A et al., (1991) *supra*). In some embodiments, the constant region of the IgG1 of an antibody described herein comprises a methionine (M) to tyrosine (Y) substitution in position 252, a serine (S) to threonine (T) substitution in position 254, and a threonine (T) to glutamic acid (E) substitution in position 256, numbered according to the EU index as in Kabat. See U.S. Pat. No. 7,658,921, which is incorporated herein by reference. This type of mutant IgG, referred to as "YTE mutant" has been shown to display fourfold increased half-life as compared to wild-type versions of the same antibody (see Dall'Acqua W F et al., (2006) *J Biol Chem* 281: 23514-24). In some embodiments, an antibody comprises an IgG constant domain comprising one, two, three or more amino acid substitutions of amino acid residues at positions 251-257, 285-290, 308-314, 385-389, and 428-436, numbered according to the EU index as in Kabat.

[000267] In some embodiments, one, two or more amino acid substitutions are introduced into an IgG constant domain Fc region to alter the effector function(s) of the anti-transferrin receptor antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Pat. Nos. 5,624,821 and 5,648,260. In some embodiments, the deletion or inactivation (through point mutations or other means) of a constant region domain can reduce Fc receptor binding of the circulating antibody thereby increasing tumor localization. See, e.g., U.S. Pat. Nos. 5,585,097 and 8,591,886 for a description of mutations that delete or inactivate the constant domain and thereby increase tumor localization. In some embodiments, one or more amino acid substitutions may be introduced into the Fc region of an antibody described herein to remove

potential glycosylation sites on Fc region, which may reduce Fc receptor binding (see, *e.g.*, Shields R L et al., (2001) J Biol Chem 276: 6591-604).

[000268] In some embodiments, one or more amino in the constant region of a muscle-targeting antibody described herein can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or (e.g., and) reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Pat. No. 6,194,551 (Idusogie et al). In some embodiments, one or more amino acid residues in the N-terminal region of the CH2 domain of an antibody described herein are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in International Publication No. WO 94/29351. In some embodiments, the Fc region of an antibody described herein is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or (e.g., and) to increase the affinity of the antibody for an Fcγ receptor. This approach is described further in International Publication No. WO 00/42072.

[000269] In some embodiments, the heavy and/or (e.g., and) light chain variable domain(s) sequence(s) of the antibodies provided herein can be used to generate, for example, CDR-grafted, chimeric, humanized, or composite human antibodies or antigen-binding fragments, as described elsewhere herein. As understood by one of ordinary skill in the art, any variant, CDR-grafted, chimeric, humanized, or composite antibodies derived from any of the antibodies provided herein may be useful in the compositions and methods described herein and will maintain the ability to specifically bind transferrin receptor, such that the variant, CDR-grafted, chimeric, humanized, or composite antibody has at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95% or more binding to transferrin receptor relative to the original antibody from which it is derived.

[000270] In some embodiments, the antibodies provided herein comprise mutations that confer desirable properties to the antibodies. For example, to avoid potential complications due to Fab-arm exchange, which is known to occur with native IgG4 mAbs, the antibodies provided herein may comprise a stabilizing ‘Adair’ mutation (Angal S., et al., “A single amino acid substitution abolishes the heterogeneity of chimeric mouse/human (IgG4) antibody,” Mol Immunol 30, 105-108; 1993), where serine 228 (EU numbering; residue 241 Kabat numbering) is converted to proline resulting in an IgG1-like hinge sequence. Accordingly, any of the antibodies may include a stabilizing ‘Adair’ mutation.

[000271] As provided herein, antibodies of this disclosure may optionally comprise constant regions or parts thereof. For example, a VL domain may be attached at its C-terminal end to a light chain constant domain like Cκ or Cλ. Similarly, a VH domain or portion thereof may be attached to all or part of a heavy chain like IgA, IgD, IgE, IgG, and IgM, and any isotype

subclass. Antibodies may include suitable constant regions (see, for example, Kabat et al., Sequences of Proteins of Immunological Interest, No. 91-3242, National Institutes of Health Publications, Bethesda, Md. (1991)). Therefore, antibodies within the scope of this may disclosure include VH and VL domains, or an antigen binding portion thereof, combined with any suitable constant regions.

ii. Muscle-Targeting Peptides

[000272] Some aspects of the disclosure provide muscle-targeting peptides as muscle-targeting agents. Short peptide sequences (e.g., peptide sequences of 5-20 amino acids in length) that bind to specific cell types have been described. For example, cell-targeting peptides have been described in Vines e., et al., A. "Cell-penetrating and cell-targeting peptides in drug delivery" *Biochim Biophys Acta* 2008, 1786: 126-38; Jarver P., et al., "In vivo biodistribution and efficacy of peptide mediated delivery" *Trends Pharmacol Sci* 2010; 31: 528-35; Samoylova T.I., et al., "Elucidation of muscle-binding peptides by phage display screening" *Muscle Nerve* 1999; 22: 460-6; U.S. Patent No. 6,329,501, issued on December 11, 2001, entitled "METHODS AND COMPOSITIONS FOR TARGETING COMPOUNDS TO MUSCLE"; and Samoylov A.M., et al., "Recognition of cell-specific binding of phage display derived peptides using an acoustic wave sensor." *Biomol Eng* 2002; 18: 269-72; the entire contents of each of which are incorporated herein by reference. By designing peptides to interact with specific cell surface antigens (e.g., receptors), selectivity for a desired tissue, e.g., muscle, can be achieved. Skeletal muscle-targeting has been investigated and a range of molecular payloads are able to be delivered. These approaches may have high selectivity for muscle tissue without many of the practical disadvantages of a large antibody or viral particle. Accordingly, in some embodiments, the muscle-targeting agent is a muscle-targeting peptide that is from 4 to 50 amino acids in length. In some embodiments, the muscle-targeting peptide is 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 amino acids in length. Muscle-targeting peptides can be generated using any of several methods, such as phage display.

[000273] In some embodiments, a muscle-targeting peptide may bind to an internalizing cell surface receptor that is overexpressed or relatively highly expressed in muscle cells, e.g. a transferrin receptor, compared with certain other cells. In some embodiments, a muscle-targeting peptide may target, e.g., bind to, a transferrin receptor. In some embodiments, a peptide that targets a transferrin receptor may comprise a segment of a naturally occurring ligand, e.g., transferrin. In some embodiments, a peptide that targets a transferrin receptor is as described in US Patent No. 6,743,893, filed 11/30/2000, "RECEPTOR-MEDIATED UPTAKE OF PEPTIDES THAT BIND THE HUMAN TRANSFERRIN RECEPTOR". In some

embodiments, a peptide that targets a transferrin receptor is as described in Kawamoto, M. et al, "A novel transferrin receptor-targeted hybrid peptide disintegrates cancer cell membrane to induce rapid killing of cancer cells." BMC Cancer. 2011 Aug 18;11:359. In some embodiments, a peptide that targets a transferrin receptor is as described in US Patent No. 8,399,653, filed 5/20/2011, "TRANSFERRIN/TRANSFERRIN RECEPTOR-MEDIATED SIRNA DELIVERY".

[000274] As discussed above, examples of muscle targeting peptides have been reported. For example, muscle-specific peptides were identified using phage display library presenting surface heptapeptides. As one example a peptide having the amino acid sequence ASSLNIA (SEQ ID NO: 138) bound to C2C12 murine myotubes *in vitro*, and bound to mouse muscle tissue *in vivo*. Accordingly, in some embodiments, the muscle-targeting agent comprises the amino acid sequence ASSLNIA (SEQ ID NO: 138). This peptide displayed improved specificity for binding to heart and skeletal muscle tissue after intravenous injection in mice with reduced binding to liver, kidney, and brain. Additional muscle-specific peptides have been identified using phage display. For example, a 12 amino acid peptide was identified by phage display library for muscle targeting in the context of treatment for DMD. See, Yoshida D., et al., "Targeting of salicylate to skin and muscle following topical injections in rats." *Int J Pharm* 2002; 231: 177-84; the entire contents of which are hereby incorporated by reference. Here, a 12 amino acid peptide having the sequence SKTFNTHPQSTP (SEQ ID NO: 139) was identified and this muscle-targeting peptide showed improved binding to C2C12 cells relative to the ASSLNIA (SEQ ID NO: 138) peptide.

[000275] An additional method for identifying peptides selective for muscle (*e.g.*, skeletal muscle) over other cell types includes *in vitro* selection, which has been described in Ghosh D., et al., "Selection of muscle-binding peptides from context-specific peptide-presenting phage libraries for adenoviral vector targeting" *J Virol* 2005; 79: 13667-72; the entire contents of which are incorporated herein by reference. By pre-incubating a random 12-mer peptide phage display library with a mixture of non-muscle cell types, non-specific cell binders were selected out. Following rounds of selection the 12 amino acid peptide TARGEHKEEELI (SEQ ID NO: 140) appeared most frequently. Accordingly, in some embodiments, the muscle-targeting agent comprises the amino acid sequence TARGEHKEEELI (SEQ ID NO: 140).

[000276] A muscle-targeting agent may an amino acid-containing molecule or peptide. A muscle-targeting peptide may correspond to a sequence of a protein that preferentially binds to a protein receptor found in muscle cells. In some embodiments, a muscle-targeting peptide contains a high propensity of hydrophobic amino acids, *e.g.* valine, such that the peptide preferentially targets muscle cells. In some embodiments, a muscle-targeting peptide has not

been previously characterized or disclosed. These peptides may be conceived of, produced, synthesized, and/or (e.g., and) derivatized using any of several methodologies, e.g. phage displayed peptide libraries, one-bead one-compound peptide libraries, or positional scanning synthetic peptide combinatorial libraries. Exemplary methodologies have been characterized in the art and are incorporated by reference (Gray, B.P. and Brown, K.C. "Combinatorial Peptide Libraries: Mining for Cell-Binding Peptides" *Chem Rev.* 2014, 114:2, 1020–1081.; Samoylova, T.I. and Smith, B.F. "Elucidation of muscle-binding peptides by phage display screening." *Muscle Nerve*, 1999, 22:4. 460-6.). In some embodiments, a muscle-targeting peptide has been previously disclosed (see, e.g. Writer M.J. et al. "Targeted gene delivery to human airway epithelial cells with synthetic vectors incorporating novel targeting peptides selected by phage display." *J. Drug Targeting*. 2004;12:185; Cai, D. "BDNF-mediated enhancement of inflammation and injury in the aging heart." *Physiol Genomics*. 2006, 24:3, 191-7.; Zhang, L. "Molecular profiling of heart endothelial cells." *Circulation*, 2005, 112:11, 1601-11.; McGuire, M.J. et al. "In vitro selection of a peptide with high selectivity for cardiomyocytes in vivo." *J Mol Biol*. 2004, 342:1, 171-82.). Exemplary muscle-targeting peptides comprise an amino acid sequence of the following group: CQAQGQLVC (SEQ ID NO: 141), CSERSMNFC (SEQ ID NO: 142), CPKTRRVPC (SEQ ID NO: 143), WLSEAGPVVTVRALRGTSW (SEQ ID NO: 144), ASSLNIA (SEQ ID NO: 138), CMQHSMRVC (SEQ ID NO: 145), and DDTRHWG (SEQ ID NO: 146). In some embodiments, a muscle-targeting peptide may comprise about 2-25 amino acids, about 2-20 amino acids, about 2-15 amino acids, about 2-10 amino acids, or about 2-5 amino acids. Muscle-targeting peptides may comprise naturally-occurring amino acids, e.g. cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, a muscle-targeting peptide may be linear; in other embodiments, a muscle-targeting peptide may be cyclic, e.g. bicyclic (see, e.g. Silvana, M.G. et al. *Mol. Therapy*, 2018, 26:1, 132–147.).

iii. Muscle-Targeting Receptor Ligands

[000277] A muscle-targeting agent may be a ligand, e.g. a ligand that binds to a receptor protein. A muscle-targeting ligand may be a protein, e.g. transferrin, which binds to an internalizing cell surface receptor expressed by a muscle cell. Accordingly, in some embodiments, the muscle-targeting agent is transferrin, or a derivative thereof that binds to a transferrin receptor. A muscle-targeting ligand may alternatively be a small molecule, e.g. a lipophilic small molecule that preferentially targets muscle cells relative to other cell types. Exemplary lipophilic small molecules that may target muscle cells include compounds

comprising cholesterol, cholesteryl, stearic acid, palmitic acid, oleic acid, oleyl, linolene, linoleic acid, myristic acid, sterols, dihydrotestosterone, testosterone derivatives, glycerine, alkyl chains, trityl groups, and alkoxy acids.

iv. Muscle-Targeting Aptamers

[000278] A muscle-targeting agent may be an aptamer, e.g. an RNA aptamer, which preferentially targets muscle cells relative to other cell types. In some embodiments, a muscle-targeting aptamer has not been previously characterized or disclosed. These aptamers may be conceived of, produced, synthesized, and/or (e.g., and) derivatized using any of several methodologies, e.g. Systematic Evolution of Ligands by Exponential Enrichment. Exemplary methodologies have been characterized in the art and are incorporated by reference (Yan, A.C. and Levy, M. "Aptamers and aptamer targeted delivery" RNA biology, 2009, 6:3, 316-20.; Germer, K. et al. "RNA aptamers and their therapeutic and diagnostic applications." Int. J. Biochem. Mol. Biol. 2013; 4: 27-40.). In some embodiments, a muscle-targeting aptamer has been previously disclosed (see, e.g. Phillippou, S. et al. "Selection and Identification of Skeletal-Muscle-Targeted RNA Aptamers." Mol Ther Nucleic Acids. 2018, 10:199-214.; Thiel, W.H. et al. "Smooth Muscle Cell-targeted RNA Aptamer Inhibits Neointimal Formation." Mol Ther. 2016, 24:4, 779-87.). Exemplary muscle-targeting aptamers include the A01B RNA aptamer and RNA Apt 14. In some embodiments, an aptamer is a nucleic acid-based aptamer, an oligonucleotide aptamer or a peptide aptamer. In some embodiments, an aptamer may be about 5-15 kDa, about 5-10 kDa, about 10-15 kDa, about 1-5 Da, about 1-3 kDa, or smaller.

v. Other Muscle-Targeting Agents

[000279] One strategy for targeting a muscle cell (e.g., a skeletal muscle cell) is to use a substrate of a muscle transporter protein, such as a transporter protein expressed on the sarcolemma. In some embodiments, the muscle-targeting agent is a substrate of an influx transporter that is specific to muscle tissue. In some embodiments, the influx transporter is specific to skeletal muscle tissue. Two main classes of transporters are expressed on the skeletal muscle sarcolemma, (1) the adenosine triphosphate (ATP) binding cassette (ABC) superfamily, which facilitate efflux from skeletal muscle tissue and (2) the solute carrier (SLC) superfamily, which can facilitate the influx of substrates into skeletal muscle. In some embodiments, the muscle-targeting agent is a substrate that binds to an ABC superfamily or an SLC superfamily of transporters. In some embodiments, the substrate that binds to the ABC or SLC superfamily of transporters is a naturally-occurring substrate. In some embodiments, the substrate that binds to the ABC or SLC superfamily of transporters is a non-naturally occurring substrate, for example, a synthetic derivative thereof that binds to the ABC or SLC superfamily of transporters.

[000280] In some embodiments, the muscle-targeting agent is a substrate of an SLC superfamily of transporters. SLC transporters are either equilibrative or use proton or sodium ion gradients created across the membrane to drive transport of substrates. Exemplary SLC transporters that have high skeletal muscle expression include, without limitation, the SATT transporter (ASCT1; SLC1A4), GLUT4 transporter (SLC2A4), GLUT7 transporter (GLUT7; SLC2A7), ATRC2 transporter (CAT-2; SLC7A2), LAT3 transporter (KIAA0245; SLC7A6), PHT1 transporter (PTR4; SLC15A4), OATP-J transporter (OATP5A1; SLC21A15), OCT3 transporter (EMT; SLC22A3), OCTN2 transporter (FLJ46769; SLC22A5), ENT transporters (ENT1; SLC29A1 and ENT2; SLC29A2), PAT2 transporter (SLC36A2), and SAT2 transporter (KIAA1382; SLC38A2). These transporters can facilitate the influx of substrates into skeletal muscle, providing opportunities for muscle targeting.

[000281] In some embodiments, the muscle-targeting agent is a substrate of an equilibrative nucleoside transporter 2 (ENT2) transporter. Relative to other transporters, ENT2 has one of the highest mRNA expressions in skeletal muscle. While human ENT2 (hENT2) is expressed in most body organs such as brain, heart, placenta, thymus, pancreas, prostate, and kidney, it is especially abundant in skeletal muscle. Human ENT2 facilitates the uptake of its substrates depending on their concentration gradient. ENT2 plays a role in maintaining nucleoside homeostasis by transporting a wide range of purine and pyrimidine nucleobases. The hENT2 transporter has a low affinity for all nucleosides (adenosine, guanosine, uridine, thymidine, and cytidine) except for inosine. Accordingly, in some embodiments, the muscle-targeting agent is an ENT2 substrate. Exemplary ENT2 substrates include, without limitation, inosine, 2',3'-dideoxyinosine, and calofarabine. In some embodiments, any of the muscle-targeting agents provided herein are associated with a molecular payload (e.g., oligonucleotide payload). In some embodiments, the muscle-targeting agent is covalently linked to the molecular payload. In some embodiments, the muscle-targeting agent is non-covalently linked to the molecular payload.

[000282] In some embodiments, the muscle-targeting agent is a substrate of an organic cation/carnitine transporter (OCTN2), which is a sodium ion-dependent, high affinity carnitine transporter. In some embodiments, the muscle-targeting agent is carnitine, mildronate, acetylcarnitine, or any derivative thereof that binds to OCTN2. In some embodiments, the carnitine, mildronate, acetylcarnitine, or derivative thereof is covalently linked to the molecular payload (e.g., oligonucleotide payload).

[000283] A muscle-targeting agent may be a protein that is protein that exists in at least one soluble form that targets muscle cells. In some embodiments, a muscle-targeting protein may be hemojuvelin (also known as repulsive guidance molecule C or hemochromatosis type 2

protein), a protein involved in iron overload and homeostasis. In some embodiments, hemojuvelin may be full length or a fragment, or a mutant with at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% sequence identity to a functional hemojuvelin protein. In some embodiments, a hemojuvelin mutant may be a soluble fragment, may lack a N-terminal signaling, and/or (e.g., and) lack a C-terminal anchoring domain. In some embodiments, hemojuvelin may be annotated under GenBank RefSeq Accession Numbers NM_001316767.1, NM_145277.4, NM_202004.3, NM_213652.3, or NM_213653.3. It should be appreciated that a hemojuvelin may be of human, non-human primate, or rodent origin.

B. Molecular Payloads

[000284] Some aspects of the disclosure provide molecular payloads, e.g., for modulating a biological outcome, e.g., the transcription of a DNA sequence, the expression of a protein, or the activity of a protein. In some embodiments, a molecular payload is linked to, or otherwise associated with a muscle-targeting agent. In some embodiments, such molecular payloads are capable of targeting to a muscle cell, e.g., via specifically binding to a nucleic acid or protein in the muscle cell following delivery to the muscle cell by an associated muscle-targeting agent. It should be appreciated that various types of muscle-targeting agents may be used in accordance with the disclosure. For example, the molecular payload may comprise, or consist of, an oligonucleotide (e.g., antisense oligonucleotide), a peptide (e.g., a peptide that binds a nucleic acid or protein associated with disease in a muscle cell), a protein (e.g., a protein that binds a nucleic acid or protein associated with disease in a muscle cell), or a small molecule (e.g., a small molecule that modulates the function of a nucleic acid or protein associated with disease in a muscle cell). In some embodiments, the molecular payload is an oligonucleotide that comprises a strand having a region of complementarity to a DMPK allele comprising a disease-associated-repeat expansion. Exemplary molecular payloads are described in further detail herein, however, it should be appreciated that the exemplary molecular payloads provided herein are not meant to be limiting.

i. Oligonucleotides

[000285] Any suitable oligonucleotide may be used as a molecular payload, as described herein. In some embodiments, the oligonucleotide may be designed to cause degradation of an mRNA (e.g., the oligonucleotide may be a gapmer, an siRNA, a ribozyme or an aptamer that causes degradation). In some embodiments, the oligonucleotide may be designed to block translation of an mRNA (e.g., the oligonucleotide may be a mixmer, an siRNA or an aptamer that blocks translation). In some embodiments, an oligonucleotide may be designed to caused degradation and block translation of an mRNA. In some embodiments, an oligonucleotide may

be a guide nucleic acid (e.g., guide RNA) for directing activity of an enzyme (e.g., a gene editing enzyme). Other examples of oligonucleotides are provided herein. It should be appreciated that, in some embodiments, oligonucleotides in one format (e.g., antisense oligonucleotides) may be suitably adapted to another format (e.g., siRNA oligonucleotides) by incorporating functional sequences (e.g., antisense strand sequences) from one format to the other format.

[000286] Examples of oligonucleotides useful for targeting DMPK are provided in US Patent Application Publication 20100016215A1, published on January 1, 2010, entitled *Compound And Method For Treating Myotonic Dystrophy*; US Patent Application Publication 20130237585A1, published July 19, 2010, *Modulation Of Dystrophia Myotonica-Protein Kinase (DMPK) Expression*; US Patent Application Publication 20150064181A1, published on March 5, 2015, entitled “*Antisense Conjugates For Decreasing Expression Of Dmpk*”; US Patent Application Publication 20150238627A1, published on August 27, 2015, entitled “*Peptide-Linked Morpholino Antisense Oligonucleotides For Treatment Of Myotonic Dystrophy*”; and US Patent Application Publication 20160304877A1, published on October 20, 2016, entitled “*Compounds And Methods For Modulation Of Dystrophia Myotonica-Protein Kinase (Dmpk) Expression*,” the contents of each of which are incorporated herein in their entireties.

[000287] Examples of oligonucleotides for promoting DMPK gene editing include US Patent Application Publication 20170088819A1, published on March 3, 2017, entitled “*Genetic Correction Of Myotonic Dystrophy Type I*”; and International Patent Application Publication WO18002812A1, published on April 1, 2018, entitled “*Materials And Methods For Treatment Of Myotonic Dystrophy Type 1 (DM1) And Other Related Disorders*,” the contents of each of which are incorporated herein in their entireties.

[000288] In some embodiments, oligonucleotides may have a region of complementarity to a sequence set forth as follows, which is an example human DMPK gene sequence (Gene ID 1760; NM_001081560.2):

```
AGGGGGGCTGGACCAAGGGGTGGGGAGAAGGGGAGGAGGCCTCGGCCGGCCGCAG
AGAGAAGTGGCCAGAGAGGCCAGGGGACAGCCAGGGACAGGCAGACATGCAGCC
AGGGCTCCAGGGCCTGGACAGGGGCTGCCAGGCCCTGTGACAGGAGGACCCCGAG
CCCCCGGCCCGGGAGGGGCCATGGTGTCTGCCTGTCCAACATGTCAGCCGAGGTGC
GGCTGAGGCGGCTCCAGCAGCTGGTGTGTTGGACCCGGGCTTCCTGGGGCTGGAGCCC
CTGCTCGACCTTCTCCTGGGCGTCCACCAGGAGCTGGGCGCCTCCGAAGTGGCCAG
GACAAGTACGTGGCCGACTTCTTGCAGTGGGCGGAGCCCATCGTGGTGAGGCTTAA
GGAGGTCCGACTGCAGAGGGACGACTTCGAGATTCTGAAGGTGATCGGACGCGGG
```

GCGTTCAGCGAGGTAGCGGTAGTGAAGATGAAGCAGACGGGCCAGGTGTATGCCAT
GAAGATCATGAACAAGTGGGACATGCTGAAGAGGGGCGAGGTGTCGTGCTTCCGTG
AGGAGAGGGACGTGTTGGTGAATGGGGACCGGCGGTGGATCACGCAGCTGCACTTC
GCCTTCCAGGATGAGAACTACCTGTACCTGGTCATGGAGTATTACGTGGGCGGGGA
CCTGCTGACACTGCTGAGCAAGTTTGGGGAGCGGATTCCGGCCGAGATGGCGCGCT
TCTACCTGGCGGAGATTGTCATGGCCATAGACTCGGTGCACCGGCTTGGCTACGTGC
ACAGGGACATCAAACCCGACAACATCCTGCTGGACCGCTGTGGCCACATCCGCCTG
GCCGACTTCGGCTCTTGCCTCAAGCTGCGGGCAGATGGAACGGTGCGGTCGCTGGT
GGCTGTGGGCACCCCAGACTACCTGTCCCCCGAGATCCTGCAGGCTGTGGGCGGTG
GGCCTGGGACAGGCAGCTACGGGCCCCGAGTGTGACTGGTGGGCGCTGGGTGTATTC
GCCTATGAAATGTTCTATGGGCAGACGCCCTTCTACGCGGATTCCACGGCGGAGAC
CTATGGCAAGATCGTCCACTACAAGGAGCACCTCTCTCTGCCGCTGGTGGACGAAG
GGGTCCCTGAGGAGGCTCGAGACTTCATTACGCGGTTGCTGTGTCCCCCGGAGACA
CGGCTGGGCGGGGTGGAGCAGGCGACTTCCGGACACATCCCTTCTTCTTTGGCCTC
GACTGGGATGGTCTCCGGGACAGCGTGCCCCCTTTACACCGGATTTCTGAAGGTGC
CACCGACACATGCAACTTCGACTTGGTGGAGGACGGGCTCACTGCCATGGAGACAC
TGTCGGACATTCGGGAAGGTGCGCCGCTAGGGGTCCACCTGCCTTTTGTGGGCTACT
CCTACTCCTGCATGGCCCTCAGGGACAGTGAGGTCCCAGGCCCCACACCCATGGAA
CTGGAGGCCGAGCAGCTGCTTGAGCCACACGTGCAAGCGCCCAGCCTGGAGCCCTC
GGTGTCCCCACAGGATGAAACAGCTGAAGTGGCAGTTCCAGCGGCTGTCCCTGCGG
CAGAGGCTGAGGCCGAGGTGACGCTGCGGGAGCTCCAGGAAGCCCTGGAGGAGGA
GGTGCTCACCCGGCAGAGCCTGAGCCGGGAGATGGAGGCCATCCGCACGGACAAC
CAGAACTTCGCCAGTCAACTACGCGAGGCAGAGGCTCGGAACCGGGACCTAGAGG
CACACGTCCGGCAGTTGCAGGAGCGGATGGAGTTGCTGCAGGCAGAGGGAGCCAC
AGCTGTACGGGGGTCCCCAGTCCCCGGGCCACGGATCCACCTTCCCATCTAGATG
GCCCCCGGCCGTGGCTGTGGGCCAGTGCCCGCTGGTGGGGCCAGGCCCCATGCAC
CGCCGCCACCTGCTGCTCCCTGCCAGGGTCCCTAGGCCTGGCCTATCGGAGGCGCTT
TCCCTGCTCCTGTTCCGCCGTTGTTCTGTCTCGTGCCGCCGCCCTGGGCTGCATTGGGT
TGGTGGCCACGCCGGCCAACTACCCGAGTCTGGCGCCGCCAGGAGCCGCCCGC
GCTCCCTGAACCCTAGAACTGTCTTCGACTCCGGGGCCCCGTTGGAAGACTGAGTGC
CCGGGGCACGGCACAGAAGCCGCGCCACCGCCTGCCAGTTCACAACCGCTCCGAG
CGTGGGTCTCCGCCAGCTCCAGTCCTGTGATCCGGGGCCCCGCCCCCTAGCGGCCGGG
GAGGGAGGGGCCGGGTCCGCGGCCGGCGAACGGGGCTCGAAGGGTCCTTGAGCC
GGGAATGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT
GCTGCTGCTGGGGGGATCACAGACCATTCTTTCTTTTCGGCCAGGCTGAGGCCCTGA

CGTGGATGGGCAAACCTGCAGGCCTGGGAAGGCAGCAAGCCGGGCCGTCCGTGTTCC
ATCCTCCACGCACCCCCACCTATCGTTGGTTCGCAAAGTGCAAAGCTTTCTTGTGCA
TGACGCCCTGCTCTGGGGAGCGTCTGGCGCGATCTCTGCCTGCTTACTCGGGAAATT
TGCTTTTGCCAAACCCGCTTTTTTCGGGGATCCCGCGCCCCCTCCTCACTTGCGCTGC
TCTCGGAGCCCCAGCCGGCTCCGCCCCGCTTCGGCGGTTTGGATATTTATTGACCTCG
TCCTCCGACTCGCTGACAGGCTACAGGACCCCCAACAACCCAATCCACGTTTTGGA
TGCACTGAGACCCCGACATTCCTCGGTATTTATTGTCTGTCCCCACCTAGGACCCCC
ACCCCGACCCCTCGCGAATAAAAGGCCCTCCATCTGCCCAAAGCTCTGGA(SEQ ID
NO: 131).

[000289] In some embodiments, oligonucleotides may have a region of complementarity to a sequence set forth as follows, which is an example mouse DMPK gene sequence (Gene ID 13400; NM_001190490.1).

GAACTGGCCAGAGAGACCCAAGGGATAGTCAGGGACGGGCAGACATGCAGCTAGG
GTTCTGGGGCCTGGACAGGGGCAGCCAGGCCCTGTGACGGGAAGACCCCGAGCTCC
GGCCCGGGGAGGGGCCATGGTGTTCCTGCCCAACATGTCAGCCGAAGTGCGGCTG
AGGCAGCTCCAGCAGCTGGTGCTGGACCCAGGCTTCCTGGGACTGGAGCCCCTGCT
CGACCTTCTCCTGGGCGTCCACCAGGAGCTGGGTGCCTCTCACCTAGCCCAGGACA
AGTATGTGGCCGACTTCTTGCACTGGGTGGAGCCCATTGCAGCAAGGCTTAAGGAG
GTCCGACTGCAGAGGGATGATTTTGAGATTTTGAAGGTGATCGGGCGTGGGGCGTT
CAGCGAGGTAGCGGTGGTGAAGATGAAACAGACGGGCCAAGTGTATGCCATGAAG
ATTATGAATAAGTGGGACATGCTGAAGAGAGGCGAGGTGTCGTGCTTCCGGGAAGA
AAGGGATGTATTAGTGAAAGGGGACCGGCGCTGGATCACACAGCTGCACTTTGCCT
TCCAGGATGAGAATACTACCTGTACCTGGTCATGGAATACTACGTGGGCGGGGACCTG
CTAACGCTGCTGAGCAAGTTTGGGGAGCGGATCCCCGCCGAGATGGCTCGCTTCTA
CCTGGCCGAGATTGTTCATGGCCATAGACTCCGTGCACCGGCTGGGCTACGTGCACA
GGGACATCAAACCAGATAACATTCTGCTGGACCGATGTGGGCACATTCGCCTGGCA
GACTTCGGCTCCTGCCTCAAACCTGCAGCCTGATGGAATGGTGAGGTCGCTGGTGGCT
GTGGGCACCCCGGACTACCTGTCTCCTGAGATTCTGCAGGCCGTTGGTGGAGGGCCT
GGGGCAGGCAGCTACGGGCCAGAGTGTGACTGGTGGGCACTGGGCGTGTTCGCCTA
TGAGATGTTCTATGGGCAGACCCCCTTCTACGCGGACTCCACAGCCGAGACATATG
CCAAGATTGTGCACTACAGGGAACACTTGTGCTGCGCTGCCGCTGGCAGACACAGTTGTC
CCCGAGGAAGCTCAGGACCTCATTCGTGGGCTGCTGTGTCTGCTGAGATAAGGCT
AGGTCGAGGTGGGGCAGACTTCGAGGGTGCCACGGACACATGCAATTTGATGTGG
TGGAGGACCGGCTCACTGCCATGGTGAGCGGGGGCGGGGAGACGCTGTCAGACAT
GCAGGAAGACATGCCCCCTTGGGGTGCGCCTGCCCTTCGTGGGCTACTCCTACTGCTG

CATGGCCTTCAGAGACAATCAGGTCCCGGACCCACCCCTATGGAAGTAGAGGCCC
TGCAGTTGCCTGTGTCAGACTTGCAAGGGCTTGACTTGACGCCCCAGTGTCCCCAC
CGGATCAAGTGGCTGAAGAGGCTGACCTAGTGGCTGTCCCTGCCCCCTGTGGCTGAG
GCAGAGACCACGGTAACGCTGCAGCAGCTCCAGGAAGCCCTGGAAGAAGAGGTTTC
TCACCCGGCAGAGCCTGAGCCGCGAGCTGGAGGCCATCCGGACCGCCAACCAGAAC
TTCTCCAGCCAACTACAGGAGGCCGAGGTCCGAAACCGAGACCTGGAGGGCGCATGT
TCGGCAGCTACAGGAACGGATGGAGATGCTGCAGGCCCCAGGAGCCGCAGCCATC
ACGGGGGTCCCCAGTCCCCGGGCCACGGATCCACCTTCCCATCTAGATGGCCCCC
GGCCGTGGCTGTGGGCCAGTGCCCGCTGGTGGGGCCAGGCCCCATGCACCGCCGTC
ACCTGCTGCTCCCTGCCAGGATCCCTAGGCCTGGCCTATCCGAGGGCGCGTTGCCTGC
TCCTGTTCGCCGCTGCTCTGGCTGCTGCCGCCACACTGGGCTGCACTGGGTTGGTGG
CCTATACCGGCGGTCTCACCCAGTCTGGTGTTCGCGGGAGCCACCTTCGCCCCCT
GAACCCTAAGACTCCAAGCCATCTTTCATTTAGGCCTCCTAGGAAGGTCGAGCGAC
CAGGGAGCGACCCAAAGCGTCTCTGTGCCATCGCGCCCCCCCCCCCCCCCCACCG
CTCCGCTCCACACTTCTGTGAGCCTGGGTCCCCACCCAGCTCCGCTCCTGTGATCCA
GGCCTGCCACCTGGCGGCCGGGAGGGAGGAACAGGGCTCGTGCCAGCACCCCTG
GTTCTGCAGAGCTGGTAGCCACCGCTGCTGCAGCAGCTGGGCATTGCGCCGACCTTG
CTTTACTAGCCCCGACGTGGATGGGCAAAGTCTGCTCAGCTCATCCGATTTCACTTTT
TCACTCTCCCAGCCATCAGTTACAAGCCATAAGCATGAGCCCCCTATTTCCAGGGAC
ATCCCATTTCCCATAGTGATGGATCAGCAAGACCTCTGCCAGCACACACGGAGTCTTT
GGCTTCGGACAGCCTCACTCCTGGGGGTTGCTGCAACTCCTTCCCCGTGTACACGTC
TGCACTCTAACAACGGAGCCACAGCTGCACTCCCCCTCCCCCAAAGCAGTGTGGG
TATTTATTGATCTTGTTATCTGACTCACTGACAGACTCCGGGACCCACGTTTTAGAT
GCATTGAGACTCGACATTCCTCGGTATTTATTGTCTGTCCCCACCTACGACCTCCACT
CCCGACCCTTGCGAATAAAATACTTCTGGTCTGCCCTAAA (SEQ ID NO: 147). In

some embodiments, an oligonucleotide may have a region of complementarity to DMPK gene sequences of multiple species, e.g., selected from human, mouse and non-human species.

[000290] In some embodiments, the oligonucleotide may have region of complementarity to a mutant form of DMPK, for example, a mutant form as reported in Botta A. et al. "The CTG repeat expansion size correlates with the splicing defects observed in muscles from myotonic dystrophy type 1 patients." J Med Genet. 2008 Oct;45(10):639-46.; and Machuca-Tzili L. et al. "Clinical and molecular aspects of the myotonic dystrophies: a review." Muscle Nerve. 2005 Jul;32(1):1-18.; the contents of each of which are incorporated herein by reference in their entireties.

[000291] In some embodiments, the oligonucleotide may target lncRNA or mRNA, *e.g.*, for degradation. In some embodiments, the oligonucleotide may target, *e.g.*, for degradation, a nucleic acid encoding a protein involved in a mismatch repair pathway, *e.g.*, MSH2, MutLalpha, MutSbeta, MutLalpha. Non-limiting examples of proteins involved in mismatch repair pathways, for which mRNAs encoding such proteins may be targeted by oligonucleotides described herein, are described in Iyer, R.R. et al., “*DNA triplet repeat expansion and mismatch repair*” *Annu Rev Biochem.* 2015;84:199-226.; and Schmidt M.H. and Pearson C.E., “Disease-associated repeat instability and mismatch repair” *DNA Repair (Amst).* 2016 Feb;38:117-26.

[000292] In some embodiments, an oligonucleotide provided herein is an antisense oligonucleotide targeting DMPK. In some embodiments, the oligonucleotide targeting is any one of the antisense oligonucleotides (*e.g.*, a Gapmer) targeting DMPK as described in US Patent Application Publication US20160304877A1, published on October 20, 2016, entitled “Compounds And Methods For Modulation Of Dystrophia Myotonica-Protein Kinase (DMPK) Expression,” incorporated herein by reference). In some embodiments, the DMPK targeting oligonucleotide targets a region of the DMPK gene sequence as set forth in Genbank accession No. NM_001081560.2 (SEQ ID NO: 131) or as set forth in Genbank accession No. NG_009784.1.

[000293] In some embodiments, the DMPK targeting oligonucleotide comprises a nucleotide sequence comprising a region complementary to a target region that is at least 10 continuous nucleotides (*e.g.*, at least 10, at least 12, at least 14, at least 16, or more continuous nucleotides) in SEQ ID NO: 131.

[000294] In some embodiments, the DMPK targeting oligonucleotide comprise a gapmer motif. “Gapmer” means a chimeric antisense compound in which an internal region having a plurality of nucleotides that support RNase H cleavage is positioned between external regions having one or more nucleotides, wherein the nucleotides comprising the internal region are chemically distinct from the nucleotide or nucleotides comprising the external regions. The internal region can be referred to as a “gap segment” and the external regions can be referred to as “wing segments.” In some embodiments, the DMPK targeting oligonucleotide comprises one or more modified nucleotides, and/or (*e.g.*, and) one or more modified internucleotide linkages. In some embodiments, the internucleotide linkage is a phosphorothioate linkage. In some embodiments, the oligonucleotide comprises a full phosphorothioate backbone. In some embodiments, the oligonucleotide is a DNA gapmer with cET ends (*e.g.*, 3-10-3; cET-DNA-cET). In some embodiments, the DMPK targeting oligonucleotide comprises one or more 6′-(S)-CH₃ biocyclic nucleotides, one or more β-D-2′-deoxyribonucleotides, and/or (*e.g.*, and) one or more 5-methylcytosine nucleotides.

[000295] In some embodiments, the DMPK targeting oligonucleotide is a gapmer having the formula 5'-X-Y-Z-3', with X and Z as wing segments and Y as the gap segment. In some embodiments, the DMPK targeting oligonucleotide is a gapmer having a 5'-4-8-4-3' formula. In some embodiments, the DMPK targeting oligonucleotide is a gapmer having a 5'-5-10-5-3' formula. In some embodiments, the DMPK targeting oligonucleotide is a gapmer having a 5'-3-10-3-3' formula. In some embodiments, the DMPK targeting oligonucleotide is a gapmer comprising one or more of 5-methylcytosine nucleotides, 2'OMe nucleotides, 2'fluoro nucleotides, LNAs, and/or (e.g., and) 2'-O-methoxyethyl (2'-O-MOE) nucleotides. In some embodiments, the DMPK targeting oligonucleotide is a gapmer comprising one or more modified internucleotide (e.g., a phosphorothioate linkage). In some embodiments, the DMPK targeting oligonucleotide is a gapmer comprising a full phosphorothioate backbone.

[000296] In some embodiments, any one of the oligonucleotides can be in salt form, e.g., as sodium, potassium, or magnesium salts.

[000297] In some embodiments, the 5' or 3' nucleoside (e.g., terminal nucleoside) of any one of the oligonucleotides described herein is conjugated to an amine group, optionally via a spacer. In some embodiments, the spacer comprises an aliphatic moiety. In some embodiments, the spacer comprises a polyethylene glycol moiety. In some embodiments, a phosphodiester linkage is present between the spacer and the 5' or 3' nucleoside of the oligonucleotide. In some embodiments, the 5' or 3' nucleoside (e.g., terminal nucleoside) of any of the oligonucleotides described herein is conjugated to a spacer that is a substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted carbocyclylene, substituted or unsubstituted heterocyclylene, substituted or unsubstituted arylene, substituted or unsubstituted heteroarylene, -O-, -N(R^A)-, -S-, -C(=O)-, -C(=O)O-, -C(=O)NR^A-, -NR^AC(=O)-, -NR^AC(=O)R^A-, -C(=O)R^A-, -NR^AC(=O)O-, -NR^AC(=O)N(R^A)-, -OC(=O)-, -OC(=O)O-, -OC(=O)N(R^A)-, -S(O)₂NR^A-, -NR^AS(O)₂-, or a combination thereof; each R^A is independently hydrogen or substituted or unsubstituted alkyl. In certain embodiments, the spacer is a substituted or unsubstituted alkylene, substituted or unsubstituted heterocyclylene, substituted or unsubstituted heteroarylene, -O-, -N(R^A)-, or -C(=O)N(R^A)₂, or a combination thereof.

[000298] In some embodiments, the 5' or 3' nucleoside of any one of the oligonucleotides described herein is conjugated to a compound of the formula -NH₂-(CH₂)_n-, wherein n is an integer from 1 to 12. In some embodiments, n is 6, 7, 8, 9, 10, 11, or 12. In some embodiments, a phosphodiester linkage is present between the compound of the formula NH₂-(CH₂)_n- and the 5' or 3' nucleoside of the oligonucleotide. In some embodiments, a compound of the formula NH₂-(CH₂)₆- is conjugated to the oligonucleotide via a reaction between 6-amino-1-hexanol (NH₂-(CH₂)₆-OH) and the 5' phosphate of the oligonucleotide.

[000299] In some embodiments, the oligonucleotide is conjugated to a targeting agent, e.g., a muscle targeting agent such as an anti-TfR antibody, e.g., via the amine group.

a. Oligonucleotide Size/Sequence

[000300] Oligonucleotides may be of a variety of different lengths, e.g., depending on the format. In some embodiments, an oligonucleotide is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 75, or more nucleotides in length. In some embodiments, the oligonucleotide is 8 to 50 nucleotides in length, 8 to 40 nucleotides in length, 8 to 30 nucleotides in length, 10 to 15 nucleotides in length, 10 to 20 nucleotides in length, 15 to 25 nucleotides in length, 21 to 23 nucleotides in lengths, etc.

[000301] In some embodiments, a complementary nucleic acid sequence of an oligonucleotide for purposes of the present disclosure is specifically hybridizable or specific for the target nucleic acid when binding of the sequence to the target molecule (e.g., mRNA) interferes with the normal function of the target (e.g., mRNA) to cause a loss of activity (e.g., inhibiting translation) or expression (e.g., degrading a target mRNA) and there is a sufficient degree of complementarity to avoid non-specific binding of the sequence to non-target sequences under conditions in which avoidance of non-specific binding is desired, e.g., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed under suitable conditions of stringency. Thus, in some embodiments, an oligonucleotide may be at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% complementary to the consecutive nucleotides of a target nucleic acid. In some embodiments a complementary nucleotide sequence need not be 100% complementary to that of its target to be specifically hybridizable or specific for a target nucleic acid.

[000302] In some embodiments, an oligonucleotide comprises region of complementarity to a target nucleic acid that is in the range of 8 to 15, 8 to 30, 8 to 40, or 10 to 50, or 5 to 50, or 5 to 40 nucleotides in length. In some embodiments, a region of complementarity of an oligonucleotide to a target nucleic acid is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length. In some embodiments, the region of complementarity is complementary with at least 8 consecutive nucleotides of a target nucleic acid. In some embodiments, an oligonucleotide may contain 1, 2 or 3 base mismatches compared to the portion of the consecutive nucleotides of target nucleic acid. In some embodiments the

oligonucleotide may have up to 3 mismatches over 15 bases, or up to 2 mismatches over 10 bases.

[000303] In some embodiments, an oligonucleotide comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides of a sequence comprising any one of SEQ ID NO: 148-383 and 621-638. In some embodiments, an oligonucleotide comprises a sequence comprising any one of SEQ ID NO: 148-383 and 621-638. In some embodiments, an oligonucleotide comprises a sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, or 97% sequence identity with at least 12 or at least 15 consecutive nucleotides of any one of SEQ ID NO: 148-383 and 621-638.

[000304] In some embodiments, an oligonucleotide comprises a sequence that targets a DMPK sequence comprising any one of SEQ ID NO: 384-619. In some embodiments, an oligonucleotide comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides (*e.g.*, consecutive nucleotides) that are complementary to a DMPK sequence comprising any one of SEQ ID NO: 384-619. In some embodiments, an oligonucleotide comprises a sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, or 97% complementary with at least 12 or at least 15 consecutive nucleotides of any one of SEQ ID NO: 384-619.

[000305] In some embodiments, the oligonucleotide is complementary (*e.g.*, at least 85% at least 90%, at least 95%, or 100%) to a target sequence of any one of the oligonucleotides provided herein (*e.g.*, the oligonucleotides listed in Table 8 or Table 17). In some embodiments, such target sequence is 100% complementary to the oligonucleotide listed in Table 8 or Table 17.

[000306] In some embodiments, any one or more of the thymine bases (T's) in any one of the oligonucleotides provided herein (*e.g.*, the oligonucleotides listed in Table 8 or Table 17) may optionally be uracil bases (U's), and/or any one or more of the U's may optionally be T's.

b. Oligonucleotide Modifications:

[000307] The oligonucleotides described herein may be modified, *e.g.*, comprise a modified sugar moiety, a modified internucleoside linkage, a modified nucleotide and/or (*e.g.*, and) combinations thereof. In addition, in some embodiments, oligonucleotides may exhibit one or more of the following properties: do not mediate alternative splicing; are not immune stimulatory; are nuclease resistant; have improved cell uptake compared to unmodified oligonucleotides; are not toxic to cells or mammals; have improved endosomal exit internally in a cell; minimizes TLR stimulation; or avoid pattern recognition receptors. Any of the modified chemistries or formats of oligonucleotides described herein can be combined with each other. For example, one, two, three, four, five, or more different types of modifications can be included within the same oligonucleotide.

[000308] In some embodiments, certain nucleotide modifications may be used that make an oligonucleotide into which they are incorporated more resistant to nuclease digestion than the native oligodeoxynucleotide or oligoribonucleotide molecules; these modified oligonucleotides survive intact for a longer time than unmodified oligonucleotides. Specific examples of modified oligonucleotides include those comprising modified backbones, for example, modified internucleoside linkages such as phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Accordingly, oligonucleotides of the disclosure can be stabilized against nucleolytic degradation such as by the incorporation of a modification, e.g., a nucleotide modification.

[000309] In some embodiments, an oligonucleotide may be of up to 50 or up to 100 nucleotides in length in which 2 to 10, 2 to 15, 2 to 16, 2 to 17, 2 to 18, 2 to 19, 2 to 20, 2 to 25, 2 to 30, 2 to 40, 2 to 45, or more nucleotides of the oligonucleotide are modified nucleotides. The oligonucleotide may be of 8 to 30 nucleotides in length in which 2 to 10, 2 to 15, 2 to 16, 2 to 17, 2 to 18, 2 to 19, 2 to 20, 2 to 25, 2 to 30 nucleotides of the oligonucleotide are modified nucleotides. The oligonucleotide may be of 8 to 15 nucleotides in length in which 2 to 4, 2 to 5, 2 to 6, 2 to 7, 2 to 8, 2 to 9, 2 to 10, 2 to 11, 2 to 12, 2 to 13, 2 to 14 nucleotides of the oligonucleotide are modified nucleotides. Optionally, the oligonucleotides may have every nucleotide except 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides modified. Oligonucleotide modifications are described further herein.

c. Modified Nucleosides

[0001] In some embodiments, the oligonucleotide described herein comprises at least one nucleoside modified at the 2' position of the sugar. In some embodiments, an oligonucleotide comprises at least one 2'-modified nucleoside. In some embodiments, all of the nucleosides in the oligonucleotide are 2'-modified nucleosides.

[0002] In some embodiments, the oligonucleotide described herein comprises one or more non-bicyclic 2'-modified nucleosides, e.g., 2'-deoxy, 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified nucleoside.

[0003] In some embodiments, the oligonucleotide described herein comprises one or more 2'-4' bicyclic nucleosides in which the ribose ring comprises a bridge moiety connecting two atoms in the ring, e.g., connecting the 2'-O atom to the 4'-C atom via a methylene (LNA) bridge, an ethylene (ENA) bridge, or a (S)-constrained ethyl (cEt) bridge. Examples of LNAs are described in International Patent Application Publication WO/2008/043753, published on April

17, 2008, and entitled “*RNA Antagonist Compounds For The Modulation Of PCSK9*”, the contents of which are incorporated herein by reference in its entirety. Examples of ENAs are provided in International Patent Publication No. WO 2005/042777, published on May 12, 2005, and entitled “*APP/ENA Antisense*”; Morita et al., *Nucleic Acid Res.*, Suppl 1:241-242, 2001; Surono et al., *Hum. Gene Ther.*, 15:749-757, 2004; Koizumi, *Curr. Opin. Mol. Ther.*, 8:144-149, 2006 and Horie et al., *Nucleic Acids Symp. Ser (Oxf)*, 49:171-172, 2005; the disclosures of which are incorporated herein by reference in their entireties. Examples of cEt are provided in US Patents 7,101,993; 7,399,845 and 7,569,686, each of which is herein incorporated by reference in its entirety.

[0004] In some embodiments, the oligonucleotide comprises a modified nucleoside disclosed in one of the following United States Patent or Patent Application Publications: US Patent 7,399,845, issued on July 15, 2008, and entitled “*6-Modified Bicyclic Nucleic Acid Analogs*”; US Patent 7,741,457, issued on June 22, 2010, and entitled “*6-Modified Bicyclic Nucleic Acid Analogs*”; US Patent 8,022,193, issued on September 20, 2011, and entitled “*6-Modified Bicyclic Nucleic Acid Analogs*”; US Patent 7,569,686, issued on August 4, 2009, and entitled “*Compounds And Methods For Synthesis Of Bicyclic Nucleic Acid Analogs*”; US Patent 7,335,765, issued on February 26, 2008, and entitled “*Novel Nucleoside And Oligonucleotide Analogues*”; US Patent 7,314,923, issued on January 1, 2008, and entitled “*Novel Nucleoside And Oligonucleotide Analogues*”; US Patent 7,816,333, issued on October 19, 2010, and entitled “*Oligonucleotide Analogues And Methods Utilizing The Same*” and US Publication Number 2011/0009471 now US Patent 8,957,201, issued on February 17, 2015, and entitled “*Oligonucleotide Analogues And Methods Utilizing The Same*”, the entire contents of each of which are incorporated herein by reference for all purposes.

[0005] In some embodiments, the oligonucleotide comprises at least one modified nucleoside that results in an increase in T_m of the oligonucleotide in a range of 1 °C, 2 °C, 3 °C, 4 °C, or 5 °C compared with an oligonucleotide that does not have the at least one modified nucleoside. The oligonucleotide may have a plurality of modified nucleosides that result in a total increase in T_m of the oligonucleotide in a range of 2 °C, 3 °C, 4 °C, 5 °C, 6 °C, 7 °C, 8 °C, 9 °C, 10 °C, 15 °C, 20 °C, 25 °C, 30 °C, 35 °C, 40 °C, 45 °C or more compared with an oligonucleotide that does not have the modified nucleoside.

[0006] The oligonucleotide may comprise a mix of nucleosides of different kinds. For example, an oligonucleotide may comprise a mix of 2'-deoxyribonucleosides or ribonucleosides and 2'-fluoro modified nucleosides. An oligonucleotide may comprise a mix of deoxyribonucleosides or ribonucleosides and 2'-O-Me modified nucleosides. An oligonucleotide may comprise a mix of 2'-fluoro modified nucleosides and 2'-O-Me modified nucleosides. An oligonucleotide may

comprise a mix of 2'-4' bicyclic nucleosides and 2'-MOE, 2'-fluoro, or 2'-O-Me modified nucleosides. An oligonucleotide may comprise a mix of non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE, 2'-fluoro, or 2'-O-Me) and 2'-4' bicyclic nucleosides (e.g., LNA, ENA, cEt).

[0007] The oligonucleotide may comprise alternating nucleosides of different kinds. For example, an oligonucleotide may comprise alternating 2'-deoxyribonucleosides or ribonucleosides and 2'-fluoro modified nucleosides. An oligonucleotide may comprise alternating deoxyribonucleosides or ribonucleosides and 2'-O-Me modified nucleosides. An oligonucleotide may comprise alternating 2'-fluoro modified nucleosides and 2'-O-Me modified nucleosides. An oligonucleotide may comprise alternating 2'-4' bicyclic nucleosides and 2'-MOE, 2'-fluoro, or 2'-O-Me modified nucleosides. An oligonucleotide may comprise alternating non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE, 2'-fluoro, or 2'-O-Me) and 2'-4' bicyclic nucleosides (e.g., LNA, ENA, cEt).

[0008] In some embodiments, an oligonucleotide described herein comprises a 5'-vinylphosphonate modification, one or more abasic residues, and/or one or more inverted abasic residues.

d. Internucleoside Linkages / Backbones

[0009] In some embodiments, oligonucleotide may contain a phosphorothioate or other modified internucleoside linkage. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages between at least two nucleotides. In some embodiments, the oligonucleotide comprises phosphorothioate internucleoside linkages between all nucleotides. For example, in some embodiments, oligonucleotides comprise modified internucleoside linkages at the first, second, and/or (e.g., and) third internucleoside linkage at the 5' or 3' end of the nucleotide sequence.

[00010] Phosphorus-containing linkages that may be used include, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'; see US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302;

5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455, 233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563, 253; 5,571,799; 5,587,361; and 5,625,050.

[00011] In some embodiments, oligonucleotides may have heteroatom backbones, such as methylene(methylimino) or MMI backbones; amide backbones (see De Mesmaeker et al. *Ace. Chem. Res.* 1995, 28:366-374); morpholino backbones (see Summerton and Weller, U.S. Pat. No. 5,034,506); or peptide nucleic acid (PNA) backbones (wherein the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone, see Nielsen et al., *Science* 1991, 254, 1497).

e. Stereospecific Oligonucleotides

[000310] In some embodiments, internucleotidic phosphorus atoms of oligonucleotides are chiral, and the properties of the oligonucleotides by adjusted based on the configuration of the chiral phosphorus atoms. In some embodiments, appropriate methods may be used to synthesize P-chiral oligonucleotide analogs in a stereocontrolled manner (e.g., as described in Oka N, Wada T, Stereocontrolled synthesis of oligonucleotide analogs containing chiral internucleotidic phosphorus atoms. *Chem Soc Rev.* 2011 Dec;40(12):5829-43.) In some embodiments, phosphorothioate containing oligonucleotides comprise nucleoside units that are joined together by either substantially all Sp or substantially all Rp phosphorothioate intersugar linkages are provided. In some embodiments, such phosphorothioate oligonucleotides having substantially chirally pure intersugar linkages are prepared by enzymatic or chemical synthesis, as described, for example, in US Patent 5,587,261, issued on December 12, 1996, the contents of which are incorporated herein by reference in their entirety. In some embodiments, chirally controlled oligonucleotides provide selective cleavage patterns of a target nucleic acid. For example, in some embodiments, a chirally controlled oligonucleotide provides single site cleavage within a complementary sequence of a nucleic acid, as described, for example, in US Patent Application Publication 20170037399 A1, published on February 2, 2017, entitled "CHIRAL DESIGN", the contents of which are incorporated herein by reference in their entirety.

f. Morpholinos

[000311] In some embodiments, the oligonucleotide may be a morpholino-based compounds. Morpholino-based oligomeric compounds are described in Dwaine A. Braasch and David R. Corey, *Biochemistry*, 2002, 41(14), 4503-4510); Genesis, volume 30, issue 3, 2001; Heasman, J., *Dev. Biol.*, 2002, 243, 209-214; Nasevicius et al., *Nat. Genet.*, 2000, 26, 216-220; Lacerra et al., *Proc. Natl. Acad. Sci.*, 2000, 97, 9591-9596; and U.S. Pat. No. 5,034,506, issued

Jul. 23, 1991. In some embodiments, the morpholino-based oligomeric compound is a phosphorodiamidate morpholino oligomer (PMO) (e.g., as described in Iverson, *Curr. Opin. Mol. Ther.*, 3:235-238, 2001; and Wang et al., *J. Gene Med.*, 12:354-364, 2010; the disclosures of which are incorporated herein by reference in their entireties).

g. Peptide Nucleic Acids (PNAs)

[000312] In some embodiments, both a sugar and an internucleoside linkage (the backbone) of the nucleotide units of an oligonucleotide are replaced with novel groups. In some embodiments, the base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, for example, an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative publication that report the preparation of PNA compounds include, but are not limited to, US patent nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen *et al.*, *Science*, 1991, 254, 1497-1500.

h. Gapmers

[000313] In some embodiments, an oligonucleotide described herein is a gapmer. A gapmer oligonucleotide generally has the formula 5'-X-Y-Z-3', with X and Z as flanking regions around a gap region Y. In some embodiments, flanking region X of formula 5'-X-Y-Z-3' is also referred to as X region, flanking sequence X, 5' wing region X, or 5' wing segment. In some embodiments, flanking region Z of formula 5'-X-Y-Z-3' is also referred to as Z region, flanking sequence Z, 3' wing region Z, or 3' wing segment. In some embodiments, gap region Y of formula 5'-X-Y-Z-3' is also referred to as Y region, Y segment, or gap-segment Y. In some embodiments, each nucleoside in the gap region Y is a 2'-deoxyribonucleoside, and neither the 5' wing region X or the 3' wing region Z contains any 2'-deoxyribonucleosides.

[000314] In some embodiments, the Y region is a contiguous stretch of nucleotides, e.g., a region of 6 or more DNA nucleotides, which are capable of recruiting an RNase, such as RNase H. In some embodiments, the gapmer binds to the target nucleic acid, at which point an RNase is recruited and can then cleave the target nucleic acid. In some embodiments, the Y region is flanked both 5' and 3' by regions X and Z comprising high-affinity modified nucleosides, e.g., one to six high-affinity modified nucleosides. Examples of high affinity modified nucleosides include, but are not limited to, 2'-modified nucleosides (e.g., 2'-MOE, 2'-O-Me, 2'-F) or 2'-4' bicyclic nucleosides (e.g., LNA, cEt, ENA). In some embodiments, the

flanking sequences X and Z may be of 1-20 nucleotides, 1-8 nucleotides, or 1-5 nucleotides in length. The flanking sequences X and Z may be of similar length or of dissimilar lengths. In some embodiments, the gap-segment Y may be a nucleotide sequence of 5-20 nucleotides, 5-15 twelve nucleotides, or 6-10 nucleotides in length.

[00012] In some embodiments, the gap region of the gapmer oligonucleotides may contain modified nucleotides known to be acceptable for efficient RNase H action in addition to DNA nucleotides, such as C4'-substituted nucleotides, acyclic nucleotides, and arabino-configured nucleotides. In some embodiments, the gap region comprises one or more unmodified internucleoside linkages. In some embodiments, one or both flanking regions each independently comprise one or more phosphorothioate internucleoside linkages (e.g., phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleotides. In some embodiments, the gap region and two flanking regions each independently comprise modified internucleoside linkages (e.g., phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleotides.

[000315] A gapmer may be produced using appropriate methods. Representative U.S. patents, U.S. patent publications, and PCT publications that teach the preparation of gapmers include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; 5,700,922; 5,898,031; 7,015,315; 7,101,993; 7,399,845; 7,432,250; 7,569,686; 7,683,036; 7,750,131; 8,580,756; 9,045,754; 9,428,534; 9,695,418; 10,017,764; 10,260,069; 9,428,534; 8,580,756; U.S. patent publication Nos. US20050074801, US20090221685; US20090286969, US20100197762, and US20110112170; PCT publication Nos. WO2004069991; WO2005023825; WO2008049085 and WO2009090182; and EP Patent No. EP2,149,605, each of which is herein incorporated by reference in its entirety.

[000316] In some embodiments, a gapmer is 10-40 nucleosides in length. For example, a gapmer may be 10-40, 10-35, 10-30, 10-25, 10-20, 10-15, 15-40, 15-35, 15-30, 15-25, 15-20, 20-40, 20-35, 20-30, 20-25, 25-40, 25-35, 25-30, 30-40, 30-35, or 35-40 nucleosides in length. In some embodiments, a gapmer is 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleosides in length.

[000317] In some embodiments, the gap region Y in a gapmer is 5-20 nucleosides in length. For example, the gap region Y may be 5-20, 5-15, 5-10, 10-20, 10-15, or 15-20 nucleosides in length. In some embodiments, the gap region Y is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleosides in length. In some embodiments, each nucleoside in the gap region Y is a 2'-deoxyribonucleoside. In some embodiments, all nucleosides in the gap

region Y are 2'-deoxyribonucleosides. In some embodiments, one or more of the nucleosides in the gap region Y is a modified nucleoside (e.g., a 2' modified nucleoside such as those described herein). In some embodiments, one or more cytosines in the gap region Y are optionally 5-methyl-cytosines. In some embodiments, each cytosine in the gap region Y is a 5-methyl-cytosines.

[000318] In some embodiments, the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of a gapmer (Z in the 5'-X-Y-Z-3' formula) are independently 1-20 nucleosides long. For example, the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) may be independently 1-20, 1-15, 1-10, 1-7, 1-5, 1-3, 1-2, 2-5, 2-7, 3-5, 3-7, 5-20, 5-15, 5-10, 10-20, 10-15, or 15-20 nucleosides long. In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) are independently 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleosides long. In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) are of the same length. In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) are of different lengths. In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) is longer than the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula). In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) is shorter than the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula).

[000319] In some embodiments, a gapmer comprises a 5'-X-Y-Z-3' of 5-10-5, 4-12-4, 3-14-3, 2-16-2, 1-18-1, 3-10-3, 2-10-2, 1-10-1, 2-8-2, 4-6-4, 3-6-3, 2-6-2, 4-7-4, 3-7-3, 2-7-2, 4-8-4, 3-8-3, 2-8-2, 1-8-1, 2-9-2, 1-9-1, 2-10-2, 1-10-1, 1-12-1, 1-16-1, 2-15-1, 1-15-2, 1-14-3, 3-14-1, 2-14-2, 1-13-4, 4-13-1, 2-13-3, 3-13-2, 1-12-5, 5-12-1, 2-12-4, 4-12-2, 3-12-3, 1-11-6, 6-11-1, 2-11-5, 5-11-2, 3-11-4, 4-11-3, 1-17-1, 2-16-1, 1-16-2, 1-15-3, 3-15-1, 2-15-2, 1-14-4, 4-14-1, 2-14-3, 3-14-2, 1-13-5, 5-13-1, 2-13-4, 4-13-2, 3-13-3, 1-12-6, 6-12-1, 2-12-5, 5-12-2, 3-12-4, 4-12-3, 1-11-7, 7-11-1, 2-11-6, 6-11-2, 3-11-5, 5-11-3, 4-11-4, 1-18-1, 1-17-2, 2-17-1, 1-16-3, 1-16-3, 2-16-2, 1-15-4, 4-15-1, 2-15-3, 3-15-2, 1-14-5, 5-14-1, 2-14-4, 4-14-2, 3-14-3, 1-13-6, 6-13-1, 2-13-5, 5-13-2, 3-13-4, 4-13-3, 1-12-7, 7-12-1, 2-12-6, 6-12-2, 3-12-5, 5-12-3, 1-11-8, 8-11-1, 2-11-7, 7-11-2, 3-11-6, 6-11-3, 4-11-5, 5-11-4, 1-18-1, 1-17-2, 2-17-1, 1-16-3, 3-16-1, 2-16-2, 1-15-4, 4-15-1, 2-15-3, 3-15-2, 1-14-5, 2-14-4, 4-14-2, 3-14-3, 1-13-6, 6-13-1, 2-13-5, 5-13-2, 3-13-4, 4-13-3, 1-12-7, 7-12-1, 2-12-6, 6-12-2, 3-12-5, 5-12-3, 1-11-8, 8-11-1, 2-11-7, 7-11-2, 3-11-6, 6-11-3, 4-11-5, 5-11-4, 1-19-1, 1-18-2, 2-18-1, 1-17-3, 3-17-1, 2-17-2, 1-16-4, 4-16-1, 2-16-3, 3-16-2, 1-15-5, 2-15-4, 4-15-2, 3-15-3, 1-14-6, 6-14-1, 2-14-5, 5-14-2, 3-14-4,

4-14-3, 1-13-7, 7-13-1, 2-13-6, 6-13-2, 3-13-5, 5-13-3, 4-13-4, 1-12-8, 8-12-1, 2-12-7, 7-12-2, 3-12-6, 6-12-3, 4-12-5, 5-12-4, 2-11-8, 8-11-2, 3-11-7, 7-11-3, 4-11-6, 6-11-4, 5-11-5, 1-20-1, 1-19-2, 2-19-1, 1-18-3, 3-18-1, 2-18-2, 1-17-4, 4-17-1, 2-17-3, 3-17-2, 1-16-5, 2-16-4, 4-16-2, 3-16-3, 1-15-6, 6-15-1, 2-15-5, 5-15-2, 3-15-4, 4-15-3, 1-14-7, 7-14-1, 2-14-6, 6-14-2, 3-14-5, 5-14-3, 4-14-4, 1-13-8, 8-13-1, 2-13-7, 7-13-2, 3-13-6, 6-13-3, 4-13-5, 5-13-4, 2-12-8, 8-12-2, 3-12-7, 7-12-3, 4-12-6, 6-12-4, 5-12-5, 3-11-8, 8-11-3, 4-11-7, 7-11-4, 5-11-6, 6-11-5, 1-21-1, 1-20-2, 2-20-1, 1-20-3, 3-19-1, 2-19-2, 1-18-4, 4-18-1, 2-18-3, 3-18-2, 1-17-5, 2-17-4, 4-17-2, 3-17-3, 1-16-6, 6-16-1, 2-16-5, 5-16-2, 3-16-4, 4-16-3, 1-15-7, 7-15-1, 2-15-6, 6-15-2, 3-15-5, 5-15-3, 4-15-4, 1-14-8, 8-14-1, 2-14-7, 7-14-2, 3-14-6, 6-14-3, 4-14-5, 5-14-4, 2-13-8, 8-13-2, 3-13-7, 7-13-3, 4-13-6, 6-13-4, 5-13-5, 1-12-10, 10-12-1, 2-12-9, 9-12-2, 3-12-8, 8-12-3, 4-12-7, 7-12-4, 5-12-6, 6-12-5, 4-11-8, 8-11-4, 5-11-7, 7-11-5, 6-11-6, 1-22-1, 1-21-2, 2-21-1, 1-21-3, 3-20-1, 2-20-2, 1-19-4, 4-19-1, 2-19-3, 3-19-2, 1-18-5, 2-18-4, 4-18-2, 3-18-3, 1-17-6, 6-17-1, 2-17-5, 5-17-2, 3-17-4, 4-17-3, 1-16-7, 7-16-1, 2-16-6, 6-16-2, 3-16-5, 5-16-3, 4-16-4, 1-15-8, 8-15-1, 2-15-7, 7-15-2, 3-15-6, 6-15-3, 4-15-5, 5-15-4, 2-14-8, 8-14-2, 3-14-7, 7-14-3, 4-14-6, 6-14-4, 5-14-5, 3-13-8, 8-13-3, 4-13-7, 7-13-4, 5-13-6, 6-13-5, 4-12-8, 8-12-4, 5-12-7, 7-12-5, 6-12-6, 5-11-8, 8-11-5, 6-11-7, or 7-11-6. The numbers indicate the number of nucleosides in X, Y, and Z regions in the 5'-X-Y-Z-3' gapmer.

[000320] In some embodiments, one or more nucleosides in the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) or the 3' wing region of a gapmer (Z in the 5'-X-Y-Z-3' formula) are modified nucleotides (e.g., high-affinity modified nucleosides). In some embodiments, the modified nucleoside (e.g., high-affinity modified nucleosides) is a 2'-modified nucleoside. In some embodiments, the 2'-modified nucleoside is a 2'-4' bicyclic nucleoside or a non-bicyclic 2'-modified nucleoside. In some embodiments, the high-affinity modified nucleoside is a 2'-4' bicyclic nucleoside (e.g., LNA, cEt, or ENA) or a non-bicyclic 2'-modified nucleoside (e.g., 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA)).

[000321] In some embodiments, one or more nucleosides in the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) are high-affinity modified nucleosides. In some embodiments, each nucleoside in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) is a high-affinity modified nucleoside. In some embodiments, one or more nucleosides in the 3' wing region of a gapmer (Z in the 5'-X-Y-Z-3' formula) are high-affinity modified nucleosides. In some embodiments, each nucleoside in the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) is a high-affinity modified nucleoside. In some embodiments, one or more nucleosides in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) are high-affinity modified

nucleosides and one or more nucleosides in the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) are high-affinity modified nucleosides. In some embodiments, each nucleoside in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) is a high-affinity modified nucleoside and each nucleoside in the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) is high-affinity modified nucleoside.

[000322] In some embodiments, the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) comprises the same high affinity nucleosides as the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula). For example, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) may comprise one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me). In another example, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) may comprise one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt). In some embodiments, each nucleoside in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) is a non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me). In some embodiments, each nucleoside in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) is a 2'-4' bicyclic nucleosides (e.g., LNA or cEt).

[000323] In some embodiments, a gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 1-7 (e.g., 1, 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein each nucleoside in X and Z is a non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and each nucleoside in Y is a 2'-deoxyribonucleoside. In some embodiments, the gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 1-7 (e.g., 1, 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein each nucleoside in X and Z is a 2'-4' bicyclic nucleosides (e.g., LNA or cEt) and each nucleoside in Y is a 2'-deoxyribonucleoside. In some embodiments, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) comprises different high affinity nucleosides as the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula). For example, the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) may comprise one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) may comprise one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt). In another example, the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) may comprise one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) may comprise one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt).

[000324] In some embodiments, a gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 1-7 (e.g., 1, 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein each nucleoside in X is a non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me), each nucleoside in Z is a 2'-4' bicyclic nucleosides (e.g., LNA or cEt), and each nucleoside in Y is a 2'-deoxyribonucleoside. In some embodiments, the gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 1-7 (e.g., 1, 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein each nucleoside in X is a 2'-4' bicyclic nucleosides (e.g., LNA or cEt), each nucleoside in Z is a non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and each nucleoside in Y is a 2'-deoxyribonucleoside.

[000325] In some embodiments, the 5' wing region of a gapmer (X in the 5'-X-Y-Z-3' formula) comprises one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt). In some embodiments, the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) comprises one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt). In some embodiments, both the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) comprise one or more non-bicyclic 2'-modified nucleosides (e.g., 2'-MOE or 2'-O-Me) and one or more 2'-4' bicyclic nucleosides (e.g., LNA or cEt).

[000326] In some embodiments, a gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 2-7 (e.g., 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein at least one but not all (e.g., 1, 2, 3, 4, 5, or 6) of positions 1, 2, 3, 4, 5, 6, or 7 in X (the 5' most position is position 1) is a non-bicyclic 2'-modified nucleoside (e.g., 2'-MOE or 2'-O-Me), wherein the rest of the nucleosides in both X and Z are 2'-4' bicyclic nucleosides (e.g., LNA or cEt), and wherein each nucleoside in Y is a 2'-deoxyribonucleoside. In some embodiments, the gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 2-7 (e.g., 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein at least one but not all (e.g., 1, 2, 3, 4, 5, or 6) of positions 1, 2, 3, 4, 5, 6, or 7 in Z (the 5' most position is position 1) is a non-bicyclic 2'-modified nucleoside (e.g., 2'-MOE or 2'-O-Me), wherein the rest of the nucleosides in both X and Z are 2'-4' bicyclic nucleosides (e.g., LNA or cEt), and wherein each nucleoside in Y is a 2'-deoxyribonucleoside. In some embodiments, the gapmer comprises a 5'-X-Y-Z-3' configuration, wherein X and Z is independently 2-7 (e.g., 2, 3, 4, 5, 6, or 7) nucleosides in length and Y is 6-10 (e.g., 6, 7, 8, 9, or 10) nucleosides in length, wherein at least one but not all (e.g., 1, 2, 3, 4, 5, or 6) of positions 1, 2, 3, 4, 5, 6, or 7 in X and at least one of

[1000327] Non-limiting examples of gapmers configurations with a mix of non-bicyclic 2'-modified nucleoside (e.g., 2'-MOE or 2'-O-Me) and 2'-4' bicyclic nucleosides (e.g., LNA or cEt) in the 5' wing region of the gapmer (X in the 5'-X-Y-Z-3' formula) and/or the 3' wing region of the gapmer (Z in the 5'-X-Y-Z-3' formula) include: BBB-(D)n-BBBAA; KKK-(D)n-KKKAA; LLL-(D)n-LLLAA; BBB-(D)n-BBBEE; KKK-(D)n-KKKEE; LLL-(D)n-LLLEE; BBB-(D)n-BBBAA; KKK-(D)n-KKKAA; LLL-(D)n-LLLAA; BBB-(D)n-BBBEE; KKK-(D)n-KKKEE; LLL-(D)n-LLLEE; BBB-(D)n-BBBAAA; KKK-(D)n-KKKAAA; LLL-(D)n-LLLAAA; BBB-(D)n-BBBEEE; KKK-(D)n-KKKEEE; LLL-(D)n-LLLEEE; BBB-(D)n-BBBAAA; KKK-(D)n-KKKAAA; LLL-(D)n-LLLAAA; BBB-(D)n-BBBEEE; KKK-(D)n-KKKEEE; LLL-(D)n-LLLEEE; BABA-(D)n-ABAB; KAKA-(D)n-AKAK; LALA-(D)n-ALAL; BEBE-(D)n-EBEB; KEKE-(D)n-EKEK; LELE-(D)n-ELEL; BABA-(D)n-ABAB; KAKA-(D)n-AKAK; LALA-(D)n-ALAL; BEBE-(D)n-EBEB; KEKE-(D)n-EKEK; LELE-(D)n-ELEL; ABAB-(D)n-ABAB; AKAK-(D)n-AKAK; ALAL-(D)n-ALAL; EBEB-(D)n-EBEB; EKEK-(D)n-EKEK; ELEL-(D)n-ELEL; ABAB-(D)n-ABAB; AKAK-(D)n-AKAK; ALAL-(D)n-ALAL; EBEB-(D)n-EBEB; EKEK-(D)n-EKEK; ELEL-(D)n-ELEL; AABB-(D)n-BBAA; BBAA-(D)n-AABB; AAKK-(D)n-KKAA; AALL-(D)n-LLAA; EEBB-(D)n-BBEE; EEKK-(D)n-KKEE; EELL-(D)n-LLEE; AABB-(D)n-BBAA; AAKK-(D)n-KKAA; AALL-(D)n-LLAA; EEBB-(D)n-BBEE; EEKK-(D)n-KKEE; EELL-(D)n-LLEE; BBB-(D)n-BBA; KKK-(D)n-KKA; LLL-(D)n-LLA; BBB-(D)n-BBE; KKK-(D)n-KKE; LLL-(D)n-LLE; BBB-(D)n-BBA; KKK-(D)n-KKA; LLL-(D)n-LLA; BBB-(D)n-BBE; KKK-(D)n-KKE; LLL-(D)n-LLE; BBB-(D)n-BBA; KKK-(D)n-KKA; LLL-(D)n-LLA; BBB-(D)n-BBE; KKK-(D)n-KKE; LLL-(D)n-LLE; ABBB-(D)n-BBBA; AKKK-(D)n-KKKA; ALLL-(D)n-LLLA; EBBB-(D)n-BBBE; EKKK-(D)n-KKKE; ELLL-(D)n-LLLE; ABBB-(D)n-BBBA; AKKK-(D)n-KKKA; ALLL-(D)n-LLLA; EBBB-(D)n-BBBE; EKKK-(D)n-KKKE; ELLL-(D)n-LLLE; ABBB-(D)n-BBBAA; AKKK-(D)n-KKKAA; ALLL-(D)n-LLLAA; EBBB-(D)n-BBBEE; EKKK-(D)n-KKKEE; ELLL-(D)n-LLLEE; ABBB-(D)n-BBBAA; AKKK-(D)n-KKKAA; ALLL-(D)n-LLLAA; EBBB-(D)n-BBBEE; EKKK-(D)n-KKKEE; ELLL-(D)n-LLLEE; AABBB-(D)n-BBB; AAKKK-(D)n-KKK; AALLL-(D)n-LLL; EEBBB-(D)n-BBB; EEKKK-(D)n-KKK; EELLL-(D)n-LLL; AABBB-(D)n-BBB; AAKKK-(D)n-KKK; AALLL-(D)n-LLL; EEBBB-(D)n-BBB; EEKKK-(D)n-KKK; EELLL-(D)n-LLL; AABBB-(D)n-BBBA; AAKKK-(D)n-KKKA; AALLL-(D)n-LLLA; EEBBB-(D)n-BBBE; EEKKK-(D)n-KKKE; EELLL-(D)n-

LLLE; AABBB-(D)_n-BBBA; AAKKK-(D)_n-KKKA; AALLL-(D)_n-LLLA; EEBBB-(D)_n-BBBE; EEKKK-(D)_n-KKKE; ELLLL-(D)_n-LLLE; ABBAABB-(D)_n-BB; AKKAAKK-(D)_n-KK; ALLAALLL-(D)_n-LL; EBBEEBB-(D)_n-BB; EKKEEKK-(D)_n-KK; ELLEELL-(D)_n-LL; ABBAABB-(D)_n-BB; AKKAAKK-(D)_n-KK; ALLAALL-(D)_n-LL; EBBEEBB-(D)_n-BB; EKKEEKK-(D)_n-KK; ELLEELL-(D)_n-LL; ABBABB-(D)_n-BBB; AKKAKK-(D)_n-KKK; ALLALLL-(D)_n-LLL; EBBEBB-(D)_n-BBB; EKKEKK-(D)_n-KKK; ELLELL-(D)_n-LLL; ABBABB-(D)_n-BBB; AKKAKK-(D)_n-KKK; ALLALL-(D)_n-LLL; EBBEBB-(D)_n-BBB; EKKEKK-(D)_n-KKK; ELLELL-(D)_n-LLL; EEK-(D)_n-EEEEEEEE; EEK-(D)_n-EEEEEEEE; EK-(D)_n-EEEEEEEE; EK-(D)_n-EEEEK; K-(D)_n-EEEKEKE; K-(D)_n-EEEKEKEE; K-(D)_n-EEKEK; EK-(D)_n-EEEKEKE; EK-(D)_n-EEKEK; EEK-(D)_n-KEEKE; EK-(D)_n-EEKEK; EK-(D)_n-KEEK; EEK-(D)_n-EEKEK; EK-(D)_n-KEEKEE; EK-(D)_n-EEKEKE; EK-(D)_n-EEKEKE; and EK-(D)_n-EEKEKE;. “A” nucleosides comprise a 2'-modified nucleoside; “B” represents a 2'-4' bicyclic nucleoside; “K” represents a constrained ethyl nucleoside (cEt); “L” represents an LNA nucleoside; and “E” represents a 2'-MOE modified ribonucleoside; “D” represents a 2'-deoxyribonucleoside; “n” represents the length of the gap segment (Y in the 5'-X-Y-Z-3' configuration) and is an integer between 1-20.

[000328] In some embodiments, any one of the gapmers described herein comprises one or more modified nucleoside linkages (e.g., a phosphorothioate linkage) in each of the X, Y, and Z regions. In some embodiments, each internucleoside linkage in the any one of the gapmers described herein is a phosphorothioate linkage. In some embodiments, each of the X, Y, and Z regions independently comprises a mix of phosphorothioate linkages and phosphodiester linkages. In some embodiments, each internucleoside linkage in the gap region Y is a phosphorothioate linkage, the 5' wing region X comprises a mix of phosphorothioate linkages and phosphodiester linkages, and the 3' wing region Z comprises a mix of phosphorothioate linkages and phosphodiester linkages.

i. Mixmers

[000329] In some embodiments, an oligonucleotide described herein may be a mixmer or comprise a mixmer sequence pattern. In general, mixmers are oligonucleotides that comprise both naturally and non-naturally occurring nucleosides or comprise two different types of non-naturally occurring nucleosides typically in an alternating pattern. Mixmers generally have higher binding affinity than unmodified oligonucleotides and may be used to specifically bind a target molecule, *e.g.*, to block a binding site on the target molecule. Generally, mixmers do not recruit an RNase to the target molecule and thus do not promote cleavage of the target molecule.

Such oligonucleotides that are incapable of recruiting RNase H have been described, for example, see WO2007/112754 or WO2007/112753.

[000330] In some embodiments, the mixmer comprises or consists of a repeating pattern of nucleoside analogues and naturally occurring nucleosides, or one type of nucleoside analogue and a second type of nucleoside analogue. However, a mixmer need not comprise a repeating pattern and may instead comprise any arrangement of modified nucleosides and naturally occurring nucleosides or any arrangement of one type of modified nucleoside and a second type of modified nucleoside. The repeating pattern, may, for instance be every second or every third nucleoside is a modified nucleoside, such as LNA, and the remaining nucleosides are naturally occurring nucleosides, such as DNA, or are a 2' substituted nucleoside analogue such as 2'-MOE or 2' fluoro analogues, or any other modified nucleoside described herein. It is recognized that the repeating pattern of modified nucleoside, such as LNA units, may be combined with modified nucleoside at fixed positions—*e.g.* at the 5' or 3' termini.

[000331] In some embodiments, a mixmer does not comprise a region of more than 5, more than 4, more than 3, or more than 2 consecutive naturally occurring nucleosides, such as DNA nucleosides. In some embodiments, the mixmer comprises at least a region consisting of at least two consecutive modified nucleoside, such as at least two consecutive LNAs. In some embodiments, the mixmer comprises at least a region consisting of at least three consecutive modified nucleoside units, such as at least three consecutive LNAs.

[000332] In some embodiments, the mixmer does not comprise a region of more than 7, more than 6, more than 5, more than 4, more than 3, or more than 2 consecutive nucleoside analogues, such as LNAs. In some embodiments, LNA units may be replaced with other nucleoside analogues, such as those referred to herein.

[000333] Mixmers may be designed to comprise a mixture of affinity enhancing modified nucleosides, such as in non-limiting example LNA nucleosides and 2'-O-Me nucleosides. In some embodiments, a mixmer comprises modified internucleoside linkages (*e.g.*, phosphorothioate internucleoside linkages or other linkages) between at least two, at least three, at least four, at least five or more nucleosides.

[000334] A mixmer may be produced using any suitable method. Representative U.S. patents, U.S. patent publications, and PCT publications that teach the preparation of mixmers include U.S. patent publication Nos. US20060128646, US20090209748, US20090298916, US20110077288, and US20120322851, and U.S. patent No. 7687617.

[000335] In some embodiments, a mixmer comprises one or more morpholino nucleosides. For example, in some embodiments, a mixmer may comprise morpholino

nucleosides mixed (*e.g.*, in an alternating manner) with one or more other nucleosides (*e.g.*, DNA, RNA nucleosides) or modified nucleosides (*e.g.*, LNA, 2'-O-Me nucleosides).

[000336] In some embodiments, mixmers are useful for splice correcting or exon skipping, for example, as reported in Touznik A., et al., *LNA/DNA mixmer-based antisense oligonucleotides correct alternative splicing of the SMN2 gene and restore SMN protein expression in type 1 SMA fibroblasts* Scientific Reports, volume 7, Article number: 3672 (2017), Chen S. et al., *Synthesis of a Morpholino Nucleic Acid (MNA)-Uridine Phosphoramidite, and Exon Skipping Using MNA/2'-O-Methyl Mixmer Antisense Oligonucleotide*, *Molecules* 2016, 21, 1582, the contents of each which are incorporated herein by reference.

j. RNA Interference (RNAi)

[000337] In some embodiments, oligonucleotides provided herein may be in the form of small interfering RNAs (siRNA), also known as short interfering RNA or silencing RNA. siRNA, is a class of double-stranded RNA molecules, typically about 20-25 base pairs in length that target nucleic acids (*e.g.*, mRNAs) for degradation via the RNA interference (RNAi) pathway in cells. Specificity of siRNA molecules may be determined by the binding of the antisense strand of the molecule to its target RNA. Effective siRNA molecules are generally less than 30 to 35 base pairs in length to prevent the triggering of non-specific RNA interference pathways in the cell via the interferon response, although longer siRNA can also be effective. In some embodiments, the siRNA molecules are 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or more base pairs in length. In some embodiments, the siRNA molecules are 8 to 30 base pairs in length, 10 to 15 base pairs in length, 10 to 20 base pairs in length, 15 to 25 base pairs in length, 19 to 21 base pairs in length, 21 to 23 base pairs in length.

[000338] Following selection of an appropriate target RNA sequence, siRNA molecules that comprise a nucleotide sequence complementary to all or a portion of the target sequence, *i.e.* an antisense sequence, can be designed and prepared using appropriate methods (see, *e.g.*, PCT Publication Number WO 2004/016735; and U.S. Patent Publication Nos. 2004/0077574 and 2008/0081791). The siRNA molecule can be double stranded (*i.e.* a dsRNA molecule comprising an antisense strand and a complementary sense strand that hybridizes to form the dsRNA) or single-stranded (*i.e.* a ssRNA molecule comprising just an antisense strand). The siRNA molecules can comprise a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense strands.

[000339] In some embodiments, the antisense strand of the siRNA molecule is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or

more nucleotides in length. In some embodiments, the antisense strand is 8 to 50 nucleotides in length, 8 to 40 nucleotides in length, 8 to 30 nucleotides in length, 10 to 15 nucleotides in length, 10 to 20 nucleotides in length, 15 to 25 nucleotides in length, 19 to 21 nucleotides in length, 21 to 23 nucleotides in lengths.

[000340] In some embodiments, the sense strand of the siRNA molecule is 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, or more nucleotides in length. In some embodiments, the sense strand is 8 to 50 nucleotides in length, 8 to 40 nucleotides in length, 8 to 30 nucleotides in length, 10 to 15 nucleotides in length, 10 to 20 nucleotides in length, 15 to 25 nucleotides in length, 19 to 21 nucleotides in length, 21 to 23 nucleotides in lengths.

[000341] In some embodiments, siRNA molecules comprise an antisense strand comprising a region of complementarity to a target region in a target mRNA. In some embodiments, the region of complementarity is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% complementary to a target region in a target mRNA. In some embodiments, the target region is a region of consecutive nucleotides in the target mRNA. In some embodiments, a complementary nucleotide sequence need not be 100% complementary to that of its target to be specifically hybridizable or specific for a target RNA sequence.

[000342] In some embodiments, siRNA molecules comprise an antisense strand that comprises a region of complementarity to a target RNA sequence and the region of complementarity is in the range of 8 to 15, 8 to 30, 8 to 40, or 10 to 50, or 5 to 50, or 5 to 40 nucleotides in length. In some embodiments, a region of complementarity is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length. In some embodiments, the region of complementarity is complementary with at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25 or more consecutive nucleotides of a target RNA sequence. In some embodiments, siRNA molecules comprise a nucleotide sequence that contains no more than 1, 2, 3, 4, or 5 base mismatches compared to the portion of the consecutive nucleotides of target RNA sequence. In some embodiments, siRNA molecules comprise a nucleotide sequence that has up to 3 mismatches over 15 bases, or up to 2 mismatches over 10 bases.

[000343] In some embodiments, siRNA molecules comprise an antisense strand comprising a nucleotide sequence that is complementary (e.g., at least 85%, at least 90%, at least 95%, or 100%) to the target RNA sequence of the oligonucleotides provided herein. In some

embodiments, siRNA molecules comprise an antisense strand comprising a nucleotide sequence that is at least 85%, at least 90%, at least 95%, or 100% identical to the oligonucleotides provided herein. In some embodiments, siRNA molecules comprise an antisense strand comprising at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25 or more consecutive nucleotides of the oligonucleotides provided herein.

[000344] Double-stranded siRNA may comprise sense and anti-sense RNA strands that are the same length or different lengths. Double-stranded siRNA molecules can also be assembled from a single oligonucleotide in a stem-loop structure, wherein self-complementary sense and antisense regions of the siRNA molecule are linked by means of a nucleic acid based or non-nucleic acid-based linker(s), as well as circular single-stranded RNA having two or more loop structures and a stem comprising self-complementary sense and antisense strands, wherein the circular RNA can be processed either in vivo or in vitro to generate an active siRNA molecule capable of mediating RNAi. Small hairpin RNA (shRNA) molecules thus are also contemplated herein. These molecules comprise a specific antisense sequence in addition to the reverse complement (sense) sequence, typically separated by a spacer or loop sequence. Cleavage of the spacer or loop provides a single-stranded RNA molecule and its reverse complement, such that they may anneal to form a dsRNA molecule (optionally with additional processing steps that may result in addition or removal of one, two, three or more nucleotides from the 3' end and/or (e.g., and) the 5' end of either or both strands). A spacer can be of a sufficient length to permit the antisense and sense sequences to anneal and form a double-stranded structure (or stem) prior to cleavage of the spacer (and, optionally, subsequent processing steps that may result in addition or removal of one, two, three, four, or more nucleotides from the 3' end and/or (e.g., and) the 5' end of either or both strands). A spacer sequence is may be an unrelated nucleotide sequence that is situated between two complementary nucleotide sequence regions which, when annealed into a double-stranded nucleic acid, comprise a shRNA.

[000345] The overall length of the siRNA molecules can vary from about 14 to about 100 nucleotides depending on the type of siRNA molecule being designed. Generally between about 14 and about 50 of these nucleotides are complementary to the RNA target sequence, i.e. constitute the specific antisense sequence of the siRNA molecule. For example, when the siRNA is a double- or single-stranded siRNA, the length can vary from about 14 to about 50 nucleotides, whereas when the siRNA is a shRNA or circular molecule, the length can vary from about 40 nucleotides to about 100 nucleotides.

[000346] An siRNA molecule may comprise a 3' overhang at one end of the molecule. The other end may be blunt-ended or have also an overhang (5' or 3'). When the siRNA molecule comprises an overhang at both ends of the molecule, the length of the overhangs may be the same or different. In one embodiment, the siRNA molecule of the present disclosure comprises 3' overhangs of about 1 to about 3 nucleotides on both ends of the molecule. In some embodiments, the siRNA molecule comprises 3' overhangs of about 1 to about 3 nucleotides on the sense strand. In some embodiments, the siRNA molecule comprises 3' overhangs of about 1 to about 3 nucleotides on the antisense strand. In some embodiments, the siRNA molecule comprises 3' overhangs of about 1 to about 3 nucleotides on both the sense strand and the antisense strand.

[000347] In some embodiments, the siRNA molecule comprises one or more modified nucleotides (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more). In some embodiments, the siRNA molecule comprises one or more modified nucleotides and/or (e.g., and) one or more modified internucleotide linkages. In some embodiments, the modified nucleotide is a modified sugar moiety (e.g. a 2' modified nucleotide). In some embodiments, the siRNA molecule comprises one or more 2' modified nucleotides, e.g., a 2'-deoxy, 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O--N-methylacetamido (2'-O--NMA). In some embodiments, each nucleotide of the siRNA molecule is a modified nucleotide (e.g., a 2'-modified nucleotide). In some embodiments, the siRNA molecule comprises one or more phosphorodiamidate morpholinos. In some embodiments, each nucleotide of the siRNA molecule is a phosphorodiamidate morpholino.

[000348] In some embodiments, the siRNA molecule contains a phosphorothioate or other modified internucleotide linkage. In some embodiments, the siRNA molecule comprises phosphorothioate internucleoside linkages. In some embodiments, the siRNA molecule comprises phosphorothioate internucleoside linkages between at least two nucleotides. In some embodiments, the siRNA molecule comprises phosphorothioate internucleoside linkages between all nucleotides. For example, in some embodiments, the siRNA molecule comprises modified internucleotide linkages at the first, second, and/or (e.g., and) third internucleoside linkage at the 5' or 3' end of the siRNA molecule.

[000349] In some embodiments, the modified internucleotide linkages are phosphorus-containing linkages. In some embodiments, phosphorus-containing linkages that may be used include, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl

phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'; see US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5, 177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563, 253; 5,571,799; 5,587,361; and 5,625,050.

[000350] Any of the modified chemistries or formats of siRNA molecules described herein can be combined with each other. For example, one, two, three, four, five, or more different types of modifications can be included within the same siRNA molecule.

[000351] In some embodiments, the antisense strand comprises one or more modified nucleotides (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more). In some embodiments, the antisense strand comprises one or more modified nucleotides and/or (e.g., and) one or more modified internucleotide linkages. In some embodiments, the modified nucleotide comprises a modified sugar moiety (e.g. a 2' modified nucleotide). In some embodiments, the antisense strand comprises one or more 2' modified nucleotides, e.g., a 2'-deoxy, 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O--N-methylacetamido (2'-O--NMA). In some embodiments, each nucleotide of the antisense strand is a modified nucleotide (e.g., a 2'-modified nucleotide). In some embodiments, the antisense strand comprises one or more phosphorodiamidate morpholinos. In some embodiments, the antisense strand is a phosphorodiamidate morpholino oligomer (PMO).

[000352] In some embodiments, antisense strand contains a phosphorothioate or other modified internucleotide linkage. In some embodiments, the antisense strand comprises phosphorothioate internucleoside linkages. In some embodiments, the antisense strand comprises phosphorothioate internucleoside linkages between at least two nucleotides. In some embodiments, the antisense strand comprises phosphorothioate internucleoside linkages between all nucleotides. For example, in some embodiments, the antisense strand comprises modified internucleotide linkages at the first, second, and/or (e.g., and) third internucleoside linkage at the 5' or 3' end of the siRNA molecule. In some embodiments, the modified internucleotide linkages are phosphorus-containing linkages. In some embodiments, phosphorus-containing linkages that may be used include, but are not limited to, phosphorothioates, chiral phosphorothioates,

phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'; see US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

[000353] Any of the modified chemistries or formats of the antisense strand described herein can be combined with each other. For example, one, two, three, four, five, or more different types of modifications can be included within the same antisense strand.

[000354] In some embodiments, the sense strand comprises one or more modified nucleotides (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more). In some embodiments, the sense strand comprises one or more modified nucleotides and/or (e.g., and) one or more modified internucleotide linkages. In some embodiments, the modified nucleotide is a modified sugar moiety (e.g. a 2' modified nucleotide). In some embodiments, the sense strand comprises one or more 2' modified nucleotides, e.g., a 2'-deoxy, 2'-fluoro (2'-F), 2'-O-methyl (2'-O-Me), 2'-O-methoxyethyl (2'-MOE), 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), 2'-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O--N-methylacetamido (2'-O--NMA). In some embodiments, each nucleotide of the sense strand is a modified nucleotide (e.g., a 2'-modified nucleotide). In some embodiments, the sense strand comprises one or more phosphorodiamidate morpholinos. In some embodiments, the antisense strand is a phosphorodiamidate morpholino oligomer (PMO). In some embodiments, the sense strand contains a phosphorothioate or other modified internucleotide linkage. In some embodiments, the sense strand comprises phosphorothioate internucleoside linkages. In some embodiments, the sense strand comprises phosphorothioate internucleoside linkages between at least two nucleotides. In some embodiments, the sense strand comprises phosphorothioate internucleoside linkages between all nucleotides. For example, in some embodiments, the sense strand comprises modified internucleotide linkages at the first, second, and/or (e.g., and) third internucleoside linkage at the 5' or 3' end of the sense strand.

[000355] In some embodiments, the modified internucleotide linkages are phosphorus-containing linkages. In some embodiments, phosphorus-containing linkages that may be used

include, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'; see US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; and 5,625,050.

[000356] Any of the modified chemistries or formats of the sense strand described herein can be combined with each other. For example, one, two, three, four, five, or more different types of modifications can be included within the same sense strand.

[000357] In some embodiments, the antisense or sense strand of the siRNA molecule comprises modifications that enhance or reduce RNA-induced silencing complex (RISC) loading. In some embodiments, the antisense strand of the siRNA molecule comprises modifications that enhance RISC loading. In some embodiments, the sense strand of the siRNA molecule comprises modifications that reduce RISC loading and reduce off-target effects. In some embodiments, the antisense strand of the siRNA molecule comprises a 2'-O-methoxyethyl (2'-MOE) modification. The addition of the 2'-O-methoxyethyl (2'-MOE) group at the cleavage site improves both the specificity and silencing activity of siRNAs by facilitating the oriented RNA-induced silencing complex (RISC) loading of the modified strand, as described in Song et al., (2017) *Mol Ther Nucleic Acids* 9:242-250, incorporated herein by reference in its entirety. In some embodiments, the antisense strand of the siRNA molecule comprises a 2'-OMe-phosphorodithioate modification, which increases RISC loading as described in Wu et al., (2014) *Nat Commun* 5:3459, incorporated herein by reference in its entirety.

[000358] In some embodiments, the sense strand of the siRNA molecule comprises a 5'-morpholino, which reduces RISC loading of the sense strand and improves antisense strand selection and RNAi activity, as described in Kumar et al., (2019) *Chem Commun (Camb)* 55(35):5139-5142, incorporated herein by reference in its entirety. In some embodiments, the sense strand of the siRNA molecule is modified with a synthetic RNA-like high affinity nucleotide analogue, Locked Nucleic Acid (LNA), which reduces RISC loading of the sense strand and further enhances antisense strand incorporation into RISC, as described in Elman et al., (2005) *Nucleic Acids Res.* 33(1): 439-447, incorporated herein by reference in its entirety. In

some embodiments, the sense strand of the siRNA molecule comprises a 5' unlocked nucleic acid (UNA) modification, which reduce RISC loading of the sense strand and improve silencing potency of the antisense strand, as described in Snead et al., (2013) *Mol Ther Nucleic Acids* 2(7):e103, incorporated herein by reference in its entirety. In some embodiments, the sense strand of the siRNA molecule comprises a 5-nitroindole modification, which decreased the RNAi potency of the sense strand and reduces off-target effects as described in Zhang et al., (2012) *Chembiochem* 13(13):1940-1945, incorporated herein by reference in its entirety. In some embodiments, the sense strand comprises a 2'-O-methyl (2'-O-Me) modification, which reduces RISC loading and the off-target effects of the sense strand, as described in Zheng et al., *FASEB* (2013) 27(10): 4017-4026, incorporated herein by reference in its entirety. In some embodiments, the sense strand of the siRNA molecule is fully substituted with morpholino, 2'-MOE or 2'-O-Me residues, and are not recognized by RISC as described in Kole et al., (2012) *Nature reviews. Drug Discovery* 11(2):125-140, incorporated herein by reference in its entirety. In some embodiments the antisense strand of the siRNA molecule comprises a 2'-MOE modification and the sense strand comprises an 2'-O-Me modification (see e.g., Song et al., (2017) *Mol Ther Nucleic Acids* 9:242-250). In some embodiments at least one (e.g., at least 2, at least 3, at least 4, at least 5, at least 10) siRNA molecule is linked (e.g., covalently) to a muscle-targeting agent. In some embodiments, the muscle-targeting agent may comprise, or consist of, a nucleic acid (e.g., DNA or RNA), a peptide (e.g., an antibody), a lipid (e.g., a microvesicle), or a sugar moiety (e.g., a polysaccharide). In some embodiments, the muscle-targeting agent is an antibody. In some embodiments, the muscle-targeting agent is an anti-transferrin receptor antibody (e.g., any one of the anti-TfR antibodies provided herein). In some embodiments, the muscle-targeting agent may be linked to the 5' end of the sense strand of the siRNA molecule. In some embodiments, the muscle-targeting agent may be linked to the 3' end of the sense strand of the siRNA molecule. In some embodiments, the muscle-targeting agent may be linked internally to the sense strand of the siRNA molecule. In some embodiments, the muscle-targeting agent may be linked to the 5' end of the antisense strand of the siRNA molecule. In some embodiments, the muscle-targeting agent may be linked to the 3' end of the antisense strand of the siRNA molecule. In some embodiments, the muscle-targeting agent may be linked internally to the antisense strand of the siRNA molecule.

k. microRNA (miRNAs)

[000359] In some embodiments, an oligonucleotide may be a microRNA (miRNA). MicroRNAs (referred to as "miRNAs") are small non-coding RNAs, belonging to a class of regulatory molecules that control gene expression by binding to complementary sites on a target

RNA transcript. Typically, miRNAs are generated from large RNA precursors (termed pri-miRNAs) that are processed in the nucleus into approximately 70 nucleotide pre-miRNAs, which fold into imperfect stem-loop structures. These pre-miRNAs typically undergo an additional processing step within the cytoplasm where mature miRNAs of 18-25 nucleotides in length are excised from one side of the pre-miRNA hairpin by an RNase III enzyme, Dicer.

[000360] As used herein, miRNAs including pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of mature miRNA. In one embodiment, the size range of the miRNA can be from 21 nucleotides to 170 nucleotides. In one embodiment the size range of the miRNA is from 70 to 170 nucleotides in length. In another embodiment, mature miRNAs of from 21 to 25 nucleotides in length can be used.

l. Aptamers

[000361] In some embodiments, oligonucleotides provided herein may be in the form of aptamers. Generally, in the context of molecular payloads, aptamer is any nucleic acid that binds specifically to a target, such as a small molecule, protein, nucleic acid in a cell. In some embodiments, the aptamer is a DNA aptamer or an RNA aptamer. In some embodiments, a nucleic acid aptamer is a single-stranded DNA or RNA (ssDNA or ssRNA). It is to be understood that a single-stranded nucleic acid aptamer may form helices and/or (e.g., and) loop structures. The nucleic acid that forms the nucleic acid aptamer may comprise naturally occurring nucleotides, modified nucleotides, naturally occurring nucleotides with hydrocarbon linkers (e.g., an alkylene) or a polyether linker (e.g., a PEG linker) inserted between one or more nucleotides, modified nucleotides with hydrocarbon or PEG linkers inserted between one or more nucleotides, or a combination of thereof. Exemplary publications and patents describing aptamers and method of producing aptamers include, e.g., Lorsch and Szostak, 1996; Jayasena, 1999; U.S. Pat. Nos. 5,270,163; 5,567,588; 5,650,275; 5,670,637; 5,683,867; 5,696,249; 5,789,157; 5,843,653; 5,864,026; 5,989,823; 6,569,630; 8,318,438 and PCT application WO 99/31275, each incorporated herein by reference.

m. Ribozymes

[000362] In some embodiments, oligonucleotides provided herein may be in the form of a ribozyme. A ribozyme (ribonucleic acid enzyme) is a molecule, typically an RNA molecule, that is capable of performing specific biochemical reactions, similar to the action of protein enzymes. Ribozymes are molecules with catalytic activities including the ability to cleave at specific phosphodiester linkages in RNA molecules to which they have hybridized, such as mRNAs, RNA-containing substrates, lncRNAs, and ribozymes, themselves.

[000363] Ribozymes may assume one of several physical structures, one of which is called a "hammerhead." A hammerhead ribozyme is composed of a catalytic core containing nine

conserved bases, a double-stranded stem and loop structure (stem-loop II), and two regions complementary to the target RNA flanking regions the catalytic core. The flanking regions enable the ribozyme to bind to the target RNA specifically by forming double-stranded stems I and III. Cleavage occurs in cis (i.e., cleavage of the same RNA molecule that contains the hammerhead motif) or in trans (cleavage of an RNA substrate other than that containing the ribozyme) next to a specific ribonucleotide triplet by a transesterification reaction from a 3', 5'-phosphate diester to a 2', 3'-cyclic phosphate diester. Without wishing to be bound by theory, it is believed that this catalytic activity requires the presence of specific, highly conserved sequences in the catalytic region of the ribozyme.

[000364] Modifications in ribozyme structure have also included the substitution or replacement of various non-core portions of the molecule with non-nucleotidic molecules. For example, Benseler et al. (J. Am. Chem. Soc. (1993) 115:8483-8484) disclosed hammerhead-like molecules in which two of the base pairs of stem II, and all four of the nucleotides of loop II were replaced with non-nucleoside linkers based on hexaethylene glycol, propanediol, bis(triethylene glycol) phosphate, tris(propanediol)bisphosphate, or bis(propanediol) phosphate. Ma et al. (Biochem. (1993) 32:1751-1758; Nucleic Acids Res. (1993) 21:2585-2589) replaced the six nucleotide loop of the TAR ribozyme hairpin with non-nucleotidic, ethylene glycol-related linkers. Thomson et al. (Nucleic Acids Res. (1993) 21:5600-5603) replaced loop II with linear, non-nucleotidic linkers of 13, 17, and 19 atoms in length.

[000365] Ribozyme oligonucleotides can be prepared using well known methods (see, e.g., PCT Publications WO9118624; WO9413688; WO9201806; and WO 92/07065; and U.S. Patents 5436143 and 5650502) or can be purchased from commercial sources (e.g., US Biochemicals) and, if desired, can incorporate nucleotide analogs to increase the resistance of the oligonucleotide to degradation by nucleases in a cell. The ribozyme may be synthesized in any known manner, e.g., by use of a commercially available synthesizer produced, e.g., by Applied Biosystems, Inc. or Milligen. The ribozyme may also be produced in recombinant vectors by conventional means. See, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory (Current edition). The ribozyme RNA sequences may be synthesized conventionally, for example, by using RNA polymerases such as T7 or SP6.

n. Guide Nucleic Acids

[000366] In some embodiments, oligonucleotides are guide nucleic acid, e.g., guide RNA (gRNA) molecules. Generally, a guide RNA is a short synthetic RNA composed of (1) a scaffold sequence that binds to a nucleic acid programmable DNA binding protein (napDNAbp), such as Cas9, and (2) a nucleotide spacer portion that defines the DNA target sequence (e.g., genomic DNA target) to which the gRNA binds in order to bring the nucleic acid programmable

DNA binding protein in proximity to the DNA target sequence. In some embodiments, the napDNAbp is a nucleic acid-programmable protein that forms a complex with (e.g., binds or associates with) one or more RNA(s) that targets the nucleic acid-programmable protein to a target DNA sequence (e.g., a target genomic DNA sequence). In some embodiments, a nucleic acid-programmable nuclease, when in a complex with an RNA, may be referred to as a nuclease:RNA complex. Guide RNAs can exist as a complex of two or more RNAs, or as a single RNA molecule.

[000367] Guide RNAs (gRNAs) that exist as a single RNA molecule may be referred to as single-guide RNAs (sgRNAs), though gRNA is also used to refer to guide RNAs that exist as either single molecules or as a complex of two or more molecules. Typically, gRNAs that exist as a single RNA species comprise two domains: (1) a domain that shares homology to a target nucleic acid (i.e., directs binding of a Cas9 complex to the target); and (2) a domain that binds a Cas9 protein. In some embodiments, domain (2) corresponds to a sequence known as a tracrRNA and comprises a stem-loop structure. In some embodiments, domain (2) is identical or homologous to a tracrRNA as provided in Jinek et al., *Science* 337:816-821 (2012), the entire contents of which is incorporated herein by reference.

[000368] In some embodiments, a gRNA comprises two or more of domains (1) and (2), and may be referred to as an extended gRNA. For example, an extended gRNA will bind two or more Cas9 proteins and bind a target nucleic acid at two or more distinct regions, as described herein. The gRNA comprises a nucleotide sequence that complements a target site, which mediates binding of the nuclease:RNA complex to said target site, providing the sequence specificity of the nuclease:RNA complex. In some embodiments, the RNA-programmable nuclease is the (CRISPR-associated system) Cas9 endonuclease, for example, Cas9 (Csn1) from *Streptococcus pyogenes* (see, e.g., “Complete genome sequence of an M1 strain of *Streptococcus pyogenes*.” Ferretti J.J., McShan W.M., Ajdic D.J., Savic D.J., Savic G., Lyon K., Primeaux C., Sezate S., Suvorov A.N., Kenton S., Lai H.S., Lin S.P., Qian Y., Jia H.G., Najar F.Z., Ren Q., Zhu H., Song L., White J., Yuan X., Clifton S.W., Roe B.A., McLaughlin R.E., *Proc. Natl. Acad. Sci. U.S.A.* 98:4658-4663 (2001); “CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III.” Deltcheva E., Chylinski K., Sharma C.M., Gonzales K., Chao Y., Pirzada Z.A., Eckert M.R., Vogel J., Charpentier E., *Nature* 471:602-607 (2011); and “A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity.” Jinek M., Chylinski K., Fonfara I., Hauer M., Doudna J.A., Charpentier E. *Science* 337:816-821 (2012), the entire contents of each of which are incorporated herein by reference.

o. Multimers

[000369] In some embodiments, molecular payloads may comprise multimers (e.g., concatemers) of 2 or more oligonucleotides connected by a linker. In this way, in some embodiments, the oligonucleotide loading of a complex/conjugate can be increased beyond the available linking sites on a targeting agent (e.g., available thiol sites on an antibody) or otherwise tuned to achieve a particular payload loading content. Oligonucleotides in a multimer can be the same or different (e.g., targeting different genes or different sites on the same gene or products thereof).

[000370] In some embodiments, multimers comprise 2 or more oligonucleotides linked together by a cleavable linker. However, in some embodiments, multimers comprise 2 or more oligonucleotides linked together by a non-cleavable linker. In some embodiments, a multimer comprises 2, 3, 4, 5, 6, 7, 8, 9, 10 or more oligonucleotides linked together. In some embodiments, a multimer comprises 2 to 5, 2 to 10 or 4 to 20 oligonucleotides linked together.

[000371] In some embodiments, a multimer comprises 2 or more oligonucleotides linked end-to-end (in a linear arrangement). In some embodiments, a multimer comprises 2 or more oligonucleotides linked end-to-end via an oligonucleotide based linker (e.g., poly-dT linker, an abasic linker). In some embodiments, a multimer comprises a 5' end of one oligonucleotide linked to a 3' end of another oligonucleotide. In some embodiments, a multimer comprises a 3' end of one oligonucleotide linked to a 3' end of another oligonucleotide. In some embodiments, a multimer comprises a 5' end of one oligonucleotide linked to a 5' end of another oligonucleotide. Still, in some embodiments, multimers can comprise a branched structure comprising multiple oligonucleotides linked together by a branching linker.

[000372] Further examples of multimers that may be used in the complexes provided herein are disclosed, for example, in US Patent Application Number 2015/0315588 A1, entitled *Methods of delivering multiple targeting oligonucleotides to a cell using cleavable linkers*, which was published on November 5, 2015; US Patent Application Number 2015/0247141 A1, entitled *Multimeric Oligonucleotide Compounds*, which was published on September 3, 2015, US Patent Application Number US 2011/0158937 A1, entitled *Immunostimulatory Oligonucleotide Multimers*, which was published on June 30, 2011; and US Patent Number 5,693,773, entitled *Triplex-Forming Antisense Oligonucleotides Having Abasic Linkers Targeting Nucleic Acids Comprising Mixed Sequences Of Purines And Pyrimidines*, which issued on December 2, 1997, the contents of each of which are incorporated herein by reference in their entireties.

ii. Small Molecules:

[000373] Any suitable small molecule may be used as a molecular payload, as described herein. In some embodiments, the small molecule is as described in US Patent Application

Publication 2016052914A1, published on February 25, 2016, entitled “*Compounds And Methods For Myotonic Dystrophy Therapy*”. Further examples of small molecule payloads are provided in Lopez-Morato M, et al., Small Molecules Which Improve Pathogenesis of Myotonic Dystrophy Type 1, (Review) Front. Neurol., 18 May 2018. For example, in some embodiments, the small molecule is an MBNL1 upregulator such as phenylbuthazone, ketoprofen, ISOX, or vorinostat. In some embodiments, the small molecule is an H-Ras pathway inhibitor such as manumycin A. In some embodiments, the small molecule is a protein kinase modulator such as Ro-318220, C16, C51, Metformin, AICAR, lithium chloride, TDZD-8 or Bio. In some embodiments, the small molecule is a plant alkaloid such as harmine. In some embodiments, the small molecule is a transcription inhibitor such as pentamidine, propamidine, heptamidine or actinomycin D. In some embodiments, the small molecule is an inhibitor of Glycogen synthase kinase 3 beta (GSK3B), for example, as disclosed in Jones K, et al., GSK3 β mediates muscle pathology in myotonic dystrophy. J Clin Invest. 2012 Dec;122(12):4461-72; and Wei C, et al., GSK3 β is a new therapeutic target for myotonic dystrophy type 1. Rare Dis. 2013; 1: e26555; and Palomo V, et al., Subtly Modulating Glycogen Synthase Kinase 3 β : Allosteric Inhibitor Development and Their Potential for the Treatment of Chronic Diseases. J Med Chem. 2017 Jun 22;60(12):4983-5001, the contents of each of which are incorporated herein by reference in their entireties. In some embodiments, the small molecule is a substituted pyrido[2,3-d]pyrimidines and pentamidine-like compound, as disclosed in Gonzalez AL, et al., In silico discovery of substituted pyrido[2,3-d]pyrimidines and pentamidine-like compounds with biological activity in myotonic dystrophy models. PLoS One. 2017 Jun 5;12(6):e0178931, the contents of which are incorporated herein by reference in its entirety. In some embodiments, the small molecule is an MBNL1 modulator, for example, as disclosed in: Zhange F, et al., A flow cytometry-based screen identifies MBNL1 modulators that rescue splicing defects in myotonic dystrophy type I. Hum Mol Genet. 2017 Aug 15;26(16):3056-3068, the contents of which are incorporated herein by reference in its entirety.

iii. Peptides

[000374] Any suitable peptide or protein may be used as a molecular payload, as described herein. A peptide or protein payload may correspond to a sequence of a protein that preferentially binds to a nucleic acid, *e.g.* a disease-associated repeat, or a protein, *e.g.* MBNL1, found in muscle cells. In some embodiments, peptides or proteins may be produced, synthesized, and/or (*e.g.*, and) derivatized using several methodologies, *e.g.* phage displayed peptide libraries, one-bead one-compound peptide libraries, or positional scanning synthetic peptide combinatorial libraries. Exemplary methodologies have been characterized in the art and are incorporated by reference (Gray, B.P. and Brown, K.C. “Combinatorial Peptide

Libraries: Mining for Cell-Binding Peptides” Chem Rev. 2014, 114:2, 1020–1081.; Samoylova, T.I. and Smith, B.F. “Elucidation of muscle-binding peptides by phage display screening.” Muscle Nerve, 1999, 22:4. 460-6.).

[000375] In some embodiments, the peptide is as described in US Patent Application 2018/0021449, published on 1/25/2018, “Antisense conjugates for decreasing expression of DMPK”. In some embodiments, the peptide is as described in Garcia-Lopez et al., “*In vivo* discovery of a peptide that prevents CUG–RNA hairpin formation and reverses RNA toxicity in myotonic dystrophy models”, PNAS July 19, 2011. 108 (29) 11866-11871. In some embodiments, the peptide or protein may target, *e.g.*, bind to, a disease-associated repeat, *e.g.* an RNA CUG repeat expansion.

[000376] In some embodiments, the peptide or protein comprises a fragment of an MBNL protein, *e.g.*, MBNL1. In some embodiments, the peptide or protein comprises at least one zinc finger. In some embodiments, the peptide or protein may comprise about 2-25 amino acids, about 2-20 amino acids, about 2-15 amino acids, about 2-10 amino acids, or about 2-5 amino acids. The peptide or protein may comprise naturally-occurring amino acids, *e.g.* cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, the peptide may be linear; in other embodiments, the peptide may be cyclic, *e.g.* bicyclic.

iv. Nucleic Acid Constructs

[000377] Any suitable gene expression construct may be used as a molecular payload, as described herein. In some embodiments, a gene expression construct may be a vector or a cDNA fragment. In some embodiments, a gene expression construct may be messenger RNA (mRNA). In some embodiments, a mRNA used herein may be a modified mRNA, *e.g.*, as described in US Patent 8,710,200, issued on April 24, 2014, entitled “*Engineered nucleic acids encoding a modified erythropoietin and their expression*”. In some embodiments, a mRNA may comprise a 5'-methyl cap. In some embodiments, a mRNA may comprise a polyA tail, optionally of up to 160 nucleotides in length. A gene expression construct may encode a sequence of a protein that preferentially binds to a nucleic acid, *e.g.* a disease-associated repeat, or a protein, *e.g.* MBNL1, found in muscle cells. In some embodiments, the gene expression construct may be expressed, *e.g.*, overexpressed, within the nucleus of a muscle cell. In some embodiments, the gene expression construct encodes a MBNL protein, *e.g.*, MBNL1. In some embodiments, the gene expression construct encodes a protein that comprises at least one zinc finger. In some embodiments, the gene expression construct encodes a protein that binds to a

disease-associated repeat. In some embodiments, the gene expression construct encodes a protein that leads to a reduction in the expression of a disease-associated repeat. In some embodiments, the gene expression construct encodes a gene editing enzyme. Additional examples of nucleic acid constructs that may be used as molecular payloads are provided in International Patent Application Publication WO2017152149A1, published on September 19, 2017, entitled, “*Closed-Ended Linear Duplex Dna For Non-Viral Gene Transfer*”; US Patent 8,853,377B2, issued on October 7, 2014, entitled, “*mRNA For Use In Treatment Of Human Genetic Diseases*”; and US Patent US8822663B2, issued on September 2, 2014, *Engineered Nucleic Acids And Methods Of Use Thereof*,” the contents of each of which are incorporated herein by reference in their entireties.

C. Linkers

[000378] Complexes described herein generally comprise a linker that connects any one of the anti-TfR antibodies described herein to a molecular payload. A linker comprises at least one covalent bond. In some embodiments, a linker may be a single bond, e.g., a disulfide bond or disulfide bridge, that connects an anti-TfR antibody to a molecular payload. However, in some embodiments, a linker may connect any one of the anti-TfR antibodies described herein to a molecular payload through multiple covalent bonds. In some embodiments, a linker may be a cleavable linker. However, in some embodiments, a linker may be a non-cleavable linker. A linker is generally stable *in vitro* and *in vivo*, and may be stable in certain cellular environments. Additionally, generally a linker does not negatively impact the functional properties of either the anti-TfR antibody or the molecular payload. Examples and methods of synthesis of linkers are known in the art (see, e.g. Kline, T. et al. “Methods to Make Homogenous Antibody Drug Conjugates.” *Pharmaceutical Research*, 2015, 32:11, 3480–3493.; Jain, N. et al. “Current ADC Linker Chemistry” *Pharm Res.* 2015, 32:11, 3526–3540.; McCombs, J.R. and Owen, S.C. “Antibody Drug Conjugates: Design and Selection of Linker, Payload and Conjugation Chemistry” *AAPS J.* 2015, 17:2, 339–351.).

[000379] A precursor to a linker typically will contain two different reactive species that allow for attachment to both the anti-TfR antibody and a molecular payload. In some embodiments, the two different reactive species may be a nucleophile and/or (e.g., and) an electrophile. In some embodiments, a linker is connected to an anti-TfR antibody via conjugation to a lysine residue or a cysteine residue of the anti-TfR antibody. In some embodiments, a linker is connected to a cysteine residue of an anti-TfR antibody via a maleimide-containing linker, wherein optionally the maleimide-containing linker comprises a maleimidocaproyl or maleimidomethyl cyclohexane-1-carboxylate group. In some embodiments, a linker is connected to a cysteine residue of an anti-TfR antibody or thiol

functionalized molecular payload via a 3-arylpropionitrile functional group. In some embodiments, a linker is connected to a lysine residue of an anti-TfR antibody. In some embodiments, a linker is connected to an anti-TfR antibody and/or (e.g., and) a molecular payload via an amide bond, a carbamate bond, a hydrazide, a triazole, a thioether, or a disulfide bond.

i. Cleavable Linkers


[000380] A cleavable linker may be a protease-sensitive linker, a pH-sensitive linker, or a glutathione-sensitive linker. These linkers are generally cleavable only intracellularly and are preferably stable in extracellular environments, e.g. extracellular to a muscle cell.

[000381] Protease-sensitive linkers are cleavable by protease enzymatic activity. These linkers typically comprise peptide sequences and may be 2-10 amino acids, about 2-5 amino acids, about 5-10 amino acids, about 10 amino acids, about 5 amino acids, about 3 amino acids, or about 2 amino acids in length. In some embodiments, a peptide sequence may comprise naturally-occurring amino acids, e.g. cysteine, alanine, or non-naturally-occurring or modified amino acids. Non-naturally occurring amino acids include β -amino acids, homo-amino acids, proline derivatives, 3-substituted alanine derivatives, linear core amino acids, N-methyl amino acids, and others known in the art. In some embodiments, a protease-sensitive linker comprises a valine-citrulline or alanine-citrulline dipeptide sequence. In some embodiments, a protease-sensitive linker can be cleaved by a lysosomal protease, e.g. cathepsin B, and/or (e.g., and) an endosomal protease.


[000382] A pH-sensitive linker is a covalent linkage that readily degrades in high or low pH environments. In some embodiments, a pH-sensitive linker may be cleaved at a pH in a range of 4 to 6. In some embodiments, a pH-sensitive linker comprises a hydrazone or cyclic acetal. In some embodiments, a pH-sensitive linker is cleaved within an endosome or a lysosome.

[000383] In some embodiments, a glutathione-sensitive linker comprises a disulfide moiety. In some embodiments, a glutathione-sensitive linker is cleaved by a disulfide exchange reaction with a glutathione species inside a cell. In some embodiments, the disulfide moiety further comprises at least one amino acid, e.g. a cysteine residue.


[000384] In some embodiments, the linker is a Val-cit linker (e.g., as described in US Patent 6,214,345, incorporated herein by reference). In some embodiments, before conjugation, the val-cit linker has a structure of:



[000385] In some embodiments, after conjugation, the val-cit linker has a structure of:




[000386] In some embodiments, the Val-cit linker is attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation). In some embodiments, before click chemistry conjugation, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) has the structure of:




wherein n is any number from 0-10. In some embodiments, n is 3.

[000387] In some embodiments, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) is conjugated (e.g., via a different chemical moiety) to a molecular payload (e.g., an oligonucleotide). In some embodiments, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) and conjugated to a molecular payload (e.g., an oligonucleotide) has the structure of (before click chemistry conjugation):



wherein n is any number from 0-10. In some embodiments, n is 3.

[000388] In some embodiments, after conjugation to a molecular payload (e.g., an oligonucleotide), the val-cit linker has a structure of:



wherein n is any number from 0-10, and wherein m is any number from 0-10. In some embodiments, n is 3 and m is 4.

ii. Non-Cleavable Linkers

[000389] In some embodiments, non-cleavable linkers may be used. Generally, a non-cleavable linker cannot be readily degraded in a cellular or physiological environment. In some embodiments, a non-cleavable linker comprises an optionally substituted alkyl group, wherein the substitutions may include halogens, hydroxyl groups, oxygen species, and other common substitutions. In some embodiments, a linker may comprise an optionally substituted alkyl, an optionally substituted alkylene, an optionally substituted arylene, a heteroarylene, a peptide sequence comprising at least one non-natural amino acid, a truncated glycan, a sugar or sugars that cannot be enzymatically degraded, an azide, an alkyne-azide, a peptide sequence comprising a LPXT sequence, a thioether, a biotin, a biphenyl, repeating units of polyethylene glycol or equivalent compounds, acid esters, acid amides, sulfamides, and/or (e.g., and) an alkoxy-amine linker. In some embodiments, sortase-mediated ligation will be utilized to covalently link an

anti-TfR antibody comprising a LPXT sequence to a molecular payload comprising a (G)_n sequence (see, e.g. Proft T. Sortase-mediated protein ligation: an emerging biotechnology tool for protein modification and immobilization. *Biotechnol Lett.* 2010, 32(1):1-10.).

[000390] In some embodiments, a linker may comprise a substituted alkylene, an optionally substituted alkenylene, an optionally substituted alkynylene, an optionally substituted cycloalkylene, an optionally substituted cycloalkenylene, an optionally substituted arylene, an optionally substituted heteroarylene further comprising at least one heteroatom selected from N, O, and S,; an optionally substituted heterocyclylene further comprising at least one heteroatom selected from N, O, and S,; an imino, an optionally substituted nitrogen species, an optionally substituted oxygen species O, an optionally substituted sulfur species, or a poly(alkylene oxide), e.g. polyethylene oxide or polypropylene oxide.

iii. Linker conjugation

[000391] In some embodiments, a linker is connected to an anti-TfR antibody and/or (e.g., and) molecular payload via a phosphate, thioether, ether, carbon-carbon, carbamate, or amide bond. In some embodiments, a linker is connected to an oligonucleotide through a phosphate or phosphorothioate group, e.g. a terminal phosphate of an oligonucleotide backbone. In some embodiments, a linker is connected to an anti-TfR antibody, through a lysine or cysteine residue present on the anti-TfR antibody.

[000392] In some embodiments, a linker is connected to an anti-TfR antibody and/or (e.g., and) molecular payload by a cycloaddition reaction between an azide and an alkyne to form a triazole, wherein the azide and the alkyne may be located on the anti-TfR antibody, molecular payload, or the linker. In some embodiments, an alkyne may be a cyclic alkyne, e.g., a cyclooctyne. In some embodiments, an alkyne may be bicyclononyne (also known as bicyclo[6.1.0]nonyne or BCN) or substituted bicyclononyne. In some embodiments, a cyclooctane is as described in International Patent Application Publication WO2011136645, published on November 3, 2011, entitled, “*Fused Cyclooctyne Compounds And Their Use In Metal-free Click Reactions*”. In some embodiments, an azide may be a sugar or carbohydrate molecule that comprises an azide. In some embodiments, an azide may be 6-azido-6-deoxygalactose or 6-azido-N-acetylgalactosamine. In some embodiments, a sugar or carbohydrate molecule that comprises an azide is as described in International Patent Application Publication WO2016170186, published on October 27, 2016, entitled, “*Process For The Modification Of A Glycoprotein Using A Glycosyltransferase That Is Or Is Derived From A β (1,4)-N-Acetylgalactosaminyltransferase*”. In some embodiments, a cycloaddition reaction between an azide and an alkyne to form a triazole, wherein the azide and the alkyne may be located on the anti-TfR antibody, molecular payload, or the linker is as described in International


Patent Application Publication WO2014065661, published on May 1, 2014, entitled, “*Modified antibody, antibody-conjugate and process for the preparation thereof*”; or International Patent Application Publication WO2016170186, published on October 27, 2016, entitled, “*Process For The Modification Of A Glycoprotein Using A Glycosyltransferase That Is Or Is Derived From A $\beta(1,4)$ -N-Acetylgalactosaminyltransferase*”.

[000393] In some embodiments, a linker further comprises a spacer, *e.g.*, a polyethylene glycol spacer or an acyl/carbomoyl sulfamide spacer, *e.g.*, a HydraSpaceTM spacer. In some embodiments, a spacer is as described in Verkade, J.M.M. et al., “*A Polar Sulfamide Spacer Significantly Enhances the Manufacturability, Stability, and Therapeutic Index of Antibody-Drug Conjugates*”, *Antibodies*, 2018, 7, 12.

[000394] In some embodiments, a linker is connected to an anti-TfR antibody and/or (*e.g.*, and) molecular payload by the Diels-Alder reaction between a dienophile and a diene/hetero-diene, wherein the dienophile and the diene/hetero-diene may be located on the anti-TfR antibody, molecular payload, or the linker. In some embodiments a linker is connected to an anti-TfR antibody and/or (*e.g.*, and) molecular payload by other pericyclic reactions, *e.g.* ene reaction. In some embodiments, a linker is connected to an anti-TfR antibody and/or (*e.g.*, and) molecular payload by an amide, thioamide, or sulfonamide bond reaction. In some embodiments, a linker is connected to an anti-TfR antibody and/or (*e.g.*, and) molecular payload by a condensation reaction to form an oxime, hydrazone, or semicarbazide group existing between the linker and the anti-TfR antibody and/or (*e.g.*, and) molecular payload.


[000395] In some embodiments, a linker is connected to an anti-TfR antibody and/or (*e.g.*, and) molecular payload by a conjugate addition reactions between a nucleophile, *e.g.* an amine or a hydroxyl group, and an electrophile, *e.g.* a carboxylic acid, carbonate, or an aldehyde. In some embodiments, a nucleophile may exist on a linker and an electrophile may exist on an anti-TfR antibody or molecular payload prior to a reaction between a linker and an anti-TfR antibody or molecular payload. In some embodiments, an electrophile may exist on a linker and a nucleophile may exist on an anti-TfR antibody or molecular payload prior to a reaction between a linker and an anti-TfR antibody or molecular payload. In some embodiments, an electrophile may be an azide, pentafluorophenyl, a silicon centers, a carbonyl, a carboxylic acid, an anhydride, an isocyanate, a thioisocyanate, a succinimidyl ester, a sulfosuccinimidyl ester, a maleimide, an alkyl halide, an alkyl pseudohalide, an epoxide, an episulfide, an aziridine, an aryl, an activated phosphorus center, and/or (*e.g.*, and) an activated sulfur center. In some embodiments, a nucleophile may be an optionally substituted alkene, an optionally substituted alkyne, an optionally substituted aryl, an optionally substituted heterocyclyl, a hydroxyl group, an amino group, an alkylamino group, an anilido group, or a thiol group.

[000396] In some embodiments, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) is conjugated to the anti-TfR antibody by a structure of:




wherein m is any number from 0-10. In some embodiments, m is 4.

[000397] In some embodiments, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) is conjugated to an anti-TfR antibody having a structure of:




wherein m is any number from 0-10. In some embodiments, m is 4.

[000398] In some embodiments, the val-cit linker attached to a reactive chemical moiety (e.g., SPAAC for click chemistry conjugation) and conjugated to an anti-TfR antibody has a structure of:




wherein n is any number from 0-10, wherein m is any number from 0-10. In some embodiments, n is 3 and/or (e.g., and) m is 4.

[000399] In some embodiments, the val-cit linker that links the antibody and the molecular payload has a structure of:




wherein n is any number from 0-10, wherein m is any number from 0-10. In some embodiments, n is 3 and/or (e.g., and) m is 4. In some embodiments, n is 3 and/or (e.g., and) m is 4. In some embodiments, X is NH (e.g., NH from an amine group of a lysine), S (e.g., S from a thiol group of a cysteine), or O (e.g., O from a hydroxyl group of a serine, threonine, or tyrosine) of the antibody.














[000400] In some embodiments, the val-cit linker used to covalently link an anti-TfR antibody and a molecular payload (e.g., an oligonucleotide) has a structure of:




wherein n is any number from 0-10, wherein m is any number from 0-10. In some embodiments, n is 3 and m is 4.

[000401] In structures formula (A), (B), (C), and (D), L1, in some embodiments, is a spacer that is a substituted or unsubstituted aliphatic, substituted or unsubstituted heteroaliphatic, substituted or unsubstituted carbocyclylene, substituted or unsubstituted heterocyclylene, substituted or unsubstituted arylene, substituted or unsubstituted heteroarylene, -O-, -N(R^A)-, -S-, -C(=O)-, -C(=O)O-, -C(=O)NR^A-, -NR^AC(=O)-, -NR^AC(=O)R^A-, -C(=O)R^A-, -NR^AC(=O)O-, -NR^AC(=O)N(R^A)-, -OC(=O)-, -OC(=O)O-, -OC(=O)N(R^A)-, -S(O)₂NR^A-, -NR^AS(O)₂-, or a combination thereof. In some embodiments, L1 is




wherein the piperazine moiety links to the oligonucleotide, wherein L2 is ,
,
,
,
,
,
,
,
,
,
,
,
,
or .

[000402] In some embodiments, L1 is:




wherein the piperazine moiety links to the oligonucleotide.

[000403] In some embodiments, L1 is .

[000404] In some embodiments, L1 is linked to a 5' phosphate of the oligonucleotide.

[000405] In some embodiments, L1 is optional (e.g., need not be present).

[000406] In some embodiments, any one of the complexes described herein has a structure of:




wherein n is 0-15 (e.g., 3) and m is 0-15 (e.g., 4).

C. Examples of Antibody-Molecular Payload Complexes


[000407] Further provided herein are non-limiting examples of complexes comprising any one the anti-TfR antibodies described herein covalently linked to any of the molecular payloads (e.g., an oligonucleotide) described herein. In some embodiments, the anti-TfR antibody (e.g., any one of the anti-TfR antibodies provided in Table 2) is covalently linked to a molecular payload (e.g., an oligonucleotide) via a linker. Any of the linkers described herein may be used. In some embodiments, if the molecular payload is an oligonucleotide, the linker is linked to the 5' end, the 3' end, or internally of the oligonucleotide. In some embodiments, the linker is linked to the anti-TfR antibody via a thiol-reactive linkage (e.g., via a cysteine in the anti-TfR antibody). In some embodiments, the linker (e.g., a Val-cit linker) is linked to the antibody (e.g., an anti-TfR antibody described herein) via an amine group (e.g., via a lysine in the antibody). In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000408] An example of a structure of a complex comprising an anti-TfR antibody covalently linked to a molecular payload via a Val-cit linker is provided below:



wherein the linker is linked to the antibody via a thiol-reactive linkage (e.g., via a cysteine in the antibody). In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000409] Another example of a structure of a complex comprising an anti-TfR antibody covalently linked to a molecular payload via a Val-cit linker is provided below:



wherein n is a number between 0-10, wherein m is a number between 0-10, wherein the linker is linked to the antibody via an amine group (e.g., on a lysine residue), and/or (e.g., and) wherein the linker is linked to the oligonucleotide (e.g., at the 5' end, 3' end, or internally). In some embodiments, the linker is linked to the antibody via a lysine, the linker is linked to the oligonucleotide at the 5' end, n is 3, and m is 4. In some embodiments, the molecular payload is an oligonucleotide comprising a sense strand and an antisense strand, and, the linker is linked to the sense strand or the antisense strand at the 5' end or the 3' end. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000410] It should be appreciated that antibodies can be linked to molecular payloads with different stoichiometries, a property that may be referred to as a drug to antibody ratios (DAR) with the “drug” being the molecular payload. In some embodiments, one molecular payload is linked to an antibody (DAR = 1). In some embodiments, two molecular payloads are linked to an antibody (DAR = 2). In some embodiments, three molecular payloads are linked to an antibody (DAR = 3). In some embodiments, four molecular payloads are linked to an antibody (DAR = 4). In some embodiments, a mixture of different complexes, each having a different DAR, is provided. In some embodiments, an average DAR of complexes in such a mixture may be in a range of 1 to 3, 1 to 4, 1 to 5 or more. DAR may be increased by conjugating molecular payloads to different sites on an antibody and/or (e.g., and) by conjugating multimers to one or more sites on antibody. For example, a DAR of 2 may be achieved by conjugating a single

molecular payload to two different sites on an antibody or by conjugating a dimer molecular payload to a single site of an antibody.

[000411] In some embodiments, the complex described herein comprises an anti-TfR antibody described herein (e.g., the 3-A4, 3-M12, and 5-H12 antibodies provided in Table 2 in an IgG or Fab form) covalently linked to a molecular payload. In some embodiments, the complex described herein comprises an anti-TfR antibody described herein (e.g., the 3-A4, 3-M12, and 5-H12 antibodies provided in Table 2 in a IgG or Fab form) covalently linked to molecular payload via a linker (e.g., a Val-cit linker). In some embodiments, the linker (e.g., a Val-cit linker) is linked to the antibody (e.g., an anti-TfR antibody described herein) via a thiol-reactive linkage (e.g., via a cysteine in the antibody). In some embodiments, the linker (e.g., a Val-cit linker) is linked to the antibody (e.g., an anti-TfR antibody described herein) via an amine group (e.g., via a lysine in the antibody). In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000412] In some embodiments, in any one of the examples of complexes described herein, the molecular payload a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000413] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a CDR-H1, a CDR-H2, and a CDR-H3 that are the same as the CDR-H1, CDR-H2, and CDR-H3 shown in Table 2; and a CDR-L1, a CDR-L2, and a CDR-L3 that are the same as the CDR-L1, CDR-L2, and CDR-L3 shown in Table 2. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000414] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 69, SEQ ID NO: 71, or SEQ ID NO: 72, and a VL comprising the amino acid sequence of SEQ ID NO: 70. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000415] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 73 or SEQ ID NO: 76, and a VL comprising the amino acid sequence of SEQ ID NO: 74. In some embodiments, the molecular

payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000416] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 73 or SEQ ID NO: 76, and a VL comprising the amino acid sequence of SEQ ID NO: 75. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000417] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 77, and a VL comprising the amino acid sequence of SEQ ID NO: 78. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000418] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 77 or SEQ ID NO: 79, and a VL comprising the amino acid sequence of SEQ ID NO: 80. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000419] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 84, SEQ ID NO: 86 or SEQ ID NO: 87 and a light chain comprising the amino acid sequence of SEQ ID NO: 85. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000420] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 88 or SEQ ID NO: 91, and a light chain comprising the amino acid sequence of SEQ ID NO: 89. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000421] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 88 or SEQ ID NO: 91, and a light chain comprising the amino acid sequence of SEQ ID NO: 90. In some embodiments, the

molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000422] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92 or SEQ ID NO: 94, and a light chain comprising the amino acid sequence of SEQ ID NO: 95. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000423] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92, and a light chain comprising the amino acid sequence of SEQ ID NO: 93. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000424] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 97, SEQ ID NO: 98, or SEQ ID NO: 99 and a VL comprising the amino acid sequence of SEQ ID NO: 85. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000425] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 or SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 89. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).


[000426] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 or SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 90. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000427] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain

comprising the amino acid sequence of SEQ ID NO: 93. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).


[000428] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked to a molecular payload, wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 or SEQ ID NO: 103 and a light chain comprising the amino acid sequence of SEQ ID NO: 95. In some embodiments, the molecular payload is a DMPK-targeting oligonucleotide (e.g., a DMPK-targeting oligonucleotide listed in Table 8 or Table 17).

[000429] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 84 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000430] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 86 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264). In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 87 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000431] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising


wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000433] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of in SEQ ID NO: 89; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000434] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of in SEQ ID NO: 90; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the


oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000435] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92 and a light chain comprising the amino acid sequence of in SEQ ID NO: 93; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000436] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 94 and a light chain comprising the amino acid sequence of in SEQ ID NO: 95; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a


gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000437] In some embodiments, the complex described herein comprises an anti-TfR antibody covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 92 and a light chain comprising the amino acid sequence of in SEQ ID NO: 95; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000438] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 69 and a VL comprising the amino acid sequence of in SEQ ID NO: 70; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000439] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 71 and a VL comprising the amino acid sequence of in SEQ ID NO: 70; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000440] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence

of SEQ ID NO: 72 and a VL comprising the amino acid sequence of in SEQ ID NO: 70; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000441] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 73 and a VL comprising the amino acid sequence of in SEQ ID NO: 74; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000442] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 73 and a VL comprising the amino acid sequence of in SEQ ID NO: 75; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000443] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of in SEQ ID NO: 74; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the


oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000444] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of in SEQ ID NO: 75; wherein the complex has the structure of:




gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000446] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 79 and a VL comprising the amino acid sequence of in SEQ ID NO: 80; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000447] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a VH comprising the amino acid sequence of SEQ ID NO: 77 and a VL comprising the amino acid sequence of in SEQ ID NO: 80; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000448] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 97 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000449] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid

sequence of SEQ ID NO: 98 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000450] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 99 and a light chain comprising the amino acid sequence of in SEQ ID NO: 85; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000451] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of in SEQ ID NO: 89; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000452] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of in SEQ ID NO: 90; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the


oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000453] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of in SEQ ID NO: 89; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000454] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of in SEQ ID NO: 90; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a


gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000455] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of in SEQ ID NO: 93; wherein the complex has the structure of:




wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).

[000456] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 103 and a light chain comprising the amino acid sequence of in SEQ ID NO: 95; wherein the complex has the structure of:



wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000457] In some embodiments, the complex described herein comprises an anti-TfR Fab covalently linked via a lysine to the 5' end of an oligonucleotide (e.g., a DMPK-targeting oligonucleotide), wherein the anti-TfR Fab comprises a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of in SEQ ID NO: 95; wherein the complex has the structure of:
















wherein n is 3 and m is 4, optionally wherein the DMPK-targeting oligonucleotide (e.g., a gapmer) comprises at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides (e.g., 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 consecutive nucleotides) of any one of the oligonucleotides listed in Table 8 or Table 17 (e.g., any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264).


[000458] In some embodiments, in any one of the examples of complexes described herein, L1 is any one of the spacers described herein.

[000459] In some embodiments, L1 is:




wherein the piperazine moiety links to the oligonucleotide, wherein L2 is , , , , , , , , , , , , , or .

[000460] In some embodiments, L1 is:



wherein the piperazine moiety links to the oligonucleotide.

[000461] In some embodiments, L1 is .

[000462] In some embodiments, L1 is linked to a 5' phosphate of the oligonucleotide.

[000463] In some embodiments, L1 is optional (e.g., need not be present).

III. Formulations

[000464] Complexes provided herein may be formulated in any suitable manner.

Generally, complexes provided herein are formulated in a manner suitable for pharmaceutical use. For example, complexes can be delivered to a subject using a formulation that minimizes degradation, facilitates delivery and/or (e.g., and) uptake, or provides another beneficial property to the complexes in the formulation. In some embodiments, provided herein are compositions comprising complexes and pharmaceutically acceptable carriers. Such compositions can be suitably formulated such that when administered to a subject, either into the immediate

environment of a target cell or systemically, a sufficient amount of the complexes enter target muscle cells. In some embodiments, complexes are formulated in buffer solutions such as phosphate-buffered saline solutions, liposomes, micellar structures, and capsids.

[000465] It should be appreciated that, in some embodiments, compositions may include separately one or more components of complexes provided herein (e.g., muscle-targeting agents, linkers, molecular payloads, or precursor molecules of any one of them).

[000466] In some embodiments, complexes are formulated in water or in an aqueous solution (e.g., water with pH adjustments). In some embodiments, complexes are formulated in basic buffered aqueous solutions (e.g., PBS). In some embodiments, formulations as disclosed herein comprise an excipient. In some embodiments, an excipient confers to a composition improved stability, improved absorption, improved solubility and/or (e.g., and) therapeutic enhancement of the active ingredient. In some embodiments, an excipient is a buffering agent (e.g., sodium citrate, sodium phosphate, a tris base, or sodium hydroxide) or a vehicle (e.g., a buffered solution, petrolatum, dimethyl sulfoxide, or mineral oil).

[000467] In some embodiments, a complex or component thereof (e.g., oligonucleotide or antibody) is lyophilized for extending its shelf-life and then made into a solution before use (e.g., administration to a subject). Accordingly, an excipient in a composition comprising a complex, or component thereof, described herein may be a lyoprotectant (e.g., mannitol, lactose, polyethylene glycol, or polyvinyl pyrrolidone), or a collapse temperature modifier (e.g., dextran, ficoll, or gelatin).

[000468] In some embodiments, a pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, administration. Typically, the route of administration is intravenous or subcutaneous.

[000469] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. In some embodiments, formulations include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, and sodium chloride in the composition. Sterile injectable solutions can be prepared by incorporating the complexes in a required amount in a selected solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization.

[000470] In some embodiments, a composition may contain at least about 0.1% of the complex, or component thereof, or more, although the percentage of the active ingredient(s) may be between about 1% and about 80% or more of the weight or volume of the total composition. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

IV. Methods of Use / Treatment

[000471] Complexes comprising a muscle-targeting agent covalently linked to a molecular payload as described herein are effective in treating myotonic dystrophy. In some embodiments, complexes are effective in treating myotonic dystrophy type 1 (DM1). In some embodiments, DM1 is associated with an expansion of a CTG trinucleotide repeat in the 3' non-coding region of DMPK. In some embodiments, the nucleotide expansions lead to toxic RNA repeats capable of forming hairpin structures that bind critical intracellular proteins, e.g., muscleblind-like proteins, with high affinity.

[000472] In some embodiments, a subject may be a human subject, a non-human primate subject, a rodent subject, or any suitable mammalian subject. In some embodiments, a subject may have myotonic dystrophy. In some embodiments, a subject has a DMPK allele, which may optionally contain a disease-associated repeat. In some embodiments, a subject may have a DMPK allele with an expanded disease-associated-repeat that comprises about 2-10 repeat units, about 2-50 repeat units, about 2-100 repeat units, about 50-1,000 repeat units, about 50-500 repeat units, about 50-250 repeat units, about 50-100 repeat units, about 500-10,000 repeat units, about 500-5,000 repeat units, about 500-2,500 repeat units, about 500-1,000 repeat units, or about 1,000-10,000 repeat units. In some embodiments, a subject is suffering from symptoms of DM1, e.g. muscle atrophy or muscle loss. In some embodiments, a subject is not suffering from symptoms of DM1. In some embodiments, subjects have congenital myotonic dystrophy.

[000473] An aspect of the disclosure includes a method involving administering to a subject an effective amount of a complex as described herein. In some embodiments, an effective amount of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload can be administered to a subject in need of treatment. In some embodiments, a pharmaceutical composition comprising a complex as described herein may be administered by a suitable route, which may include intravenous administration, e.g., as a bolus or by continuous infusion over a period of time. In some embodiments, intravenous administration may be performed by intramuscular,

intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, or intrathecal routes. In some embodiments, a pharmaceutical composition may be in solid form, aqueous form, or a liquid form. In some embodiments, an aqueous or liquid form may be nebulized or lyophilized. In some embodiments, a nebulized or lyophilized form may be reconstituted with an aqueous or liquid solution.

[000474] Compositions for intravenous administration may contain various carriers such as vegetable oils, dimethylactamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, and polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like). For intravenous injection, water soluble antibodies can be administered by the drip method, whereby a pharmaceutical formulation containing the antibody and a physiologically acceptable excipients is infused. Physiologically acceptable excipients may include, for example, 5% dextrose, 0.9% saline, Ringer's solution or other suitable excipients. Intramuscular preparations, e.g., a sterile formulation of a suitable soluble salt form of the antibody, can be dissolved and administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose solution.

[000475] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload is administered via site-specific or local delivery techniques. Examples of these techniques include implantable depot sources of the complex, local delivery catheters, site specific carriers, direct injection, or direct application.

[000476] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload is administered at an effective concentration that confers therapeutic effect on a subject. Effective amounts vary, as recognized by those skilled in the art, depending on the severity of the disease, unique characteristics of the subject being treated, e.g. age, physical conditions, health, or weight, the duration of the treatment, the nature of any concurrent therapies, the route of administration and related factors. These related factors are known to those in the art and may be addressed with no more than routine experimentation. In some embodiments, an effective concentration is the maximum dose that is considered to be safe for the patient. In some embodiments, an effective concentration will be the lowest possible concentration that provides maximum efficacy.

[000477] Empirical considerations, e.g. the half-life of the complex in a subject, generally will contribute to determination of the concentration of pharmaceutical composition that is used for treatment. The frequency of administration may be empirically determined and adjusted to maximize the efficacy of the treatment.

[000478] Generally, for administration of any of the complexes described herein, an initial candidate dosage may be about 1 to 100 mg/kg, or more, depending on the factors described above, e.g. safety or efficacy. In some embodiments, a treatment will be administered once. In some embodiments, a treatment will be administered daily, biweekly, weekly, bimonthly, monthly, or at any time interval that provide maximum efficacy while minimizing safety risks to the subject. Generally, the efficacy and the treatment and safety risks may be monitored throughout the course of treatment.

[000479] In some embodiments, an initial candidate dosage is about 1-50, 1-25, 1-10, 1-5, 5-100, 5-50, 5-25, 5-10, 10-100, 10-75, 10-50, 10-25, 10-20, 25-100, 25-75, or 25-50 mg/kg. In some embodiments, an initial candidate dosage is about 1-20, 1-15, 1-10, 1-5, 1-3, 1-2, 5-20, 5-15, or 5-10 mg/kg. In some embodiments, an initial candidate dosage is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, or 20 mg/kg.

[000480] The efficacy of treatment may be assessed using any suitable methods. In some embodiments, the efficacy of treatment may be assessed by evaluation of observation of symptoms associated with DM1, e.g. muscle atrophy or muscle weakness, through measures of a subject's self-reported outcomes, e.g. mobility, self-care, usual activities, pain/discomfort, and anxiety/depression, or by quality-of-life indicators, e.g. lifespan.

[000481] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein is administered to a subject at an effective concentration sufficient to inhibit activity or expression of a target gene by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% relative to a control, e.g. baseline level of gene expression prior to treatment.

[000482] In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1-5, 1-10, 5-15, 10-20, 15-30, 20-40, 25-50, or more days. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15, 20, or 24 weeks. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1-5, 1-10, 2-5, 2-10, 4-8, 4-12, 5-10, 5-12, 5-15, 8-12, 8-15, 10-12, 10-15, 10-20, 12-15, 12-20, 15-20, or 15-25 weeks. In

some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject is sufficient to inhibit activity or expression of a target gene for at least 1, 2, 3, 4, 5, or 6 months.

[000483] In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject persists or remains in the subject for at least 1-5, 1-10, 5-15, 10-20, 15-30, 20-40, 25-50, or more days. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject persists or remains in the subject for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15, 20, or 24 weeks. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject persists or remains in the subject for at least 1-5, 1-10, 2-5, 2-10, 4-8, 4-12, 5-10, 5-12, 5-15, 8-12, 8-15, 10-12, 10-15, 10-20, 12-15, 12-20, 15-20, or 15-25 weeks. In some embodiments, a single dose or administration of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein to a subject persists or remains in the subject for at least 1, 2, 3, 4, 5, or 6 months.

[000484] In some embodiments, multiple doses or administrations of a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein are delivered to a subject. In some embodiments, multiple doses of a pharmaceutical composition comprise delivering 2, 3, 4, 5, 6, 7, 8, 9, or 10 doses to a subject. In some embodiments, multiple doses of a pharmaceutical composition comprise delivering a dose to a subject every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 weeks. In some embodiments, multiple doses of a pharmaceutical composition comprise delivering a dose to a subject once every 4 weeks. In some embodiments, multiple doses of a pharmaceutical composition comprise delivering a dose to a subject once every 1-10, 2-5, 2-10, 4-8, 4-12, 5-10, 5-12, 5-15, 8-12, 8-16, 10-12, 10-15, 10-20, 12-15, 12-20, 15-20, or 15-25 weeks. In some embodiments, multiple doses of a pharmaceutical composition comprise delivering a dose to a subject on a biweekly (*i.e.*, every two weeks), bimonthly (*i.e.*, every two months), or quarterly schedule (*i.e.*, every twelve weeks).

[000485] In some embodiments, a single dose or administration is about 1-50, 1-25, 1-10, 1-15, 1-5, 5-100, 5-50, 5-25, 5-10, 10-100, 10-75, 10-50, 10-25, 10-20, 25-100, 25-75, or 25-50 mg/kg. In some embodiments, a single dose or administration is about 1-20, 1-15, 1-10, 1-5, 1-

3, 1-2, 5-20, 5-15, or 5-10 mg/kg. In some embodiments, a single dose or administration is about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 20 mg/kg.

[000486] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein is delivered to a subject every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 weeks. In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein is delivered to a subject every 1-10, 2-5, 2-10, 4-8, 4-12, 5-10, 5-12, 5-15, 8-12, 8-16, 9-15, 10-12, 10-14, 10-15, 10-20, 11-13, 11-15, 12-15, 12-16, 12-20, 15-20, or 15-25 weeks. In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein is delivered to a subject on a biweekly (*i.e.*, every two weeks), bimonthly (*i.e.*, every two months), or quarterly schedule (*i.e.*, every twelve weeks).

[000487] In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein at a concentration of 1-15 mg/kg of RNA is delivered to a subject every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 weeks. In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein at a concentration of 1-15 mg/kg of RNA is delivered to a subject every 1-10, 2-5, 2-10, 4-8, 4-12, 5-10, 5-12, 5-15, 8-12, 8-16, 9-15, 10-12, 10-14, 10-15, 10-20, 11-13, 11-15, 12-15, 12-16, 12-20, 15-20, or 15-25 weeks. In some embodiments, a pharmaceutical composition that comprises a complex comprising a muscle-targeting agent covalently linked to a molecular payload described herein at a concentration of 1-15 mg/kg of RNA is delivered to a subject on a biweekly (*i.e.*, every two weeks), bimonthly (*i.e.*, every two months), or quarterly schedule (*i.e.*, every twelve weeks).

[000488] In some embodiments, a pharmaceutical composition may comprise more than one complex comprising a muscle-targeting agent covalently linked to a molecular payload. In some embodiments, a pharmaceutical composition may further comprise any other suitable therapeutic agent for treatment of a subject, e.g. a human subject having DM1. In some embodiments, the other therapeutic agents may enhance or supplement the effectiveness of the complexes described herein. In some embodiments, the other therapeutic agents may function to treat a different symptom or disease than the complexes described herein.

EXAMPLES

Example 1: Targeting DMPK with transfected antisense oligonucleotides

[000489] A gapmer antisense oligonucleotide that targets both wild-type and mutant alleles of DMPK (ASO300) was tested *in vitro* for its ability to reduce expression levels of DMPK in an immortalized cell line. Briefly, Hepa 1-6 cells were transfected with ASO300 (100 nM) formulated with Lipofectamine 2000. DMPK expression levels were evaluated 72 hours following transfection. A control experiment was also performed in which vehicle (phosphate-buffered saline) was delivered to Hepa 1-6 cells in culture and the cells were maintained for 72 hours. As shown in FIG. 1, it was found that ASO300 reduced DMPK expression levels by ~90% compared with controls.

Example 2: Targeting DMPK with a muscle-targeting complex

[000490] A muscle-targeting complex was generated comprising the DMPK ASO used in Example 1 (ASO300) covalently linked, via a cathepsin cleavable linker, to DTX-A-002 (RI7 217 Fab), an anti-transferrin receptor antibody.

[000491] Briefly, a maleimidocaproyl-L-valine-L-citrulline-p-aminobenzyl alcohol p-nitrophenyl carbonate (MC-Val-Cit-PABC-PNP) linker molecule was coupled to NH₂-C₆-ASO300 using an amide coupling reaction. Excess linker and organic solvents were removed by gel permeation chromatography. The purified Val-Cit-linker-ASO300 was then coupled to a thiol on the anti-transferrin receptor antibody (DTX-A-002).

[000492] The product of the antibody coupling reaction was then subjected to hydrophobic interaction chromatography (HIC-HPLC). FIG. 2A shows a resulting HIC-HPLC chromatogram, in which fractions B7-C2 of the chromatogram (denoted by vertical lines) contained antibody-oligonucleotide complexes (referred to as DTX-C-008) comprising one or two DMPK ASO molecules covalently attached to DTX-A-002, as determined by SDS-PAGE. These HIC-HPLC fractions were combined and densitometry following additional purification confirmed that this sample of DTX-C-008 complexes had an average ASO to antibody ratio (DAR) of 1.48. SDS-PAGE analysis demonstrated that 86.4% of this sample of DTX-C-008 complexes comprised DTX-A-002 linked to either one or two DMPK ASO molecules (FIG. 2B).

[000493] Using the same methods as described above, a control complex was generated comprising the DMPK ASO used in Example 1 (ASO300) covalently linked via a Val-Cit linker to an IgG2a (Fab) antibody (DTX-C-007).

[000494] The purified RI7 217 Fab antibody-ASO complex (DTX-C-008) was then tested for cellular internalization and inhibition of DMPK. Hepa 1-6 cells, which have relatively high expression levels of transferrin receptor, were incubated in the presence of vehicle control, DTX-C-008 (100 nM), or DTX-C-007 (100 nM) for 72 hours. After the 72 hour incubation, the

cells were isolated and assayed for expression levels of DMPK (FIG. 3). Cells treated with the DTX-C-008 demonstrated a reduction in DMPK expression by ~65% relative to the cells treated with the vehicle control. Meanwhile, cells treated with the DTX-C-007 had DMPK expression levels comparable to the vehicle control (no reduction in DMPK expression). These data indicate that the anti-transferrin receptor antibody of the DTX-C-008 enabled cellular internalization of the complex, thereby allowing the DMPK ASO to inhibit expression of DMPK.

Example 3: Targeting DMPK in mouse muscle tissues with a muscle-targeting complex

[000495] The muscle-targeting complex described in Example 2, RI7 217 Fab antibody-ASO complex DTX-C-008, was tested for inhibition of DMPK in mouse tissues. C57BL/6 wild-type mice were intravenously injected with a single dose of a vehicle control, naked ASO300 (3 mg/kg of ASO), DTX-C-008 (3 mg/kg of ASO, corresponding to 20 mg/kg antibody conjugate), or DTX-C-007 IgG2a Fab antibody-ASO complex (3 mg/kg of ASO, corresponding to 20 mg/kg antibody conjugate). Naked ASO300, the DMPK ASO as described in Example 1, was used as a control. Each experimental condition was replicated in three individual C57BL/6 wild-type mice. Following a seven-day period after injection, the mice were euthanized and segmented into isolated tissue types. Individual tissue samples were subsequently assayed for expression levels of DMPK (FIGs. 4A-4E and 5A-5B).

[000496] Mice treated with the DTX-C-008 complex demonstrated a reduction in DMPK expression in a variety of skeletal, cardiac, and smooth muscle tissues. For example, as shown in FIGs 4A-4E, DMPK expression levels were significantly reduced in gastrocnemius (50% reduction), heart (30% reduction), esophagus (45% reduction), tibialis anterior (47% reduction), and soleus (31% reduction) tissues, relative to the mice treated with the vehicle control. Meanwhile, mice treated with the DTX-C-007 complex had DMPK expression levels comparable to the vehicle control mice and mice treated with naked ASO300 (no reduction in DMPK expression) for all assayed muscle tissue types.

[000497] Mice treated with the DTX-C-008 complex demonstrated no change in DMPK expression in non-muscle tissues such as spleen and brain tissues (FIGs. 5A and 5B).

[000498] These data indicate that the anti-transferrin receptor antibody of the DTX-C-008 enabled cellular internalization of the complex into muscle-specific tissues in an *in vivo* mouse model, thereby allowing the DMPK ASO to inhibit expression of DMPK. These data further demonstrate that the DTX-C-008 complex is capable of specifically targeting muscle tissues.

Example 4: Targeting DMPK in mouse muscle tissues with a muscle-targeting complex

[000499] The muscle-targeting complex described in Example 2, RI7 217 Fab antibody-ASO complex DTX-C-008, was tested for dose-dependent inhibition of DMPK in mouse tissues. C57BL/6 wild-type mice were intravenously injected with a single dose of a vehicle control (phosphate-buffered saline, PBS), naked ASO300 (10 mg/kg of ASO), DTX-C-008 (3 mg/kg or 10 mg/kg of ASO, wherein 3 mg/kg corresponds to 20 mg/kg antibody conjugate), or DTX-C-007 IgG2a Fab antibody-ASO complex (3 mg/kg or 10 mg/kg of ASO, wherein 3 mg/kg corresponds to 20 mg/kg antibody conjugate). Naked ASO300, the DMPK ASO as described in Example 1, was used as a control. Each experimental condition was replicated in five individual C57BL/6 wild-type mice. Following a seven-day period after injection, the mice were euthanized and segmented into isolated tissue types. Individual tissue samples were subsequently assayed for expression levels of DMPK (FIGs. 6A-6F).

[000500] Mice treated with the DTX-C-008 complex demonstrated a reduction in DMPK expression in a variety of skeletal muscle tissues. As shown in FIGs 6A-6F, DMPK expression levels were significantly reduced in tibialis anterior (58% and 75% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively), soleus (55% and 66% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively), extensor digitorum longus (EDL) (52% and 72% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively), gastrocnemius (55% and 77% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively), heart (19% and 35% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively), and diaphragm (53% and 70% reduction for 3 mg/kg and 10 mg/kg DTX-C-008, respectively) tissues, relative to the mice treated with the vehicle control. Notably, all assayed muscle tissue types experienced dose-dependent inhibition of DMPK, with greater reduction in DMPK levels at 10 mg/kg antibody conjugate relative to 3 mg/kg antibody conjugate.

[000501] Meanwhile, mice treated with the control DTX-C-007 complex had DMPK expression levels comparable to the vehicle control (no reduction in DMPK expression) for all assayed muscle tissue types. These data indicate that the anti-transferrin receptor antibody of the DTX-C-008 enabled cellular internalization of the complex into muscle-specific tissues in an *in vivo* mouse model, thereby allowing the DMPK ASO to inhibit expression of DMPK. These data further demonstrate that the DTX-C-008 complex is capable of specifically targeting muscle tissues for dose-dependent inhibition of DMPK.

Example 5: Targeting DMPK in cynomolgus monkey muscle tissues with a muscle-targeting complex

[000502] A muscle-targeting complex comprising ASO300 (DTX-C-012), was generated and purified using methods described in Example 2. DTX-C-012 is a complex comprising a

human anti-transferrin receptor antibody covalently linked, via a cathepsin cleavable Val-Cit linker, to ASO300. The anti-TfR antibody used in DTX-C-012 is cross-reactive with cynomolgus and human TfR1. Following HIC-HPLC purification and additional purification, densitometry confirmed that DTX-C-012 had an average ASO to antibody ratio of 1.32, and SDS-PAGE revealed a purity of 92.3%.

[000503] DTX-C-012 was tested for inhibition of DMPK in male cynomolgus monkey tissues. Male cynomolgus monkeys (19-31 months; 2-3 kg) were intravenously injected with a single dose of a saline control, naked ASO300 (10 mg/kg of ASO), or DTX-C-012 (10 mg/kg of ASO) on Day 0. Each experimental condition was replicated in three individual male cynomolgus monkeys. On Day 7 after injection, tissue biopsies (including muscle tissues) were collected. DMPK mRNA expression levels, ASO detection assays, serum clinical chemistries, tissue histology, clinical observations, and body weights were analyzed. The monkeys were euthanized on Day 14.

[000504] Significant knockdown (KD) of DMPK mRNA expression using DTX-C-012 was observed in soleus, deep flexor, and masseter muscles relative to saline control, with 39% KD, 62% KD, and 41% KD, respectively (FIGs. 7A-7C). Robust knockdown of DMPK mRNA expression by DTX-C-012 was further observed in gastrocnemius (62% KD; FIG. 7D), EDL (29% KD; FIG. 7E), tibialis anterior muscle (23% KD; FIG. 7F), diaphragm (54% KD; FIG. 7G), tongue (43% KD; FIG. 7H), heart muscle (36% KD; FIG. 7I), quadriceps (58% KD; FIG. 7J), bicep (51% KD; FIG. 7K), and deltoid muscles (47% KD; FIG. 7L). Knockdown of DMPK mRNA expression by DTX-C-012 in smooth muscle was also observed in the intestine, with 63% KD at jejunum-duodenum ends (FIG. 8A) and 70% KD in ileum (FIG. 8B). Notably, naked DMPK ASO300 (*i.e.*, not linked to a muscle-targeting agent) had minimal effects on DMPK expression levels relative to the vehicle control (*i.e.*, little or no reduction in DMPK expression) for most assayed muscle tissue types. Monkeys treated with the DTX-C-012 complex demonstrated no change in DMPK expression in most non-muscle tissues, such as kidney, brain, and spleen tissues (FIGs. 9A-9D). Additional tissues were examined, as depicted in FIG. 10, which shows normalized DMPK mRNA tissue expression levels across several tissue types in cynomolgus monkeys. (N=3 male cynomolgus monkeys)

[000505] Prior to euthanization, all monkeys were tested for reticulocyte levels, platelet levels, hemoglobin expression, alanine aminotransferase (ALT) expression, aspartate aminotransferase (AST) expression, and blood urea nitrogen (BUN) levels on days 2, 7, and 14 after dosing. As shown in FIG. 12, monkeys dosed with antibody-oligonucleotide complex had normal reticulocyte levels, platelet levels and hemoglobin expression throughout the length of the experiment. Monkeys dosed with DTX-C-012 also had normal alanine aminotransferase

(ALT) expression, aspartate aminotransferase (AST) expression, and blood urea nitrogen (BUN) levels throughout the length of the experiment. These data show that a single dose of a complex comprising ASO300 is safe and tolerated in cynomolgus monkeys.

[000506] These data demonstrate that the anti-transferrin receptor antibody of the DTX-C-012 complex enabled cellular internalization of the complex into muscle-specific tissues in an *in vivo* cynomolgus monkey model, thereby allowing the DMPK ASO (ASO300) to inhibit expression of DMPK. These data further demonstrate that the DTX-C-012 complex is capable of specifically targeting muscle tissues for dose-dependent inhibition of DMPK without substantially impacting non-muscle tissues. This is direct contrast with the limited ability of naked DMPK ASO300 (not linked to a muscle-targeting agent), to inhibit expression of DMPK in muscle tissues of an *in vivo* cynomolgus monkey model.

Example 6: Targeting DMPK in mouse muscle tissues with a muscle-targeting complex

[000507] The RI7 217 Fab antibody-ASO muscle-targeting complex described in Example 2, DTX-C-008, was tested for time-dependent inhibition of DMPK in mouse tissues. C57BL/6 wild-type mice were intravenously injected with a single dose of a vehicle control (saline), naked ASO300 (10 mg/kg of ASO), or DTX-C-008 (10 mg/kg of ASO) and euthanized after a prescribed period of time, as described in Table 13. Following euthanization, the mice were segmented into isolated tissue types and tissue samples were subsequently assayed for expression levels of DMPK (FIGs. 11A-11B).

Table 13. Experimental conditions

Group	Dosage	Days after injection before euthanization	Number of mice
1	Vehicle (saline)	3 days	3
2	Vehicle (saline)	7 days	3
3	Vehicle (saline)	14 days	3
4	Vehicle (saline)	28 days	3
5	ASO300	3 days	3
6	ASO300	7 days	3
7	ASO300	14 days	3
8	ASO300	28 days	3
9	DTX-C-008	3 days	3
10	DTX-C-008	7 days	3
11	DTX-C-008	14 days	3
12	DTX-C-008	28 days	3

[000508] Mice treated with the DTX-C-008 complex demonstrated approximately 50% reduction in DMPK expression in gastrocnemius (FIG. 11A) and tibialis anterior (FIG. 11B) muscles for all of Groups 9-12 (3-28 days between injection and euthanization), relative to

vehicle. Mice treated with the naked ASO300 oligonucleotide did not demonstrate significant reduction in DMPK expression.

[000509] These data indicate that the DTX-C-008 complex was capable of providing persistent reduction in DMPK expression for up to 28 days following dosage of mice with said DTX-C-008 complex.

Example 7: Evaluation of antisense oligonucleotides that target DMPK in immortalized myoblasts

[000510] Two hundred and thirty-six oligonucleotides for targeting DMPK were generated using *in silico* analysis. Each individual oligonucleotide was evaluated for their ability to target DMPK *in cellulo* at two doses – 0.5 nM (low dose) and 50 nM (high dose).

[000511] Briefly, DM1 C15 immortalized myoblasts were cultured in T-75 flasks until near confluency (~80% confluent). Myoblasts were then disrupted with trypsin and seeded into 96-well microplates at a density of 50,000 cells/well. Cells were allowed to recover overnight before the growth media was washed out and replaced with a no-serum media to induce differentiation into myotubes. Differentiation proceeded for seven days prior to treatment with DMPK-targeting oligonucleotides.

[000512] On day seven following induction of differentiation, DM1 C15 myotubes were transfected with an individual oligonucleotide using 0.3 μ L of Lipofectamine MessengerMax per well. All oligonucleotides were tested at both 0.5 nM and 50 nM final concentrations in biological triplicates. After treatment with oligonucleotides, cells were incubated for 72 hours prior to being harvested for total RNA. cDNA was synthesized from the total RNA extracts and qPCR was performed to determine expression levels of DMPK in technical quadruplicate. All qPCR data were analyzed using a traditional $\Delta\Delta$ CT method and were normalized to a plate-based negative control that comprised cells treated with vehicle control (0.3 μ L/well Lipofectamine MessengerMax without any oligonucleotide). Results from these experiments are shown in Table 8. ‘Normalized DMPK Remaining’ for each antisense oligonucleotide in Table 8 refers to the expression level of DMPK in cell treated with said antisense oligonucleotide relative to the negative control that comprised cells treated with vehicle control (wherein the expression level of the negative control has been normalized to equal 1.00)

[000513] The majority of tested DMPK-targeting antisense oligonucleotides demonstrated a reduction in DMPK expression in differentiated myotubes at both the low and high dose concentrations (0.5 nM and 50 nM, respectively). These data demonstrate that the antisense oligonucleotides shown in Table 8 are capable of targeting DMPK *in cellulo*, suggesting that

muscle-targeting complexes comprising these antisense oligonucleotides would be capable of targeting DMPK in muscle tissues *in vivo*.

Table 8. Ability of DMPK-targeting antisense oligonucleotides to reduce expression of DMPK *in cellulo*

Antisense Oligonucleotide Sequence	SEQ ID NO:	DMPK Target Sequence	SEQ ID NO:	0.5 nM		50 nM	
				Normalized DMPK Remaining	Percent DMPK Reduction	Normalized DMPK Remaining	Percent DMPK Reduction
GGACGGCCCCG GCUUGCUGCC	148	GGCAGCAAGC CGGGCCGTCC	384	0.42	58.25	0.31	69.30
GGGCCCCGAU CACAGGACUG	149	CAGTCCTGTG ATCCGGGCCC	385	0.42	57.97	0.38	61.96
CAAACUUGC CAGCAGUGUC	150	GACACTGCTG AGCAAGTTTG	386	0.69	31.45	0.46	53.93
AAACUUGCUC AGCAGUGUCA	151	TGACACTGCT GAGCAAGTTT	387	0.69	30.85	0.49	50.69
CGGAUGGCCU CCAUCUCCCG	152	CGGGAGATGG AGGCCATCCG	388	0.71	28.92	0.44	55.57
CUCGGCCGGA AUCCGCUCCC	153	GGGAGCGGAT TCCGGCCGAG	389	0.71	28.64	0.35	64.75
UCUCGGCCGG AAUCCGCUCC	154	GGAGCGGATT CCGGCCGAGA	390	0.72	27.88	0.33	67.46
UGCUCAGCAG UGUCAGCAGG	155	CCTGCTGACA CTGCTGAGCA	391	0.73	27.08	0.34	65.78
UUGUCGGGUU UGAUGUCCCU	156	AGGGACATCA AACCCGACAA	392	0.66	34.16	0.44	55.56
GUUGCGGGUU UGAUGUCCC	157	GGGACATCAA ACCCGACAAC	393	0.67	33.31	0.39	61.07
UCCGCCAGGU AGAAGCGCGC	158	GCGCGCTTCT ACCTGGCGGA	394	0.72	27.99	0.20	80.06
CAUGGCAUAC ACCUGGCCCG	159	CGGGCCAGGT GTATGCCATG	395	0.68	31.63	0.26	74.03
AACUUGCUC GCAGUGUCAG	160	CTGACACTGC TGAGCAAGTT	396	0.80	19.81	0.47	52.64
CAGCUGCGUG AUCCACCGCC	161	GGCGGTGGAT CACGCAGCTG	397	0.81	19.03	0.32	68.34
CGA AUGUCCG ACAGUGUCUC	162	GAGACACTGT CGGACATTCG	398	0.60	40.21	0.36	64.42
GAAGUCGGCC AGGCGGAUGU	163	ACATCCGCCT GGCCGACTTC	399	0.82	18.36	0.56	44.04
UGUCGGGUUU GAUGUCCUG	164	CAGGGACATC AAACCCGACA	400	0.70	30.09	0.32	68.14
GGAUGGCCUC CAUCUCCCG	165	CCGGGAGATG GAGGCCATCC	401	0.75	24.93	0.39	60.77
AGGAUGUUGU CGGGUUUGAU	166	ATCAAACCCG ACAACATCCT	402	0.76	24.19	0.61	39.48
GUCGGGUUUG AUGUCCUGU	167	ACAGGGACAT CAAACCCGAC	403	0.71	28.89	0.36	64.15
AAUACUCCAU GACCAGGUAC	168	GTACCTGGTC ATGGAGTATT	404	0.71	28.86	0.48	52.07
CUUGUUAUG AUCUUAUGG	169	CCATGAAGAT CATGAACAAG	405	0.84	16.06	0.51	49.47
UCAGUGCAUC CAAACGUGG	170	CCACGTTTGG ATGCACTGA	406	0.84	15.76	0.58	42.06
CUGUCCCGGA GACCAUCCCA	171	TGGGATGGTC TCCGGGACAG	407	0.64	35.85	0.49	50.78

GGGCCUGGGA CCUCACUGUC	172	GACAGTGAGG TCCCAGGCC	408	0.63	37.19	0.23	76.81
CCCACGAAU ACUCCAUGAC	173	GTCATGGAGT ATTACGTGGG	409	0.72	28.21	0.54	45.94
CUCUGCCGCA GGGACAGCCG	174	CGGCTGTCCCT GCGGCAGAG	410	0.63	37.09	0.06	93.59
CUGUGCACGU AGCCAAGCCG	175	CGGCTTGGCT ACGTGCACAG	411	0.74	25.67	0.30	70.10
UGCCCAUCCA CGUCAGGGCC	176	GGCCCTGACG TGGATGGGCA	412	0.86	13.63	0.67	33.09
AGCGCCUCCG AUAGGCCAGG	177	CCTGGCCTATC GGAGGCGCT	413	0.79	21.19	0.38	61.91
UGUGCACGUA GCCAAGCCGG	178	CCGGCTTGGC TACGTGCACA	414	0.75	24.74	0.25	75.09
GACCAGGUAC AGGUAGUUCU	179	AGAACTACCT GTACCTGGTC	415	0.57	42.85	0.29	70.95
CCAUCUCGGC CGGAAUCCGC	180	GCGGATTCCG GCCGAGATGG	416	0.79	20.50	0.40	59.76
CAUCUCGGCC GGAAUCCGCU	181	AGCGGATTCC GGCCGAGATG	417	0.80	20.21	0.41	59.40
UUGCCAUAGG UCUCCGCCGU	182	ACGGCGGAGA CCTATGGCAA	418	0.64	36.30	0.40	60.12
ACAGCGGUCC AGCAGGAUGU	183	ACATCCTGCT GGACCGCTGT	419	0.80	19.94	0.45	55.14
AAAGCGCCUC CGAUAGGCCA	184	TGGCCTATCG GAGGCGCTTT	420	0.80	19.89	0.38	62.04
GCCAAAGAAG AAGGGAUGUG	185	CACATCCCTTC TTCTTTGGC	421	0.75	24.87	0.44	56.19
CACGUAUAC UCCAUGACCA	186	TGGTCATGGA GTATTACGTG	422	0.76	24.40	0.54	46.50
AUCUCGGCCG GAAUCCGCUC	187	GAGCGGATTC CGGCCGAGAT	423	0.88	11.61	0.34	65.98
GCUUCAUCUU CACUACCGCU	188	AGCGGTAGTG AAGATGAAGC	424	0.69	31.44	0.48	51.78
GCCAUCUCGG CCGGAAUCCG	189	CGGATTCGGG CCGAGATGGC	425	0.81	18.56	0.14	86.39
CAGGACAGC CGCUGGAACU	190	AGTTCCAGCG GCTGTCCCTG	426	0.68	32.09	0.41	58.84
AUGACAAUCU CCGCCAGGUA	191	TACCTGGCGG AGATTGTCAT	427	0.58	42.38	0.40	60.47
GGCCAUGACA AUCUCCGCCA	192	TGGCGGAGAT TGTCATGGCC	428	0.58	42.38	0.25	75.00
AUACUCCAUG ACCAGGUACA	193	TGTACCTGGTC ATGGAGTAT	429	0.77	23.07	0.43	56.84
GCCUCUGCCU CGCGUAGUUG	194	CAACTACGCG AGGCAGAGGC	430	0.65	35.38	0.19	81.18
GAAUGUCCGA CAGUGUCUCC	195	GGAGACACTG TCGGACATTC	431	0.70	30.09	0.37	63.41
CGUCCAUCU GCCCCGAGCU	196	AGCTGCGGGC AGATGGAACG	432	0.66	33.74	0.31	68.72
CCUUGUAGUG GACGAUCUUG	197	CAAGATCGTC CACTACAAGG	433	0.83	17.20	0.34	65.91
AUCUCCGCCA GGUAGAAGCG	198	CGCTTCTACCT GGCGGAGAT	434	0.58	42.37	0.35	65.50
CUCAGGCUCU GCCGGGUGAG	199	CTCACCCGGC AGAGCCTGAG	435	0.70	30.13	0.37	63.07
UGCUUCAUCU UCACUACCGC	200	GCGGTAGTGA AGATGAAGCA	436	0.71	28.82	0.40	60.24
GCAGGAUGUU GUCGGGUUUG	201	CAAACCCGAC AACATCCTGC	437	0.56	44.39	0.22	78.03

GGCCUCAGCC UCUGCCGCAG	202	CTGCGGCAGA GGCTGAGGCC	438	0.80	20.12	0.29	71.28
UGUUGUCGGG UUUGAUGUCC	203	GGACATCAAA CCCGACAACA	439	0.79	21.00	0.58	42.19
CCACGUAAUA CUCCAUGACC	204	GGTCATGGAG TATTACGTGG	440	0.79	20.84	0.50	50.06
CCGUUCCAUC UGCCCGCAGC	205	GCTGCGGGCA GATGGAACGG	441	0.68	31.74	0.23	77.46
UUCCCCAGUA AGCAGGCAGA	206	TCTGCCTGCTT ACTCGGGAA	442	0.69	31.49	0.50	49.81
UGAUCUUAU GGCAUACACC	207	GGTGTATGCC ATGAAGATCA	443	0.72	27.70	0.10	89.68
AGGGACAGCC GCUGGAACTG	208	CAGTTCCAGC GGCTGTCCCT	444	0.71	28.72	0.55	45.34
GGGUUUGAUG UCCUGUGCA	209	TGCACAGGGA CATCAAACCC	445	0.60	40.12	0.37	62.61
UGACAAUCUC CGCCAGGUAG	210	CTACCTGGCG GAGATTGTCA	446	0.61	38.86	0.33	66.56
CACAGCGGUC CAGCAGGAUG	211	CATCCTGCTG GACCGCTGTG	447	0.93	6.62	0.40	59.58
GCGUAGAAGG GCGUCUGCCC	212	GGGCAGACGC CCTTCTACGC	448	0.60	39.53	0.22	77.91
CUCAGCCUCU GCCGCAGGGA	213	TCCCTGCGGC AGAGGCTGAG	449	0.82	17.86	0.20	79.58
GUCUCAGUGC AUCCAAAACG	214	CGTTTTGGATG CACTGAGAC	450	0.81	18.85	0.54	46.13
GGACGAUCUU GCCAUAGGUC	215	GACCTATGGC AAGATCGTCC	451	0.70	29.82	0.51	48.97
UCAGCAGUGU CAGCAGGUCC	216	GGACCTGCTG ACACTGCTGA	452	0.67	33.46	0.39	61.11
GCUCCUGGGC GGCGCCAGAC	217	GTCTGGCGCC GCCCAGGAGC	453	0.91	8.52	0.21	78.79
AGCAGGAUGU UGUCGGGUUU	218	AAACCCGACA ACATCCTGCT	454	0.59	41.05	0.26	74.02
AUCCGCUCCU GCAACUGCCG	219	CGGCAGTTGC AGGAGCGGAT	455	0.87	12.80	0.60	40.06
AGGAGCAGGG AAAGCGCCUC	220	GAGGCGCTTT CCCTGCTCCT	456	0.67	33.24	0.38	62.37
ACACCUGGCC CGUCUGCUUC	221	GAAGCAGACG GGCCAGGTGT	457	0.67	33.00	0.45	55.40
CCCAGCGCCC ACCAGUCACA	222	TGTGACTGGT GGGCGCTGGG	458	0.62	37.93	0.32	67.82
GCUCCUCUG CCUGCAGCAA	223	TTGCTGCAGG CAGAGGGAGC	459	0.74	26.41	0.30	70.15
GCUCAGGCUC UGCCGGGUGA	224	TCACCCGGCA GAGCCTGAGC	460	0.74	25.69	0.39	60.71
UUGAUGUCCC UGUGCACGUA	225	TACGTGCACA GGGACATCAA	461	0.74	25.67	0.45	55.13
GCCUCAGCCU CUGCCGCAGG	226	CCTGCGGCAG AGGCTGAGGC	462	0.84	16.37	0.54	46.42
GGUAGUUCUC AUCCUGGAAG	227	CTTCCAGGAT GAGAACTACC	463	0.75	25.48	0.44	56.15
CAGCGCCAC CAGUCACACU	228	AGTGTGACTG GTGGGCGCTG	464	0.63	37.28	0.35	64.93
CCCAAACUUG CUCAGCAGUG	229	CACTGCTGAG CAAGTTTGGG	465	0.63	37.02	0.38	61.78
CUUGCCAUAG GUCUCCGCCG	230	CGGCGGAGAC CTATGGCAAG	466	0.73	27.04	0.29	71.05
UACACCGGC CCGUCUGCUU	231	AAGCAGACGG GCCAGGTGTA	467	0.69	31.10	0.43	57.43

CCAGCGCCCA CCAGUCACAC	232	GTGTGACTGG TGGGCGCTGG	468	0.64	36.17	0.29	70.96
GGCCUCAGCC UGGCCGAAAG	233	CTTTCGGCCA GGCTGAGGCC	469	0.86	14.49	0.35	64.80
AAUCUCCGCC AGGUAGAAGC	234	GCTTCTACCTG GCGGAGATT	470	0.64	35.85	0.35	65.27
AUGGCAUACA CCUGGCCCGU	235	ACGGGCCAGG TGTATGCCAT	471	0.86	14.31	0.50	49.63
CCAUGACAAU CUCCGCCAGG	236	CCTGGCGGAG ATTGTCATGG	472	0.65	34.53	0.24	76.46
UCCCCAAACU UGCUCAGCAG	237	CTGCTGAGCA AGTTTGGGGA	473	0.94	5.73	0.55	44.67
GAUGUUGUCG GGUUUGAUGU	238	ACATCAAACC CGACAACATC	474	0.90	10.06	0.58	42.42
GUUUGCCCAU CCACGUCAGG	239	CCTGACGTGG ATGGGCAAAC	475	0.66	34.36	0.46	54.49
CGGACGGCCC GGCUUGCUGC	240	GCAGCAAGCC GGGCCGTCCG	476	0.95	5.42	0.70	30.41
CUCCGCCAGG UAGAAGCGCG	241	CGCGCTTCTAC CTGGCGGAG	477	0.70	30.22	0.22	78.14
GUACAGGUAG UUCUACUCCU	242	AGGATGAGAA CTACCTGTAC	478	0.68	31.52	0.34	65.57
AGGGCGUCUG CCCAUAGAAC	243	GTTCTATGGG CAGACGCCCT	479	0.87	13.23	0.41	58.98
UGGCCACAGC GGUCCAGCAG	244	CTGCTGGACC GCTGTGGCCA	480	0.70	29.59	0.31	69.44
CGUAGUUGAC UGGCGAAGUU	245	AACTTCGCCA GTCAACTACG	481	0.75	25.26	0.38	61.52
UCUGCCGCAG GGACAGCCGC	246	GCGGCTGTCC CTGCGGCAGA	482	0.77	22.97	0.18	82.10
AAGCGCCUCC GAUAGGCCAG	247	CTGGCCTATC GGAGGCGCTT	483	0.91	8.91	0.56	43.93
GACAGAACAA CGGCGAACAG	248	CTGTTCCCGT TGTTCTGTC	484	0.79	21.41	0.30	70.49
GCUCAGCAGU GUCAGAGGU	249	ACCTGCTGAC ACTGCTGAGC	485	0.71	29.18	0.27	73.46
AUGAUCUUCA UGGCAUACAC	250	GTGTATGCCA TGAAGATCAT	486	0.87	12.76	0.60	39.97
UUUGCCCAUC CACGUCAGGG	251	CCCTGACGTG GATGGGCAAA	487	0.67	32.79	0.41	59.36
ACUUGCUCAG CAGUGUCAGC	252	GCTGACACTG CTGAGCAAGT	488	0.72	27.84	0.39	60.71
UGAUGUCCCU GUGCACGUAG	253	CTACGTGCAC AGGGACATCA	489	0.79	20.58	0.41	59.00
AAAUACCGAG GAAUGUCGGG	254	CCCGACATTC CTCGGTATTT	490	0.89	11.25	0.49	50.91
GGCGAAUACA CCCAGCGCCC	255	GGGCGCTGGG TGTATTCGCC	491	0.80	19.77	0.31	68.72
AGACAAUAAA UACCGAGGAA	256	TTCCTCGGTAT TTATTGTCT	492	0.71	29.37	0.52	48.20
CCCGUCUGCU UCAUCUUCAC	257	GTGAAGATGA AGCAGACGGG	493	0.80	20.31	0.56	43.97
CUGCCUGCAG CAACUCCAUC	258	GATGGAGTTG CTGCAGGCAG	494	0.77	23.10	0.53	46.69
CCUCAGCCUC UGCCGCAGGG	259	CCCTGCGGCA GAGGCTGAGG	495	0.89	10.87	0.45	55.22
GUGUCCGGAA GUCGCCUGCU	260	AGCAGGCGAC TTCCGGACAC	496	0.77	22.99	0.26	73.65
UGCACGUGUG GCUCAAGCAG	261	CTGCTTGAGC CACACGTGCA	497	0.89	10.81	0.36	64.18

GACAAUAAAU ACCGAGGAAU	262	ATTCTCGGTA TTTATTGTC	498	0.71	28.97	0.52	47.51
GCCAUGACAA UCUCCGCCAG	263	CTGGCGGAGA TTGTCATGGC	499	0.69	30.96	0.19	81.00
GCUGUCCCGG AGACCAUCCC	264	GGGATGGTCT CCGGGACAGC	500	0.77	22.57	0.34	66.27
CAUGACCAGG UACAGGUAGU	265	ACTACCTGTA CCTGGTCATG	501	0.81	19.39	0.41	59.09
AGCGCCACC AGUCACACUC	266	GAGTGTGACT GGTGGGCGCT	502	0.70	30.36	0.36	63.67
UCUCAGUGCA UCCAAAACGU	267	ACGTTTTGGAT GCACTGAGA	503	0.89	10.88	0.49	51.34
UUUGGGCAGA UGGAGGGCCU	268	AGGCCCTCCA TCTGCCAAA	504	0.65	35.14	0.30	70.00
GAUGUCCUG UGCAGUAGC	269	GCTACGTGCA CAGGGACATC	505	0.81	18.99	0.38	62.46
CAGCAGUGUC AGCAGGUCCC	270	GGGACCTGCT GACACTGCTG	506	0.74	25.67	0.48	51.97
CAUGACAAUC UCCGCCAGGU	271	ACCTGGCGGA GATTGTCATG	507	0.71	29.45	0.29	70.52
ACUUGUUCAU GAUCUUAUG	272	CATGAAGATC ATGAACAAGT	508	0.75	25.47	0.47	52.89
GUGGAAUCCG CGUAGAAGGG	273	CCCTTCTACGC GGATTCCAC	509	0.69	30.55	0.51	49.34
UGGCCAUGAC AAUCUCCGCC	274	GGCGGAGATT GTCATGGCCA	510	0.70	30.46	0.27	72.55
GGGACAGACA AUAAAUACCG	275	CGGTATTTATT GTCTGTCCC	511	0.73	27.19	0.49	50.50
CCGCUCCCCA AACUUGCUCA	276	TGAGCAAGTT TGGGGAGCGG	512	1.00	0.28	0.43	56.82
CGGCUCAGGC UCUGCCGGGU	277	ACCCGGCAGA GCCTGAGCCG	513	0.82	17.97	0.31	69.03
GGCUCCUGGG CGGCGCCAGA	278	TCTGGCGCCG CCCAGGAGCC	514	1.00	0.05	0.04	96.23
UUUCCCGAGU AAGCAGGCAG	279	CTGCCTGCTTA CTCGGGA	515	0.79	20.69	0.55	44.89
GGAUGUUGUC GGGUUUGAUG	280	CATCAAACCC GACAACATCC	516	0.96	4.26	0.59	40.81
CAGGUAGUUC UCAUCCUGGA	281	TCCAGGATGA GAACTACCTG	517	0.74	25.92	0.23	76.71
UGCCCAUAGA ACAUUUCAUA	282	TATGAAATGT TCTATGGGCA	518	0.92	7.67	0.65	34.56
UAGUUCUCAU CCUGGAAGGC	283	GCCTTCCAGG ATGAGAACTA	519	0.83	16.83	0.56	43.88
AUGUCCUGU GCACGUAGCC	284	GGCTACGTGC ACAGGGACAT	520	0.83	16.78	0.51	49.29
CGGGCCCCGA UCACAGGACU	285	AGTCCTGTGA TCCGGGCCCCG	521	0.83	17.45	0.33	67.11
UGGACGAUCU UGCCAUAGGU	286	ACCTATGGCA AGATCGTCCA	522	0.81	19.20	0.57	42.52
GUUGGCCGGC GUGGGCCACC	287	GGTGGCCAC GCCGGCCAAC	523	1.02	-1.82	0.56	43.57
CUCAGUGCAU CCAAAACGUG	288	CACGTTTTGG ATGCACTGAG	524	0.92	7.65	0.46	54.26
UCGAAGUUGC AUGUGUCGGU	289	ACCGACACAT GCAACTTCGA	525	0.77	22.96	0.42	58.15
UGGAACACGG ACGGCCCGGC	290	GCCGGGCCGT CCGTGTTCCA	526	1.02	-1.90	0.39	60.96
CCGAGAGCAG CGCAAGUGAG	291	CTCACTTGCGC TGCTCTCGG	527	0.84	16.13	0.59	40.93

UCCUGCAACU GCCGGACGUG	292	CACGTCCGGC AGTTGCAGGA	528	0.84	16.06	0.55	44.61
UCACCAACAC GUCCCUCUCC	293	GGAGAGGGAC GTGTTGGTGA	529	0.53	47.12	0.16	84.09
UGCCUGCAGC AACUCCAUCC	294	GGATGGAGTT GCTGCAGGCA	530	0.86	13.99	0.50	49.75
UUGGCCGGCG UGGGCCACCA	295	TGGTGGCCCA CGCCGGCCAA	531	1.03	-3.19	0.56	44.37
GAGCCUCUGC CUCGCGUAGU	296	ACTACGCGAG GCAGAGGCTC	532	0.81	18.77	0.22	77.78
AAGGGCGUCU GCCCAUAGAA	297	TTCTATGGGC AGACGCCCTT	533	0.87	13.15	0.65	34.56
ACAGACAAUA AAUACCGAGG	298	CCTCGGTATTT ATTGTCTGT	534	1.04	-3.95	0.26	74.02
GGACAGACAA UAAAUACCGA	299	TCGGTATTTAT TGTCGTGCC	535	0.77	22.57	0.47	52.51
ACGUGUGCCU CUAGGUCCCCG	300	CGGGACCTAG AGGCACACGT	536	0.84	16.47	0.22	77.73
GGCACGAGAC AGAACAACGG	301	CCGTTGTTCTG TCTCGTGCC	537	0.84	16.10	0.32	68.01
UGACCAGGUA CAGGUAGUUC	302	GAACCTACCTG TACCTGGTCA	538	0.78	22.00	0.36	63.73
CUCUGCCGGG UGAGCACCUC	303	GAGGTGCTCA CCCGGCAGAG	539	0.75	25.25	0.26	74.36
GACAAUCUCC GCCAGGUAGA	304	TCTACCTGGC GGAGATTGTC	540	0.76	23.70	0.50	49.82
UCUCCGCCAG GUAGAAGCGC	305	GCGCTTCTACC TGGCGGAGA	541	0.80	19.59	0.33	66.52
CUCUGCCUCG CGUAGUUGAC	306	GTCAACTACG CGAGGCAGAG	542	0.83	16.61	0.09	91.21
CUUUGGGCAG AUGGAGGGCC	307	GGCCCTCCAT CTGCCCAAAG	543	0.72	28.06	0.33	67.50
ACAGGUAGUU CUCAUCCUGG	308	CCAGGATGAG AACTACCTGT	544	0.79	20.51	0.15	85.36
CCAAACUUGC TCAGCAGUGU	309	ACACTGCTGA GCAAGTTTGG	545	0.76	23.64	0.42	57.70
UCGGGUUUGA UGUCCUGUG	310	CACAGGGACA TCAAACCCGA	546	0.78	22.49	0.43	57.16
GGCUUGCUGC CUUCCAGGC	311	GCCTGGGAAG GCAGCAAGCC	547	1.06	-6.32	0.52	48.15
UACAGGUAGU UCUCAUCCUG	312	CAGGATGAGA ACTACCTGTA	548	0.80	19.83	0.27	72.51
UUGCCCAUCC ACGUCAGGGC	313	GCCCTGACGT GGATGGGCAA	549	0.78	22.23	0.33	67.15
AGGUACAGGU AGUUCUCAUC	314	GATGAGAACT ACCTGTACCT	550	0.81	18.68	0.41	58.92
GACAGACAAU AAAUACCGAG	315	CTCGGTATTTA TTGTCTGTC	551	0.82	18.26	0.62	38.07
UAGAACAUUU CAUAGGCGAA	316	TTCGCCTATGA AATGTTCTA	552	0.80	20.23	0.56	43.67
AGGGCCUUUU AUUCGCGAGG	317	CCTCGCGAAT AAAAGGCCCT	553	0.86	13.63	0.34	66.43
GCCUCGCGUA GUUGACUGGC	318	GCCAGTCAAC TACGCGAGGC	554	0.87	12.98	0.09	91.10
CCAGCAGGAU GUUGUCGGGU	319	ACCCGACAAC ATCCTGCTGG	555	0.60	40.29	0.10	89.59
GUAGUUGACU GGCGAAGUUC	320	GAACCTCGCC AGTCAACTAC	556	0.93	7.50	0.55	45.33
UGCGGAUGGC CUCCAUCUCC	321	GGAGATGGAG GCCATCCGCA	557	0.60	40.15	0.16	84.43

ACAAUCUCCG CCAGGUAGAA	322	TTCTACCTGGC GGAGATTGT	558	0.81	19.09	0.50	49.75
GCGAAUACAC CCAGCGCCCA	323	TGGGCGCTGG GTGTATTTCGC	559	0.93	6.94	0.30	69.72
GUAGUUCUCA UCCUGGAAGG	324	CCTTCCAGGA TGAGAACTAC	560	0.93	7.43	0.45	55.09
GGCUCAGGCU CUGCCGGGUG	325	CACCCGGCAG AGCCTGAGCC	561	0.93	7.38	0.34	65.82
CCAUUCACCA ACACGUCCCU	326	AGGGACGTGT TGGTGAATGG	562	0.61	39.26	0.13	86.83
ACCAGGUACA GGUAGUUCUC	327	GAGAACTACC TGTACCTGGT	563	0.84	16.09	0.23	76.96
CTGCAGUUUG CCCAUCCACG	328	CGTGGATGGG CAAACCTGCAG	564	1.11	-10.69	0.40	60.08
UUGUUCAUGA UCUUAUGGC	329	GCCATGAAGA TCATGAACAA	565	0.86	14.13	0.55	45.23
UUGAUGUCCC UGUGCACGU	330	ACGTGCACAG GGACATCAAA	566	0.93	6.92	0.57	43.07
GCGGUCCAGC AGGAUGUUGU	331	ACAACATCCT GCTGGACCGC	567	0.61	38.84	0.16	83.64
GUCUAUGGCC AUGACAAUCU	332	AGATTGTCAT GGCCATAGAC	568	1.11	-11.00	0.27	73.11
GGAGCAGGGA AAGCGCCUCC	333	GGAGGCGCTT TCCCTGCTCC	569	0.79	21.46	0.12	88.35
UGCCUCGCGU AGUUGACUGG	334	CCAGTCAACT ACGCGAGGCA	570	0.89	11.03	0.12	88.02
GCGGAUGGCC UCCAUCUCCC	335	GGGAGATGGA GGCCATCCGC	571	0.79	21.25	0.28	71.77
UUUCAUAGGC GAAUACACCC	336	GGGTGTATTC GCCTATGAAA	572	0.94	5.56	0.47	53.28
GCCUGUCAGC GAGUCGGAGG	337	CCTCCGACTC GCTGACAGGC	573	0.89	10.81	0.24	75.67
CCACUUCAGC UGUUUCAUCC	338	GGATGAAACA GCTGAAGTGG	574	0.78	22.40	0.36	64.20
CAUCCGCUCC UGCAACUGCC	339	GGCAGTTGCA GGAGCGGATG	575	0.79	21.04	0.23	76.81
UCUAGGGUUC AGGGAGCGCG	340	CGCGCTCCCT GAACCCTAGA	576	0.78	21.81	0.17	83.22
CACCAACACG UCCCUUCCU	341	AGGAGAGGGA CGTGTTGGTG	577	0.62	37.51	0.18	81.57
CAGGAGCAGG GAAAGCGCCU	342	AGGCGCTTTC CCTGCTCCTG	578	0.88	12.48	0.48	51.82
CAAUCUCCGC CAGGUAGAAG	343	CTTCTACCTGG CGGAGATTG	579	0.84	15.95	0.51	49.25
AUGUUGUCGG GUUUGAUGUC	344	GACATCAAAC CCGACAACAT	580	0.83	16.93	0.47	52.83
CCAUCCGCUC CUGCAACUGC	345	GCAGTTGCAG GAGCGGATGG	581	0.80	19.53	0.28	71.62
GCGUCACCUC GGCCUCAGCC	346	GGCTGAGGCC GAGGTGACGC	582	0.80	20.02	0.19	81.27
GAGGGCCUUU UAUUCGCGAG	347	CTCGCGAATA AAAGGCCCTC	583	0.92	8.23	0.38	62.21
AGCGGCAGAG AGAGGUGCUC	348	GAGCACCTCT CTCTGCCGCT	584	0.80	19.75	0.09	90.71
CAUCCAAAAC GUGGAUUGGG	349	CCCAATCCAC GTTTTGGATG	585	0.81	19.12	0.22	77.98
UUGGGCAGAU GGAGGGCCUU	350	AAGGCCCTCC ATCTGCCCAA	586	0.81	19.08	0.22	78.39
CCUCUGCCUC GCGUAGUUGA	351	TCAACTACGC GAGGCAGAGG	587	0.93	7.39	0.15	85.33

ACAGAACAAC GGCGAACAGG	352	CCTGTTTCGCCG TTGTTCTGT	588	0.98	2.07	0.44	55.96
CAGGAUGUUG UCGGGUUUGA	353	TCAAACCCGA CAACATCCTG	589	0.83	17.17	0.21	79.31
CGGCCUCAGC CUCUGCCGCA	354	TGCGGCAGAG GCTGAGGCCG	590	0.93	6.71	0.40	60.06
CAGCAGGAUG UUGUCGGGUU	355	AACCCGACAA CATCCTGCTG	591	0.66	34.18	0.15	84.54
GCAGAGAGAG GUGCUCUUG	356	CAAGGAGCAC CTCTCTCTGC	592	0.83	17.29	0.14	85.95
UCCAGUUGCA UGGGUGUGGG	357	CCCACACCCA TGGAAGTGA	593	0.84	15.66	0.22	78.48
CCUCAGCCUG GCCGAAAGAA	358	TTCTTTCGGCC AGGCTGAGG	594	0.83	16.83	0.36	63.99
GGGCCUUUUA UUCGCGAGGG	359	CCCTCGCGAA TAAAAGGCC	595	0.95	5.11	0.49	50.65
GUCGGCCAGG CGGAUGUGGC	360	GCCACATCCG CCTGGCCGAC	596	0.85	15.35	0.25	74.59
GCUUGCUGCC UUCCCAGGCC	361	GGCCTGGGAA GGCAGCAAGC	597	0.99	1.14	0.19	81.01
GGUCCAGCAG GAUGUUGCG	362	CGACAACATC CTGCTGGACC	598	0.68	31.78	0.20	79.93
CGAGACCAU CCCAGUCGAG	363	CTCGACTGGG ATGGTCTCCG	599	0.86	14.08	0.20	79.93
UCUGCCUCGC GUAGUGACU	364	AGTCAACTAC GCGAGGCAGA	600	0.96	3.53	0.13	86.86
AGGUAGUUCU CAUCCUGGAA	365	TTCCAGGATG AGAACTACCT	601	0.93	7.36	0.37	62.62
UCCUUGUAGU GGACGAUCUU	366	AAGATCGTCC ACTACAAGGA	602	0.87	12.96	0.15	84.87
GCAUCCAAAA CGUGGAUUGG	367	CCAATCCACG TTTTGGATGC	603	0.97	2.54	0.27	72.69
GUCCAGCAGG AUGUGUCGG	368	CCGACAACAT CCTGCTGGAC	604	0.70	30.00	0.17	82.64
AGCUCCCGCA GCGUACCUC	369	GAGGTGACGC TGCGGGAGCT	605	0.86	13.72	0.20	80.40
CGAGAGCAGC GCAAGUGAGG	370	CCTCACTTGCG CTGCTCTCG	606	1.02	-2.19	0.63	37.11
CAGGGAAAGC GCCUCCGAUA	371	TATCGGAGGC GCTTTCCTG	607	0.89	11.10	0.08	91.59
AUUUCAUAGG CGAAUACACC	372	GGTGTATTCG CCTATGAAAT	608	1.05	-4.54	0.56	44.15
UCGGCCAGGC GGAUGUGGCC	373	GGCCACATCC GCCTGGCCGA	609	0.73	26.53	0.17	83.04
AAGGGAUGUG UCCGGAAGUC	374	GACTTCCGGA CACATCCCTT	610	0.90	10.37	0.26	73.52
CUUGUAGUGG ACGAUCUUGC	375	GCAAGATCGT CCACTACAAG	611	0.76	24.09	0.11	89.16
AGUCGGCCAG GCGGAUGUGG	376	CCACATCCGC CTGGCCGACT	612	0.94	6.15	0.33	67.44
GCCUCAGCCU GGCCGAAAGA	377	TCTTTCGGCCA GGCTGAGGC	613	1.05	-4.82	0.37	63.11
AGCGUACCU CGGCCUCAGC	378	GCTGAGGCCG AGGTGACGCT	614	0.78	22.10	0.35	64.70
CAGCGGCAGA GAGAGGUGCT	379	AGCACCTCTCT CTGCCGCTG	615	0.96	4.49	0.14	86.00
CCAGCGGCAG AGAGAGGUGC	380	GCACCTCTCTC TGCCGCTGG	616	0.97	3.23	0.15	84.55
UUGUAGUGGA CGAUCUUGCC	381	GGCAAGATCG TCCACTACAA	617	0.83	17.22	0.19	81.05

AGGGAAAGCG CCUCCGAUAG	382	CTATCGGAGG CGCTTTCCT	618	1.01	-1.12	0.25	75.50
GGGAAAGCGC CUCCGAUAGG	383	CCTATCGGAG GCGCTTCCC	619	0.90	10.02	0.23	76.79

Example 8: Selected antisense oligonucleotides provided dose-dependent reduction in DMPK expression in immortalized myoblasts

[000514] Eighteen oligonucleotides from Example 7 were selected to be evaluated for their ability to reduce DMPK expression in a dose-responsive manner. DM1 C15 myoblasts were prepared as in Example 7 to yield differentiated myotubes in 96-well microplates. After seven days of differentiation, cells were transfected with individual oligonucleotides using Lipofectamine MessengerMax. Each oligonucleotide was tested in triplicate at concentrations of 0.046 nM, 0.137 nM, 0.412 nM, 1.235 nM, 3.704 nM, 11.11 nM, 33.33 nM, and 100 nM by 3-fold serial dilutions using 0.3 μ L of Lipofectamine MessengerMax per well.

[000515] Following addition of oligonucleotide, cells were incubated for 72 hours prior to harvesting for total RNA. cDNA was synthesized from the total RNA extracts and qPCR was performed to determine expression levels of DMPK using a commercially available Taqman probeset in technical quadruplicate. All qPCR data were analyzed using a traditional $\Delta\Delta$ CT method and were normalized to a plate-based negative control that comprised of cells treated with vehicle control (0.3 μ L/well Lipofectamine MessengerMax without any oligonucleotide). Data for each oligonucleotide to was fit to sigmoidal curve in order to determine an effective concentration of each oligonucleotide that provided a half-maximal response (EC-50). Results from these experiments are shown in Table 9.

[000516] Each of the eighteen antisense oligonucleotides selected for dose-dependent experimentation were capable of dose-dependently reducing DMPK in differentiated myotubes. Further, each of the tested antisense oligonucleotides reduced DMPK with EC-50 values below 25 nM. For example, antisense oligonucleotides comprising SEQ ID NOs: 264, 215, 222, 190, and 212 resulted in EC-50 values of 3.27 nM, 3.59 nM, 5.45 nM, 6.04 nM, and 24.59 nM, respectively. These data demonstrate that the antisense oligonucleotides shown in Table 9 are capable of dose-dependent reduction of DMPK *in cellulo*, suggesting that muscle-targeting complexes comprising these antisense oligonucleotides would be capable of targeting DMPK in muscle tissues *in vivo*.

Table 9. Ability of DMPK-targeting antisense oligonucleotides to reduce expression of DMPK in dose-dependent manner *in cellulo*

				Results
--	--	--	--	----------------

Antisense Oligonucleotide Sequence	SEQ ID NO:	DMPK Target Sequence	SEQ ID NO:	EC-50 (nM)	Percent DMPK reduction at 100 nM
GCAGGAUGUUGUCGGGUUUG	201	CAAACCCGACAA CATCCTGC	437	0.1679	89.77
AGCAGGAUGUUGUCGGGUUU	218	AAACCCGACAAC ATCCTGCT	454	0.2266	85.81
GCGUAGAAGGGCGUCUGCCC	212	GGGCAGACGCCC TTCTACGC	448	24.59	95.13
CCCAGCGCCCACCAGUCACA	222	TGTGACTGGTGG GCGCTGGG	458	5.454	63.69
CCAUCUCGGCCGGAUCCGC	180	GCGGATTCCGGC CGAGATGG	416	0.44	95.42
CGUCCAUCUGCCCGCAGCU	196	AGTGCGGGCAG ATGGAACG	432	0.19	89.97
CAGGGACAGCCGCUGGAACU	190	AGTTCCAGCGGC TGTCCCTG	426	6.04	90.59
CAUGGCAUACACCUGGCCCG	159	CGGGCCAGGTGT ATGCCATG	395	0.42	75.28
GCUUCAUCUUCACUACCGCU	188	AGCGGTAGTGAA GATGAAGC	424	0.03	64.06
GAAUGUCCGACAGUGUCC	195	GGAGACACTGTC GGACATTC	431	0.07	97.23
GGACGAUCUUGCCAUAUGUC	215	GACCTATGGCAA GATCGTCC	451	3.59	92.18
GCUGUCCCGGAGACCAUCCC	264	GGGATGGTCTCC GGGACAGC	500	3.27	93.07
GACAGAACAACGGCGAACAG	248	CTGTTCGCCGTT GTTCTGTC	484	0.08	94.32
UGUUGUCGGGUUUGAUGUCC	203	GGACATCAAACC CGACAACA	439	0.21	93.95
CGAAUGUCCGACAGUGUCUC	162	GAGACACTGTCTG GACATTCTG	398	0.18	95.93
GGGCCUGGGACCUCACUGUC	172	GACAGTGAGGTC CCAGGCCC	408	0.07	90.58
CUCUGCCGACAGGACAGCCG	174	CGGCTGTCCCTG CGGCAGAG	410	0.42	93.66
UUGCCAUAGGUCUCCGCGCU	182	ACGGCGGAGACC TATGGCAA	418	0.37	93.70

Example 9: Targeting DMPK in mouse muscle tissues with a muscle-targeting complex

[000517] The RI7 217 Fab antibody-ASO muscle-targeting complex described in Example 2, DTX-C-008, was tested for time-dependent inhibition of DMPK in mouse tissues *in vivo*. C57BL/6 wild-type mice were intravenously injected with a single dose of a vehicle control (phosphate-buffered saline (PBS)), naked antisense oligonucleotide (ASO300) (10 mg/kg of ASO), DTX-C-007 IgG2a Fab antibody-ASO control complex (10 mg/kg of ASO), or DTX-C-008 (10 mg/kg of ASO) on Day 0 and euthanized after a prescribed period of time, as described in Table 10. One group of mice in each experimental condition was subjected to a second dose (multi-dose groups) at four weeks (Day 28). Following euthanization, the mice were segmented into isolated tissue types and samples of tibialis anterior and gastrocnemius muscle tissues were subsequently assayed for expression levels of DMPK (FIGs. 13A-13B).

Table 10. Experimental conditions

Group	Dosage	Weeks after injection before euthanization	Number of mice
1	Single dose of vehicle (PBS)	2	5
2	Single dose of vehicle (PBS)	4	5
3	Single dose of vehicle (PBS)	8	3
4	Multi-dose of vehicle (PBS)	8	2
5	Single dose of vehicle (PBS)	12	5
6	Single dose of ASO300	2	5
7	Single dose of ASO300	4	5
8	Single dose of ASO300	8	5
9	Multi-dose of ASO300	8	5
10	Single dose of ASO300	12	5
11	Single dose of DTX-C-007 control	2	5
12	Single dose of DTX-C-007 control	4	5
13	Single dose of DTX-C-007 control	8	5
14	Multi-dose of DTX-C-007 control	8	5
15	Single dose of DTX-C-007 control	12	5
16	Single dose of DTX-C-008	2	5
17	Single dose of DTX-C-008	4	5
18	Single dose of DTX-C-008	8	5
19	Multi-dose of DTX-C-008	8	5
20	Single dose of DTX-C-008	12	5

[000518] Mice treated with the RI7 217 Fab antibody-ASO DTX-C-008 complex demonstrated about 50-60% reduction in DMPK expression in tibialis anterior muscle (FIG. 13A) and about 30-50% reduction in DMPK expression in gastrocnemius muscle (FIG. 13B) for all of Groups 16-20 (2-12 weeks between injection and euthanization), relative to vehicle. These data show that a single dose of the muscle-targeting complex DTX-C-008 reduces expression of DMPK for at least twelve weeks following administration of the complex.

[000519] In contrast, mice treated with the naked antisense oligonucleotide or the control complex did not demonstrate significant inhibition of DMPK expression across all experimental groups and tissues.

[000520] These data demonstrate that a muscle-targeting complex as described herein is capable of providing persistent inhibition of DMPK expression *in vivo* for up to 12 weeks following a single dose or administration of said muscle targeting complex.

Example 10: A muscle-targeting complex can target gene expression in the nucleus

[000521] The RI7 217 Fab antibody-ASO muscle-targeting complex as described in Example 2, DTX-C-008, was tested for inhibition of nuclear-retained DMPK RNA in mouse muscle tissues. The mice used for this Example have been engineered to express a human mutant DMPK gene (schematic shown in FIG. 14A) – DMPK with a 350 CTG repeat region and

a downstream G-for-C single-nucleotide polymorphism. As shown in FIG. 14A, the human mutant DMPK RNA is retained in the nucleus, while the mouse wild-type DMPK RNA is located in the cytoplasm and the nucleus.

[000522] Mice were intravenously injected with a single dose of a vehicle control (saline), an IgG2a Fab-ASO control complex DTX-C-007 (10 mg/kg of ASO), naked ASO300 (10 mg/kg of ASO), or DTX-C-008 (10 mg/kg of ASO) and euthanized after 14 days. Six mice were treated in each experimental condition. Following euthanization, the mice were segmented into isolated tissue types and tissue samples were subsequently assayed for expression levels of mutant and wild-type DMPK (FIG. 14B).

[000523] Mice treated with the muscle-targeting RI7 217 Fab antibody-ASO complex DTX-C-008 demonstrated statistically significant reduction in both nuclear-retained mutant DMPK and wild-type DMPK. (p-value <0.05). These data demonstrate that a muscle-targeting complex as described herein is capable of targeting DMPK in the nucleus.

Example 11: A muscle-targeting complex reverses myotonia in HSA^{LR} mouse model

[000524] A muscle-targeting complex (DTX-Actin) was generated comprising an antisense oligonucleotide (ASO) that targets actin covalently linked to DTX-A-002 (RI7 217 Fab), an anti-transferrin receptor antibody.

[000525] The actin-targeting ASO is an MOE 5-10-5 gapmer that comprises: 5'-NH₂-(CH₂)₆-dA*oC*oC*oA*oT*oT*dT*dT*dC*dT*dT*dC*dC*dA*dC*dA*oG*oG*oG*oC*oT-3 (SEQ ID NO: 620); wherein '*' represents a PS linkage; 'd' represents a deoxynucleic acid; and 'o' represents a 2'-MOE.

[000526] DTX-Actin was then tested for its ability to reduce target gene expression (hACTA1) and reduce myotonia in HSA^{LR} mice, a mouse model that has a functional myotonia phenotype similar to that observed in human DM1 patients. Details of the HSA^{LR} mouse model are as described in Mankodi, A. et al. Science. 289: 1769, 2000. HSA^{LR} mice were intravenously injected with a single dose of PBS or DTX-Actin (either 10 mg/kg or 20 mg/kg ASO equivalent). Each of these three experimental conditions were replicated in two individual mice. On Day 14 after injection, mice were euthanized and specific muscles were collected – quadriceps (quad), gastrocnemius (gastroc) and tibialis anterior (TA). The muscle tissues were analyzed for expression of hACTA1. DTX-Actin demonstrated reduction of hACTA1 expression in all three muscle tissues relative to vehicle control (FIG. 15A).

[000527] On Day 14 after injection, and prior to the euthanasia and tissue collection described above, electromyography (EMG) was performed on specific muscles. EMG myotonic discharges were graded by a blinded examiner on a 4-point scale: 0, no myotonia; 1, occasional

myotonic discharge in less than 50% of needle insertions; 2, myotonic discharge in greater than 50% of needle insertions; and 3: myotonic discharge with nearly every insertion. DTX-Actin demonstrated reduction in graded myotonia in all three muscle tissues relative to vehicle control (FIG. 15B). Mice treated with DTX-Actin at a dose of 20 mg/kg ASO equivalent demonstrated little-to-no myotonia in quadriceps and gastrocnemius muscles.

[000528] These data demonstrate that a single dose of a muscle-targeting complex is capable of gene-specific targeting and reduction in functional myotonia in the HSA^{LR} mice, a mouse model that has a functional myotonia phenotype similar to that observed in human DM1 patients.

Example 12: A muscle-targeting complex can functionally correct arrhythmia in a DM1 mouse model

[000529] The RI7 217 Fab antibody-ASO muscle-targeting complex as described in Example 2, DTX-C-008, was tested for its ability to functionally correct arrhythmia in a DM1 mouse model. The mice used for this Example are the offspring of mice expressing myosin heavy chain reverse tet transactivator (MHCrtTA) and mice expressing a mutant form of human DMPK (CUG₉₆₀). FIG. 16A shows the genomic structure of the mutant transgene.

[000530] Doxycycline containing chow (2 g doxycycline/kg chow, Bio-Serv) was provided to the mice beginning at postnatal day 1, initially through the nursing dam and subsequently through chow after weaning, to induce selective expression of mutant DMPK in the heart. All mice were maintained on chow containing doxycycline throughout the entire course of the study except the “off Dox Control” group. At 12 weeks of age, all mice underwent a baseline pre-dose ECG evaluation. Mice were then treated intravenously with a single dose of either vehicle (saline), naked ASO300 (10 mg/kg), DTX-C-008 (10 mg/kg ASO equivalent) or DTX-C-008 (20 mg/kg ASO equivalent). Following baseline pre-dose ECG evaluation mice in the “off Dox Control” group were switched to chow without doxycycline. Post dose ECG evaluations were performed in all mice 7 and 14 days after treatment, or in the case of the “off Dox Control” group 7 and 14 days after reversion to chow without doxycycline. For each of the ECG spectra, QRS and QTc intervals were measured.

[000531] In this model, mice treated with doxycycline exhibit prolongation of QRS and QTC intervals driven by expression of mutant DMPK in the heart, similar to those reported in DM1 patients, and consistent with increased propensity for cardiac arrhythmia. Removal of doxycycline for the diet in the “Off Dox Control” group turns off expression of mutant DMPK, resulting in a normalization of QRS (FIG. 16B) and QTC (FIG. 16C) intervals. Mice maintained on doxycycline and treated with 20 mg/kg of the muscle-targeting complex DTX-C-

008 demonstrated statistically significant reduction in their QTc intervals after 14 days despite continued expression of mutant DMPK in the heart (FIG. 16C). This reduction in QTc intervals represents a correction in cardiac arrhythmia in a DM1 mouse model. These data demonstrate that a muscle-targeting complex as described herein is capable of providing a phenotypic and therapeutic benefit in a DM1 model.

Example 13: A muscle-targeting complex can target DMPK and correct DM1-related genetic splicing

[000532] In isolated muscles cells derived from human DM1 patients, the muscle-targeting complex as described in Example 5, DTX-C-012, was tested for reduction of DMPK expression and subsequent correction of splicing defects in Bin1, a downstream gene of DMPK.

[000533] Briefly, patient cells were seeded at a density of 10k cells/well before being allowed to recover overnight. Cells were then treated with PBS (vehicle control), naked ASO300, or DTX-C-012 (500 nM; equivalent to 55.5 nM ASO). Cells were allowed to differentiate for 14 days. Expression levels of DMPK and %Bin1 exon-11 inclusion were determined on Days 10, 11, 12, 13, and 14 post differentiation.

[000534] Treatment of DM1 patient cells with the DTX-C-012 complex leads to reduction of DMPK levels as early as Day 10 post differentiation (FIG. 17A). Treatment of DM1 patient cells with the DTX-C-012 complex also leads to a statistically significant time-dependent change in Bin1 splicing (FIG. 17B). (**p<0.01, ***p<0.001). These data demonstrate that a muscle-targeting complex as described herein is capable of providing phenotypic and therapeutic benefit (increased correction of DM1 gene-specific splicing) in a DM1 model.

Example 14: Selected antisense oligonucleotides provided dose-dependent reduction in DMPK expression in DM1 and NHP myotubes

[000535] The antisense oligonucleotides listed in Table 9 were further assessed to identify oligos that are safe in vivo (e.g., as indicated by low immunogenicity as measured by cytokine induction), and further based on manufacturability and secondary structure considerations. Three antisense oligonucleotides from Table 9, (SEQ ID NO: 212, DMPK-ASO-1), (SEQ ID NO: 222, DMPK-ASO-2), and (SEQ ID NO: 180, DMPK-ASO-3) were selected. These oligonucleotides were then further evaluated for their ability to reduce DMPK expression in DM1 myotubes and NHP myotubes in a dose-responsive manner. Naked ASO300 was used as control. Each of the antisense oligonucleotides were capable of dose-dependently reducing DMPK in DM1 and NHP myotubes (see FIGs. 18A-18C and FIGs. 19A-19B, respectively).

[000536] These data demonstrate that these antisense oligonucleotides are safe *in vivo* and are capable of dose-dependent reduction of DMPK *in cellulo*, suggesting that muscle-targeting complexes comprising these antisense oligonucleotides would be capable of targeting DMPK in muscle tissues *in vivo*.

Example 15: Binding Affinity of selected anti-TfR1 antibodies to human TfR1

[000537] Selected anti-TfR1 antibodies were tested for their binding affinity to human TfR1 for measurement of K_a (association rate constant), K_d (dissociation rate constant), and K_D (affinity). Two known anti-TfR1 antibodies were used as control, 15G11 and OKT9. The binding experiment was performed on Carterra LSA at 25°C. An anti-mouse IgG and anti-human IgG antibody "lawn" was prepared on a HC30M chip by amine coupling. 59 IgGs (58 mouse mAbs and 1 human mAb) were captured on the chip. Dilution series of hTfR1, cyTfR1, and hTfR2 were injected to the chip for binding (starting from 1000 nM, 1:3 dilution, 8 concentrations).

[000538] Binding data were referenced by subtracting the responses from a buffer analyte injection and globally fitting to a 1:1 Langmuir binding model for estimate of K_a (association rate constant), K_d (dissociation rate constant), and K_D (affinity) using the Carterra™ Kinetics software. 5-6 concentrations were used for curve fitting.

[000539] The result showed that the mouse mAbs demonstrated binding to hTfR1 with K_D values ranging from 13 pM to 50 nM. A majority of the mouse mAbs had K_D values in the single digit nanomolar to sub-nanomolar range. The tested mouse mAbs showed cross-reactive binding to cyTfR1 with K_D values ranging from 16 pM to 22 nM.

[000540] K_a , K_d , and K_D values of anti-TfR1 antibodies are provided in Table 11.

Table 11. K_a , K_d , and K_D values of anti-TfR1 antibodies

Name	K_D (M)	K_a (M)	K_d (M)
ctrl-15G11	2.83E-10	3.70E+05	1.04E-04
ctrl-OKT9 mIgG	5.36E-10	7.74E+05	4.15E-04
3-A04	4.36E-10	4.47E+05	1.95E-04
3-M12	7.68E-10	1.66E+05	1.27E-04
5-H12	2.08E-07	6.67E+04	1.39E-02

Example 16: Conjugation of anti-TfR1 antibodies with oligonucleotides

[000541] Complexes containing an anti-TfR1 antibody covalently conjugated to ASO300 were generated. First, Fab fragments of anti-TfR antibody clones 3-A4, 3-M12, and 5-H12 were prepared by cutting the mouse monoclonal antibodies with an enzyme in or below the hinge

region of the full IgG followed by partial reduction. The Fabs were confirmed to be comparable to mAbs in avidity or affinity.

[000542] Muscle-targeting complexes was generated by covalently linking the anti-TfR mAbs to the ASO300 via a cathepsin cleavable linker. Briefly, a Bicyclo[6.1.0]nonyne-PEG3-L-valine-L-citrulline-pentafluorophenyl ester (BCN-PEG3-Val-Cit-PFP) linker molecule was coupled to ASO300 through a carbamate bond. Excess linker and organic solvents were removed by tangential flow filtration (TFF). The purified Val-Cit-linker-ASO was then coupled to an azide functionalized anti-transferrin receptor antibody generated through modifying ϵ -amine on lysine with Azide-PEG4-PFP. A positive control muscle-targeting complex was also generated using 15G11.

[000543] The product of the antibody coupling reaction was then subjected to two purification methods to remove free antibody and free payload: 1) hydrophobic interaction chromatography (HIC-HPLC), and 2) Size exclusion chromatography (SEC). The HIC column utilized a decreasing salt gradient to separate free antibody from conjugate. During SEC, fractionation was performed based on A260/A280 traces to specifically collect conjugated material. Concentrations of the conjugates were determined by either Nanodrop A280 or BCA protein assay (for antibody) and Quant-It Ribogreen assay (for payload). Corresponding drug-antibody ratios (DARs) were calculated. DARs ranged between 0.8 and 2.0, and were standardized so that all samples receive equal amounts of payload.

[000544] The purified complexes were then tested for cellular internalization and inhibition of the target gene, DMPK. Non-human primate (NHP) or DM1 (donated by DM1 patients) cells were grown in 96-well plates and differentiated into myotubes for 7 days. Cells were then treated with escalating concentrations (0.5 nM, 5 nM, 50 nM) of each complex for 72 hours. Cells were harvested, RNA was isolated, and reverse transcription was performed to generate cDNA. qPCR was performed using TaqMan kits specific for Ppib (control) and DMPK on the QuantStudio 7. The relative amounts of remaining DMPK transcript in treated vs non-treated cells was calculated and the results are shown in Table 12.

[000545] The results demonstrated that the anti-TfR1 antibodies are able to target muscle cells, be internalized by the muscle cells with the molecular payload (ASO300), and that the molecular payload is able to target and knockdown the target gene (DMPK).

Table 12. Binding Affinity of anti-TfR1 Antibodies and Efficacy of Conjugates

Clone Name	huTfR1 Avg K_D (M) (antibody alone)	cyTfR1 Avg K_D (M) (antibody alone)	% knockdown of DMPK in NHP cells using	% knockdown of DMPK in cells from human DM1 patients using
------------	--	--	--	--

			Antibody-DMPK ASO conjugate	Antibody-DMPK ASO conjugate
15G11 (control)	8.0E-10	1.0E-09	36	46
3-A4	4.36E-10	2.32E-09	77	70
3-M12	7.68E-10	5.18E-09	77	52
5-H12	2.02316E-07	1.20E-08	88	57

[000546] Interestingly, the DMPK knockdown results showed a lack of correlation between the binding affinity of the anti-TfR to transferrin receptor and efficacy in delivering a DMPK-targeting ASO to cells for DMPK knockdown. Surprisingly, the anti-TfR antibodies provided herein (e.g., at least 3-A4, 3-M12, and 5-H12) demonstrated superior activity in delivering a payload (e.g., DMPK ASO) to the target cells and achieving the biological effect of the molecular payload (e.g., DMPK knockdown) in either cyno cells or human DM1 patient cells, compared to the control antibody 15G11, despite the comparable binding affinity (or, in certain instances, such as 5-H12, lower binding affinity) to human or cyno transferrin receptor between these antibodies and the control antibody 15G11.

[000547] Top attributes such as high huTfR1 affinity, >50% knockdown of DMPK in NHP and DM1 patient cell line, identified epitope binding with 3 unique sequences, low/no predicted PTM sites, and good expression and conjugation efficiency led to the selection of the top 3 clones for humanization, 3-A4, 3-M12, and 5-H12.

Example 17: Humanized anti-TfR1 antibodies

[000548] The anti-TfR antibodies shown in Table 2 were subjected to humanization and mutagenesis to reduce manufacturability liabilities. The humanized variants were screened and tested for their binding properties and biological actives. Humanized variants of anti-TfR1 heavy and light chain variable regions (5 variants each) were designed using Composite Human Technology. Genes encoding Fabs having these heavy and light chain variable regions were synthesized, and vectors were constructed to express each humanized heavy and light chain variant. Subsequently, each vector was expressed on a small scale and the resultant humanized anti-TfR1 Fabs were analyzed. Humanized Fabs were selected for further testing based upon several criteria including Biacore assays of antibody affinity for the target antigen, relative expression, percent homology to human germline sequence, and the number of MHC class II predicted T cell epitopes (determined using iTope™ MCH class II in silico analysis).

[000549] Potential liabilities were identified within the parental sequence of some antibodies by introducing amino acid substitutions in the heavy chain and light chain variable regions. These substitutions were chosen based on relative expression levels, iTope™ score and relative K_D from Biacore single cycle kinetics analysis. The humanized variants were tested and

variants were selected initially based upon affinity for the target antigen. Subsequently, the selected humanized Fabs were further screened based on a series of biophysical assessments of stability and susceptibility to aggregation and degradation of each analyzed variant, shown in **Table 14** and **Table 15**. The selected Fabs were analyzed for their properties binding to TfR1 by kinetic analysis. The results of these analyses are shown in **Table 16**. For conjugates shown in Table 14 and Table 15, the selected humanized Fabs were conjugated to a DMPK-targeting oligonucleotide ASO300. The selected Fabs are thermally stable, as indicated by the comparable binding affinity to human and cyno TfR1 after been exposed to high temperature (40 °C) for 9 days, compared to before the exposure (see Table 16).

Table 14. Biophysical assessment data for humanized anti-TfR Fabs

Variant	3M12 (VH3/Vk2)	3M12 (VH3/Vk3)	3M12 (VH4/Vk2)	3M12 (VH4/Vk3)	3A4 (VH3-N54T/Vk4)
Binding Affinity (Biacore d0)	395 pM	345 pM	396 pM	341 pM	3.09 nM
Binding Affinity (Biacore d25)	567 pM	515 pM	510 pM	486 pM	3.01 nM
Fab binding affinity ELISA (human/cyno TfR1)	0.8 nM/9.9 nM	0.6 nM/4.7 nM	0.4 nM/1.4 nM	0.5 nM/2.2 nM	2.6 nM/156 nM*
Conjugate binding affinity ELISA (human/cyno TfR1)	2.2 nM/2.9 nM	N/A	N/A	1.7 nM/2.1 nM	2.8 nM/4.7 nM
...					
Variant	3A4 (VH3-N54S/Vk4)	3A4 (VH3/Vk4)	5H12 (VH5-C33Y/Vk3)	5H12 (VH5-C33D/Vk4)	5H12 (VH4-C33Y/Vk4)
Binding Affinity (Biacore d0)	1.34 nM	1.5 nM	627 pM	991 pM	626 pM
Binding Affinity (Biacore d25)	1.39 nM	1.35 nM	1.07 nM	3.01 nM	1.33 nM
Fab binding affinity ELISA (human/cyno TfR1)	1.6 nM/398 nM*	1.5 nM/122 nM*	6.3 nM/2.1 nM	6.0 nM/3.5 nM	2.8 nM/3.3 nM
Conjugate binding affinity ELISA (human/cyno TfR1)	2.9 nM/7.8 nM	2.8 nM/7.6 nM	33.4 nM/2.3 nM	110 nM/10.2 nM	23.7 nM/3.3 nM

*Regains cyno binding after conjugation;

Table 15. Thermal Stability for humanized anti-TfR Fabs and conjugates

Variant	3M12 (VH3/Vk2)	3M12 (VH3/Vk3)	3M12 (VH4/Vk2)	3M12 (VH4/Vk3)	3A4 (VH3-N54T/Vk4)
Binding affinity hTfR1 d0 (nM)	0.8	0.6	0.4	0.5	2.6
Binding affinity hTfR1 d9 (nM)	0.98	1.49	0.50	0.28	0.40
Binding affinity cyno TfR1 d0 (nM)	9.9	4.7	1.4	2.2	156
Binding affinity cyno TfR1 d9 (nM)	19.51	15.58	5.01	16.40	127.50

DMPK oligo conjugate binding to hTfR1 (nM)	1.14	N/A	N/A	1.18	2.22
DMPK oligo conjugate binding to cyno TfR1 (nM)	2.26	N/A	N/A	1.85	5.12
...					
Variant	3A4 (VH3-N54S/Vk4)	3A4 (VH3/Vk4)	5H12 (VH5-C33Y/Vk3)	5H12 (VH5-C33D/Vk4)	5H12 (VH4-C33Y/Vk4)
Criteria					
Binding affinity hTfR1 d0 (nM)	1.6	1.5	6.3	6	2.8
Binding affinity hTfR1 d9 (nM)	0.65	0.46	71.90	92.34	1731.00
Binding affinity cyno TfR1 d0 (nM)	398	122	2.1	3.5	3.3
Binding affinity cyno TfR1 d9 (nM)	248.30	878.40	0.69	0.63	0.26
DMPK oligo conjugate binding to hTfR1 (nM)	2.71	2.837	N/A	110.5	13.9
DMPK oligo conjugate binding to cyno TfR1 (nM)	4.1	7.594	N/A	10.18	13.9

Table 16. Kinetic analysis of humanized anti-TfR Fabs binding to TfR1

Humanized anti-TfR Fabs	k_a (1/Ms)	k_d (1/s)	K_D (M)	R_{MAX}	χ^2 (RU ²)
3A4 (VH3/Vk4)	7.65E+10	1.15E+02	1.50E-09	48.0	0.776
3A4 (VH3-N54S/Vk4)	4.90E+10	6.56E+01	1.34E-09	49.4	0.622
3A4 (VH3-N54T/Vk4)	2.28E+05	7.05E-04	3.09E-09	61.1	1.650
3M12 (VH3/Vk2)	2.64E+05	1.04E-04	3.95E-10	78.4	0.037
3M12 (VH3/Vk3)	2.42E+05	8.34E-05	3.45E-10	91.1	0.025
3M12 (VH4/Vk2)	2.52E+05	9.98E-05	3.96E-10	74.8	0.024
3M12 (VH4/Vk3)	2.52E+05	8.61E-05	3.41E-10	82.4	0.030
5H12 (VH5-C33D/Vk4)	6.78E+05	6.72E-04	9.91E-10	49.3	0.093
5H12 (VH5-C33Y/Vk3)	1.95E+05	1.22E-04	6.27E-10	68.5	0.021
5H12 (VH5-C33Y/Vk4)	1.86E+05	1.17E-04	6.26E-10	75.2	0.026

Binding of humanized anti-TfR1 Fabs to TfR1 (ELISA)

[000550] To measure binding of humanized anti-TfR antibodies to TfR1, ELISAs were conducted. High binding, black, flat bottom, 96 well plates (Corning# 3925) were first coated with 100 μ L/well of recombinant huTfR1 at 1 μ g/mL in PBS and incubated at 4°C overnight. Wells were emptied and residual liquid was removed. Blocking was conducted by adding 200 μ L of 1%BSA (w/w) in PBS to each well. Blocking was allowed to proceed for 2 hours at room temperature on a shaker at 300 rpm. After blocking, liquid was removed and wells were washed three times with 300 μ L of TBST. Anti-TfR1 antibodies were then added in 0.5% BSA/TBST in triplicate in an 8 point serial dilution (dilution range 5 μ g/mL – 5 ng/mL). A positive control and isotype controls were also included on the ELISA plate. The plate was incubated at room temperature on an orbital shaker for 60 minutes at 300 rpm, and the plate was washed three times with 300 μ L of TBST. Anti-(H+L)IgG-A488 (1:500) (Invitrogen #A11013) was diluted in

0.5% BSA in TBST, and 100 μ L was added to each well. The plate was then allowed to incubate at room temperature for 60 minutes at 300 rpm on orbital shaker. The liquid was removed and the plate was washed four times with 300 μ L of TBST. Absorbance was then measured at 495 nm excitation and 50 nm emission (with a 15 nm bandwidth) on a plate reader. Data was recorded and analyzed for EC₅₀. The data for binding to human TfR1 (hTfR1) for the humanized 3M12, 3A4 and 5H12 Fabs are shown in **FIG. 21A**, **21C**, and **21E**, respectively. ELISA measurements were conducted using cynomolgus monkey (*Macaca fascicularis*) TfR1 (cTfR1) according to the same protocol described above for hTfR1, and results are shown in **FIG. 21B**, **21D**, and **21F**.

[000551] Results of these two sets of ELISA analyses for binding of the humanized anti-TfR Fabs to hTfR1 and cTfR1 demonstrate that humanized 3M12 Fabs show consistent binding to both hTfR1 and cTfR1, and that humanized 3A4 Fabs show decreased binding to cTfR1 relative to hTfR1.

[000552] Antibody-oligonucleotide conjugates were prepared using six humanized anti-TfR Fabs, each of which were conjugated to a DMPK targeting oligonucleotide ASO300. Conjugation efficiency and down-stream purification were characterized, and various properties of the product conjugates were measured. The results demonstrate that conjugation efficiency was robust across all 10 variants tested, and that the purification process (hydrophobic interaction chromatography followed by hydroxyapatite resin chromatography) were effective. The purified conjugates showed a >97% purity as analyzed by size exclusion chromatography.

[0006] Several humanized Fabs were tested in cellular uptake experiments to evaluate TfR1-mediated internalization. To measure such cellular uptake mediated by antibodies, humanized anti-TfR Fab conjugates were labeled with Cypher5e, a pH-sensitive dye. Rhabdomyosarcoma (RD) cells were treated for 4 hours with 100 nM of the conjugates, trypsinized, washed twice, and analyzed by flow cytometry. Mean Cypher5e fluorescence (representing uptake) was calculated using Attune NxT software. As shown in FIG. 22, the humanized anti-TfR Fabs show similar or greater endosomal uptake compared to a positive control anti-TfR1 Fab. Similar internalization efficiencies were observed for different oligonucleotide payloads. An anti-mouse TfR antibody was used as the negative control. Cold (non-internalizing) conditions abrogated the fluorescence signal of the positive control antibody-conjugate (data not shown), indicating that the positive signal in the positive control and humanized anti-TfR Fab-conjugates is due to internalization of the Fab-conjugates.

[000553] Conjugates of six humanized anti-TfR Fabs were also tested for binding to hTfR1 and cTfR1 by ELISA, and compared to the unconjugated forms of the humanized Fabs. Results demonstrate that humanized 3M12 and 5H12 Fabs maintain similar levels of hTfR1 and

cTfR1 binding after conjugation relative to their unconjugated forms (3M12, **FIG. 23A** and **23B**; 5H12, **FIG. 23E** and **23F**). Interestingly, 3A4 clones show improved binding to cTfR1 after conjugation relative to their unconjugated forms (**FIG. 23C** and **23D**).

[000554] As used in this Example, the term ‘unconjugated’ indicates that the antibody was not conjugated to an oligonucleotide.

Example 18. Knockdown of DMPK mRNA level facilitated by oligonucleotides conjugates *in vitro*

[000555] DMPK-targeting oligonucleotides (e.g., ASO) were tested in rhabdomyosarcoma (RD) cells for knockdown of DMPK transcript expression. RD cells were cultured in a growth medium of DMEM with glutamine, supplemented with 10% FBS and penicillin/streptomycin until nearly confluent. Cells were then seeded into a 96 well plate at 20K cells per well and were allowed to recover for 24 hours. Cells were then treated with free DMPK-targeting oligonucleotides or by transfection of the oligonucleotides using 0.3 μ L per well of Lipofectamine MessengerMAX transfection reagent. After 3 days, total RNA was collected from cells, cDNA was synthesized and DMPK expression was measured by qPCR.

[000556] Results in FIG. 24 show that DMPK expression level was reduced in cells treated with each given DMPK-targeting oligonucleotide, relative to expression in PBS-treated cells. Several DMPK-oligonucleotides showed dose-dependent reduction of DMPK expression level. In FIG. 24, DMPK-ASO-1 has the sequence GCGUAGAAGGGCGUCUGCCC (SEQ ID NO: 212). DMPK-ASO-2 has the sequence CCCAGCGCCCACCAGUCACA (SEQ ID NO: 222). DMPK-ASO-3 has the sequence CCAUCUCGGCCGGAUCCGC (SEQ ID NO: 180). ASO300 was also used in this experiment.

Example 19. Knockdown of DMPK mRNA level facilitated by antibody-oligonucleotide conjugates *in vitro*

[000557] Conjugates containing humanized anti-TfR Fabs 3M12(VH3/Vk2), 3M-12 (VH4/Vk3), and 3A4(VH3-N54S/Vk4) were conjugated to a DMPK-targeting oligonucleotide (ASO300) and were tested in rhabdomyosarcoma (RD) cells for knockdown of DMPK transcript expression. Antibodies were conjugated to ASO300 via the linker shown in Formula (C).

[000558] RD cells were cultured in a growth medium of DMEM with glutamine, supplemented with 10% FBS and penicillin/streptomycin until nearly confluent. Cells were then seeded into a 96 well plate at 20K cells per well and were allowed to recover for 24 hours. Cells were then treated with the conjugates for 3 days. Total RNA was collected from cells, cDNA was synthesized and DMPK expression was measured by qPCR.

[000559] Results in FIG. 25 show that DMPK expression level was reduced in cells treated with each indicated conjugate, relative to expression in PBS-treated cells, indicating that the humanized anti-TfR Fabs are able to mediate the uptake of the DMPK-targeting oligonucleotide by the RD cells and that the internalized DMPK-targeting oligonucleotide are effective in knocking down DMPK mRNA level.

Example 20. Splicing correction and functional efficacy in HSA-LR mouse model of DM1

[000560] Correction of splicing in the HSA-LR mouse model of DM1 was demonstrated with conjugates containing anti-TfR antibodies conjugated to oligonucleotides target human skeletal actin (ACTA1). The anti-TfR1 antibody used in this study was RI7 217. The oligonucleotide targeting ACTA1 is an MOE 5-10-5 gapmer that comprises: 5'-NH₂-(CH₂)₆-dA*oC*oC*oA*oT*oT*dT*dT*dC*dT*dT*dC*dC*dA*dC*dA*oG*oG*oG*oC*oT-3 (SEQ ID NO: 620); wherein '*' represents a PS linkage; 'd' represents a deoxynucleic acid; and 'o' represents a 2'-MOE.

[000561] The HSA-LR DM1 mouse model is a well-validated model of DM1 that exhibits pathologies that are very similar to human DM1 patients. The HSA-LR model uses the human skeletal actin (ACTA1) promoter to drive expression of CUG long repeats (LR). In this model, toxic DMPK RNA accumulates within the nucleus and sequesters proteins responsible for splicing, such as Muscleblind-like protein (MBNL), resulting in mis-splicing of multiple RNAs, including CLCN1 (chloride channel), ATP2a1 (calcium channel), and others. This mis-splicing causes the mice to also exhibit myotonia which is a hallmark of the DM1 clinical presentation in humans.

[000562] The anti-TfR-oligonucleotide conjugate delivered intravenously has been shown to have activity in dose-dependent correction of splicing in multiple RNAs and multiple muscles and was well tolerated by HSA-LR mice. In this study, the ability of the conjugates to correct splicing in more than 30 different RNAs was evaluated. In DM1, significant RNA mis-splicing of these RNAs reduces the ability of multiple muscles to function. The RNAs monitored are critical for contraction and relaxation of muscle in HSA-LR mice. Dose-dependent correction of splicing was observed.

[000563] FIG. 26 shows results for Atp2a1, which encodes a calcium channel and contributes to muscle contraction and relaxation. The X-axis represents splice derangement with 1.00 representing severe mis-splicing and 0.00 representing a wild type (WT) splice pattern. Progression from right to left in the figure represents a correction of splicing. The Y-axis represents the percent of the gene spliced in (PSI). Severe mis-splicing of ATP2a1 is caused by exclusion of exon 22 in the ATP2a1 RNA. WT splicing reflects near complete inclusion of exon

22. Results demonstrate that the conjugate corrected splicing of ATP2a1 in a dose-dependent manner in the gastrocnemius muscle.

[000564] Data for the more than 30 different RNAs that were tested in this study are shown in FIGs. 27, 35A, and 35B. Similar dose-dependent correction of splicing was achieved for all of the tested RNAs in gastrocnemius muscle (FIG. 27), tibialis anterior (FIG. 35A), and quadriceps (FIG. 35B). For some of these RNAs, correction of splicing is reflected by a decrease in PSI, as in FIG. 26, and for other RNAs correction is reflected by an increase in PSI.

[000565] Similar dose-dependent improvements in splicing within the set of RNAs were observed in the quadriceps and tibialis anterior muscles, after treatment with the conjugate. FIG. 28 shows composite levels of splicing derangement observed for saline and different doses of the Ab-ASO across the more than 30 RNAs that were tested in each muscle type. Doses of 10mg/kg and 20mg/kg were administered in this study.

[000566] In addition to reductions in splicing derangement across multiple genes in several muscles in the HSA-LR model, disease modification was observed in the HSA-LR model. The results in FIG. 29 show that almost complete reversal of myotonia was achieved after a single dose of the conjugate. The severity of myotonia on a four-point scale was evaluated 14 days following dosing with saline (PBS), naked oligonucleotide, or the conjugate. Grade 0 indicates no myotonia was observed, grade 1 indicates myotonic discharge was measured by electromyography (EMG) in less than 50% of needle insertions, grade 2 indicates myotonic discharge was measured in greater than 50% of needle insertions and grade 3 indicates myotonic discharge was measured with nearly every needle insertion.

Example 21. DMPK-targeting PMOs

[000567] Additional DMPK targeting oligonucleotides (PMOs) were designed and tested for their activity in reducing DMPK expression in primary human myotubes. Wild-type primary myoblasts were cultured in PromoCell Skeletal Muscle growth medium with 5% FBS and penicillin/streptomycin until nearly confluent. Cells were then seeded into a 96 well plate at 50K cells per well and allowed to recover for 24 hours. Cells were then differentiated in a differentiation medium of DMEM with glutamine and penicillin/streptomycin for 7 days. Cells were then treated with the unconjugated PMO for 3 days. Total RNA was collected from cells, cDNA was synthesized and DMPK expression was measured by qPCR. The sequences of the PMOs and their activity in knocking down DMPK in vitro are shown in Table 17.

[000568] As used in this Example, the term ‘unconjugated’ indicates that the oligonucleotide was not conjugated to an antibody.”

Table 17. DMPK-targeting PMOs and activity in knocking down DMPK in vitro

PMO Sequence	SEQ ID NO	Target region	% DMPK Knockdown in Primary Human Myotubes
CAGGTGACAGTTCAGGTGCAG	624	Intron 1-2	59%
TCCACCCTGACTCCAGGTGAC	625	Intron 1-2	16%
GAGAAGGAAATAAGACCCAGTT	626	Intron 1-2	14%
CCTTCTCTCTGCCTCTCAGCTT	627	Intron 1-2	58%
CCACCCTCTGTCTGTCTCC	628	Intron 1-2	64%
TCCGCTGGGTGGTGGGAAAAGAA	629	Exon 2	9%
ATGGGCTCCGCTGGGTGGTGG	630	Exon 2	11%
ACGATGGGCTCCGCTGGG	631	Exon 2	60%
CCATCCTTGGGCAGAGACCT	632	Intron 4-5	41%
ATGACCAGGTACTGAGAAGGG	633	Exon 5	33%
AGGTACTGAGAAGGGTTCGTC	634	Exon 5	40%
TAGGGACCTGCGGAGAGGGCGA	635	Exon 15	35%
GCCTAGGGACCTGCGGAGAG	636	Exon 15	62%
GCCTTTTATTCGCGAGGGTTCGG	637	polyA	73%
TGGAGGGCCTTTTATTCGCGAGG	638	polyA	66%
TAGGCACTACCCACTGCAAGA	621	Exon 1	69%
CGGAGCTACCAAGGTAGTTCT	622	Intron 4-5	73%
AGGGCAGTGCTTACCTGAGGG	623	Intron 9-10	57%

Example 22. Serum stability of the linker linking the anti-TfR antibody and the molecular payload

[000569] Oligonucleotides which were linked to antibodies in examples were conjugated via a cleavable linker shown in Formula (C). It is important that the linker maintain stability in serum and provide release kinetics that favor sufficient payload accumulation in the targeted muscle cell. This serum stability is important for systemic intravenous administration, stability of the conjugated oligonucleotide in the bloodstream, delivery to muscle tissue and internalization of the therapeutic payload in the muscle cells. The linker has been confirmed to facilitate precise conjugation of multiple types of payloads (including ASOs, siRNAs and PMOs) to Fabs. This flexibility enabled rational selection of the appropriate type of payload to address the genetic basis of each muscle disease. Additionally, the linker and conjugation chemistry allowed the optimization of the ratio of payload molecules attached to each Fab for each type of payload, and enabled rapid design, production and screening of molecules to enable use in various muscle disease applications.

[000570] FIG. 20 shows serum stability of the linker *in vivo*, which was comparable across multiple species over the course of 72 hours after intravenous dosing. At least 75% stability was measured in each case at 72 hours after dosing.

Example 23. *In vivo* activity of anti-TfR conjugates in hTfR1 mice

[000571] In DM1, the higher than normal number of CUG repeats form large hairpin loops that remain trapped in the nucleus, forming nuclear foci that bind splicing proteins and inhibit the ability of splicing proteins to perform their normal function. When toxic nuclear DMPK levels are reduced, the nuclear foci are diminished, releasing splicing proteins, allowing restoration of normal mRNA processing, and potentially stopping or reversing disease progression.

[000572] The *in vivo* activity of conjugates containing an anti-TfR Fab (control, 3M12 VH3/VK2, 3M12 VH4/VK3, 3A4 VH3 N54S/VK4) conjugated to the DMPK-targeting oligonucleotide (ASO300) in reducing DMPK mRNA level in multiple muscle tissues following systemic intravenous administration in mice was evaluated.

[000573] Male and female C57BL/6 mice where one TfR1 allele was replaced with a human TFR1 allele were administered between the ages of 5 and 15 weeks according to the dosing schedule outlined in Table 18 and in FIG. 30A. Mice were sacrificed 14 days after the first injection and selected muscles collected as indicated in Table 19.

Table 18							
Group	Animal No.	Treatment Antibody	Treatment Oligo	Dose Level (mg/kg)	Dose Volume (mL/kg)	Dosing Regimen	Terminal Time Point
1	4	Vehicle	NA	0	10	Day 0 and Day 7 by IV	Day 14
2	4	NA	ASO300	10	5.0		
3	4	control anti-TfR Fab	ASO300		10.2		
4	4	3M12 VH3/VK2	ASO300		11.5		
5	4	3M12 VH4/VK3	ASO300		10.1		
6	4	3A4 VH3 N54S/VK4	ASO300		10.7		

Table 19	
Tissue	Storage
Gastrocnemius	Right leg of each animal stored in RNALater at -80 °C
Tibialis Anterior	One leg (R) of each animal stored in RNALater at -80 °C
Heart	Dissect transversally and store the apex in RNALater at -80 °C
Diaphragm	Split in half and collect one half in RNALater at -80 °C

[000574] Total RNA was extracted on a Maxwell Rapid Sample Concentrator (RSC) Instrument using kits provided by the manufacturer (Promega). Purified RNA was reverse-transcribed and levels of *Dmpk* and *Ppib* transcripts determined by qRT-PCR with specific TaqMan assays (ThermoFisher). Log fold changes in *Dmpk* expression were calculated according to the $2^{-\Delta\Delta CT}$ method using *Ppib* as the reference gene and mice injected with vehicle as the control group. Statistical significance in differences of *Dmpk* expression between control mice and mice administered with the conjugates were determined by one-way ANOVA with Dunnet's correction for multiple comparisons. As shown in FIGs. 30B-30E, the tested conjugates showed robust activity in reducing DMPK mRNA level *in vivo* in various muscle tissues.

Example 24. *In vitro* activity of anti-TfR conjugates in patient-derived cells

[000575] An *in vitro* experiment was conducted to determine the activities of anti-TfR conjugates in reducing DMPK mRNA expression, correcting BIN1 splicing, and reducing nuclear foci in CM-DM1-32F primary cells expressing a mutant DMPK mRNA containing 380 CUG repeats. The CM-DM1-32F primary cell is an immortalized myoblastic cell line isolated from a DM1 patient (CL5 cells; Described in Arandel et al., *Dis Model Mech.* 2017 Apr 1; 10(4): 487–497). Conjugate 1 contains an anti-TfR mAb conjugated to DMPK-targeting oligonucleotide (ASO300). Conjugate 2 contains an anti-TfR Fab conjugated to DMPK ASO-1 (GCGUAGAAGGGCGUCUGCCC; SEQ ID NO: 212).

[000576] CL5 cells were seeded at a density of 156,000 cells/cm², allowed to recover for 24 hours, transferred to differentiation media to induce myotube formation, as described (Arandel et al. *Dis Model Mech.* (2017) 10(4):487-497) and subsequently exposed to conjugate 1 and conjugate 2 at a payload concentration of 500 nM. Parallel cultures exposed to vehicle PBS served as controls. Cells were harvested after 10 days of culture.

[000577] For analysis of gene expression, cells were collected with Qiazol for total RNA extraction with a Qiagen miRNAeasy kit. Purified RNA was reverse-transcribed and levels of *DMPK*, *PPIB*, *BIN1* transcripts and of the *BIN1* mRNA isoform containing exon 11 determined by qRT-PCR with specific TaqMan assays (ThermoFisher). Log fold changes in *DMPK* expression were calculated according to the $2^{-\Delta\Delta CT}$ method using *PPIB* as the reference gene and cells exposed to vehicle as the control group. Log fold changes in the levels *BIN1* isoform containing exon 11 were calculated according to the $2^{-\Delta\Delta CT}$ method using *BIN1* as the reference gene and cells exposed to vehicle as the control group.

[000578] To measure the area of mutant DMPK nuclear foci, cells were fixed in 4% formalin, permeabilized with 0.1% Triton X-100 and hybridized at 70 °C with a CAG peptide-

nucleic acid probe conjugated to the Cy5 fluorophore. After multiple washes in hybridization buffer and 2xSSC solution, nuclei were counterstained with DAPI. Images were collected at a 400x magnification by confocal microscopy and foci area measured as the area of Cy5 signal contained within the area of DAPI signal. Data were expressed as foci area corrected for nuclear area.

[000579] The results show that a single dose of the conjugates containing an anti-TfR (IgG or Fab) conjugated to a DMPK-targeting oligonucleotide (ASO300 or DMPK ASO-1 (SEQ ID NO: 212)) resulted reduced mutant DMPK expression (FIG. 31A), corrected BIN1 splicing (FIG. 31B), and reduced nuclear foci by approximately 40% (FIG. 31C).

Example 25. Characterization of binding activities of anti-TfR Fab 3M12 VH4/Vk3

[000580] In vitro studies were performed to test the specificity of anti-TfR Fab 3M12 VH4/Vk3 for human and cynomolgus monkey TfR1 binding and to confirm its selectivity for human TfR1 vs TfR2. The binding affinity of anti-TfR Fab 3M12 VH4/Vk3 to TfR1 from various species was determined using an enzyme-linked immunosorbent assay (ELISA). Serial dilutions of the Fab were added to plates precoated with recombinant human, cynomolgus monkey, mouse, or rat TfR1. After a short incubation, binding of the Fab was quantified by addition of a fluorescently tagged anti-(H+L) IgG secondary antibody and measurement of fluorescence intensity at 495nm excitation and 520nm emission. The Fab showed strong binding affinity to human and cynomolgus monkey TfR1, and no detectable binding of mouse or rat TfR1 was observed (FIG. 32). Surface plasmon resonance (SPR) measurements were also conducted, and results are shown in Table 20. The K_d of the Fab against the human TfR1 receptor was calculated to be 7.68×10^{-10} M and against the cynomolgus monkey TfR1 receptor was calculated to be 5.18×10^{-9} M.

Table 20. Kinetic analysis of anti-TfR Fab 3M12 VH4/Vk3 binding to human and cynomolgus monkey TfR1 or human TfR2, measured using surface plasmon resonance

Target	Anti-TfR Fab 3M12 VH4/Vk3				
	K_d (M)	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	R_{max}	R_{es} SD
Human TfR1	7.68E-10	1.66E+05	1.27E-04	1.11E+02	3.45E+00
Cyno TfR1	5.18E-09	9.19E+04	4.76E-04	1.87E+02	6.24E+00
Human TfR2	ND	ND	ND	ND	ND

ND = No detectable binding by SPR (10pM – 100 uM)

[000581] To test for cross-reactivity of anti-TfR Fab 3M12 VH4/Vk3 to human TfR2, an ELISA was performed. Recombinant human TfR2 protein was plated overnight at $2 \mu g/mL$ and was blocked for 1 hour with 1% bovine serum albumin (BSA) in PBS. Serial dilutions of the Fab or a positive control anti-TfR2 antibody were added in 0.5% BSA/TBST for 1 hour. After

washing, anti-(H+L) IgG-A488 (Invitrogen #MA5-25932) fluorescent secondary antibody was added at a 1:500 dilution in 0.5% BSA/TBST and the plate was incubated for 1 hour. Relative fluorescence was measured using a Biotek Synergy plate reader at 495nm excitation and 520nm emission. No binding of anti-TfR Fab 3M12 VH4/Vk3 to hTfR2 was observed (FIG. 33).

Example 26. Serum stability of anti-TfR Fab-ASO conjugate

[000582] Anti-TfR Fab VH4/Vk3 was conjugated to a control antisense oligonucleotide (ASO) via a linker as shown in Formula (C) and the resulting conjugate was tested for stability of the linker conjugating the Fab to the ASO. Serum stability was measured by incubating fluorescently labeled conjugate in PBS or in rat, mouse, cynomolgus monkey, or human serum and measuring relative fluorescence intensity over time, with higher fluorescence indicating more conjugate remaining intact. FIG. 34 shows serum stability was similar across multiple species and remained high after 72 hours.

Example 27. Effect of a single dose of an anti-mouse TfR Fab conjugated to an oligonucleotide (ASO) against the human *ACTA1* mRNA in HSA^{LR} mice, a model of DM1.

[000583] Six- to 8-week-old homozygous HSA^{LR} mice were allocated randomly to one of five treatment groups and treated with vehicle, a naked oligonucleotide (ASO) targeting human *ACTA1* mRNA at a dose of 10 mg/kg or 20 mg/kg, or conjugates containing anti-TfR RI7217 Fab conjugated to the ASO (Ab-ASO) at a dose equivalent to 10 mg/kg or 20 mg/kg of ASO.

[000584] Twenty-eight days after intravenous injections of vehicle, ASO, or Ab-ASO, EMG myotonic discharges in quadriceps (FIG. 36A), gastrocnemius (FIG. 36B), and tibialis anterior (FIG. 36C), were graded by blinded examiner on a 4-point scale in which 0 indicated no myotonia; 1 indicated occasional myotonic discharge in less than 50% of needle insertions; 2 indicated myotonic discharge in greater than 50% of needle insertions; and 3 indicated myotonic discharge with nearly every insertion. A single dose of the conjugate, but not naked ASO, dose-dependently reversed myotonia in the HSA^{LR} DM1 model.

[000585] Statistical significance of differences between vehicle-treated group and each experimental arm was determined by Kruskal-Wallis test with Dunn's multiple comparisons test. Data are reported as means $\pm$ S.E.M.; * $p < 0.05$, ** $p < 0.01$.

[000586] Additionally, fourteen days and twenty-eight days after the treatment mice were sacrificed and quadriceps (quad), gastrocnemius (gastroc) and tibialis anterior (TA) muscles were collected and analyzed for expression of *ACTA1*. Knockdown (KD) of *ACTA1* expression using the Ab-ASO conjugate was observed in quadriceps (quad), gastrocnemius (gastroc) and tibialis anterior (TA) muscles relative to PBS control 14 days following a single

administered dose, whereas naked ASO did not facilitate ACTA1 suppression in the same timeframe (FIG 37). Measurement of ACTA1 in muscle 28 days following the treatment showed that the, the Ab-ASO conjugate reduced ACTA1 expression relative to vehicle control at both tested doses (10 mg/kg ASO equivalent and 20 mg/kg ASO equivalent), whereas the same doses of naked ASO did not facilitate ACTA1 suppression (FIGs. 38A-38C; * $p < 0.05$, ** $p < 0.01$).

ADDITIONAL EMBODIMENTS

1. A complex comprising a muscle-targeting agent covalently linked to a molecular payload configured for inhibiting expression or activity of a DMPK allele comprising a disease-associated-repeat, wherein the muscle-targeting agent specifically binds to an internalizing cell surface receptor on muscle cells,

wherein the muscle targeting agent is a humanized antibody that binds to a transferrin receptor, wherein the antibody comprises:

(i) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 69; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(ii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 71; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(iii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 72; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(iv) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 74;

(v) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 75;

(vi) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 74;

(vii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 75;

(viii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 78;

(ix) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 79; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 80; or

(x) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 80.

2. The complex of embodiment 1, wherein the antibody comprises:

(i) a VH comprising the amino acid sequence of SEQ ID NO: 69 and a VL comprising the amino acid sequence of SEQ ID NO: 70;

(ii) a VH comprising the amino acid sequence of SEQ ID NO: 71 and a VL comprising the amino acid sequence of SEQ ID NO: 70;

(iii) a VH comprising the amino acid sequence of SEQ ID NO: 72 and a VL comprising the amino acid sequence of SEQ ID NO: 70;

(iv) a VH comprising the amino acid sequence of SEQ ID NO: 73 and a VL comprising the amino acid sequence of SEQ ID NO: 74;

(v) a VH comprising the amino acid sequence of SEQ ID NO: 73 and a VL comprising the amino acid sequence of SEQ ID NO: 75;

(vi) a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of SEQ ID NO: 74;

(vii) a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of SEQ ID NO: 75;

(viii) a VH comprising the amino acid sequence of SEQ ID NO: 77 and a VL comprising the amino acid sequence of SEQ ID NO: 78;

(ix) a VH comprising the amino acid sequence of SEQ ID NO: 79 and a VL comprising the amino acid sequence of SEQ ID NO: 80; or

(x) a VH comprising the amino acid sequence of SEQ ID NO: 77 and a VL comprising the amino acid sequence of SEQ ID NO: 80.

3. The complex of embodiment 1 or embodiment 2, wherein the antibody is selected from the group consisting of a full-length IgG, a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, a scFv, and a Fv.

4. The complex of embodiment 3, wherein the antibody is a full-length IgG, optionally wherein the full-length IgG comprises a heavy chain constant region of the isotype IgG1, IgG2, IgG3, or IgG4.
5. The complex of embodiment 4, wherein the antibody comprises:
 - (i) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 84; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;
 - (ii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 86; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;
 - (iii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 87; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;
 - (iv) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 88; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;
 - (v) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 88; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;
 - (vi) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 91; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;
 - (vii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 91; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;
 - (viii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 92; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 93;
 - (ix) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 94; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95; or
 - (x) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 92; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95.
6. The complex of embodiment 3, wherein the antibody is a Fab fragment.

7. The complex of embodiment 6, wherein the antibody comprises:

(i) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 97; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(ii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 98; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 99; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iv) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(v) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(vi) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(vii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(viii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 93;

(ix) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 103; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95; or

(x) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95.

8. The complex of embodiment 6 or embodiment 7, wherein the antibody comprises:

(i) a heavy chain comprising the amino acid sequence of SEQ ID NO: 97; and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(ii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 98; and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 99; and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iv) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100; and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(v) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100; and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(vi) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101; and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(vii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101; and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(viii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102; and a light chain comprising the amino acid sequence of SEQ ID NO: 93;

(ix) a heavy chain comprising the amino acid sequence of SEQ ID NO: 103; and a light chain comprising the amino acid sequence of SEQ ID NO: 95; or

(x) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102; and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

9. The complex of any one of embodiments 1 to 8, wherein the equilibrium dissociation constant (K_D) of binding of the antibody to the transferrin receptor is in a range from 10^{-11} M to 10^{-6} M.

10. The complex of any one of embodiments 1 to 9, wherein the antibody does not specifically bind to the transferrin binding site of the transferrin receptor and/or wherein the antibody does not inhibit binding of transferrin to the transferrin receptor.

11. The complex of any one of embodiments 1 to 10, wherein the antibody is cross-reactive with extracellular epitopes of two or more of a human, non-human primate and rodent transferrin receptor.

12. The complex of any one of embodiments 1 to 11, wherein the complex is configured to promote transferrin receptor mediated internalization of the molecular payload into a muscle cell.

13. The complex of any one of embodiments 1 to 12, wherein the antibody is a chimeric antibody, optionally wherein the chimeric antibody is a humanized monoclonal antibody.

14. The complex of any one of embodiments 1 to 13, wherein the antibody is in the form of a ScFv, Fab fragment, Fab' fragment, F(ab')₂ fragment, or Fv fragment.

15. The complex of any one of embodiments 1 to 14, wherein the molecular payload is an oligonucleotide.
16. The complex of embodiment 15, wherein the oligonucleotide comprises at least 15 consecutive nucleotides of a sequence comprising any one of SEQ ID NOs: 148-383 and 621-638.
17. The complex of embodiment 16, wherein the oligonucleotide comprises a sequence comprising any one of SEQ ID NOs: 148-383 and 621-638.
18. The complex of embodiment 17, wherein the oligonucleotide comprises a sequence comprising any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264.
- 18.1. The complex of embodiment 18, wherein the oligonucleotide comprises a sequence comprising any one of SEQ ID NOs: 180, 212, and 222.
19. The complex of any one of embodiments 1-14, wherein the oligonucleotide comprises a region of complementarity to any one of SEQ ID NO: 384-619.
20. The complex of embodiment 19, wherein the oligonucleotide comprises a region of complementarity to at least 15 consecutive nucleotides of any one of SEQ ID NO: 384-619.
21. The complex of any one of embodiments 15 to 20, wherein the oligonucleotide comprises a region of complementarity to the DMPK allele comprising the disease-associated-repeat expansion.
22. The complex of any one of embodiments 1 to 14, wherein the molecular payload is a polypeptide.
23. The complex of embodiment 22, wherein the polypeptide is a muscleblind-like (MBNL) polypeptide.
24. The complex of any one of embodiments 15 to 21, wherein the oligonucleotide comprises an antisense strand that hybridizes, in a cell, with a wild-type DMPK mRNA transcript encoded by the allele, wherein the DMPK mRNA transcript comprises repeating units of a CUG trinucleotide sequence.
25. The complex of any one of embodiments 15 to 21, wherein the oligonucleotide comprises an antisense strand that hybridizes, in a cell, with a mutant DMPK mRNA transcript encoded by the allele, wherein the DMPK mRNA transcript comprises repeating units of a CUG trinucleotide sequence.
26. The complex of any one of embodiments 1 to 25, wherein the disease-associated-repeat is 38 to 200 repeating units in length.

27. The complex of embodiment 26, wherein the disease-associated-repeat is associated with late onset myotonic dystrophy.
28. The complex of any one of embodiments 1 to 25, wherein the disease-associated-repeat is 100 to 10,000 repeat units in length.
29. The complex of embodiment 28, wherein the disease-associated-repeat is associated with congenital myotonic dystrophy.
30. The complex of any one of embodiments 15 to 21 and 24 to 29, wherein the oligonucleotide comprises at least one modified internucleotide linkage.
31. The complex of embodiment 30, wherein the at least one modified internucleotide linkage is a phosphorothioate linkage.
32. The complex of embodiment 31, wherein the oligonucleotide comprises phosphorothioate linkages in the Rp stereochemical conformation and/or in the Sp stereochemical conformation.
33. The complex of embodiment 32, wherein the oligonucleotide comprises phosphorothioate linkages that are all in the Rp stereochemical conformation or that are all in the Sp stereochemical conformation.
34. The complex of any one of embodiments 15 to 21 and 24 to 33, wherein the oligonucleotide comprises one or more modified nucleotides.
35. The complex of embodiment 34, wherein the one or more modified nucleotides are 2'-modified nucleotides.
36. The complex of any one of embodiments 15 to 21 and 24 to 35, wherein the oligonucleotide is a gapmer oligonucleotide that directs RNase H-mediated cleavage of a DMPK mRNA transcript in a cell.
37. The complex of embodiment 36, wherein the gapmer oligonucleotide comprises a central portion of 5 to 15 deoxyribonucleotides flanked by wings of 2 to 8 modified nucleotides.
38. The complex of embodiment 37, wherein the modified nucleotides of the wings are 2'-modified nucleotides.
39. The complex of any one of embodiments 15 to 21 and 24 to 35, wherein the oligonucleotide is a mixmer oligonucleotide.
40. The complex of embodiment 39, wherein the mixmer oligonucleotide inhibits binding of muscleblind-like protein 1, muscleblind-like protein 2, or muscleblind-like protein 3 to the DMPK mRNA transcript.
41. The complex of embodiment 39 or 40, wherein the mixmer oligonucleotide comprises two or more different 2' modified nucleotides.

42. The complex of any one of embodiments 15 or 21 and 24 to 35, wherein the oligonucleotide is an RNAi oligonucleotide that promotes RNAi-mediated cleavage of the DMPK mRNA transcript.
43. The complex of embodiment 42, wherein the RNAi oligonucleotide is a double-stranded oligonucleotide of 19 to 25 nucleotides in length.
44. The complex of embodiment 42 or 43, wherein the RNAi oligonucleotide comprises at least one 2' modified nucleotide.
45. The complex of any one of embodiments 35, 38, 41, or 44, wherein each 2' modified nucleotide is selected from the group consisting of: 2'-O-methyl, 2'-fluoro (2'-F), 2'-O-methoxyethyl (2'-MOE), and 2', 4'-bridged nucleotides.
46. The complex of embodiment 34, wherein the one or more modified nucleotides are bridged nucleotides.
47. The complex of any one of embodiment 35, 38, 41, or 44, wherein at least one 2' modified nucleotide is a 2',4'-bridged nucleotide selected from: 2',4'-constrained 2'-O-ethyl (cEt) and locked nucleic acid (LNA) nucleotides.
48. The complex of any one of embodiments 15 to 21 and 24 to 35, wherein the oligonucleotide comprises a guide sequence for a genome editing nuclease.
49. The complex of any one of embodiments 15 to 21 and 24 to 35, wherein the oligonucleotide is phosphorodiamidite morpholino oligomer.
50. The complex of any one of embodiments 1 to 49, wherein the muscle-targeting agent is covalently linked to the molecular payload via a cleavable linker.
51. The complex of embodiment 50, wherein the cleavable linker is selected from: a protease-sensitive linker, pH-sensitive linker, and glutathione-sensitive linker.
52. The complex of embodiment 51, wherein the cleavable linker is a protease-sensitive linker.
53. The complex of embodiment 52, wherein the protease-sensitive linker comprises a sequence cleavable by a lysosomal protease and/or an endosomal protease.
54. The complex of embodiment 52, wherein the protease-sensitive linker comprises a valine-citrulline dipeptide sequence.
55. The complex of embodiment 51, wherein the linker is pH-sensitive linker that is cleaved at a pH in a range of 4 to 6.
56. The complex of any one of embodiments 1 to 49, wherein the muscle-targeting agent is covalently linked to the molecular payload via a non-cleavable linker.
57. The complex of embodiment 56, wherein the non-cleavable linker is an alkane linker.

58. The complex of any of embodiments 2 to 57, wherein the antibody comprises a non-natural amino acid to which the oligonucleotide is covalently linked.

59. The complex of any of embodiments 2 to 57, wherein the antibody is covalently linked to the oligonucleotide via conjugation to a lysine residue or a cysteine residue of the antibody.

60. The complex of embodiment 59, wherein the antibody is conjugated to the cysteine via a maleimide-containing linker, optionally wherein the maleimide-containing linker comprises a maleimidocaproyl or maleimidomethyl cyclohexane-1-carboxylate group.

61. The complex of any one of embodiments 2 to 60, wherein the antibody is a glycosylated antibody that comprises at least one sugar moiety to which the oligonucleotide is covalently linked.

62. The complex of embodiment 61, wherein the sugar moiety is a branched mannose.

63. The complex of embodiment 61 or 62, wherein the antibody is a glycosylated antibody that comprises one to four sugar moieties each of which is covalently linked to a separate oligonucleotide.

64. The complex of embodiment 61, wherein the antibody is a fully-glycosylated antibody.

65. The complex of embodiment 61, wherein the antibody is a partially-glycosylated antibody.

66. The complex of embodiment 65, wherein the partially-glycosylated antibody is produced via chemical or enzymatic means.

67. The complex of embodiment 65, wherein the partially-glycosylated antibody is produced in a cell, cell that is deficient for an enzyme in the N- or O- glycosylation pathway.

68. A method of delivering a molecular payload to a cell expressing transferrin receptor, the method comprising contacting the cell with the complex of any one of embodiments 1 to 67.

69. A method of inhibiting activity of DMPK in a cell, the method comprising contacting the cell with the complex of any one of embodiments 1 to 67 in an amount effective for promoting internalization of the molecular payload to the cell.

70. The method of embodiment 69, wherein the cell is *in vitro*.

71. The method of embodiment 69, wherein the cell is in a subject.

72. The method of embodiment 71, wherein the subject is a human.

73. The method of any one of embodiments 69 to 72, wherein the complex inhibits the expression of DMPK.

74. The method of any one of embodiments 69 to 73, wherein the cell is contacted with a single dose of the complex.

75. The method of embodiment 74, wherein a single dose of the complex inhibits the expression of DMPK for at least two, four, eight, or twelve weeks.

76. The method of embodiment 75, wherein the complex inhibits the expression of DMPK by at least 30%, 40%, 50%, or 60% relative to a control.

77. The method of embodiment 75 or 76, the complex inhibits the expression of DMPK in muscle tissues by 40-60% for at least 12 weeks following administration of the single dose, relative to a control

78. A method of treating a subject having an expansion of a disease-associated-repeat of a DMPK allele that is associated with myotonic dystrophy, the method comprising administering to the subject an effective amount of the complex of any one of embodiments 1 to 67.

79. The method of embodiment 78, wherein the disease-associated-repeat comprises repeating units of a trinucleotide sequence.

80. The method of embodiment 78, wherein the trinucleotide sequence is a CTG trinucleotide sequence.

81. The method of any one of embodiments 78 to 80, wherein the disease-associated-repeat is 38 to 200 repeating units in length.

82. The method of 81, wherein the disease-associated-repeat is associated with late onset myotonic dystrophy.

83. The method of any one of embodiments 78 to 80, wherein the disease-associated-repeat is 100 to 10,000 repeating units in length.

84. The method of 83, wherein the disease-associated-repeat is associated with congenital myotonic dystrophy.

85. The method of any one of embodiments 78 to 84, wherein administration of the complex results in inhibition of the expression of DMPK in muscle tissues.

86. The method of any one of embodiments 78 to 85, wherein the complex is intravenously administered to the subject.

87. The method of any one of embodiments 78 to 86, wherein an effective amount of the complex comprises 1-15 mg/kg of RNA.

88. The method of any one of embodiments 78 to 87, wherein the complex is administered to the subject in a single dose.

89. The method of embodiment 88, wherein administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues for at least two, four, eight, or twelve weeks.

90. The method of embodiment 89, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues by at least 30%, 40%, 50%, or 60% relative to a control.

91. The method of embodiment 89 or 90, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues for at least 12 weeks following administration of the single dose.

92. The method of embodiment 91, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues by 40-60%, relative to a control, for at least 12 weeks following administration of the single dose.

93. The method of embodiment 89 or 90, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues for a duration of time in the range of 4-8, 5-10, 8-12, 10-14, or 8-16 weeks following administration of the single dose.

94. The method of embodiment 93, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues by 40-60%, relative to a control, for a duration of time in the range of 4-8, 5-10, 8-12, 10-14, or 8-16 weeks following administration of the single dose.

95. The method of embodiment 89 or 90, wherein the administration of a single dose of the complex results in inhibition of the expression of DMPK in muscle tissues by 40-60%, relative to a control, at 12 weeks following administration of the single dose.

96. The method of any one of embodiments 78 to 87, wherein the complex is administered to the subject in a single dose once every 4-8, 5-10, 8-12, or 8-16 weeks.

97. The method of embodiment 96, wherein the complex is administered to the subject in a single dose once every 12 weeks.

98. The method of any one of embodiments 88 to 97, wherein the single dose comprises the complex at a concentration of 1-15 mg/kg of RNA.

99. The method of embodiment 98, wherein the single dose comprises the complex at a concentration of 10 mg/kg of RNA.

100. A method of treating a subject having an expansion of a disease-associated-repeat of a DMPK allele that is associated with myotonic dystrophy, the method comprising administering the complex of any one of embodiments 1 to 67 to the subject,

wherein the administration results in inhibition of DMPK expression in muscle tissues by 40-60%, relative to a control, for a duration of time in the range of 4-8, 5-10, 8-12, 10-14, or 8-16 weeks following administration of the complex.

101. A method of inhibiting DMPK expression in a subject, the method comprising administering the complex of any one of embodiments 1 to 67,

wherein the administration results in inhibition of DMPK expression in muscle tissues by 40-60%, relative to a control, for a duration of time in the range of 4-8, 5-10, 8-12, 10-14, or 8-16 weeks following administration of the complex.

102. The method of embodiment 100 or 101, wherein the molecular payload is an oligonucleotide.

103. The method of embodiment 102, wherein the concentration of the complex is 1-15 mg/kg of RNA.

104. A complex comprising an anti-transferrin receptor (TfR) antibody covalently linked to a molecular payload configured for reducing expression or activity of DMPK, wherein the anti-TfR antibody comprises:

(i) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 75;

(ii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 69; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(iii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 71; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(iv) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 72; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 70;

(v) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 74;

(vi) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 75;

(vii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 74;

(viii) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 78;

(ix) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 79; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 80; or

(x) a heavy chain variable region (VH) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 85% identical to SEQ ID NO: 80.

105. A complex comprising an anti-transferrin receptor (TfR) antibody covalently linked to a molecular payload configured for reducing expression or activity of DMPK, wherein the anti-TfR antibody has undergone pyroglutamate formation resulting from a post-translational modification.

EQUIVALENTS AND TERMINOLOGY

[000587] The disclosure illustratively described herein suitably can be practiced in the absence of any element or elements, limitation or limitations that are not specifically disclosed herein. Thus, for example, in each instance herein any of the terms “comprising”, “consisting essentially of”, and “consisting of” may be replaced with either of the other two terms. The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention that in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the disclosure. Thus, it should be understood that although the present disclosure has been specifically disclosed by preferred embodiments, optional features, modification and variation of the concepts herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this disclosure.

[000588] In addition, where features or aspects of the disclosure are described in terms of Markush groups or other grouping of alternatives, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group or other group.

[000589] It should be appreciated that, in some embodiments, sequences presented in the sequence listing may be referred to in describing the structure of an oligonucleotide or other nucleic acid. In such embodiments, the actual oligonucleotide or other nucleic acid may have one or more alternative nucleotides (e.g., an RNA counterpart of a DNA nucleotide or a DNA counterpart of an RNA nucleotide) and/or (e.g., and) one or more modified nucleotides and/or (e.g., and) one or more modified internucleotide linkages and/or (e.g., and) one or more other modification compared with the specified sequence while retaining essentially same or similar complementary properties as the specified sequence.

[000590] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (*i.e.*, meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (*e.g.*, “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[000591] Embodiments of this invention are described herein. Variations of those embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description.

[000592] The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS

What is claimed is:

1. A complex comprising an anti-transferrin receptor (TfR) antibody covalently linked to a molecular payload configured for reducing expression or activity of DMPK, wherein the anti-TfR antibody comprises:
 - (i) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 75;
 - (ii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 69; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;
 - (iii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 71; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;
 - (iv) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 72; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 70;
 - (v) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 74;
 - (vi) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 73; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 75;
 - (vii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 76; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 74;
 - (viii) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 78;
 - (ix) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 79; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 80; or

(x) a heavy chain variable region (VH) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 77; and/or a light chain variable region (VL) comprising an amino acid sequence at least 95% identical to SEQ ID NO: 80.

2. The complex of claim 1, wherein the antibody comprises:

(i) a VH comprising the amino acid sequence of SEQ ID NO: 76 and a VL comprising the amino acid sequence of SEQ ID NO: 75;

(ii) a VH comprising an amino acid sequence of SEQ ID NO: 69 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(iii) a VH comprising an amino acid sequence of SEQ ID NO: 71 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(iv) a VH comprising an amino acid sequence of SEQ ID NO: 72 and a VL comprising an amino acid sequence of SEQ ID NO: 70;

(v) a VH comprising an amino acid sequence of SEQ ID NO: 73 and a VL comprising an amino acid sequence of SEQ ID NO: 74;

(vi) a VH comprising an amino acid sequence of SEQ ID NO: 73 and a VL comprising an amino acid sequence of SEQ ID NO: 75;

(vii) a VH comprising an amino acid sequence of SEQ ID NO: 76 and a VL comprising an amino acid sequence of SEQ ID NO: 74;

(viii) a VH comprising an amino acid sequence of SEQ ID NO: 77 and a VL comprising an amino acid sequence of SEQ ID NO: 78;

(ix) a VH comprising an amino acid sequence of SEQ ID NO: 79 and a VL comprising an amino acid sequence of SEQ ID NO: 80; or

(x) a VH comprising an amino acid sequence of SEQ ID NO: 77 and a VL comprising an amino acid sequence of SEQ ID NO: 80.

3. The complex of claim 1 or claim 2, wherein the antibody is selected from the group consisting of a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, an scFv, an Fv, and a full-length IgG.

4. The complex of claim 3, wherein the antibody is a Fab fragment.

5. The complex of claim 4, wherein the antibody comprises:

(i) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(ii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 97; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 98; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(iv) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 99; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 85;

(v) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(vi) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 100; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 90;

(vii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 101; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 89;

(viii) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 93;

(ix) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 103; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95; or

(x) a heavy chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 102; and/or a light chain comprising an amino acid sequence at least 85% identical to SEQ ID NO: 95.

6. The complex of claim 4 or claim 5, wherein the antibody comprises:

(i) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(ii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 97 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 98 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(iv) a heavy chain comprising the amino acid sequence of SEQ ID NO: 99 and a light chain comprising the amino acid sequence of SEQ ID NO: 85;

(v) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(vi) a heavy chain comprising the amino acid sequence of SEQ ID NO: 100 and a light chain comprising the amino acid sequence of SEQ ID NO: 90;

(vii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 101 and a light chain comprising the amino acid sequence of SEQ ID NO: 89;

(viii) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 93;

(ix) a heavy chain comprising the amino acid sequence of SEQ ID NO: 103 and a light chain comprising the amino acid sequence of SEQ ID NO: 95; or

(x) a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 95.

7. The complex of any one of claims 1 to 6, wherein the antibody does not specifically bind to the transferrin binding site of the transferrin receptor and/or wherein the antibody does not inhibit binding of transferrin to the transferrin receptor.

8. The complex of any one of claims 1 to 7, wherein the antibody is cross-reactive with extracellular epitopes of two or more of a human, non-human primate and rodent transferrin receptor.

9. The complex of any one of claims 1 to 8, wherein the complex is configured to promote transferrin receptor mediated internalization of the molecular payload into a muscle cell.

10. The complex of any one of claims 1 to 9, wherein the molecular payload is an oligonucleotide.

11. The complex of claim 10, wherein the oligonucleotide comprises at least 15 consecutive nucleotides of SEQ ID NOs: 148-383 and 621-638, wherein any one or more of the thymidine bases (T's) in the oligonucleotide may optionally be a uridine base (U) and/or any one or more of the U's may optionally be a T.

12. The complex of claim 11, wherein the oligonucleotide comprises a sequence comprising any one of SEQ ID NOs: 159, 162, 172, 174, 180, 182, 188, 190, 195, 196, 201, 203, 212, 215, 218, 222, 248, and 264, wherein any one or more of the U's in the oligonucleotide may optionally be a T.

13. The complex of claim 10, wherein the oligonucleotide comprises a region of complementarity to at least 15 consecutive nucleotides of any one of SEQ ID NO: 384-619.
14. The complex of any one of claims 10 to 13, wherein the oligonucleotide mediates RNase H-mediated cleavage of a DMPK mRNA transcript.
15. The complex of any one of claims 10 to 14, wherein the oligonucleotide comprises a 5'-X-Y-Z-3' formula, wherein X and Z are flanking regions comprising one or more 2'-modified nucleosides selected from the group consisting of: 2'-O-methyl, 2'-fluoro, 2'-O-methoxyethyl, and 2',4'- bridged nucleosides, and wherein Y is a gap region and each nucleoside in Y is a 2'-deoxyribonucleoside.
16. The complex of any one of claims 10 to 15, wherein the oligonucleotide comprises one or more phosphorothioate internucleoside linkages.
17. The complex of any one of claims 1 to 16, wherein the antibody is covalently linked to the molecular payload via a cleavable linker.
18. The complex of claim 17, wherein the cleavable linker comprises a valine-citrulline sequence.
19. The complex of any of claims 1 to 18, wherein the antibody is covalently linked to the molecular payload via conjugation to a lysine residue or a cysteine residue of the antibody.
20. The complex of any one of claims 1 to 19, wherein reducing expression comprises reducing RNA levels of DMPK, optionally wherein the reduced RNA levels are in the nucleus of a cell, optionally wherein the cell is a muscle cell.
21. The complex of any one of claims 1 to 20, wherein the DMPK is encoded from an allele comprising a disease-associated repeat.
22. A method of reducing DMPK expression in a cell, the method comprising contacting the cell with the complex of any one of claims 1 to 21 in an effective amount for promoting internalization of the molecular payload in the cell, optionally wherein the cell is a muscle cell.
23. A method of treating a subject having an expansion of a disease-associated-repeat of a DMPK allele that is associated with myotonic dystrophy, the method comprising administering to the subject an effective amount of the complex of any one of claims 1 to 21.
24. The method of claim 23, wherein the disease-associated-repeat comprises repeating units of a CTG trinucleotide sequence.

25. The method of claim 23 or claim 24, wherein the complex is intravenously administered to the subject.