CLAIMS

1. Use of a combination of (a) valacyclovir or a salt or solvate thereof and (b) famciclovir or a salt or solvate thereof in the preparation of a medicament for the treatment of herpes simplex virus (HSV) infection, prophylaxis, or pre-exposure prophylaxis against HSV infection.

2. Use according to Claim 1, in which the medicament comprises:

(a) 0.5 mg/kg body weight to 1000 mg/kg body weight of (a);

(b) 5 mg/kg body weight to 50 mg/kg body weight of (a);

(c) 25 mg/kg body weight to 100 mg/kg body weight of (a);

(d) 50 mg/kg body weight to 250 mg/kg body weight of (a); or

(e) 100 mg/kg body weight to 500 mg/kg body weight of (a).

3. Use according to Claim 1 or 2, in which the medicament comprises:

(a) 0.5 mg/kg body weight to 1000 mg/kg body weight of (b);

(b) 5 mg/kg body weight to 50 mg/kg body weight of (b);

(c) 25 mg/kg body weight to 100 mg/kg body weight of (b);

(d) 50 mg/kg body weight to 250 mg/kg body weight of (b); or

(e) 100 mg/kg body weight to 500 mg/kg body weight of (b).

4. Use according to Claim 1, 2 or 3, in which the medicament comprises (a) and (b) in a dose ratio in mg/kg of 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, or 1:5.

5. Use according to any preceding claim, in which the medicament comprises:

(a) 5 mg/kg body weight to 15 mg/kg body weight of (a) and 2.5 mg/kg body weight to 7.5 mg/kg body weight of (b);

(b) 10 mg/kg body weight to 50 mg/kg body weight of (a) and 5 mg/kg body weight to 25 mg/kg body weight of (b);

(c) 50 mg/kg body weight to 150 mg/kg body weight of (a) and 20 mg/kg body weight to 50 mg/kg body weight of (b);

(d) 100 mg/kg body weight to 150 mg/kg body weight of (a) and 50 mg/kg body weight to 100 mg/kg body weight of (b);

(e) 150 mg/kg body weight to 500 mg/kg body weight of (a) and 100 mg/kg body weight to 250 mg/kg body weight of (b); or

(f) 250 mg/kg body weight to 1000 mg/kg body weight of (a) and 150 mg/kg body weight to 500 mg/kg body weight of (b).

6. Use according to any preceding claim, in which the medicament comprises from greater than 100 mg/kg to 700 mg/kg body weight of valacyclovir and from greater than 100 mg/kg to 500 mg/kg body weight of famciclovir.

7. Use according to any preceding claim, in which the medicament comprises:

(a) 1 mg to 8000 mg of valacyclovir or a salt or solvate thereof and 1 mg to 8000 mg famciclovir or a salt or solvate thereof; or

(b) 125 mg to 2000 mg of valacyclovir or a salt or solvate thereof and 60 mg to 1000 mg famciclovir or a salt or solvate thereof.

8. Use according to any preceding claim, in which the medicament is formulated for oral administration, or comprises an oral tablet, oral capsule, or oral solution, or is formulated for administration once daily or twice daily, or is formulated in sustained-release form, or is formulated in parenteral administration form.

9. Use according to any preceding claim, in which the medicament additionally comprises a therapeutically effective amount of an additional antiviral agent or an HIV antiviral agent, or is formulated for a female and additionally comprises a therapeutically effective amount of a contraceptive.

10. Use according to Claim 9, in which the additional antiviral agent comprises a helicase-primase inhibitor, amenamevir, N-methancocarbathymidine (N-MCT), tenofovir, emtricitabine, lamivudine, interfereon, ribavirin, boceprevir, telaprevir, simeprevir, sofosbuvir, ledipasvir, tenofovir, ombitasvir, paritaprevir, penciclovir, pritelivir, brincidofovir, cidofovir, ganciclovir, valganciclovir, valomaciclovir, or ritonavir.

11. Use according to any preceding claim, in which the HSV is HSV-1.

12. Use according to any of Claims 1 to 11, in which the HSV is HSV-2.