KCNT1 INHIBITORS AND METHODS OF USE

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to and the benefit of U.S. Provisional Patent Application Number 62/842,849 filed May 3, 2019 and U.S. Provisional Patent Application Number 62/982,864 filed February 28, 2020, the contents of each of which are incorporated herein by reference in their entirety.


BACKGROUND

[0002] KCNT1 encodes sodium-activated potassium channels known as Slack (Sequence like a calcium-activated K^+ channel). These channels are found in neurons throughout the brain and can mediate a sodium-activated potassium current I_{KNa}. This delayed outward current can regulate neuronal excitability and the rate of adaption in response to maintained stimulation. Abnormal Slack activity have been associated with development of early onset epilepsies and intellectual impairment. Accordingly, pharmaceutical compounds that selectively regulate sodium-activated potassium channels, e.g., abnormal KCNT1, abnormal I_{KNa}, are useful in treating a neurological disease or disorder or a disease or condition related to excessive neuronal excitability and/or KCNT1 gain-of-function mutations.

SUMMARY

[0003] Described herein are compounds and compositions useful for preventing and/or treating a disease, disorder, or condition, e.g., a neurological disease or disorder, a disease, disorder, or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene, for example, KCNT1.


[0004] In one example, the present disclosure features a pharmaceutical composition comprising a compounds of Formula (I):



(I),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein, and a pharmaceutically acceptable excipient.

[0004a] In one aspect, the present disclosure refers to a pharmaceutical composition comprising a compound of Formula **I**:



(I),

or a pharmaceutically acceptable salt thereof, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be replaced with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, $-C(O)OR_8$, $C(O)N(R_9)_2$, C_{1-6}alkylene-CN, -CN, $-S(O)_2-C_{1-6}$alkyl, C_{1-6}alkylene-$S(O)_2-C_{1-6}$alkyl, $-S(O)_2-N(R_9)_2$, $-OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, $-N(R_9)_2$, $-NR_9-SO_2-C_{1-6}$alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, $-C(O)OR_8$, $-C(O)N(R_9)_2$,-$NR_{10}C(O)-R_{11}$, -CN, $-S(O)_2-C_{1-6}$alkyl, $-S(O)_2-N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and $-N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;


each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N;

and a pharmaceutically acceptable excipient.


[0005] In another example, the present disclosure features a compound of Formula (II):



(II),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein.

[0005a] In one aspect, the present disclosure refers to a compound of Formula **II**:



(**II**),

or a pharmaceutically acceptable salt thereof, wherein

one or two selected from X, Y, Z, Y', and Z' are N, and the others are CH, wherein the hydrogen of CH may be replaced with R_5;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy optionally substituted with C_{3-7}cycloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, -$N(R_9)_2$, -$C(O)OR_8$, -$C(O)N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$-$N(R_9)_2$, -OC(O)C_{1-6}alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -$N(R_9)_2$, -NR_9-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -$C(O)OR_8$, -$C(O)N(R_9)_2$,-$NR_{10}C(O)$-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is selected from the group consisting of hydrogen, C_{1-6}alkyl, and C_{3-10}cycloalkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to

form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl; and

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl.


[0006] In another example, the present disclosure features a compound of Formula (III):



(III),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein.

[0006a] In one aspect, the present disclosure refers to a compound of Formula **III**:



(**III**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is C_{1-6}alkyl or C_{3-7}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-7}cycloalkyl is optionally substituted with one or more halogen or C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -$C(O)OR_8$, -$C(O)N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, -$OC(O)C_{1-6}$alkyl, and -O-C_{1-3}cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -$C(O)OR_8$, -$C(O)N(R_9)_2$,-$NR_{10}C(O)$-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;


each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and


n is selected from the group consisting of 0, 1, 2, and 3;

wherein the compound is not:


,



, or a pharmaceutically acceptable salt thereof.

[0007] In another example, the present disclosure features a compound of Formula (IV):


(IV),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein.

[0007a] In one aspect, the present disclosure refers to a compound of Formula **IV**:


(**IV**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is selected from the group consisting of $C_{1\text{-}6}$alkyl, and $C_{3\text{-}7}$cycloalkyl, wherein the $C_{1\text{-}6}$alkyl or $C_{3\text{-}7}$cycloalkyl is optionally substituted with one or more substituents independently selected from halogen and $C_{1\text{-}6}$alkoxy;

R_{12} is $C_{1\text{-}6}$alkyl optionally substituted with one or more halogen or $C_{1\text{-}6}$alkoxy;

R_2 is hydrogen or $C_{1\text{-}4}$alkyl;

R_3 is selected from the group consisting of $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$ cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from $C_{1\text{-}6}$alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a $C_{3\text{-}7}$cycloalkylene or 3-7 membered heterocyclene; wherein the $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$ cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, $C_{3\text{-}7}$cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with R_7;

R_5 is selected from the group consisting of halogen, $C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$alkoxy, $C_{1\text{-}6}$haloalkyl, $C_{1\text{-}6}$haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -$C_{1\text{-}6}$alkylene-OH, OH, -$C(O)OR_8$, -$C(O)N(R_9)_2$, -$C_{1\text{-}6}$alkylene-CN, -CN, -$S(O)_2$-$C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$alkylene-$S(O)_2$-$C_{1\text{-}6}$alkyl, -$S(O)_2$- $N(R_9)_2$, -$OC(O)C_{1\text{-}6}$alkyl, and -O-$C_{3\text{-}10}$cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, $C_{1\text{-}6}$alkoxy, -OH, -O-($C_{1\text{-}6}$alkylene)-phenyl, $C_{3\text{-}10}$cycloalkyl, -$C(O)OR_8$, -$C(O)N(R_9)_2$,-$NR_{10}C(O)$-R_{11}, -CN, -$S(O)_2$-$C_{1\text{-}6}$alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of $C_{1\text{-}6}$alkyl, halogen, -OH, $C_{1\text{-}6}$alkoxy, and -$N(R_9)_2$;

R_8 is hydrogen or $C_{1\text{-}6}$alkyl;


each R_9 is independently selected from the group consisting of hydrogen, $C_{1\text{-}6}$alkyl, and -($C_{1\text{-}6}$alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents selected from halogen and -OH;

each R_{10} is independently hydrogen or $C_{1\text{-}6}$alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;


wherein the compound is not:



,




,



, or a pharmaceutically acceptable salt thereof.


[0008] In another example, the present disclosure features a compound of Formula (V):



(V),

Or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein.

[0008a] In one aspect, the present disclosure refers to a compound of Formula **V**:



(**V**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is phenyl or 3-10 membered heteroaryl, wherein the phenyl or 3-10 membered heteroaryl is optionally substituted with one or more substituents independently selected from halogen and C_{1-6}alkoxy,

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -$C(O)OR_8$, -$C(O)N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, -$OC(O)C_{1-6}$alkyl, and -O-C_{3-10}cycloalkyl;

R_7 is selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-7}cycloalkyl, -$C(O)OR_8$, -$C(O)N(R_9)_2$,-$NR_{10}C(O)$-R_{11}, -CN, -$S(O)_z$-C_{1-6}alkyl, -$S(O)_z$-$N(R_9)_2$, 3-7 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with

one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;


wherein the compound is not:




,



,




,



, or a pharmaceutically acceptable salt thereof.


[0009] In another example, the present disclosure features a compound of Formula (VI):



(VI),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein.

[0009a] In one aspect, the present disclosure refers to a compound of Formula **VI**:



(**VI**),

or a pharmaceutically acceptable salt thereof, wherein

R_{13} is $C_{1\text{-}6}$alkyl or $C_{3\text{-}10}$cycloalkyl, wherein the $C_{1\text{-}6}$alkyl or $C_{3\text{-}10}$cycloalkyl is optionally substituted with phenyl;

R_{14} is hydrogen;

R_{15} is $C_{1\text{-}6}$alkyl or hydrogen;

R_{16} is $C_{1\text{-}6}$alkyl optionally substituted with one or more halogen, $C_{1\text{-}6}$alkoxy, $C_{3\text{-}10}$ cycloalkyl, or phenyl;


each R_{17} is independently selected from the group consisting of halogen, $C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$haloalkyl, $C_{1\text{-}6}$alkylene-$N(R_{20})_2$, $C_{1\text{-}6}$alkylene-O-$C_{3\text{-}10}$cycloalkyl, $C_{1\text{-}6}$alkoxy optionally substituted with $C_{3\text{-}7}$cycloalkyl, $C_{1\text{-}6}$haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or $C_{1\text{-}6}$alkoxy, 3-10 membered heteroaryl, -$C_{1\text{-}6}$alkylene-OH, $C_{1\text{-}6}$alkylene-$C_{1\text{-}6}$alkoxy, OH, -$N(R_{20})_2$, -C(O)OR_{19}, -C(O)$N(R_{20})_2$, -$C_{1\text{-}6}$alkylene-CN, -CN, -$S(O)_2$-$C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$alkylene-$S(O)_2$-$C_{1\text{-}6}$alkyl, -$S(O)_2$- $N(R_{20})_2$, -OC(O)$C_{1\text{-}6}$alkyl, -O-$C_{3\text{-}10}$cycloalkyl optionally substituted with one or more halogen or $C_{1\text{-}6}$alkyl, and $C_{3\text{-}10}$cycloalkyl optionally substituted with one or more substituents selected from halogen, $C_{1\text{-}6}$alkyl, and $C_{1\text{-}6}$alkoxy;

p is selected from the group consisting of 1, 2, and 3;

R_{19} is selected from the group consisting of hydrogen, $C_{1\text{-}6}$alkyl, and $C_{3\text{-}10}$cycloalkyl; and

each R_{20} is independently hydrogen or $C_{1\text{-}6}$alkyl.


[00010] In another example, the present disclosure features a pharmaceutical composition comprising a compound of Formula (VII):




(VII),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined herein, and a pharmaceutically acceptable excipient.


[00010a] In one aspect, the present disclosure refers to a pharmaceutical composition comprising a compound of Formula **VII**:



(**VII**), wherein the compound is a compound of Formula **VII-a** or Formula **VII-b**:



(**VII-a**),



(**VII-b**),

or a pharmaceutically acceptable salt thereof, wherein

R_{23} is a tert-butyl;

R_{24} is hydrogen;

R_{25} is hydrogen; and

R_{27} is independently selected from the group consisting of halogen and $C_{1\text{-}6}$alkoxy;


and a pharmaceutically acceptable excipient.

[00011] In another example, the present disclosure provides a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of formula (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a))) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[00011a] In one aspect, the present disclosure refers to a pharmaceutical composition comprising a compound disclosed herein or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

[00012] In another example, the present disclosure provides a method of treating a neurological disease or disorder, wherein the method comprises administering to a subject in need thereof a compound disclosed herein (e.g., a compound of formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient).

[00012a] In one aspect, the present disclosure refers to use of a compound of Formula **I**:



(**I**),

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a neurological disease or disorder, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be replaced with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or

more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, $-C(O)OR_8$, $C(O)N(R_9)_2$, C_{1-6}alkylene-CN, -CN, $-S(O)_2-C_{1-6}$alkyl, C_{1-6}alkylene-$S(O)_2-C_{1-6}$alkyl, $-S(O)_2-N(R_9)_2$, $-OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, $-N(R_9)_2$, $-NR_9-SO_2-C_{1-6}$alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, $-C(O)OR_8$, $-C(O)N(R_9)_2$,-$NR_{10}C(O)-R_{11}$, -CN, $-S(O)_2-C_{1-6}$alkyl, $-S(O)_2-N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and $-N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N.

[00013] In another example, the present disclosure provides a method of treating a disease or condition associated with excessive neuronal excitability, wherein the method comprises administering to a subject in need thereof a compound disclosed herein (e.g., a compound of formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-

b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient).

[00014] In another example, the present disclosure provides a method of treating a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the method comprises administering to a subject in need thereof a compound disclosed herein (e.g., a compound of formula ((I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of formula ((I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient).

[00015] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is epilepsy, an epilepsy syndrome, or an encephalopathy.

[00015a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is epilepsy, an epilepsy syndrome, or an encephalopathy.

[00016] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a genetic or pediatric epilepsy or a genetic or pediatric epilepsy syndrome.

[00016a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a genetic or pediatric epilepsy or a genetic or pediatric epilepsy syndrome.

[00017] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a cardiac dysfunction.

[00017a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a cardiac dysfunction.

[00018] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome,

developmental and epileptic encephalopathy, Lennox Gastaut syndrome, seizures (e.g., Generalized tonic clonic seizures, Asymmetric Tonic Seizures), leukodystrophy, leukoencephalopathy, intellectual disability, Multifocal Epilepsy, Drug resistant epilepsy, Temporal lobe epilepsy, cerebellar ataxia).

[00018a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy, Lennox Gastaut syndrome, seizures (e.g., Generalized tonic clonic seizures, Asymmetric Tonic Seizures), leukodystrophy, leukoencephalopathy, intellectual disability, Multifocal Epilepsy, Drug resistant epilepsy, Temporal lobe epilepsy, or cerebellar ataxia).

[00019] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of cardiac arrhythmia, sudden unexpected death in epilepsy, Brugada syndrome, and myocardial infarction.

[00019a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of cardiac arrhythmia, sudden unexpected death in epilepsy (SUDEP), Brugada syndrome, and myocardial infarction.

[00020] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine, etc).

[00020a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine).

[00021] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a muscle disorder (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity).

[00021a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament

for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a muscle disorder (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity).

[00022] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from itch and pruritis, ataxia and cerebellar ataxias.

[00022a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from itch and pruritis, ataxia and cerebellar ataxias.

[00023] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia).

[00023a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition of any one of disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia).

[00024] In some examples, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from the group consisting of learning disorders, Fragile X, neuronal plasticity, and autism spectrum disorders.

[00024a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from the group consisting of learning disorders, Fragile X, neuronal plasticity, and autism spectrum disorders.

[00025] In some examples, the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of epileptic encephalopathy with SCN1A, SCN2A, SCN8A mutations, early infantile epileptic encephalopathy, Dravet syndrome, Dravet syndrome with SCN1A

mutation, generalized epilepsy with febrile seizures, intractable childhood epilepsy with generalized tonic-clonic seizures, infantile spasms, benign familial neonatal-infantile seizures, SCN2A epileptic encephalopathy, focal epilepsy with SCN3A mutation, cryptogenic pediatric partial epilepsy with SCN3A mutation, SCN8A epileptic encephalopathy, sudden unexpected death in epilepsy, Rasmussen encephalitis, malignant migrating partial seizures of infancy, autosomal dominant nocturnal frontal lobe epilepsy, sudden expected death in epilepsy (SUDEP), KCNQ2 epileptic encephalopathy, and KCNT1 epileptic encephalopathy.

[00025a] In one aspect, the present disclosure refers to use of a compound disclosed herein or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of epileptic encephalopathy with SCN1A, SCN2A, SCN8A mutations, early infantile epileptic encephalopathy, Dravet syndrome, Dravet syndrome with SCN1A mutation, generalized epilepsy with febrile seizures, intractable childhood epilepsy with generalized tonic-clonic seizures, infantile spasms, benign familial neonatal-infantile seizures, SCN2A epileptic encephalopathy, focal epilepsy with SCN3A mutation, cryptogenic pediatric partial epilepsy with SCN3A mutation, SCN8A epileptic encephalopathy, sudden unexpected death in epilepsy (SUDEP), Rasmussen encephalitis, malignant migrating partial seizures of infancy, autosomal dominant nocturnal frontal lobe epilepsy, KCNQ2 epileptic encephalopathy, and KCNT1 epileptic encephalopathy.

[00026] Other objects and advantages will become apparent to those skilled in the art from a consideration of the ensuing **Detailed Description**, **Examples**, and **Claims**.

DETAILED DESCRIPTION OF THE INVENTION

[00027] As generally described herein, the present invention provides compounds and compositions useful for preventing and/or treating a disease, disorder, or condition described herein, e.g., a disease, disorder, or condition associated with excessive neuronal excitability, and/or a disease, disorder, or condition associated with gain-of-function mutations in KCNT1. Exemplary diseases, disorders, or conditions include epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy, and Lennox Gastaut syndrome, seizures, leukodystrophy, leukoencephalopathy, intellectual disability, Multifocal Epilepsy, Generalized tonic clonic seizures, Drug resistant epilepsy, Temporal lobe epilepsy, cerebellar ataxia, Asymmetric Tonic Seizures) and cardiac dysfunctions (e.g., cardiac arrhythmia, Brugada syndrome, sudden unexpected death in epilepsy, myocardial infarction), pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine, etc), muscle disorders (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity), itch and pruritis, ataxia and cerebellar ataxias, psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia), learning disorders, Fragile X, neuronal plasticity, and autism spectrum disorders.

Definitions

Chemical Definitions

[00028] Definitions of specific functional groups and chemical terms are described in more detail below. The chemical elements are identified in accordance with the Periodic

Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in Thomas Sorrell, *Organic Chemistry*, University Science Books, Sausalito, 1999; Smith and March, *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; and Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

[00029] Compounds described herein can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, e.g., enantiomers and/or diastereomers. For example, the compounds described herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. Isomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred isomers can be prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, *Stereochemistry of Carbon Compounds* (McGraw–Hill, NY, 1962); and Wilen, *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972). The invention additionally encompasses compounds described herein as individual isomers substantially free of other isomers, and alternatively, as mixtures of various isomers.

[00030] As used herein a pure enantiomeric compound is substantially free from other enantiomers or stereoisomers of the compound (*i.e.*, in enantiomeric excess). In other words, an "S" form of the compound is substantially free from the "R" form of the compound and is, thus, in enantiomeric excess of the "R" form. The term "enantiomerically pure" or "pure enantiomer" denotes that the compound comprises more than 75% by weight, more than 80% by weight, more than 85% by weight, more than 90% by weight, more than 91% by weight, more than 92% by weight, more than 93% by weight, more than 94% by weight, more than 95% by weight, more than 96% by weight, more than 97% by weight, more than 98% by weight, more than 98.5% by weight, more than 99% by weight, more than 99.2% by weight, more than 99.5% by weight, more than 99.6% by weight, more than 99.7% by weight, more than 99.8% by weight or more than 99.9% by weight, of the enantiomer. In certain

embodiments, the weights are based upon total weight of all enantiomers or stereoisomers of the compound.

[00031] In the compositions provided herein, an enantiomerically pure compound can be present with other active or inactive ingredients. For example, a pharmaceutical composition comprising enantiomerically pure R–compound can comprise, for example, about 90% excipient and about 10% enantiomerically pure R–compound. In certain embodiments, the enantiomerically pure R–compound in such compositions can, for example, comprise, at least about 95% by weight R–compound and at most about 5% by weight S–compound, by total weight of the compound. For example, a pharmaceutical composition comprising enantiomerically pure S–compound can comprise, for example, about 90% excipient and about 10% enantiomerically pure S–compound. In certain embodiments, the enantiomerically pure S–compound in such compositions can, for example, comprise, at least about 95% by weight S–compound and at most about 5% by weight R–compound, by total weight of the compound. In certain embodiments, the active ingredient can be formulated with little or no excipient or carrier.

[00032] Compound described herein may also comprise one or more isotopic substitutions. For example, H may be in any isotopic form, including ^{1}H, ^{2}H (D or deuterium), and ^{3}H (T or tritium); C may be in any isotopic form, including ^{12}C, ^{13}C, and ^{14}C; O may be in any isotopic form, including ^{16}O and ^{18}O; F may be in any isotopic form, including ^{18}F and ^{19}F; and the like.

[00033] The following terms are intended to have the meanings presented therewith below and are useful in understanding the description and intended scope of the present invention. When describing the invention, which may include compounds and pharmaceutically acceptable salts thereof, pharmaceutical compositions containing such compounds and methods of using such compounds and compositions, the following terms, if present, have the following meanings unless otherwise indicated. It should also be understood that when described herein any of the moieties defined forth below may be substituted with a variety of substituents, and that the respective definitions are intended to include such substituted moieties within their scope as set out below. Unless otherwise stated, the term "substituted" is to be defined as set out below. It should be further understood that the terms "groups" and "radicals" can be considered interchangeable when used herein. The articles "a" and "an" may be used herein to refer to one or to more than one (*i.e.* at least one) of the grammatical objects of the article. By way of example "an analogue" means one analogue or more than one analogue.

[00034] When a range of values is listed, it is intended to encompass each value and sub–range within the range. For example, "$C_{1–6}$ alkyl" is intended to encompass, C_1, C_2, C_3, C_4, C_5, C_6, $C_{1–6}$, $C_{1–5}$, $C_{1–4}$, $C_{1–3}$, $C_{1–2}$, $C_{2–6}$, $C_{2–5}$, $C_{2–4}$, $C_{2–3}$, $C_{3–6}$, $C_{3–5}$, $C_{3–4}$, $C_{4–6}$, $C_{4–5}$, and $C_{5–6}$ alkyl.

[00035] As used herein, "alkyl" refers to a radical of a straight–chain or branched saturated hydrocarbon group, e.g., having 1 to 20 carbon atoms ("$C_{1–20}$ alkyl"). In some embodiments, an alkyl group has 1 to 10 carbon atoms ("$C_{1–10}$ alkyl"). In some embodiments, an alkyl group has 1 to 9 carbon atoms ("$C_{1–9}$ alkyl"). In some embodiments, an alkyl group has 1 to 8 carbon atoms ("$C_{1–8}$ alkyl"). In some embodiments, an alkyl group has 1 to 7 carbon atoms ("$C_{1–7}$ alkyl"). In some embodiments, an alkyl group has 1 to 6 carbon atoms ("$C_{1–6}$ alkyl"). In some embodiments, an alkyl group has 1 to 5 carbon atoms ("$C_{1–5}$ alkyl"). In some embodiments, an alkyl group has 1 to 4 carbon atoms ("$C_{1–4}$ alkyl"). In some embodiments, an alkyl group has 1 to 3 carbon atoms ("$C_{1–3}$ alkyl"). In some embodiments, an alkyl group has 1 to 2 carbon atoms ("C_{1-2} alkyl"). In some embodiments, an alkyl group has 1 carbon atom ("C_1 alkyl"). Examples of $C_{1–6}$ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, hexyl, and the like.

[00036] As used herein, "alkenyl" refers to a radical of a straight–chain or branched hydrocarbon group having from 2 to 20 carbon atoms, one or more carbon–carbon double bonds (e.g., 1, 2, 3, or 4 carbon–carbon double bonds), and optionally one or more carbon–carbon triple bonds (e.g., 1, 2, 3, or 4 carbon–carbon triple bonds) ("$C_{2–20}$ alkenyl"). In certain embodiments, alkenyl does not contain any triple bonds. In some embodiments, an alkenyl group has 2 to 10 carbon atoms ("$C_{2–10}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 9 carbon atoms ("$C_{2–9}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 8 carbon atoms ("$C_{2–8}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 7 carbon atoms ("$C_{2–7}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 6 carbon atoms ("$C_{2–6}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 5 carbon atoms ("$C_{2–5}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 4 carbon atoms ("$C_{2–4}$ alkenyl"). In some embodiments, an alkenyl group has 2 to 3 carbon atoms ("$C_{2–3}$ alkenyl"). In some embodiments, an alkenyl group has 2 carbon atoms ("C_2 alkenyl"). The one or more carbon–carbon double bonds can be internal (such as in 2–butenyl) or terminal (such as in 1–butenyl). Examples of $C_{2–4}$ alkenyl groups include ethenyl (C_2), 1–propenyl (C_3), 2–propenyl (C_3), 1–butenyl (C_4), 2–butenyl (C_4), butadienyl (C_4), and the like. Examples of $C_{2–6}$ alkenyl groups include the aforementioned $C_{2–4}$ alkenyl groups as well as pentenyl (C_5),

pentadienyl (C_5), hexenyl (C_6), and the like. Additional examples of alkenyl include heptenyl (C_7), octenyl (C_8), octatrienyl (C_8), and the like.

[00037] As used herein, "alkynyl" refers to a radical of a straight–chain or branched hydrocarbon group having from 2 to 20 carbon atoms, one or more carbon–carbon triple bonds (e.g., 1, 2, 3, or 4 carbon–carbon triple bonds), and optionally one or more carbon–carbon double bonds (e.g., 1, 2, 3, or 4 carbon–carbon double bonds) ("$C_{2–20}$ alkynyl"). In certain embodiments, alkynyl does not contain any double bonds. In some embodiments, an alkynyl group has 2 to 10 carbon atoms ("$C_{2–10}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 9 carbon atoms ("$C_{2–9}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 8 carbon atoms ("$C_{2–8}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 7 carbon atoms ("$C_{2–7}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 6 carbon atoms ("$C_{2–6}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 5 carbon atoms ("$C_{2–5}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 4 carbon atoms ("$C_{2–4}$ alkynyl"). In some embodiments, an alkynyl group has 2 to 3 carbon atoms ("$C_{2–3}$ alkynyl"). In some embodiments, an alkynyl group has 2 carbon atoms ("C_2 alkynyl"). The one or more carbon–carbon triple bonds can be internal (such as in 2–butynyl) or terminal (such as in 1–butynyl). Examples of $C_{2–4}$ alkynyl groups include, without limitation, ethynyl (C_2), 1–propynyl (C_3), 2–propynyl (C_3), 1–butynyl (C_4), 2–butynyl (C_4), and the like. Examples of $C_{2–6}$ alkenyl groups include the aforementioned $C_{2–4}$ alkynyl groups as well as pentynyl (C_5), hexynyl (C_6), and the like. Additional examples of alkynyl include heptynyl (C_7), octynyl (C_8), and the like.

[00038] As used herein, "alkylene," "alkenylene," and "alkynylene," refer to a divalent radical of an alkyl, alkenyl, and alkynyl group respectively. When a range or number of carbons is provided for a particular "alkylene," "alkenylene," or "alkynylene," group, it is understood that the range or number refers to the range or number of carbons in the linear carbon divalent chain. "Alkylene," "alkenylene," and "alkynylene," groups may be substituted or unsubstituted with one or more substituents as described herein.

[00039] As used herein, "aryl" refers to a radical of a monocyclic or polycyclic (e.g., bicyclic or tricyclic) 4n+2 aromatic ring system (e.g., having 6, 10, or 14 π electrons shared in a cyclic array) having 6–14 ring carbon atoms and zero heteroatoms provided in the aromatic ring system ("$C_{6–14}$ aryl"). In some embodiments, an aryl group has six ring carbon atoms ("C_6 aryl"; e.g., phenyl). In some embodiments, an aryl group has ten ring carbon atoms ("C_{10} aryl"; e.g., naphthyl such as 1–naphthyl and 2–naphthyl). In some embodiments, an aryl group has fourteen ring carbon atoms ("C_{14} aryl"; e.g., anthracyl).

"Aryl" also includes ring systems wherein the aryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the radical or point of attachment is on the aryl ring, and in such instances, the number of carbon atoms continue to designate the number of carbon atoms in the aryl ring system. Typical aryl groups include, but are not limited to, groups derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, as-indacene, s-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, and trinaphthalene. Particularly aryl groups include phenyl, naphthyl, indenyl, and tetrahydronaphthyl.


[00040] As used herein, "heteroaryl" refers to a radical of a 5–10 membered monocyclic or bicyclic 4n+2 aromatic ring system (e.g., having 6 or 10 electrons shared in a cyclic array) having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen and sulfur ("5–10 membered heteroaryl"). In heteroaryl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. Heteroaryl bicyclic ring systems can include one or more heteroatoms in one or both rings. "Heteroaryl" includes ring systems wherein the heteroaryl ring, as defined above, is fused with one or more carbocyclyl or heterocyclyl groups wherein the point of attachment is on the heteroaryl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heteroaryl ring system. "Heteroaryl" also includes ring systems wherein the heteroaryl ring, as defined above, is fused with one or more aryl groups wherein the point of attachment is either on the aryl or heteroaryl ring, and in such instances, the number of ring members designates the number of ring members in the fused (aryl/heteroaryl) ring system. Bicyclic heteroaryl groups wherein one ring does not contain a heteroatom (e.g., indolyl, quinolinyl, carbazolyl, and the like) the point of attachment can be on either ring, i.e., either the ring bearing a heteroatom (e.g., 2–indolyl) or the ring that does not contain a heteroatom (e.g., 5–indolyl).

[00041] In some embodiments, a heteroaryl group is a 5–10 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5–10 membered heteroaryl"). In some embodiments, a heteroaryl group is a 5–8 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the

aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5–8 membered heteroaryl"). In some embodiments, a heteroaryl group is a 5–6 membered aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms provided in the aromatic ring system, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5–6 membered heteroaryl"). In some embodiments, the 5–6 membered heteroaryl has 1–3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heteroaryl has 1–2 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heteroaryl has 1 ring heteroatom selected from nitrogen, oxygen, and sulfur.

[00042] Exemplary 5–membered heteroaryl groups containing one heteroatom include, without limitation, pyrrolyl, furanyl and thiophenyl. Exemplary 5–membered heteroaryl groups containing two heteroatoms include, without limitation, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, and isothiazolyl. Exemplary 5–membered heteroaryl groups containing three heteroatoms include, without limitation, triazolyl, oxadiazolyl, and thiadiazolyl. Exemplary 5–membered heteroaryl groups containing four heteroatoms include, without limitation, tetrazolyl. Exemplary 6–membered heteroaryl groups containing one heteroatom include, without limitation, pyridinyl. Exemplary 6–membered heteroaryl groups containing two heteroatoms include, without limitation, pyridazinyl, pyrimidinyl, and pyrazinyl. Exemplary 6–membered heteroaryl groups containing three or four heteroatoms include, without limitation, triazinyl and tetrazinyl, respectively. Exemplary 7–membered heteroaryl groups containing one heteroatom include, without limitation, azepinyl, oxepinyl, and thiepinyl. Exemplary 5,6–bicyclic heteroaryl groups include, without limitation, indolyl, isoindolyl, indazolyl, benzotriazolyl, benzothiophenyl, isobenzothiophenyl, benzofuranyl, benzoisofuranyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzoxadiazolyl, benzthiazolyl, benzisothiazolyl, benzthiadiazolyl, indolizinyl, and purinyl. Exemplary 6,6–bicyclic heteroaryl groups include, without limitation, naphthyridinyl, pteridinyl, quinolinyl, isoquinolinyl, cinnolinyl, quinoxalinyl, phthalazinyl, and quinazolinyl.

[00043] Examples of representative heteroaryls include the following:



wherein each Z is selected from carbonyl, N, NR^{65}, O, and S; and R^{65} is independently hydrogen, C_1-C_8 alkyl, C_3-C_{10} carbocyclyl, 4-10 membered heterocyclyl, C_6-C_{10} aryl, and 5-10 membered heteroaryl.

[00044] As used herein, "carbocyclyl" or "carbocyclic" refers to a radical of a non–aromatic cyclic hydrocarbon group having from 3 to 10 ring carbon atoms ("C_{3-10} carbocyclyl") and zero heteroatoms in the non–aromatic ring system. In some embodiments, a carbocyclyl group has 3 to 8 ring carbon atoms ("C_{3-8} carbocyclyl"). In some embodiments, a carbocyclyl group has 3 to 6 ring carbon atoms ("C_{3-6} carbocyclyl"). In some embodiments, a carbocyclyl group has 5 to 10 ring carbon atoms ("C_{5-10} carbocyclyl"). Exemplary C_{3-6} carbocyclyl groups include, without limitation, cyclopropyl (C_3),cyclobutyl (C_4), cyclobutenyl (C_4), cyclopentyl (C_5), cyclopentenyl (C_5), cyclohexyl (C_6), cyclohexenyl (C_6), cyclohexadienyl (C_6), and the like. Exemplary C_{3-8} carbocyclyl groups include, without limitation, the aforementioned C_{3-6} carbocyclyl groups as well as cycloheptyl (C_7), cycloheptenyl (C_7), cycloheptadienyl (C_7), cycloheptatrienyl (C_7), cyclooctyl (C_8), cyclooctenyl (C_8), bicyclo[2.2.1]heptanyl (C_7), bicyclo[2.2.2]octanyl (C_8), and the like. Exemplary C_{3-10} carbocyclyl groups include, without limitation, the aforementioned C_{3-8} carbocyclyl groups as well as cyclononyl (C_9), cyclononenyl (C_9), cyclodecyl (C_{10}), cyclodecenyl (C_{10}), octahydro–1*H*–indenyl (C_9), decahydronaphthalenyl (C_{10}), spiro[4.5]decanyl (C_{10}), and the like. As the foregoing examples illustrate, in certain embodiments, the carbocyclyl group is either monocyclic ("monocyclic carbocyclyl") or contain a fused, bridged or spiro ring system such as a bicyclic system ("bicyclic carbocyclyl") and can be saturated or can be partially unsaturated. "Carbocyclyl" also includes ring systems wherein the carbocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups wherein the point of attachment is on the carbocyclyl ring, and

in such instances, the number of carbons continue to designate the number of carbons in the carbocyclic ring system.

[00045] The term "cycloalkyl" refers to a monovalent saturated cyclic, bicyclic, or bridged cyclic (e.g., adamantyl) hydrocarbon group of 3-12, 3-8, 4-8, or 4-6 carbons, referred to herein, e.g., as "C_{4-8}cycloalkyl," derived from a cycloalkane. Exemplary cycloalkyl groups include, but are not limited to, cyclohexanes, cyclopentanes, cyclobutanes and cyclopropanes. Unless specified otherwise, cycloalkyl groups are optionally substituted at one or more ring positions with, for example, alkanoyl, alkoxy, alkyl, haloalkyl, alkenyl, alkynyl, amido, amidino, amino, aryl, arylalkyl, azido, carbamate, carbonate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, imino, ketone, nitro, phosphate, phosphonato, phosphinato, sulfate, sulfide, sulfonamido, sulfonyl or thiocarbonyl. Cycloalkyl groups can be fused to other cycloalkyl, aryl, or heterocyclyl groups. In certain embodiments, the cycloalkyl group is not substituted, i.e., it is unsubstituted.

[00046] As used herein, "heterocyclyl" or "heterocyclic" refers to a radical of a 3– to 10–membered non–aromatic ring system having ring carbon atoms and 1 to 4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, sulfur, boron, phosphorus, and silicon ("3–10 membered heterocyclyl"). In heterocyclyl groups that contain one or more nitrogen atoms, the point of attachment can be a carbon or nitrogen atom, as valency permits. A heterocyclyl group can either be monocyclic ("monocyclic heterocyclyl") or a fused, bridged or spiro ring system such as a bicyclic system ("bicyclic heterocyclyl"), and can be saturated or can be partially unsaturated. Heterocyclyl bicyclic ring systems can include one or more heteroatoms in one or both rings. "Heterocyclyl" also includes ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more carbocyclyl groups wherein the point of attachment is either on the carbocyclyl or heterocyclyl ring, or ring systems wherein the heterocyclyl ring, as defined above, is fused with one or more aryl or heteroaryl groups, wherein the point of attachment is on the heterocyclyl ring, and in such instances, the number of ring members continue to designate the number of ring members in the heterocyclyl ring system.

[00047] In some embodiments, a heterocyclyl group is a 5–10 membered non–aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, sulfur, boron, phosphorus, and silicon ("5–10 membered heterocyclyl"). In some embodiments, a heterocyclyl group is a 5–8 membered non–aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms,

wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5–8 membered heterocyclyl"). In some embodiments, a heterocyclyl group is a 5–6 membered non–aromatic ring system having ring carbon atoms and 1–4 ring heteroatoms, wherein each heteroatom is independently selected from nitrogen, oxygen, and sulfur ("5–6 membered heterocyclyl"). In some embodiments, the 5–6 membered heterocyclyl has 1–3 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heterocyclyl has 1–2 ring heteroatoms selected from nitrogen, oxygen, and sulfur. In some embodiments, the 5–6 membered heterocyclyl has one ring heteroatom selected from nitrogen, oxygen, and sulfur.

[00048] Exemplary 3–membered heterocyclyl groups containing one heteroatom include, without limitation, azirdinyl, oxiranyl, thiorenyl. Exemplary 4–membered heterocyclyl groups containing one heteroatom include, without limitation, azetidinyl, oxetanyl and thietanyl. Exemplary 5–membered heterocyclyl groups containing one heteroatom include, without limitation, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothiophenyl, dihydrothiophenyl, pyrrolidinyl, dihydropyrrolyl and pyrrolyl–2,5–dione. Exemplary 5–membered heterocyclyl groups containing two heteroatoms include, without limitation, dioxolanyl, oxasulfuranyl, disulfuranyl, and oxazolidin-2-one. Exemplary 5–membered heterocyclyl groups containing three heteroatoms include, without limitation, triazolinyl, oxadiazolinyl, and thiadiazolinyl. Exemplary 6–membered heterocyclyl groups containing one heteroatom include, without limitation, piperidinyl, tetrahydropyranyl, dihydropyridinyl, and thianyl. Exemplary 6–membered heterocyclyl groups containing two heteroatoms include, without limitation, piperazinyl, morpholinyl, dithianyl, dioxanyl. Exemplary 6–membered heterocyclyl groups containing two heteroatoms include, without limitation, triazinanyl. Exemplary 7–membered heterocyclyl groups containing one heteroatom include, without limitation, azepanyl, oxepanyl and thiepanyl. Exemplary 8–membered heterocyclyl groups containing one heteroatom include, without limitation, azocanyl, oxecanyl and thiocanyl. Exemplary 5-membered heterocyclyl groups fused to a C_6 aryl ring (also referred to herein as a 5,6-bicyclic heterocyclic ring) include, without limitation, indolinyl, isoindolinyl, dihydrobenzofuranyl, dihydrobenzothienyl, benzoxazolinonyl, and the like. Exemplary 6-membered heterocyclyl groups fused to an aryl ring (also referred to herein as a 6,6-bicyclic heterocyclic ring) include, without limitation, tetrahydroquinolinyl, tetrahydroisoquinolinyl, and the like.

[00049] As used herein, "heterocylene" refers to a divalent radical of a heterocycle.

[00050] "Hetero" when used to describe a compound or a group present on a compound means that one or more carbon atoms in the compound or group have been replaced by a nitrogen, oxygen, or sulfur heteroatom. Hetero may be applied to any of the hydrocarbyl groups described above such as alkyl, e.g., heteroalkyl; carbocyclyl, e.g., heterocyclyl; aryl, e.g,. heteroaryl; and the like having from 1 to 5, and particularly from 1 to 3 heteroatoms.

[00051] As used herein, "cyano" refers to -CN.

[00052] As used herein, "halo" or "halogen" refers to fluoro (F), chloro (Cl), bromo (Br) and iodo (I). In certain embodiments, the halo group is either fluoro or chloro.

[00053] As used herein, "haloalkyl" refers to an alkyl group substituted with one or more halogen atoms.

[00054] As used herein, "nitro" refers to $-NO_2$.

[00055] As used herein, "oxo" refers to -C=O.

[00056] In general, the term "substituted", whether preceded by the term "optionally" or not, means that at least one hydrogen present on a group (e.g., a carbon or nitrogen atom) is replaced with a permissible substituent, e.g., a substituent which upon substitution results in a stable compound, e.g., a compound which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, or other reaction. Unless otherwise indicated, a "substituted" group has a substituent at one or more substitutable positions of the group, and when more than one position in any given structure is substituted, the substituent is either the same or different at each position.

[00057] Nitrogen atoms can be substituted or unsubstituted as valency permits, and include primary, secondary, tertiary, and quarternary nitrogen atoms. Exemplary nitrogen atom substitutents include, but are not limited to, hydrogen, –OH, $–OR^{aa}$, $–N(R^{cc})_2$, –CN, $–C(=O)R^{aa}$, $–C(=O)N(R^{cc})_2$, $–CO_2R^{aa}$, $–SO_2R^{aa}$, $–C(=NR^{bb})R^{aa}$, $–C(=NR^{cc})OR^{aa}$, $–C(=NR^{cc})N(R^{cc})_2$, $–SO_2N(R^{cc})_2$, $–SO_2R^{cc}$, $–SO_2OR^{cc}$, $–SOR^{aa}$, $–C(=S)N(R^{cc})_2$, $–C(=O)SR^{cc}$, $–C(=S)SR^{cc}$, $–P(=O)_2R^{aa}$, $–P(=O)(R^{aa})_2$, $–P(=O)_2N(R^{cc})_2$, $–P(=O)(NR^{cc})_2$, $C_{1–10}$ alkyl, $C_{1–10}$ perhaloalkyl, $C_{2–10}$ alkenyl, $C_{2–10}$ alkynyl, $C_{3–10}$ carbocyclyl, 3–14 membered heterocyclyl, $C_{6–14}$ aryl, and 5–14 membered heteroaryl, or two R^{cc} groups attached to a nitrogen atom are joined to form a 3–14 membered heterocyclyl or 5–14 membered heteroaryl ring, wherein each alkyl, alkenyl, alkynyl, carbocyclyl, heterocyclyl, aryl, and heteroaryl is independently substituted with 0, 1, 2, 3, 4, or 5 R^{dd} groups, and wherein R^{aa}, R^{bb}, R^{cc} and R^{dd} are as defined above.

[00058] These and other exemplary substituents are described in more detail in the **Detailed Description**, **Examples**, and **Claims**. The invention is not intended to be limited in any manner by the above exemplary listing of substituents.

Other Definitions

[00059] The term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, Berge *et al.*, describes pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences* (1977) 66:1–19, and Gould, Salt selection for basic drugs, International Journal of Pharmaceutics, 33 (1986) 201-217. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2–hydroxy–ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2–naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3–phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p–toluenesulfonate, undecanoate, valerate salts, and the like. Pharmaceutically acceptable salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate, and aryl sulfonate.

[00060] As used herein, a "subject" to which administration is contemplated includes, but is not limited to, humans (i.e., a male or female of any age group, e.g., a pediatric subject (e.g, infant, child, adolescent) or adult subject (e.g., young adult, middle–aged adult or senior adult)) and/or a non-human animal, e.g., a mammal such as primates (e.g., cynomolgus monkeys, rhesus monkeys), cattle, pigs, horses, sheep, goats, rodents, cats, and/or dogs. In certain embodiments, the subject is a human. In certain embodiments, the subject is a non-human animal. The terms "human," "patient," and "subject" are used interchangeably herein.

[00061] Disease, disorder, and condition are used interchangeably herein.

[00062] As used herein, and unless otherwise specified, the terms "treat," "treating" and "treatment" contemplate an action that occurs while a subject is suffering from the specified disease, disorder or condition, which reduces the severity of the disease, disorder or condition, or retards or slows the progression of the disease, disorder or condition (also "therapeutic treatment").

[00063] In general, the "effective amount" of a compound refers to an amount sufficient to elicit the desired biological response. As will be appreciated by those of ordinary skill in this art, the effective amount of a compound of the invention may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the disease being treated, the mode of administration, and the age, weight, health, and condition of the subject.

[00064] As used herein, and unless otherwise specified, a "therapeutically effective amount" of a compound is an amount sufficient to provide a therapeutic benefit in the treatment of a disease, disorder or condition, or to delay or minimize one or more symptoms associated with the disease, disorder or condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment of the disease, disorder or condition. The term "therapeutically effective amount" can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease or condition, or enhances the therapeutic efficacy of another therapeutic agent.

[00065] In an alternate embodiment, the present invention contemplates administration of the compounds of the present invention or a pharmaceutically acceptable salt or a pharmaceutically acceptable composition thereof, as a prophylactic before a subject begins to suffer from the specified disease, disorder or condition. As used herein, "prophylactic treatment" contemplates an action that occurs before a subject begins to suffer from the specified disease, disorder or condition. As used herein, and unless otherwise specified, a


"prophylactically effective amount" of a compound is an amount sufficient to prevent a disease, disorder or condition, or one or more symptoms associated with the disease, disorder or condition, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the disease, disorder or condition. The term "prophylactically effective amount" can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[00066] As used herein, a "disease or condition associated with a gain-of-function mutation in KCNT1" refers to a disease or condition that is associated with, is partially or completely caused by, or has one or more symptoms that are partially or completely caused by, a mutation in KCNT1 that results in a gain-of-function phenotype, i.e. an increase in activity of the potassium channel encoded by KCNT1 resulting in an increase in whole cell current.

[00067] As used herein, a "gain-of-function mutation" is a mutation in KCNT1 that results in an increase in activity of the potassium channel encoded by KCNT1. Activity can be assessed by, for example, ion flux assay or electrophysiology (e.g. using the whole cell patch clamp technique). Typically, a gain-of-function mutation results in an increase of at least or about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 125%, 150%, 175%, 200%, 225%, 250%, 275%, 300%, 325%, 350%, 375%, 400% or more compared to the activity of a potassium channel encoded by a wild-type KCNT1.

Compounds and Compositions

[00068] In one aspect, the present disclosure provides a compound of Formula **I**:



(I),

or a pharmaceutically acceptable salt thereof, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be substituted with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with

one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, $-C(O)OR_8$, $C(O)N(R_9)_2$, C_{1-6}alkylene-CN, -CN, $-S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, $-S(O)_2$-$N(R_9)_2$, $-OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, $-N(R_9)_2$, $-NR_9$-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, $-C(O)OR_8$, $-C(O)N(R_9)_2$,$-NR_{10}C(O)$-R_{11}, -CN, $-S(O)_2$-C_{1-6}alkyl, $-S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or

more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N.

[00069] In another aspect, the present disclosure provides a pharmaceutical composition comprising a compound of Formula **I**:



(I),

or a pharmaceutically acceptable salt thereof, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be substituted with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, -C(O)OR_8, C(O)$N(R_9)_2$, C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$-$N(R_9)_2$, -OC(O)C_{1-6}alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -$N(R_9)_2$, -NR_9-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)$N(R_9)_2$,-NR_{10}C(O)-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;



each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N;

and a pharmaceutically acceptable excipient.

[00070] In some embodiments, the compound is a compound of Formula **I-I** or Formula **I-II**:



(**I-I**) or

(**I-II**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00071] In some embodiments, one of X, Y, Z, Y', and Z' is N and the other four are CH.


[00072] In some embodiments, two of X, Y, Z, Y', and Z' are N and the other three are CH.

[00073] In some embodiments, the compound is a compound of Formula **I-a**:


(**I-a**),


or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00074] In some embodiments, the compound is a compound of Formula **I-b**:


(**I-b**),


or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00075] In some embodiments, the compound is a compound of Formula **I-c**:


(**I-c**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00076] In some embodiments, the compound is a compound of Formula **I-d**:



(**I-d**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00077] In some embodiments, the compound is a compound of Formula **I-e**:



(**I-e**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00078] In some embodiments, the compound is a compound of Formula **I-f**:



(**I-f**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00079] In some embodiments, the compound is a compound of Formula **I-g**:



(**I-g**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00080] In some embodiments, the compound is a compound of Formula **I-h**:



(**I-h**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00081] In some embodiments, the compound is a compound of Formula **I-i**:



(**I-i**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00082] In some embodiments, the compound is a compound of Formula **I-j**:



(I-j),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00083] In some embodiments, the compound is a compound of Formula **I-k**:



(I-k),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00084] In some embodiments, the compound is a compound of Formula **I-l**:



(I-l),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00085] In some embodiments, the compound is a compound of Formula **I-m**:



(I-m),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00086] In some embodiments, the compound is a compound of Formula **I-n**:



(I-n),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00087] In some embodiments, the compound is a compound of Formula **I-o**:



(I-o),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00088] In some embodiments, the compound is a compound of Formula **I-p**:



(**I-p**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00089] In some embodiments, the compound is a compound of Formula **I-q**:



(**I-q**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[00090] In some embodiments, the compound is a compound of Formula **I-r**:



(**I-r**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00091] In some embodiments, the compound is a compound of Formula **I-s**:



(**I-s**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[00092] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_2 is hydrogen. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_2 is methyl.

[00093] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is hydrogen. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is C_{1-6}alkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is selected from the group consisting of methyl, ethyl, and isopropyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-

l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is methyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is ethyl.

[00094] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 is C_{1-6}alkyl substituted with C_{1-6}alkoxy, -OH, or -C(O)OR_8.

[00095] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_4 is hydrogen.

[00096] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_3 and R_4 are taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene. In some embodiments, the C_{3-7}cycloalkylene is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. In some embodiments, the 3-7 membered heterocyclene is selected from the group consisting of oxetanyl, tetrahydrofuranyl, and tetrahydropyranyl.

[00097] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, C_{3-10}cycloalkyl, O-C_{3-10}cycloalkyl, -OH, -CN, $N(R_9)_2$, and -C(O)OR_8.

[00098] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is methyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is halogen. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is -F. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is -Cl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is methoxy. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is -CF_3. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g),

(I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is $-CHF_2$. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is $-C(O)OR_8$. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), each R_5 is cyclopropyl, cyclobutyl, or cyclopentyl.

[00099] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_5 is each independently selected from the group consisting of cyclopropyl, $-OCH_2CH_3$, $-OCH_2-CHF_2$, -O-cyclopropyl, -O-isopropyl, $-NHCH_3$, $-N(CH_3)_2$, and $-CH_2OCH_3$.

[000100] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 1. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 2.

[000101] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 1 and R_5 is at the *meta*-position. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 2 and the two R_5 are at the *ortho*- and *para*- positions. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 2 and the two R_5 are at the *meta*- and *para*- positions. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), n is 2 and the two R_5 are at the *meta*-positions.

[000102] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is selected from the group consisting of C_{1-6}alkyl optionally substituted with C_{1-6}alkoxy, $N(R_9)_2$, $C(O)N(R_9)_2$, C_{3-7}cycloalkyl, pyridyl, tetrahydropyranyl, or phenyl, C_{1-6}haloalkyl, C_{3-7}cycloalkyl, phenyl optionally substituted with halogen, and pyridyl optionally substituted with halogen.

[000103] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is C_{1-6}alkyl.

[000104] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is methyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is ethyl.

[000105] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{1\text{-}6}$haloalkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $-CH_2-CHF_2$. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $-CHF_2$.

[000106] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{3\text{-}7}$cycloalkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is cyclopropyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is cyclobutyl.

[000107] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{1\text{-}6}$alkyl substituted with $C_{1\text{-}6}$alkoxy. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{1\text{-}6}$alkyl substituted with methoxy.

[000108] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{1\text{-}6}$alkyl substituted with $C_{3\text{-}7}$cycloalkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is $C_{1\text{-}6}$alkyl substituted with cyclopropyl.

[000109] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_1 is phenyl substituted with halogen.

[000110] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is selected from the group consisting of $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered

heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy.

[000111] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}haloalkyl, C_{3-7}cycloalkyl, and phenyl optionally substituted with halogen.

[000112] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is C_{3-7}cycloalkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is cyclopropyl.

[000113] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is C_{1-6}alkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is ethyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is methyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is t-butyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is isopropyl.


[000114] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is C_{1-6}haloalkyl. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is $-CF_3$. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is $-CHF_2$.

[000115] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), R_{12} is phenyl optionally substituted with -F.

[000116] In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), x is

1. In some embodiments of formula (I) (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), x is 2.

[000117] In another aspect, the present disclosure provides a compound of Formula **II**:



(**II**),

or a pharmaceutically acceptable salt thereof, wherein

one or two selected from X, Y, Z, Y', and Z' are N, and the others are CH, wherein the hydrogen of CH may be substituted with R_5;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy optionally substituted with C_{3-7}cycloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally

substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, -$N(R_9)_2$, -$C(O)OR_8$, -$C(O)N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$-$N(R_9)_2$, -$OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -$N(R_9)_2$, -NR_9-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -$C(O)OR_8$, -$C(O)N(R_9)_2$,-$NR_{10}C(O)$-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is selected from the group consisting of hydrogen, C_{1-6}alkyl, and C_{3-10}cycloalkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl; and

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl.

[000118] In some embodiments, two of X, Y, Z, Y', and Z' are N and the other three are CH.


[000119] In some embodiments, the compound is a compound of Formula **II-a**:



(II-a),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000120] In some embodiments, the compound is a compound of Formula **II-b**:



(**II-b**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000121] In some embodiments, the compound is a compound of Formula **II-c**:



(**II-c**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000122] In some embodiments, the compound is a compound of Formula **II-d:**



(**II-d**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000123] In some embodiments, the compound is a compound of Formula **II-e**:



(**II-e**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000124] In some embodiments, the compound is a compound of Formula **II-f**:



(**II-f**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000125] In some embodiments, the compound is a compound of Formula **II-g**:



(**II-g**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000126] In some embodiments, the compound is a compound of Formula **II-h**:



(II-h),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000127] In some embodiments, the compound is a compound of Formula **II-i**:



(II-i),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000128] In some embodiments, the compound is a compound of Formula **II-j**:



(II-j),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000129] In some embodiments, the compound is a compound of Formula **II-k**:



(II-k),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000130] In some embodiments, the compound is a compound of Formula **II-l**:



(II-l),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000131] In some embodiments, the compound is a compound of Formula **II-m**:



(II-m),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000132] In some embodiments, the compound is a compound of Formula **II-n**:



(II-n),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[000133] In some embodiments, the compound is a compound of Formula **II-p**:



(II-p),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000134] In some embodiments, the compound is a compound of Formula **II-q**:



(II-q),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.


[000135] In some embodiments, the compound is a compound of Formula **II-r**:




(II-r),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above.

[000136] In some embodiments, the compound is a compound of Formula **II-k1** or Formula **II-k2**:



(II-k1)



(II-k2),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined as above.

[000137] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 1. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 2.

[000138] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 1 and R_5 is at the *meta*-position. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 2 and the two R_5 are at the *ortho*- and *para*- positions. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 2 and the two R_5 are at the *meta*- and *para*- positions. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), n is 2 and the two R_5 are at the *meta*-positions.

[000139] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), R_3 and R_4 are taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene. In some embodiments, the C_{3-7}cycloalkylene is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. In some embodiments, the 3-7 membered heterocyclene is selected from the group consisting of oxetanyl, tetrahydrofuranyl, and tetrahydropyranyl.

[000140] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), R_4 is hydrogen.

[000141] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), R_2 is hydrogen.

[000142] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), or (II-r)), R_2 is methyl.

[000143] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_3 is C_{1-6}alkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_3 is methyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_3 is ethyl.

[000144] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_3 is C_{1-6}alkyl substituted with C_{1-6}alkoxy, -OH, or -C(O)OR_8.

[000145] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_3 is hydrogen.

[000146] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), each R_5 is independently selected from the group consisting of C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, C_{3-10}cycloalkyl, O-C_{3-10}cycloalkyl, -CN, C_{1-6}alkylene-C_{1-6}alkoxy, C_{1-6}alkylene-N(R_9)$_2$, N(R_9)$_2$, and -C(O)OR_8.

[000147] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is C_{1-6}alkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is methyl.

[000148] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is halogen. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is -F. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is -Cl.

[000149] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is C_{1-6}haloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is CF_3. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is CF_2H.

[000150] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is C_{1-6}alkoxy. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c),

(II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is methoxy.

[000151] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is C_{3-10}cycloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is cyclopropyl.

[000152] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_5 is $-C(O)OR_8$.

[000153] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is selected from the group consisting of C_{1-6}alkyl optionally substituted with C_{1-6}alkoxy, $N(R_9)_2$, $C(O)N(R_9)_2$, C_{3-7}cycloalkyl, pyridyl, tetrahydropyranyl, or phenyl, C_{1-6}haloalkyl, C_{3-7}cycloalkyl, phenyl, and pyridyl.

[000154] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{1-6}alkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is methyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is ethyl.

[000155] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{1-6}haloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is $-CH_2-CHF_2$. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is $-CHF_2$.

[000156] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{3-7}cycloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is cyclopropyl, cyclobutyl, or cyclopentyl.

[000157] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is phenyl substituted with halogen.

[000158] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{1-6}alkyl substituted with C_{1-6}alkoxy. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{1-6}alkyl substituted with $N(R_9)_2$. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_1 is C_{1-6}alkyl substituted with cyclopropyl, cyclobutyl, or cyclopentyl.

[000159] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), x is 0 or 1.

[000160] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), x is 1.

[000161] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}haloalkyl, and phenyl optionally substituted with halogen.

[000162] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is C_{1-6}alkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is methyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is ethyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is t-butyl.

[000163] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is C_{1-6}haloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-


q), (II-r), (II-k1), or (II-k2)), R_{12} is CF_3. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is -CHF_2.

[000164] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is C_{3-7}cycloalkyl. In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), R_{12} is cyclopropyl.

[000165] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)),

[000166] In some embodiments of formula (II) (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), x is 0.

[000167] In another aspect, the present disclosure provides a compound of Formula **III**:



(III),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is C_{1-6}alkyl or C_{3-7}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-7}cycloalkyl is optionally substituted with one or more halogen or C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)N(R_9)$_2$, -C_{1-6}alkylene-CN, -CN, -S(O)$_2$-C_{1-6}alkyl, C_{1-6}alkylene-S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, -OC(O)C_{1-6}alkyl, and -O-C_{1-3}cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)N(R_9)$_2$,-NR_{10}C(O)-R_{11}, -CN, -S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;


each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;


wherein the compound is not:





, or a pharmaceutically acceptable salt thereof.

[000168] In another aspect, the present disclosure provides a compound of Formula **IV**:



(**IV**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is selected from the group consisting of C_{1-6}alkyl, and C_{3-7}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-7}cycloalkyl is optionally substituted with one or more substituents independently selected from halogen and C_{1-6}alkoxy;

R_{12} is C_{1-6}alkyl optionally substituted with one or more halogen or C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}haloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)N(R_9)$_2$, -C_{1-6}alkylene-CN, -CN, -S(O)$_2$-C_{1-6}alkyl, C_{1-6}alkylene-S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, -OC(O)C_{1-6}alkyl, and -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)N(R_9)$_2$,-NR_{10}C(O)-R_{11}, -CN, -S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;




each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and


n is selected from the group consisting of 0, 1, 2, and 3;

wherein the compound is not:

, or a pharmaceutically acceptable salt thereof.

[000169] In another aspect, the present disclosure provides a compound of Formula **V**:



(V),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is phenyl or 3-10 membered heteroaryl, wherein the phenyl or 3-10 membered heteroaryl is optionally substituted with one or more substituents independently selected from halogen and C_{1-6}alkoxy,

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)$N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, -OC(O)C_{1-6}alkyl, and -O-C_{3-10}cycloalkyl;

R_7 is selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-7}cycloalkyl, -C(O)OR_8, -C(O)N(R_9)$_2$,-NR_{10}C(O)-R_{11}, -CN, -S(O)$_z$-C_{1-6}alkyl, -S(O)$_z$- N(R_9)$_2$, 3-7 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;


wherein the compound is not:




, ,



,



,



, or a pharmaceutically acceptable salt thereof.

[000170] In some embodiments of formula (III), (IV), or (V), R_2 is hydrogen.

[000171] In some embodiments of formula (III), (IV), or (V), R_3 is C_{1-6}alkyl. In some embodiments of formula (III), (IV), or (V), R_3 is methyl. In some embodiments of formula (III), (IV), or (V), R_3 is ethyl.

[000172] In some embodiments of formula (III), (IV), or (V) , R_3 is C_{1-6}alkyl substituted with C_{1-6}alkoxy, -OH, or -C(O)OR_8.

[000173] In some embodiments of formula (III), (IV), or (V), R_4 is hydrogen.

[000174] In some embodiments of formula (III), (IV), or (V), R_3 and R_4 are taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene.

[000175] In some embodiments of formula (III), (IV), or (V), R_5 is each independently selected from the group consisting of cyclopropyl, -OCH_2CH_3, -OCH_2 -CHF_2, -O-cyclopropyl, -O-isopropyl, -$NHCH_3$, -$N(CH_3)_2$, and -CH_2OCH_3;

[000176] In some embodiments of formula (III), (IV), or (V), each R_5 is methyl.

[000177] In some embodiments of formula (III), (IV), or (V), each R_5 is halogen. In some embodiments of formula (III), (IV), or (V), each R_5 is -F. In some embodiments of formula (III), (IV), or (V), each R_5 is -Cl.

[000178] In some embodiments of formula (III), (IV), or (V), each R_5 is methoxy.

[000179] In some embodiments of formula (III), (IV), or (V), each R_5 is -CF_3.

[000180] In some embodiments of formula (III), (IV), or (V), each R_5 is -CHF_2.

[000181] In some embodiments of formula (III), (IV), or (V), each R_5 is -C(O)OR_8.

[000182] In some embodiments of formula (III), (IV), or (V), n is 1.

[000183] In some embodiments of formula (III), (IV), or (V), n is 2.

[000184] In some embodiments of formula (III), (IV), or (V), n is 1 and R_5 is at the *meta*-position. In some embodiments of formula (III), (IV), or (V), n is 2 and the two R_5 are at the *ortho*- and *para*- positions. In some embodiments of formula (III), (IV), or (V), n is 2 and the two R_5 are at the *meta*- and *para*- positions. In some embodiments of formula (III), (IV), or (V), n is 2 and the two R_5 are at the *meta*-positions.

[000185] In some embodiments of formula (III), (IV), or (V), R_1 is C_{1-6}alkyl. In some embodiments of formula (III), (IV), or (V), R_1 is methyl. In some embodiments of formula (III), (IV), or (V), R_1 is ethyl.

[000186] In some embodiments of formula (III), (IV), or (V), R_1 is C_{1-6}haloalkyl. In some embodiments of formula (III), (IV), or (V), R_1 is $-CH_2-CHF_2$. In some embodiments of formula (III), (IV), or (V), R_1 is $-CHF_2$.

[000187] In some embodiments of formula (III), (IV), or (V), R_1 is C_{3-7}cycloalkyl. In some embodiments of formula (III), (IV), or (V), R_1 is cyclopropyl.

[000188] In some embodiments of formula (III), (IV), or (V), R_1 is phenyl substituted with halogen.


[000189] In some embodiments of formula (III), (IV), or (V), R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy.

[000190] In some embodiments of formula (III), (IV), or (V), R_{12} is C_{3-7}cycloalkyl. In some embodiments of formula (III), (IV), or (V), R_{12} is cyclopropyl.

[000191] In some embodiments of formula (III), (IV), or (V), R_{12} is C_{1-6}alkyl. In some embodiments of formula (III), (IV), or (V), R_{12} is ethyl. In some embodiments of formula (III), (IV), or (V), R_{12} is methyl. In some embodiments of formula (III), (IV), or (V), R_{12} is t-butyl.

[000192] In some embodiments of formula (III), (IV), or (V), R_{12} is C_{1-6}haloalkyl. In some embodiments of formula (III), (IV), or (V), R_{12} is $-CF_3$. In some embodiments of formula (III), (IV), or (V), R_{12} is $-CHF_2$.

[000193] In another aspect, the present disclosure features a compound of Formula **VI**:



(VI),

or a pharmaceutically acceptable salt thereof, wherein

R_{13} is C_{1-6}alkyl or C_{3-10}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-10}cycloalkyl is optionally substituted with phenyl;

R_{14} is hydrogen;

R_{15} is C_{1-6}alkyl or hydrogen;

R_{16} is C_{1-6}alkyl optionally substituted with one or more halogen, C_{1-6}alkoxy, C_{3-10} cycloalkyl, or phenyl;

each R_{17} is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_{20})_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy optionally substituted with C_{3-7}cycloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, -$N(R_{20})_2$, -$C(O)OR_{19}$, -$C(O)N(R_{20})_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_{20})_2$, -$OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

p is selected from the group consisting of 1, 2, and 3;

R_{19} is selected from the group consisting of hydrogen, C_{1-6}alkyl, and C_{3-10}cycloalkyl;

each R_{20} is independently hydrogen or C_{1-6}alkyl; and

each R_{21} is independently hydrogen or C_{1-6}alkyl.

[000194] In some embodiments of formula (VI), R_{13} is selected from the group consisting of ethyl, tert-butyl, sec-butyl, iso-propyl, benzyl, and cyclopentyl.

[000195] In some embodiments of formula (VI), R_{15} is hydrogen.

[000196] In some embodiments of formula (VI), R_{16} is C_{1-6}alkyl.

[000197] In some embodiments of formula (VI), R_{16} is methyl or ethyl.

[000198] In some embodiments of formula (VI), p is 1 or 2.

[000199] In some embodiments of formula (VI), the compound is a compound of Formula **VI-a**:



(**VI-a**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined above for Formula VI.

[000200] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, C_{1-6}alkylene-C_{1-6}alkoxy, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen, or C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy.

[000201] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is cyclopropyl optionally substituted with C_{1-6}alkyl or C_{1-6}alkoxy.

[000202] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is cyclopropyl optionally substituted with methyl or methoxy.

[000203] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is C_{1-6}alkyl.

[000204] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is methyl.

[000205] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is C_{1-6}alkoxy, C_{1-6}alkylene-C_{1-6}alkoxy, or C_{1-6}haloalkoxy.


[000206] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is -$OCH(CH_3)_2$, -OCH_3, -OCH_2CH_3, O-CH_2CHF_2, or -CH_2OCH_3.

[000207] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is C_{1-6}haloalkyl.

[000208] In some embodiments of formula (VI) (e.g., (VI-a)), R_{17} is CHF_2 or CF_3.

[000209] In some embodiments of formula (VI) (e.g., (VI-a)), the compound is selected from the group consisting of Compound Nos. 272, 247, 262, 273, 274, 275, 276, 277, 284, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, and 296 in the Examples, or a pharmaceutically acceptable salt thereof.

[000210] In another aspect, the present disclosure features a compound of Formula **VII**:



(VII),

or a pharmaceutically acceptable salt thereof, wherein

W is N or CH;

R_{23} is C_{1-6}alkyl;

R_{24} is hydrogen;


R_{25} is C_{1-6}alkyl or hydrogen;

R_{26} is C_{1-6}alkyl optionally substituted with one or more halogen or C_{1-6}alkoxy;

each R_{27} is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; and

p is selected from the group consisting of 1, 2, and 3.

[000211] In another aspect, the present disclosure features a pharmaceutical composition comprising a compound of Formula **VII**:



(VII),

or a pharmaceutically acceptable salt thereof, wherein

W is N or CH;

R_{23} is C_{1-6}alkyl;

R_{24} is hydrogen;

R_{25} is C_{1-6}alkyl or hydrogen;


R_{26} is C_{1-6}alkyl optionally substituted with one or more halogen or C_{1-6}alkoxy;

each R_{27} is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; and


p is selected from the group consisting of 1, 2, and 3;

and a pharmaceutically acceptable excipient.

[000212] In some embodiments of formula (VII), the compound is a compound of Formula VII-a or Formula VII-b:



(VII-a),



(VII-b),

or a pharmaceutically acceptable salt thereof.

[000213] In some embodiments of formula (VII), p is 1.

[000214] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), R_{23} is tert-butyl.

[000215] In some embodiments of formula (VII), R_{25} is hydrogen.

[000216] In some embodiments of formula (VII), R_{26} is methyl.

[000217] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), R_{27} is halogen, C_{1-6}alkyl, or C_{1-6}alkoxy.

[000218] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), R_{27} is fluoro.

[000219] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), R_{27} is OCH_3.

[000220] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), R_{27} is methyl.

[000221] In some embodiments of formula (VII) (e.g., (VII-a) or (VII-b)), the compound is selected from the group consisting of Compound Nos. 281, 282, 283, and 285 in the Examples, or a pharmaceutically acceptable salt thereof.

[000222] In typical embodiments, the present invention is intended to encompass the compounds disclosed herein, and the pharmaceutically acceptable salts, tautomeric forms, polymorphs, and prodrugs of such compounds. In some embodiments, the present invention includes a pharmaceutically acceptable addition salt, a pharmaceutically acceptable ester, a solvate (e.g., hydrate) of an addition salt, a tautomeric form, all polymorphs including polymorphs of hydrates and solvates, an enantiomer, a mixture of enantiomers, a diastereomer, a mixture of diastereomers, a stereoisomer or mixture of stereoisomers (pure or as a racemic or non-racemic mixture) of a compound described herein, e.g., a compound of formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)).


[000223] In any and all aspects, in some embodiments, the compound of Formula (I), (II), (III), (IV), or (V) is selected from the group consisting of Compound Nos. 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237 238, 239, 240, 241, 242, 243, 244, 245, 246, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 263, 264, 265, 266, 267, 268, 269, 270, 271, 278, 279, 280, 297, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, and 310 in the Examples, or a pharmaceutically acceptable salt thereof.

General Synthetic Schemes

[000224] Exemplary methods for preparing compounds described herein are illustrated in the following synthetic schemes. These schemes are given for the purpose of illustrating the invention, and should not be regarded in any manner as limiting the scope or the spirit of the invention.


[000225] The synthetic route illustrated in Scheme 1 depicts an exemplary procedure for preparing carboxylic acid intermediates **D** and **G**. In the first step, compound **A** is reacted with hydrazine **B** to form ethyl pyrazole-5-carboxylate **C**. Then, hydrolysis of **C** provides carboxylic acid **D**. Carboxylic acid **D** may be coupled with amine **E** to form **F**, which is then hydrolyzed to yield carboxylic acid **G**.

SCHEME 1




[000226] The synthetic route illustrated in Scheme 2 depicts an exemplary procedure for preparing amine-substituted oxadiazole intermediate **L**. In the first step, nitrile **H** is treated with hydroxylamine to provide *N*-hydroxyimidamide **I**. Then, cabonyldiimidazole (CDI)-mediated cyclization of **I** with glycine **J** affords oxadiazole **K**. Deprotection of **K** under acidic conditions provides amine-substituted oxadiazole intermediate **L**.

SCHEME 2




[000227] The synthetic route illustrated in Scheme 3 depicts an exemplary procedure for preparing **M** (a compound of formula I). Coupling of amine-substituted oxadiazole amine **L** with carboxylic acid **D** using standard peptide coupling procedures (e.g., HOBt, and EDCI in dichloromethane in the presence of DIPEA) provides compound **M** (a compound of formula I).

SCHEME 3



[000228] As illustrated in Scheme 4, *N*-hydroxyimidamide **I** may undergo CDI-mediated cyclization with carboxylic acid **G** to afford compound M (a compound of formula I).

SCHEME 4



Methods of Treatment

[000229] The compounds and compositions described above and herein can be used to treat a neurological disease or disorder or a disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1). Exemplary diseases, disorders, or conditions include epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, developmental and epileptic encephalopathy (DEE), early infantile epileptic encephalopathy (EIEE), generalized epilepsy, focal epilepsy, multifocal epilepsy, temporal lobe epilepsy, Ohtahara syndrome, early myoclonic encephalopathy and Lennox Gastaut syndrome, drug resistant epilepsy, seizures (e.g., frontal lobe seizures, generalized tonic clonic seizures, asymmetric tonic seizures, focal seizures), leukodystrophy, hypomyelinating leukodystrophy, leukoencephalopathy, and sudden unexpected death in epilepsy, cardiac dysfunctions (e.g., cardiac arrhythmia, Brugada syndrome, myocardial infarction), pulmonary vasculopathy / hemorrhage, pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine, etc), muscle disorders (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity), itch and pruritis, movement disorders (e.g., ataxia and cerebellar ataxias), psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia, attention-deficit hyperactivity disorder), neurodevelopmental disorder, learning disorders, intellectual disability, Fragile X, neuronal plasticity, and autism spectrum disorders.

[000230] In some embodiments, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from EIMFS, ADNFLE and West syndrome. In some embodiments, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy and Lennox Gastaut syndrome. In some embodiments, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is seizure. In some embodiments, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from cardiac arrhythmia, Brugada syndrome, and myocardial infarction.

[000231] In some embodiments, the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from the group consisting of the learning disorders, Fragile X, intellectual function, neuronal plasticity, psychiatric disorders, and autism spectrum disorders.

[000232] Accordingly, the compounds and compositions thereof can be administered to a subject with a neurological disease or disorder or a disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene such as KCNT1 (e.g., EIMFS, ADNFLE, West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy, and Lennox Gastaut syndrome, seizures, cardiac arrhythmia, Brugada syndrome, and myocardial infarction).

[000233] EIMFS is a rare and debilitating genetic condition characterized by an early onset (before 6 months of age) of almost continuous heterogeneous focal seizures, where seizures appear to migrate from one brain region and hemisphere to another. Patients with EIMFS are generally intellectually impaired, non-verbal and non-ambulatory. While several genes have been implicated to date, the gene that is most commonly associated with EIMFS is KCNT1. Several de novo mutations in KCNT1 have been identified in patients with EIMFS, including V271F, G288S, R428Q, R474Q, R474H, R474C, I760M, A934T, P924L, G243S, H257D, A259D, R262Q, Q270E, L274I, F346L, C377S, R398Q, P409S, A477T, F502V, M516V, Q550del, K629E, K629N, I760F, E893K, M896K, R933G, R950Q, K1154Q (Barcia et al. (2012) Nat Genet. 44: 1255-1260; Ishii et al. (2013) Gene 531:467-471; McTague et al. (2013) Brain. 136: 1578-1591; Epi4K Consortium & Epilepsy Phenome/Genome Project. (2013) Nature 501:217-221; Lim et al. (2016) Neurogenetics; Ohba et al. (2015) Epilepsia 56:el21-el28; Zhou et al. (2018) Genes Brain Behav. e12456; Moller et al. (2015) Epilepsia. e114-20; Numis et al. (2018) Epilepsia. 1889-1898; Madaan et al. Brain Dev. 40(3):229-232; McTague et al. (2018) Neurology. 90(1):e55-e66; Kawasaki et al. (2017) J Pediatr. 191:270-274; Kim et al. (2014) Cell Rep. 9(5):1661-1672; Ohba et al. (2015) Epilepsia. 56(9):e121-8; Rizzo et al. (2016) Mol Cell Neurosci. 72:54-63; Zhang et al. (2017) Clin Genet. 91(5):717-724; Mikati et al. (2015) Ann Neurol. 78(6):995-9; Baumer et al. (2017) Neurology. 89(21):2212; Dilena et al. (2018) Neurotherapeutics. 15(4):1112-1126). These mutations are gain-of-function, missense mutations that are dominant (i.e. present on only one allele) and result in change in function of the encoded potassium channel that causes a marked increase in whole cell current when tested in Xenopus oocyte or mammalian

expression systems (see e.g. Milligan et al. (2015) Ann Neurol. 75(4): 581-590; Barcia et al. (2012) Nat Genet. 44(11): 1255-1259; and Mikati et al. (2015) Ann Neurol. 78(6): 995-999).

[000234] ADNFLE has a later onset than EIMFS, generally in mid-childhood, and is generally a less severe condition. It is characterized by nocturnal frontal lobe seizures and can result in psychiatric, behavioural and cognitive disabilities in patients with the condition. While ADNFLE is associated with genes encoding several neuronal nicotinic acetylcholine receptor subunits, mutations in the KCNT1 gene have been implicated in more severe cases of the disease (Heron et al. (2012) Nat Genet. 44: 1188-1190). Functional studies of the mutated KCNT1 genes associated with ADNFLE indicated that the underlying mutations (M896I, R398Q, Y796H and R928C) were dominant, gain-of-function mutations (Milligan et al. (2015) Ann Neurol. 75(4): 581-590; Mikati et al. (2015) Ann Neurol. 78(6): 995-999).

[000235] West syndrome is a severe form of epilepsy composed of a triad of infantile spasms, an interictal electroencephalogram (EEG) pattern termed hypsarrhythmia, and mental retardation, although a diagnosis can be made one of these elements is missing. Mutations in KCNT1, including G652V and R474H, have been associated with West syndrome (Fukuoka et al. (2017) Brain Dev 39:80-83 and Ohba et al. (2015) Epilepsia 56:el21-el28). Treatment targeting the KCNT1 channel suggests that these mutations are gain-of-function mutations (Fukuoka et al. (2017) Brain Dev 39:80-83).

[000236] In one aspect, the present invention features a method of treating treat a disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene such as KCNT1 (for example, epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy (DEE), and Lennox Gastaut syndrome, seizures, leukodystrophy, leukoencephalopathy, intellectual disability, Multifocal Epilepsy, Generalized tonic clonic seizures, Drug resistant epilepsy, Temporal lobe epilepsy, cerebellar ataxia, Asymmetric Tonic Seizures) and cardiac dysfunctions (e.g., cardiac arrhythmia, Brugada syndrome, sudden unexpected death in epilepsy, myocardial infarction), pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine, etc), muscle disorders (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity), itch and pruritis, ataxia and cerebellar ataxias, psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia), learning disorders, Fragile X, neuronal plasticity, and autism spectrum disorders) comprising administering to a subject in need thereof a compound disclosed herein

(e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof) or a pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient).

[000237] In some examples, the subject presenting with a disease or condition that may be associated with a gain-of-function mutation in KCNT1 is genotyped to confirm the presence of a known gain-of-function mutation in KCNT1 prior to administration of the compounds and compositions thereof. For example, whole exome sequencing can be performed on the subject. Gain-of-function mutations associated with EIMFS may include, but are not limited to, V271F, G288S, R428Q, R474Q, R474H, R474C, I760M, A934T, P924L, G243S, H257D, A259D, R262Q, Q270E, L274I, F346L, C377S, R398Q, P409S, A477T, F502V, M516V, Q550del, K629E, K629N, I760F, E893K, M896K, R933G, R950Q, and K1154Q. Gain-of-function mutations associated with ADNFLE may include, but are not limited to, M896I, R398Q, Y796H, R928C, and G288S. Gain-of-function mutations associated with West syndrome may include, but are not limited to, G652V and R474H. Gain-of-function mutations associated with temporal lobe epilepsy may include, but are not limited to, R133H and R565H. Gain-of-function mutations associated with Lennox-Gastaut may include, but are not limited to, R209C. Gain-of-function mutations associated with seizures may include, but are not limited to, A259D, G288S, R474C, R474H. Gain-of-function mutations associated with leukodystrophy may include, but are not limited to, G288S and Q906H. Gain-of-function mutations associated with Multifocal Epilepsy may include, but are not limited to, V340M. Gain-of-function mutations associated with EOE may include, but are not limited to, F346L and A934T. Gain-of-function mutations associated with Early-onset epileptic encephalopathies (EOEE) may include, but are not limited to, R428Q. Gain-of-function mutations associated with developmental and epileptic encephalopathies may include, but are not limited to, F346L, R474H, and A934T. Gain-of-

function mutations associated with epileptic encephalopathies may include, but are not limited to, L437F, Y796H, P924L, R961H. Gain-of-function mutations associated with Early Infantile Epileptic Encephalopathy (EIEE) may include, but are not limited to, M896K. Gain-of-function mutations associated with drug resistent epilepsy and generalized tonic-clonic seizure may include, but are not limited to, F346L. Gain-of-function mutations associated with migrating partial seizures of infancy may include, but are not limited to, R428Q. Gain-of-function mutations associated with Leukoencephalopathy may include, but are not limited to, F932I. Gain-of-function mutations associated with NFLE may include, but are not limited to, A934T and R950Q. Gain-of-function mutations associated with Ohtahara syndrome may include, but are not limited to, A966T. Gain-of-function mutations associated with infantile spasms may include, but are not limited to, P924L. Gain-of-function mutations associated with Brugada Syndrome may include, but are not limited to, R1106Q. Gain-of-function mutations associated with Brugada Syndrome may include, but are not limited to, R474H.

[000238] In other examples, the subject is first genotyped to identify the presence of a mutation in KCNT1 and this mutation is then confirmed to be a gain-of-function mutation using standard in vitro assays, such as those described in Milligan et al. (2015) Ann Neurol. 75(4): 581-590. Typically, the presence of a gain-of-function mutation is confirmed when the expression of the mutated KCNT1 allele results an increase in whole cell current compared to the whole cell current resulting from expression of wild-type KCNT1 as assessed using whole-cell electrophysiology (such as described in Milligan et al. (2015) Ann Neurol. 75(4): 581-590; Barcia et al. (2012) Nat Genet. 44(11): 1255-1259; Mikati et al. (2015) Ann Neurol. 78(6): 995-999; or Rizzo et al. Mol Cell Neurosci. (2016) 72:54-63). This increase of whole cell current can be, for example, an increase of at least or about 50%, 100%, 150%, 200%, 250%, 300%, 350%, 400% or more. The subject can then be confirmed to have a disease or condition associated with a gain-of-function mutation in KCNT1.

[000239] In particular examples, the subject is confirmed as having a KCNT1 allele containing a gain-of-function mutation (e.g. V271F, G288S, R398Q, R428Q, R474Q, R474H, R474C, G652V, I760M, Y796H, M896I, P924L, R928C or A934T).

[000240] The compounds disclosed herein (e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k) or a pharmaceutically acceptable salt thereof) or the pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound

disclosed herein (e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient) can also be used therapeutically for conditions associated with excessive neuronal excitability where the excessive neuronal excitability is not necessarily the result of a gain-of-function mutation in KCNT1. Even in instances where the disease is not the result of increased KCNT1 expression and/or activity, inhibition of KCNT1 expression and/or activity can nonetheless result in a reduction in neuronal excitability, thereby providing a therapeutic effect. Thus, the compounds disclosed herein (e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof) or the pharmaceutical composition disclosed herein (e.g., a pharmaceutical composition comprising a compound disclosed herein (e.g., a compound of Formula (I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) or a pharmaceutically acceptable salt thereof), and a pharmaceutically acceptable excipient) can be used to treat a subject with conditions associated with excessive neuronal excitability, for example, epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy, and Lennox Gastaut syndrome, seizures) or cardiac dysfunctions (e.g., cardiac arrhythmia, Brugada syndrome, myocardial infarction), regardless of whether or not the disease or disorder is associated with a gain-of-function mutation in KCNT1.

Pharmaceutical Compositions and Routes of Administration

[000241] Compounds provided in accordance with the present invention, e.g., a compound of formula ((I), (e.g., (I-I), (I-II), (I-a), (I-b), (I-c), (I-d), (I-e), (I-f), (I-g), (I-h), (I-

i), (I-j), (I-k), (I-l), (I-m), (I-n), (I-o), (I-p), (I-q), (I-r), or (I-s)), (II), (e.g., (II-a), (II-b), (II-c), (II-d), (II-e), (II-f), (II-g), (II-h), (II-i), (II-j), (II-k), (II-l), (II-m), (II-n), (II-o), (II-p), (II-q), (II-r), (II-k1), or (II-k2)), (III), (IV), (V), (VI), (e.g., (VI-a)), or (VII), (e.g., (VII-a) or (VII-b)) are usually administered in the form of pharmaceutical compositions. This invention therefore provides pharmaceutical compositions that contain, as the active ingredient, one or more of the compounds described, or a pharmaceutically acceptable salt or ester thereof, and one or more pharmaceutically acceptable excipients, carriers, including inert solid diluents and fillers, diluents, including sterile aqueous solution and various organic solvents, permeation enhancers, solubilizers and adjuvants. The pharmaceutical compositions may be administered alone or in combination with other therapeutic agents. Such compositions are prepared in a manner well known in the pharmaceutical art (see, e.g., Remington's Pharmaceutical Sciences, Mace Publishing Co., Philadelphia, Pa. 17th Ed. (1985); and Modern Pharmaceutics, Marcel Dekker, Inc. 3rd Ed. (G. S. Banker & C. T. Rhodes, Eds.)

[000242] The pharmaceutical compositions may be administered in either single or multiple doses by any of the accepted modes of administration of agents having similar utilities, for example as described in those patents and patent applications incorporated by reference, including rectal, buccal, intranasal and transdermal routes, by intra-arterial injection, intravenously, intraperitoneally, parenterally, intramuscularly, subcutaneously, orally, topically, as an inhalant, or via an impregnated or coated device such as a stent, for example, or an artery-inserted cylindrical polymer.

[000243] One mode for administration is parenteral, particularly by injection. The forms in which the novel compositions of the present invention may be incorporated for administration by injection include aqueous or oil suspensions, or emulsions, with sesame oil, corn oil, cottonseed oil, or peanut oil, as well as elixirs, mannitol, dextrose, or a sterile aqueous solution, and similar pharmaceutical vehicles. Aqueous solutions in saline are also conventionally used for injection, but less preferred in the context of the present invention. Ethanol, glycerol, propylene glycol, liquid polyethylene glycol, and the like (and suitable mixtures thereof), cyclodextrin derivatives, and vegetable oils may also be employed. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like.

[000244] Sterile injectable solutions are prepared by incorporating a compound according to the present invention in the required amount in the appropriate solvent with various other ingredients as enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[000245] Oral administration is another route for administration of compounds in accordance with the invention. Administration may be via capsule or enteric coated tablets, or the like. In making the pharmaceutical compositions that include at least one compound described herein, the active ingredient is usually diluted by an excipient and/or enclosed within such a carrier that can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be in the form of a solid, semi-solid, or liquid material (as above), which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, sterile injectable solutions, and sterile packaged powders.

[000246] Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, sterile water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl and propylhydroxy-benzoates; sweetening agents; and flavoring agents.

[000247] The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art. Controlled release drug delivery systems for oral administration include osmotic pump systems and dissolutional systems containing polymer-coated reservoirs or drug-polymer matrix formulations. Examples of controlled release systems are given in U.S. Pat. Nos. 3,845,770; 4,326,525; 4,902,514; and 5,616,345. Another formulation for use in the methods of the present invention employs transdermal delivery

devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Pat. Nos. 5,023,252, 4,992,445 and 5,001,139. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

[000248] The compositions are preferably formulated in a unit dosage form. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient (e.g., a tablet, capsule, ampoule). The compounds are generally administered in a pharmaceutically effective amount. Preferably, for oral administration, each dosage unit contains from 1 mg to 2 g of a compound described herein, and for parenteral administration, preferably from 0.1 to 700 mg of a compound a compound described herein. It will be understood, however, that the amount of the compound actually administered usually will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered and its relative activity, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

[000249] For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules.

[000250] The tablets or pills of the present invention may be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action, or to protect from the acid conditions of the stomach. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer that serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric

layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

[000251] Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described supra. Preferably, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably pharmaceutically acceptable solvents may be nebulized by use of inert gases. Nebulized solutions may be inhaled directly from the nebulizing device or the nebulizing device may be attached to a facemask tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions may be administered, preferably orally or nasally, from devices that deliver the formulation in an appropriate manner.

[000252] In some embodiments, a pharmaceutical composition comprising a disclosed compound, or pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

EXAMPLES

[000253] In order that the invention described herein may be more fully understood, the following examples are set forth. The synthetic and biological examples described in this application are offered to illustrate the compounds, pharmaceutical compositions and methods provided herein and are not to be construed in any way as limiting their scope.

[000254] The compounds provided herein can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless otherwise stated. Optimal reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization.

[000255] Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection are well known in the art. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and P. G. M. Wuts, *Protecting Groups in Organic Synthesis*, Second Edition, Wiley, New York, 1991, and references cited therein.

[000256] The compounds provided herein may be isolated and purified by known standard procedures. Such procedures include recrystallization, filtration, flash chromatography, trituration, high pressure liquid chromatography (HPLC), or supercritical fluid chromatography (SFC). Note that flash chromatography may either be performed manually or via an automated system. The compounds provided herein may be characterized by known standard procedures, such as nuclear magnetic resonance spectroscopy (NMR) or liquid chromatography mass spectrometry (LCMS). NMR chemical shifts are reported in part per million (ppm) and are generated using methods well known to those of skill in the art.


List of abbreviations

TEA	triethylamine
NaH	sodium hydride
THF	tetrahydrofuran
DMF	N,N-dimethylformamide
DCM	dichloromethane
EtOH	ethanol
ACN	acetonitrile
MeCN	acetonitrile
CH_3CN	acetonitrile
MeONa	sodium methoxide
CbzCl	benzyl chloroformate
MsCl	methanesulfonyl chloride
AcOH	acetic acid
TFA	trifluoroacetic acid
TFAA	trifluoroacetic anhydride
Py	pyridine
MeI	methyl iodide
LiOH	lithium hydroxide
$MeNH_2$	methylamine
$Cu(OAc)_2$	copper(II)acetate
DCC	N,N'-dicyclohexylcarbodiimide
DCE	dichloroethane
i-PrOH	isopropyl alcohol
EtOAc	ethyl acetate

HOBt	hydroxybenzotriazole
HATU	hexafluorophosphate azabenzotriazole tetramethyl uranium
DEA	diethanolamine
$MeNHNH_2$	methylhydrazine
EtONa	sodium ethoxide
SFC	supercritical fluid chemistry
DIPEA	N,N-diisopropylethylamine
EDCI	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
Boc	tert-butyloxycarbonyl
DMSO	dimethylsulfoxide
Boc-L-Ala-OH	N-(tert-butoxycarbonyl)-L-alanine
IPA	isopropyl alcohol
Ac_2O	acetic anhydride
PTFE	polytetrafluoroethylene
$Pd(dppf)Cl_2$	[1,1′-bis(diphenylphosphino)ferrocene]dichloropalladium(II)
DMS	dimethyl sulphate
T_3P	propylphosphonic anhydride solution
PE	petroleum ether
LCMS	liquid chromatography–mass spectrometry

Example 1. Synthesis of 192 and 191

a) Synthesis of 191



Synthesis of N'-hydroxy-3-methylbenzimidamide (A-3)

[000257] To a solution of 3-methylbenzonitrile (10 g, 85.36 mmol) in ethanol (200 mL) was added hydroxylamine hydrochloride (17.8 g, 256 mmol) and DIPEA (8.63 mL, 85.36 mmol). The reaction mixture was heated at 70 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (100 mL) and extracted with EtOAc (2 x 100 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-3** (9.3 g).

Synthesis of tert-butyl (R)-(1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-268)

[000258] To a solution of compound **A-3** (2.0 g, 13.32 mmol) in 1,4-dioxane (60 mL) was added (2R)-2-(tert-butoxycarbonylamino)propanoic acid (2.52 g, 13.32 mmol) and DCC (3.02 g, 14.65 mmol). The reaction mixture heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 40 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-268** (1.1 g, 3.8 mmol, 28% yield). **LCMS**: 302.1 (M-H), Rt 2.57 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (R)-1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-269)


[000259] To a stirred solution of compound **A-268** (500 mg, 1.65 mmol) in DCM (2.5 mL) was added TFA (1.5 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 8 h. The reaction mixture was concentrated and treated with saturated $NaHCO_3$ solution (10 mL). The mixture was extracted with EtOAc (2 x 20 mL) and washed with brine (20 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-269** (188 mg). The compound was used for the next step without further purification.


Synthesis of (R)-3-cyclopropyl-1-methyl-N-(1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (191)

[000260] To a stirred solution of compound **A-269** (188 mg, 0.92 mmol) in THF (5.0 mL) was added 3-cyclopropyl-1-methyl-1H-pyrazole-5-carboxylic acid (179 mg, 1.08 mmol) followed by T3P (1.17 mL, 1.97 mmol) and TEA (0.41 mL, 2.95 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (20 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by

column chromatography on silica gel with 30% EtOAc/PE to afford **191** (105 mg, 0.29 mmol, 30% yield) as a solid. **HPLC**: Rt 4.71 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 352.1 (M+H), Rt 2.36 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.79 min, SFC column: LUX A1; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 4.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 7.91-7.88 (m, 2H), 7.42-7.34 (m, 2H), 6.59 (d, 1H), 6.32 (s, 1H), 5.64-5.57 (m, 1H), 4.12 (s, 3H), 2.45 (s, 3H), 1.97-1.92 (m, 1H), 1.75 (d, 3H), 0.99-0.94 (m, 2H), 0.78-0.74 (m, 2H).

b) Synthesis of 192





Synthesis of tert-butyl (S)-(1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-270)

[000261] To a solution of compound **A-3** (1.0 g, 6.66 mmol) in 1,4-dioxane (60 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.37 g, 7.22 mmol) and DCC (1.51 g, 7.32 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (20 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-270** (0.80 g, 2.62 mmol, 39% yield).

LCMS: 302.1 (M-H), Rt 2.56 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm ; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H**

NMR (400 MHz, DMSO-d_6): δ 7.82-7.78 (m, 2H), 7.47-7.40 (m, 2H), 5.00-4.93 (m, 1H), 2.40 (s, 3H), 1.51 (d, 3H), 1.40 (s, 9H).


Synthesis of (S)-1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-271)

[000262] To a stirred solution of compound **A-270** (400 mg, 1.32 mmol) in DCM (2.4 mL) was added TFA (2.1 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 8 h. The reaction mixture was concentrated and treated with saturated $NaHCO_3$ solution (10 mL). The mixture was extracted with EtOAc (2 x 20 mL) and washed with brine (20 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-271** (200 mg). The compound was used for the next step without further purification.

Synthesis of (S)-3-cyclopropyl-1-methyl-N-(1-(3-(m-tolyl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (192)

[000263] To a stirred solution of A-271 (200 mg, 0.98 mmol) in THF (10.0 mL) was added 3-cyclopropyl-1-methyl-1H-pyrazole-5-carboxylic acid (179 mg, 1.08 mmol) followed by T3P (1.17 mL, 1.97 mmol) and TEA (0.41 mL, 20.52 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (20 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 30% EtOAc/PE to afford **192** (80 mg, 0.22 mmol, 23% yield) as a solid. **HPLC**: Rt 4.68 min, 99.7%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 352.3 (M+H), Rt 2.37 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.56 min, SFC column: LUX A1; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 4.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 7.91-7.88 (m, 2H), 7.42-7.34 (m, 2H), 6.58 (d, 1H), 6.32 (s, 1H), 5.62-5.58 (m, 1H), 4.12 (s, 3H), 2.45 (s, 3H), 1.97-1.93 (m, 1H), 1.75 (d, 3H), 0.99-0.94 (m, 2H), 0.78-0.74 (m, 2H).

Example 2. Synthesis of 3 and 4



[000264] **A-6:** A mixture of ethyl 5-cyclopropyl-1-methyl-pyrazole-3-carboxylate (1.4 g, 7.21 mmol) and NaOH (576.64 mg, 14.42 mmol) in ethanol (20 mL) and water (10 mL) was stirred at 50 °C for 2 hours. After cooling to room temperature, the reaction mixture was concentrated. Then the residue was diluted with H_2O (20 mL) and acidified with HCl (1M) to pH ~ 6, and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (2000 mg) as a solid. **LCMS** R_t = 0.46 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_{11}N_2O_2$ $[M+H]^+$ 167.1, found 167.0.

[000265] **A-7:** A mixture of 5-cyclopropyl-1-methyl-pyrazole-3-carboxylic acid (300 mg, 1.81 mmol), 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (432.72 mg, 1.81 mmol), TEA (913.38 mg, 9.03 mmol), EDCI (1038.21 mg, 5.42 mmol) and HOBt (731.84 mg, 5.42 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The reaction was quenched with the addition of sat. NH_4Cl (30 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 55-85 %B over 7 min) to give the product (300 mg, 0.85 mmol, 47% yield) as a solid. **LCMS** R_t = 1.11 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. for $C_{19}H_{22}N_5O_2$ $[M+H]^+$ 352.2, found 352.1.


[000266] **3 & 4:** Analytical SFC (Daicel CHIRALPAK AD-3 (50 mm x 3 mm, 3 μm), mobile phase: A: CO_2 B:ethanol (0.05% DEA), gradient: from 5% to 40% of B in 2.5 min and hold 40% for 0.35 min, then from 40% to 5% of B for 0.15 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 1.37 min and 1.61 min. The product was purified by SFC (Daicel CHIRALPAK AD-3 (50 mm x 3 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 25% B; 12 min run; 7 injections, Rt of peak 1 = 5.74 min, Rt of

peak 2 = 10.1 min) to give the enantiomer 1, randomly assigned as **3** (73.33 mg, 0.21 mmol, 24% yield) (Rt = 1.37 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **4** (77.03 mg, 0.22 mmol, 26% yield) **(**Rt = 1.61 min in analytical SFC) as a solid.

3: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.95 - 7.83 (m, 2H), 7.43 - 7.29 (m, 3H), 6.41 (s, 1H), 5.71 - 5.60 (m, 1H), 3.94 (s, 3H), 2.43 (s, 3H), 1.78 - 1.67 (m, 4H), 1.07 - 0.96 (m, 2H), 0.75 - 0.65 (m, 2H). **LCMS** R_t = 1.14 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{22}N_5O_2$ $[M+H]^+$ 352.2, found 352.1.

4: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.94 - 7.84 (m, 2H), 7.44 - 7.29 (m, 3H), 6.41 (s, 1H), 5.73 - 5.59 (m, 1H), 3.94 (s, 3H), 2.43 (s, 3H), 1.79 - 1.67 (m, 4H), 1.07 - 0.97 (m, 2H), 0.74 - 0.66 (m, 2H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{22}N_5O_2$ $[M+H]^+$ 352.2, found 352.1.

Example 3. Synthesis of 5




[000267] **A-10:** A mixture of 3-fluorobenzonitrile (500 mg, 4.13 mmol), hydroxylamine hydrochloride (860.66 mg, 12.39 mmol) and NaOH (495.42 mg, 12.39 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 12 hours to give a mixture. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (600 mg) as a solid. **LCMS** R_t = 0.46 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8FN_2O$ $[M+H]^+$ 155.1, found 154.8.

[000268] **5:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (170 mg, 0.72 mmol) and CDI (127.8 mg, 0.79 mmol) in DMF (3 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N'-hydroxy-benzamidine (110.44 mg, 0.72 mmol) was added. The reaction mixture was then stirred at 110 °C for 2 hours. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by Prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = water

(0.05% NH_4OH) and B = CH_3CN; 49-79% B over 8 min) to give the product (23.03 mg, 64.8 µmol, 9% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (dd, 1H), 7.81 - 7.76 (m, 1H), 7.52-7.46 (m, 1H), 7.26 - 7.19 (m, 1H), 6.52 (d, 1H), 6.31 (s, 1H), 5.64 - 5.54 (m, 1H), 4.10 (s, 3H), 1.98 - 1.87 (m, 1H), 1.75 (d, 3H), 1.00 - 0.91 (m, 2H), 0.78 - 0.72 (m, 2H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_2$ $[M+H]^+$356.1, found 356.0.

Example 4. Synthesis of 6




[000269] **A-12:** A mixture of 3-chlorobenzonitrile (500 mg, 3.63 mmol), hydroxylamine hydrochloride (757.69 mg, 10.9 mmol) and NaOH (436.14 mg, 10.9 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude. The crude product was purified by silica gel column (EtOAc in PE = 0% to 60%) to give the product (70 mg, 410.3 µmol, 11% yield) as a solid. **LCMS** R_t = 0.15 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8ClN_2O$ $[M+H]^+$ 171.0, found 170.9.

[000270] **6:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (105 mg, 0.44 mmol) and CDI (78.94 mg, 0.49 mmol) in DMF (3 mL) was stirred at 15 °C for 1 hour and then 3-chloro-N'-hydroxy-benzamidine (67.95 mg, 0.40 mmol) was added. The reaction mixture was stirred at 110 °C for 2 hours. After cooling to room temperature, the mixture was diluted with water (20 mL), extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered. The filtrate was concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters Xbridge (150 mm x 25 mm, 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 53-83% B over 8 min) to give the product (21.27 mg, 56.9 µmol, 13% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.09 (t, 1H),8.05 - 7.92 (m, 1H), 7.53 - 7.48 (m, 1H), 7.47 - 7.41 (m, 1H), 6.51 (d, 1H), 6.31 (s, 1H), 5.63 - 5.52 (m, 1H), 4.10 (s, 3H), 1.98 - 1.89 (m, 1H), 1.75 (d, 3H), 0.99 - 0.92 (m, 2H), 0.78 - 0.71 (m, 2H).

LCMS R_t = 1.31 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$ 372.1, found 372.0.

Example 5. Synthesis of 7




A-13 A-14 7

[000271] **A-14:** A mixture of 2,6-difluorobenzonitrile (500 mg, 3.59 mmol), hydroxylamine hydrochloride (749.35 mg, 10.78 mmol) and NaOH (431.34 mg, 10.78 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 2 hours. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (600 mg) as a solid. **LCMS** R_t = 0.26 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_7F_2N_2O$ $[M+H]^+$ 173.0, found 172.8

[000272] **7:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (150 mg, 0.63 mmol) and CDI (112.77 mg, 0.70 mmol) in DMF (3 mL) was stirred at 15 °C for 1 hour and then 2,6-difluoro-N'-hydroxy-benzamidine (119.71 mg, 0.70 mmol) was added. The reaction was then stirred at 110 °C for 2 hours. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered. The filtrate was concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters Xbridge (150 mm x 25mm, 5 μm), A = H_2O (0.05% NH_4OH v/v) and B = CH_3CN; 41-71% B over 8 min) to give the product (52.31 mg, 140.1 μmol, 22% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.55 - 7.45 (m, 1H), 7.07 (t, 2H), 6.60 (d, 1H), 6.27 (s, 1H), 5.69 - 5.60 (m, 1H), 4.09 (s, 3H), 1.96 - 1.86 (m, 1H), 1.76 (d, 3H), 0.96 - 0.88 (m, 2H), 0.74 - 0.66 (m, 2H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_2N_5O$ $[M+H]^+$ 374.1, found 374.0.

Example 6. Synthesis of 8



[000273] **A-16:** A mixture of 4-fluorobenzonitrile (500 mg, 4.13 mmol), hydroxylamine hydrochloride (860.66 mg, 12.39 mmol) and NaOH (495.42 mg, 12.39 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH, then diluted with H_2O (15 mL). The mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (300 mg) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.70 (s, 1H), 7.74 - 7.56 (m, 2H), 7.19 - 6.98 (m, 2H), 4.88 (s, 2H).


[000274] **A-17:** A mixture of 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (500 mg, 3.01 mmol), DIPEA (2.63 mL, 15.04 mmol), EDCI (865.18 mg, 4.51 mmol), HOBt (813.15 mg, 6.02 mmol) and methyl 2-aminopropanoate hydrochloride (419.97 mg, 3.01 mmol) in DCM (15 mL) was stirred 20 °C at for 16 hours. The reaction mixture was diluted with sat. NH_4Cl (20 mL). The mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (500 mg) as an oil. **LCMS** R_t = 0.72 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{18}N_3O_3$ $[M+H]^+$ 252.1, found 252.0.

[000275] **A-18:** To a solution of methyl 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoate (500 mg, 1.99 mmol) in THF (4 mL) was slowly added a solution of $LiOH.H_2O$ (166.99 mg, 3.98 mmol) in water (4 mL). The resulting mixture was stirred at 20 °C for 2 hours. The mixture was concentrated to remove THF. To the residue

was added 1N HCl to adjust the pH = 2, and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (350 mg) as oil. **LCMS** R_t = 0.67 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{16}N_3O_3$ $[M+H]^+$ 238.1, found 238.0.

[000276] **8:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (150 mg, 0.63 mmol) and CDI (112.77 mg, 0.70 mmol) in DMF (10 mL) was stirred at 15 °C for 1 hour and then 4-fluoro-N'-hydroxy-benzamidine (97.45 mg, 0.63 mmol) was added. The reaction mixture was then stirred at 110 °C for 16 hours. After cooling to room temperature, the mixture was diluted with H_2O (20 mL) and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Agela DuraShell (150 mm x 25 mm, 5 μm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-80% B over 8.5 minutes) to give the product (40.74 mg, 114.60 μmol, 18% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.15 - 8.00 (m, 2H), 7.23 - 7.13 (m, 2H), 6.53 (d, 1H), 6.30 (s, 1H), 5.65 - 5.51 (m, 1H), 4.10 (s, 3H), 1.98 - 1.87 (m, 1H), 1.74 (d, 3H), 1.03 - 0.88 (m, 2H), 0.81 - 0.68 (m, 2H). **LCMS** R_t = 1.15 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_2$ $[M+H]^+$ 356.1, found 356.0.


Example 7. Synthesis of 9



[000277] **A-20:** A mixture of 4-chlorobenzonitrile (494.5 mg, 3.59 mmol), hydroxylamine hydrochloride (749.35 mg, 10.78 mmol) and NaOH (431.34 mg, 10.78 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 2 hours. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (600 mg) as a solid. **LCMS** R_t = 0.43 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8ClN_2O$ $[M+H]^+$ 171.0, found 170.8.


[000278] **9:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (150 mg, 0.63 mmol) and CDI (112.77 mg, 0.70 mmol) in DMF (3 mL) was stirred at 15 °C for 1 hour and then 4-chloro-N'-hydroxy-benzamidine (118.64 mg, 0.70 mmol) was added. The reaction mixture was stirred at 110 °C for 2 hours. After cooling to room temperature, the mixture was diluted with water (20 mL), extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4 and filtered. The filtrate was concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm), A = water (0.05% NH_4OH) and B = CH_3CN; 49-79% B over 8 min) to give the product (53.21 mg, 143.1 µmol, 23% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.06 - 7.97 (m, 2H), 7.50 - 7.44 (m, 2H), 6.52 (d, 1H), 6.30 (s, 1H), 5.58 (quin, 1H), 4.09 (s, 3H), 1.98 - 1.60 (m, 1H), 1.74 (d, 3H), 1.00 - 0.91 (m, 2H), 0.78 - 0.70 (m, 2H). **LCMS** R_t = 1.20 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$372.1, found 372.0.

Example 8. Synthesis of 10



[000279] A mixture of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol), 5-isopropyl-2-methyl-pyrazole-3-carboxylic acid (70.17 mg, 0.42 mmol), HOBt (112.75 mg, 0.83 mmol), EDCI (119.96 mg, 0.63 mmol) and TEA (0.29 mL, 2.09 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The mixture was concentrated, diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Kromasil (150 mm x 30 mm, 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 55-85% B over 8 min) to give the product (44.51 mg, 0.13 mmol, 30% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.92 - 7.84 (m, 2H), 7.42 - 7.30 (m, 2H), 6.63 (d, 1H), 6.46 (s, 1H), 5.66 - 5.56 (m, 1H), 4.12 (s, 3H), 3.05 - 2.94 (m, 1H), 2.43 (s, 3H), 1.75 (d, 3H), 1.28 (d, 6H). **LCMS** R_t = 1.30 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.1.


Example 9. Synthesis of 11




A-5 11

[000280] A mixture of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol), 2-tert-butyl-5-methyl-pyrazole-3-carboxylic acid (76.02 mg, 0.42 mmol), HOBt (112.75 mg, 0.83 mmol), EDCI (119.96 mg, 0.63 mmol) and TEA (0.29 mL, 2.09 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The mixture was concentrated, diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Kromasil (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 55-85% B over 8 min) to give the product (30.81 mg, 0.08 mmol, 20% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.96 - 7.83 (m, 2H), 7.44 (d, 1H), 7.39 - 7.29 (m, 2H), 6.58 (s, 1H), 5.71 - 5.60 (m, 1H), 2.48 (s, 3H), 2.47 (s, 3H), 1.76 (d, 3H), 1.67 (s, 9H). **LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{26}N_5O_2$ $[M+H]^+$ 368.20, found 368.2.

Example 10. Synthesis of 12



A-19b A-19c




A-18 12

[000281] **A-19c:** A mixture of 3-(trifluoromethyl)benzonitrile (500 mg, 2.92 mmol), hydroxylamine hydrochloride (609.13 mg, 8.77 mmol) and NaOH (350.63 mg, 8.77 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (300 mg) as a solid. **^{1}H**

NMR (400MHz, $CDCl_3$) δ_H = 7.91 (s, 1H), 7.83 (d, 1H), 7.70 (d, 1H), 7.59 - 7.50 (m, 1H), 4.93 (br s, 2H).

[000282] **12:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (150 mg, 0.63 mmol) and CDI (112.77 mg, 0.70 mmol) in DMF (10 mL) was stirred at 15 °C for 1 hour and then N'-hydroxy-3-(trifluoromethyl)benzamidine (129.07 mg, 0.63 mmol) was added. The reaction mixture was then stirred at 110 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (20 mL), and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 μm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 20-50% B over 9 min) to give the product (26.89 mg, 65.60 mmol, 10% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.36 (s, 1H), 8.28 (d, 1H), 7.79 (d, 1H), 7.68 - 7.61 (m, 1H), 6.52 (d, 1H), 6.32 (s, 1H), 5.67 - 5.54 (m, 1H), 4.10 (s, 3H), 1.99 - 1.88 (m, 1H), 1.76 (d, 3H), 1.00 - 0.90 (m, 2H), 0.81 - 0.68 (m, 2H). **LCMS** R_t = 1.23 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{19}F_3N_5O_2$ $[M+H]^+$ 406.1, found 406.0.

Example 11. Synthesis of 13




A-5 13

[000283] A mixture of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol), 5-tert-butyl-2-methyl-pyrazole-3-carboxylic acid (76.02 mg, 0.42 mmol), HOBt (112.75 mg, 0.83 mmol), EDCI (119.96 mg, 0.63 mmol) and TEA (0.29 mL, 2.09 mmol) in DCM (20 mL) was stirred at 15 °C for 16 hours. The mixture was concentrated, diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Kromasil (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 60-90% B over 8 min) to give the product (61 mg, 0.17 mmol, 39% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.94 - 7.84 (m, 2H), 7.41 - 7.31 (m, 2H), 6.60 - 6.54 (m, 1H), 6.48 (s, 1H), 5.65 - 5.57 (m, 1H), 4.13 (s, 3H), 2.43 (s, 3H), 1.76 (d, 3H), 1.33 (s, 9H). **LCMS** R_t =

1.37 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{26}N_5O_2$ $[M+H]^+$ 368.20, found 368.1.

Example 12. Synthesis of 14 and 15



[000284] **A-21a:** A mixture of benzonitrile (370.67 mg, 3.59 mmol), hydroxylamine hydrochloride (749.35 mg, 10.78 mmol) and NaOH (431.34 mg, 10.78 mmol) in ethanol (6 mL) and water (2 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the reaction mixture was concentrated to remove most of EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (500 mg, 3.43 mmol, 95% yield) as a solid. **LCMS** R_t = 0.29 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_9N_2O$ $[M+H]^+$ 137.0, found 136.8.

[000285] **A-22:** A mixture of 2-[(5-cyclopropyl-2-methyl-pyrazole-3-carbonyl)amino]propanoic acid (200 mg, 0.84 mmol) and CDI (150.35 mg, 0.93 mmol) in DMF (3 mL) was stirred at 15 °C for 1 hour. Then N'-hydroxybenzamidine (126.25 mg, 0.93 mmol) was added and then the reaction mixture was then stirred at 110 °C for 2 hours to give a mixture. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 µm), A = water (0.05% NH_4OH v/v) and B = CH_3CN; 45-75% B over 8 min) to give the product (170 mg, 495.1 µmol, 59% yield) as a solid. **LCMS** R_t = 0.87 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{18}H_{20}N_5O_2$ $[M+H]^+$ 338.2, found 338.1.


[000286] **14 & 15:** Analytical SFC (Daicel CHIRACEL OJ-H (150 mm x 4.6 mm, 5 µm), mobile phase: A: CO_2 B: methanol (0.05% DEA,) gradient: hold 5% for 0.5 min, then

from 5% to 40% of B in 3.5 min and hold 40% for 2.5 min, then 5% of B for 1.5 min, flow rate: 3 mL/min, column temp: 40 °C.) showed two peaks at 3.40 min and 3.73 min. The product was separated by SFC (Daicel CHIRALPAK AS (250 mm x 30 mm, 5 μm); A = CO_2 and B = MeOH (0.1% NH_3H_2O); 35 °C; 50 mL/min; 20% B; 9 min run; 7 injections, Rt of peak 1 = 5.87 min, Rt of peak 2 = 7.39 min) to give the enantiomer 1, randomly assigned as **14** (11.44 mg, 33.9 μmol, 7% yield) (Rt = 3.40 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **15** (31.75 mg, 94.1 μmol, 19% yield) (Rt = 3.73 min in analytical SFC) as a solid.

14: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 8.08 (dd, 2H), 7.56 - 7.46 (m, 3H), 6.56 (d, 1H), 6.31 (s, 1H), 5.59 (quin, 1H), 4.10 (s, 3H), 1.98 - 1.87 (m, 1H), 1.75 (d, 3H), 0.99 - 0.90 (m, 2H), 0.77 - 0.68 (m, 2H). **LCMS** R_t = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{20}N_5O_2$ $[M+H]^+$ 338.2, found 338.0.

15: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 8.08 (dd, , 2H), 7.57 - 7.45 (m, 3H), 6.56 (d, 1H), 6.31 (s, 1H), 5.59 (quin, 1H), 4.10 (s, 3H), 1.99 - 1.88 (m, 1H), 1.75 (d, 3H), 1.00 - 0.91 (m, 2H), 0.79 - 0.69 (m, 2H). **LCMS** R_t = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{20}N_5O_2$ $[M+H]^+$ 338.2, found 338.0.

Example 13. Synthesis of 16 and 17




[000287] To a mixture of 2,5-dimethylpyrazole-3-carboxylic acid (58.46 mg, 420 μmol), EDCI (119.96 mg, 630 μmol), TEA (0.29 mL, 2.09 mmol) and HOBt (112.75 mg, 830 μmol) in DCM (20 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 420 μmol), and the mixture was stirred at 15 °C for 16 hours. The mixture was concentrated and diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The organic layer was washed brine (10 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 37-67% B over 8 min) to give **A-23** (65 mg, 195.7 μmol, 47% yield) as an oil. **LCMS** R_t = 0.78 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{17}H_{20}N_5O_2$ $[M+H]^+$ 326.15, found 326.7. Analytical SFC (Daicel CHIRALPAK IC-3 (150 x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and

hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.61 min and 4.64 min. The product was separated by SFC (Daicel CHIRALPAK IC (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 65 mL/min; 30% B; 7 min run; 4 injections, Rt of peak 1 = 3.75 min, Rt of peak 2 = 5.50 min) to give enantiomer 1, randomly assigned as **16** (12.35 mg, 38 µmol, 19% yield) (Rt = 3.61 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **17** (11.51 mg, 35.4 µmol, 18% yield) (Rt =4.64 min in analytical SFC) as a solid.

16: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.97 - 7.83 (m, 2H), 7.44 - 7.31 (m, 2H), 6.60 (d, 1H), 6.44 (s, 1H), 5.60 (quin, 1H), 4.12 (s, 3H), 2.44 (s, 3H), 2.30 (s, 3H), 1.75 (d, 3H). **LCMS** R_t = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{20}N_5O_2$ $[M+H]^+$ 326.15, found 325.9.

17: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.93 - 7.85 (m, 2H), 7.43 - 7.31 (m, 2H), 6.59 (d, 1H), 6.44 (s, 1H), 5.60 (quin, 1H), 4.12 (s, 3H), 2.44 (s, 3H), 2.30 (s, 3H), 1.75 (d, 3H). **LCMS** R_t = 1.17 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{20}N_5O_2$ $[M+H]^+$ 326.15, found 326.0.

Example 14. Synthesis of 19 and 20



A-5 A-27 19 20

[000288] **A-27:** To a solution of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (150 mg, 630 µmol) and 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (242.94 mg, 1.25 mmol), HATU (475.88 mg, 1.25 mmol) in DMF (10 mL) was added DIPEA (0.44 mL, 2.5 mmol) and the reaction mixture was stirred at 15 °C for 3 hours. The mixture was diluted with H_2O (40 mL) and then extracted with EtOAc (40 mL x 3). The combined organic layer was washed brine (40 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN;


65-75% B over 8 min) to give the product (200 mg, 527.2 µmol, 84% yield) as an oil. **LCMS** R_t = 3.97 min in 7.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}F_3N_5O_2$ $[M+H]^+$ 380.13, found 380.0.

[000289] **19 & 20:** Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 0.5 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C.) showed two peaks at 2.53 min and 3.11 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 35 °C; 50 mL/min; 25% B; 7 min run; 6 injections, Rt of peak 1 = 3.45 min, Rt of peak 2 = 4.40 min) to give the enantiomer 1, randomly assigned as **19** (55.59 mg, 146.5 µmol, 28% yield) (Rt = 2.53 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **20** (61.98 mg, 163 µmol, 31% yield) (Rt =3.11 min in analytical SFC) as a solid.

19: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.91 - 7.85 (m, 2H), 7.42 - 7.30 (m, 2H), 6.93 (s, 1H), 6.71 (br d, 1H), 5.60 (quin, 1H), 4.24 (s, 3H), 2.44 (s, 3H), 1.77 (d, 3H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}F_3N_5O_2$ $[M+H]^+$ 380.13, found 379.9.

20: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.92 - 7.84 (m, 2H), 7.43 - 7.32 (m, 2H), 6.93 (s, 1H), 6.71 (br d, 1H), 5.60 (quin, 1H), 4.24 (s, 3H), 2.44 (s, 3H), 1.77 (d, 3H). **LCMS** R_t = 1.34 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}F_3N_5O_2$ $[M+H]^+$ 380.13, found 379.9.

Example 15. Synthesis of 21




A-5 **21**


[000290] A mixture of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol), 5-(difluoromethyl)-2-methyl-pyrazole-3-carboxylic acid (73.48 mg, 0.42 mmol), HOBt (112.75 mg, 0.83 mmol), EDCI (119.96 mg, 0.63 mmol) and TEA (0.29 mL, 2.09 mmol) in DCM (20 mL) was stirred at 15 °C for 16 hours. The mixture was concentrated, diluted with H_2O (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and

concentrated to give the crude product. The crude product was purified by prep-HPLC (Kromasil (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 53-83% B over 8 min) to give the product (96 mg, 0.26 mmol, 63% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ_H = 9.40 (d, 1H), 7.87 - 7.76 (m, 2H), 7.51 - 7.37 (m, 2H), 7.28 (s, 1H), 7.06 (t, 1H), 5.49 - 5.40 (m, 1H), 4.09 (s, 3H), 2.39 (s, 3H), 1.67 (d, 3H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.14, found 362.1.

Example 16. Synthesis of 22



A-29a A-29b A-29



22


[000291] **A-29b:** To a mixture of Cs_2CO_3 (4649.64 mg, 14.27 mmol) and 2-bromopropane (1755.24 mg, 14.27 mmol) in DMF (15 mL) was added methyl 3-methyl-1H-pyrazole-5-carboxylate (1000 mg, 7.14 mmol), and then the reaction mixture was stirred at 100 °C for 2.5 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (100 mL) and then extracted with EtOAc (150 mL x 2). The combined organic layer was washed with brine (150 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE= 0% to 5%) to give the product (620 mg, 3.24 mmol, 45% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 6.58 (s, 1H), 5.46 (spt, 1H), 3.86 (s, 3H), 2.29 (s, 3H), 1.48 (d, 6H). **LCMS** R_t = 3.60 min in 7.0 min chromatography, 0-60AB, MS ESI calcd. for $C_9H_{15}N_2O_2$ $[M+H]^+$ 183.11, found 182.9.

[000292] **A-29:** To a solution of methyl 2-isopropyl-5-methyl-pyrazole-3-carboxylate (300 mg, 1.65 mmol) in ethanol (5mL) was added a solution of NaOH (65.85 mg, 1.65 mmol) in water (5 mL). The mixture was stirred at 15 °C for 2 hours. The reaction mixture was concentrated to give a residue. The residue was diluted with H_2O (20 mL) and washed

with EtOAc (20 mL x 1). The aqueous phase was acidified with 1N HCl (20 mL) to pH = 1. The mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (70 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (200 mg) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 6.72 (s, 1H), 5.50 - 5.40 (m, 1H), 2.37 - 2.26 (m, 3H), 1.50 (d, 6H).

[000293] **22:** To a mixture of 2-isopropyl-5-methyl-pyrazole-3-carboxylic acid (140.33 mg, 0.83 mmol), HATU (317.25 mg, 0.83 mmol), DIPEA (0.29 mL, 1.67 mmol) in DCM (8 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol) and the mixture was stirred at 15 °C for 12 hours. The mixture was concentrated and diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The organic layer was washed brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 37-67% B over 8 min) to give the product (35.39 mg, 99.7 µmol, 24% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.93 - 7.82 (m, 2H), 7.42 - 7.33 (m, 2H), 6.59 (br d, 1H), 6.39 (s, 1H), 5.58 (quin, 1H), 5.40 (spt, 1H), 2.43 (s, 3H), 2.32 (s, 3H), 1.75 (d, 3H), 1.48 (dd, 6H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.2.

Example 17. Synthesis of 23 and 24



6 23 24


[000294] Analytical SFC (Daicel CHIRALPAK AD-3 (150 mm × 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C.) showed two peaks at 3.76 min and 4.09 min. The product was separated by SFC (Daicel CHIRALPAK AS (250 mm x 30 mm, 5 µm); A = CO_2 and B = MeOH (0.1% NH_3H_2O); 38 °C; 50 mL/min; 20% B; 10 min run; 8 injections, Rt of peak 1 = 7.2 min, Rt of peak 2 = 8.5 min) to give the enantiomer 1, randomly assigned as **23** (2.31 mg, 6.2 µmol, 13% yield) (Rt = 3.76 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **24** (2.02 mg, 5.4 µmol, 11% yield) (Rt =4.09 min in analytical SFC) as a solid.

23: ^{1}H NMR (400MHz, CD_3CN) δ_H = 8.10 - 8.00 (m, 1H), 8.00 - 7.94 (m, 1H), 7.61 - 7.40 (m, 3H), 6.46 (s, 1H), 5.43 (quin, 1H), 3.96 (s, 3H), 1.90 - 1.84 (m, 1H), 1.69 (d, 3H), 0.91 -

0.86 (m, 2H), 0.68 - 0.63 (m, 2H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$ 372.1, found 372.0.

24: ^{1}H NMR (400MHz, CD_3CN) δ_H = 8.09 - 8.01 (m, 1H), 8.00 -7.95 (m, 1H), 7.62 - 7.43 (m, 3H), 6.46 (s, 1H), 5.43 (quin, 1H), 3.96 (s, 3H), 1.91 - 1.83 (m, 1H), 1.69 (d, 3H), 0.92 - 0.86 (m, 2H), 0.69 - 0.62 (m, 2H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$ 372.1, found 371.9.

Example 18. Synthesis of 25 and 26




[000295] Analytical SFC (Regis, (S,S) Whelk-O1 (250 mm x 4.6 mm, 5 μm), mobile phase: A: CO_2 B: methanol (0.05% DEA), gradient: hold 5% for 0.5 min, then from 5% to 40% of B in 3.5 min and hold 40% for 2.5 min, then 5% of B for 1.5 min, flow rate: 3 mL/min, column temp: 35 °C) showed two peaks at 3.51 min and 4.72 min. The product was separated by SFC (Daicel CHIRALPAK AS (250 mm x 30 mm, 5 μm); A = CO_2 and B = MeOH (0.1% NH_3H_2O); 38 °C; 50 mL/min; 20% B; 9 min run; 5 injections, Rt of peak 1 = 5.9 min, Rt of peak 2 =7.5 min) to give the enantiomer 1, randomly assigned as **25** (8.68 mg, 23.3 μmol, 17% yield) (Rt = 3.51 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **26** (7.31 mg, 19.7 μmol, 15% yield) (Rt = 4.72 min in analytical SFC) as a solid.

[000296] **25: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.06 - 7.99 (m, 2H), 7.51 - 7.44 (m, 2H), 6.53 (d, 1H), 6.30 (s, 1H), 5.59 (quin, 1H), 4.09 (s, 3H), 2.00 - 1.89 (m, 1H), 1.74 (d, 3H), 0.99 - 0.92 (m, 2H), 0.78 - 0.71 (m, 2H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$ 372.1, found 372.0.

[000297] **26: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.08 - 7.97 (m, 2H), 7.51 - 7.43 (m, 2H), 6.52 (d, 1H), 6.30 (s, 1H), 5.58 (quin, 1H), 4.10 (s, 3H), 1.99 - 1.87 (m, 1H), 1.74 (d, 3H), 1.00 - 0.91 (m, 2H), 0.78 - 0.70 (m, 2H). **LCMS** R_t = 1.27 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}ClN_5O_2$ $[M+H]^+$ 372.1, found 372.1.

Example 19. Synthesis of 29 and 30




13 29 30

[000298] The product was analyzed by SFC (Daicel CHIRALCEL OJ-3 (100 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 4.5 min and hold 40% for 0.5 min, then 5% of B for 1 min, flow rate: 2.8 mL/min, column temp: 40 °C) showed two peaks at 1.91 min and 2.22 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 25% B; 8 min run; 10 injections, Rt of peak 1 = 5.38 min, Rt of peak 2 = 6.62 min) to give the enantiomer 1, randomly assigned as **29** (12.72 mg, 0.03 mmol, 22% yield) (Rt = 1.91 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned 30 (15.52 mg, 0.04 mmol, 27 % yield) (Rt = 2.22 min in analytical SFC) as a solid.

29: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.93 - 7.84 (m, 2H), 7.43 - 7.30 (m, 2H), 6.57 (d, 1H), 6.48 (s, 1H), 5.61 (quin, 1H), 4.13 (s, 3H), 2.44 (s, 3H), 1.76 (d, 3H), 1.33 (s, 9H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{26}N_5O_2$ $[M+H]^+$ 368.20, found 368.2.

30: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.93 - 7.85 (m, 2H), 7.42 - 7.31 (m, 2H), 6.56 (d, 1H), 6.47 (s, 1H), 5.61 (quin, 1H), 4.13 (s, 3H), 2.44 (s, 3H), 1.76 (d, 3H), 1.33 (s, 9H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{26}N_5O_2$ $[M+H]^+$ 368.20, found 368.2.

Example 20. Synthesis of 31 and 32



22 31 32

[000299] Analytical SFC (Regis (R,R) Whelk-01 (100 mm × 4.6 mm, 5 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 3.68 min and 4.40 min.

[000300] The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = ethanol (0.1% NH_3H_2O); 38 °C; 60 mL/min; 60% B; 10 min run; 3 injections, Rt of peak 1 = 7.9 min, Rt of peak 2 = 5.5 min) to give the enantiomer 1, randomly

assigned as **31** (9.92 mg, 28.1 µmol, 33% yield) (Rt = 3.68 min in analytical SFC) as oil and the enantiomer 2, randomly assigned as **32** (9.16 mg, 25.9 µmol, 30% yield) (Rt = 4.40 min in analytical SFC) as oil.

31

^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.12 (d, 1H), 7.85 - 7.74 (m, 2H), 7.48 - 7.36 (m, 2H), 6.66 (s, 1H), 5.43 - 5.27 (m, 2H), 2.38 (s, 3H), 2.18 (s, 3H), 1.63 (d, 3H), 1.36 - 1.27 (m, 6H).


LCMS R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.1.

32

^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.13 (d, 1H), 7.85 - 7.76 (m, 2H), 7.49 - 7.37 (m, 2H), 6.66 (s, 1H), 5.45 - 5.26 (m, 2H), 2.39 (s, 3H), 2.19 (s, 3H), 1.64 (d, 3H), 1.37 - 1.28 (m, 6H).

LCMS R_t = 1.27 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.0.

Example 21. Synthesis of 33




[000301] **A-31:** To a solution of methyl 3-methyl-1H-pyrazole-5-carboxylate (800 mg, 5.71 mmol) in MeCN (20 mL) was added Cs_2CO_3 (3719.71 mg, 11.42 mmol), sodium 2-chloro-2,2-difluoro-acetate (1740.66 mg, 11.42 mmol) and 18-crown-6 (301.77 mg, 1.14 mmol) and the reaction mixture was stirred at 90 °C for 2 hours under N_2. After cooling to room temperature, the mixture was filtered, and the filtrate was concentrated to give a residue. The residue was diluted with H_2O (50 mL) and extracted with EtOAc (40 mL x 2). The combined organic phase was washed with brine (60 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5.5% to 40%) to give the product (120

mg, 631.1 μmol, 11 % yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.03 (t, 1H), 6.77 (s, 1H), 3.92 (s, 3H), 2.36 (s, 3H).

[000302] **A-32:** To a solution of methyl 2-(difluoromethyl)-5-methyl-pyrazole-3-carboxylate (120 mg, 631.1 μmol) in ethanol (2 mL) was added a solution of NaOH (25.24 mg, 631.1 μmol) in water (2 mL). The mixture was stirred at 20 °C for 2 hours. The reaction mixture was then diluted with H_2O (15 mL) and washed with EtOAc (20 mL, discarded). The aqueous phase was acidified with 1N HCl (20 mL) to adjust th pH = 1 and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (70 mL), dried over Na_2SO_4, filtered and concentrated to give the product (100 mg, 567.8 μmol, 90% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 14.15 (br s, 1H), 8.13 (t, 1H), 6.86 (s, 1H), 2.26 (s, 3H).

[000303] **33:** To a mixture of 2-(difluoromethyl)-5-methyl-pyrazole-3-carboxylic acid (100 mg, 570 μmol), HATU (380.71 mg, 1 mmol), DIPEA (0.35 mL, 2 mmol) in DCM (8 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (120.mg, 500 μ mol) and the mixture was stirred at 15 °C for 2 hours. The mixture was concentrated and diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The combined organic layer was washed brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 53-83% B over 8 min) to give the product (109.62 mg, 303.4 μmol, 61% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.55 (d, 1H), 8.19 (t, 1H), 7.85 - 7.75 (m, 2H), 7.49 - 7.37 (m, 2H), 7.02 (s, 1H), 5.44 (quin, 1H), 2.39 (s, 3H), 2.28 (s, 3H), 1.66 (d, 3H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.14, found 362.0.

Example 22. Synthesis of 34 and 35




[000304] Analytical SFC (CHIRALCEL OJ-3 (100 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 4.5 min and hold 40% for 0.5 min, then 5% of B for 1 min, flow rate: 2.8 mL/min, column temp: 40 °C) showed two peaks at 2.01 min and 2.31 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min;

25% B; 8 min run; 10 injections, Rt of peak 1 = 5.37 min, Rt of peak 2 = 6.75 min) to give the enantiomer 1, randomly assigned as (5.56 mg, 0.02 mmol, 13% yield) (Rt = 2.01 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as (10.86 mg, 0.03 mmol, 26% yield) (Rt = 2.31 min in analytical SFC) as a solid.

34: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.92 - 7.86 (m, 2H), 7.41 - 7.32 (m, 2H), 6.60 (d, 1H), 6.46 (s, 1H), 5.61 (quin, 1H), 4.13 (s, 3H), 3.04 - 2.95 (m, 1H), 2.44 (s, 3H), 1.76 (d, 3H), 1.29 (d, 6H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.1.

35: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.93 - 7.85 (m, 2H), 7.42 - 7.31 (m, 2H), 6.58 (d, 1H), 6.45 (s, 1H), 5.61 (quin, 1H), 4.13 (s, 3H), 3.05 - 2.94 (m, 1H), 2.44 (s, 3H), 1.76 (d, 3H), 1.29 (d, 6H). **LCMS** R_t = 1.27 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.1.

Example 23. Synthesis of 36 and 37




[000305] Analytical SFC (Daicel CHIRALCEL OJ-3 (100 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 4.5 min and hold 40% for 2.5 min, then 5% of B for 1 min, flow rate: 2.8 mL/min, column temp: 40 °C.) showed two peaks at 1.94 min and Rt = 2.38 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 50 mL/min; 20% B; 9 min run; 11 injections, Rt of peak 1 = 5.5 min, Rt of peak 2 = 7.6 min) to give the enantiomer 1, randomly assigned as **36** (16.17 mg, 44.4 µmol, 17% yield) (Rt = 1.94 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **37** (28.82 mg, 79.5 µmol, 30% yield) (Rt = 2.38 min in analytical SFC) as a solid.

36: ^{1}H NMR (400MHz, DMSO-*d*$_6$) δ_H = 9.38 (d, 1H), 7.85 - 7.77 (m, 2H), 7.48 - 7.39 (m, 2H), 7.27 (s, 1H), 7.06 (t, 1H), 5.44 (quin, 1H), 4.09 (s, 3H), 2.40 (s, 3H), 1.66 (d, 3H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.1, found 362.3.


37: ^{1}H NMR (400MHz, DMSO-*d*$_6$) δ_H = 9.39 (d, 1H), 7.84 - 7.78 (m, 2H), 7.48 - 7.39 (m, 2H), 7.27 (s, 1H), 7.06 (t, 1H), 5.44 (quin, 1H), 4.09 (s, 3H), 2.39 (s, 3H), 1.66 (d, 3H).

LCMS R_t = 1.23 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.1, found 362.1.

Example 24. Synthesis of 38 and 39



A-5 A-34



38 39

[000306] **A-34:** A mixture of 5-ethyl-2-methyl-pyrazole-3-carboxylic acid (69.46 mg, 0.45 mmol), HOBt (135.3 mg, 1 mmol), Et_3N (0.35 mL, 2.5 mmol), EDCI (143.95 mg, 0.75 mmol) and 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (120 mg, 0.5 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours under N_2. The reaction was quenched with the addition of sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Agela DuraShell (150 mm x 25 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 34-74% B over 8 min) to give the product (88 mg, 0.26 mmol, 52% yield) as a solid.


[000307] **38 & 39:** Analytical SFC (Column: Daicel CHIRALCEL IC-3 (150 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.42 min and Rt = 4.26 min. The product was separated by SFC (Daicel CHIRALCEL (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH; 38 °C; 50 mL/min; 30% B; 7 min run; 4 injections, Rt of peak 1 = 3.75 min, Rt of peak 2 = 5.4 min) to give the enantiomer 1, randomly assigned as **38** (24.43 mg, 0.07 mmol, 41% yield) (Rt = 3.42 min in analytical SFC) as oil and the enantiomer 2, randomly assigned as **39** (27.23 mg, 0.08 mmol, 45% yield) (Rt = 4.26 min in analytical SFC) as an oil.

38: ^{1}H NMR (400MHz, CD_3CN) δ_H = 7.87 (s, 1H), 7.83 (d, 1H), 7.49 (br d, 1H), 7.44 - 7.37 (m, 2H), 6.61 (s, 1H), 5.50 - 5.37 (m, 1H), 3.99 (s, 3H), 2.60 (q , 2H), 2.41 (s, 3H), 1.69 (d,

3H), 1.21 (t, 3H). **LCMS** R_t = 1.25 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{22}N_5O_2$ $[M+H]^+$ 340.2, found 340.1.

39: ^{1}H NMR (400MHz, CD_3CN) δ_H = 7.87 (s, 1H), 7.83 (d, 1H), 7.48 (br d, 1H), 7.44 - 7.37 (m, 2H), 6.61 (s, 1H), 5.49 - 5.39 (m, 1H), 3.99 (s, 3H), 2.60 (q, 2H), 2.41 (s, 3H), 1.69 (d, 3H), 1.21 (t, 3H). **LCMS** R_t = 1.25 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{22}N_5O_2$ $[M+H]^+$ 340.2, found 340.1.

Example 25. Synthesis of 40 and 41



[000308] **A-36:** To a mixture of Cs_2CO_3 (1310 mg, 4.02 mmol) and 2-bromopropane (494.61 mg, 4.02 mmol) in DMF (6 mL) was added ethyl 4-methyl-1H-pyrazole-5-carboxylate (310 mg, 2.01 mmol), and then the reaction mixture was stirred at 100 °C for 2 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (40 mL) and then extracted with EtOAc (30 mL x 2). The combined organic layer was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE= 0% to 5% to 20%) to give the product (190 mg, 0.97 mmol, 48% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.26 (s, 1H), 4.65 - 4.50 (m, 1H), 4.40 (q, *2*H), 2.28 (s, 3H), 1.50 (d, 6H), 1.40 (t, 3H).

[000309] **A-37:** To a solution of ethyl 1-isopropyl-4-methyl-pyrazole-3-carboxylate (190 mg, 0.97 mmol) in ethanol (9 mL) was added a solution of NaOH (116.18 mg, 2.9 mmol) in water (9 mL) and the mixture was stirred at 50 °C for 3 hours. After cooling to room temperature, the reaction mixture was concentrated to give a residue. The residue was diluted with H_2O (30 mL) and washed with EtOAc (20 mL x 1). The aqueous phase was acidified with 1N HCl (20 mL) to adjust the pH = 1 and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (70 mL), dried over Na_2SO_4, filtered and concentrated to give the product (130 mg, 0.76 mmol, 78% yield) as a solid. **LCMS** R_t =

0.65 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_8H_{13}N_2O_2$ $[M+H]^+$ 169.09, found 168.9.


[000310] **A-38:** To a mixture of 1-isopropyl-4-methyl-pyrazole-3-carboxylic acid (130 mg, 0.77 mmol), HATU (317.25 mg, 0.83 mmol), DIPEA (0.29 mL, 1.67 mmol) in DCM (8 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (100 mg, 0.42 mmol) and the mixture was stirred at 15 °C for 2 hours. The mixture was concentrated and diluted with H_2O (10 mL), then extracted with EtOAc (20 mL x 2). The combined organic layer was washed brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 59-89% B over 8 min) to give the product (150 mg, 412.9 µ mol, 99% yield) as an oil. **LCMS** R_t = 0.93 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.2.

[000311] **40 & 41:** Analytical SFC (Regis (S,S) Whelk-O1 (250 mm x 4.6 mm, 5 µm), mobile phase: A: CO_2 B:ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.46 min and 5.05 min. The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = methanol (0.1% DEA) ; 38 °C; 60 mL/min; 40% B; 12 min run; 4 injections, Rt of peak 1 = 6.2 min, Rt of peak 2 = 9.0 min) to give the enantiomer 1, randomly assigned as **41** (47.98 mg, 135.8 µmol, 32% yield) (Rt = 3.46 min in analytical SFC) as oil and enantiomer 2, randomly assigned as **40** (51.51 mg, 145.7 µmol, 34 % yield) (Rt = 5.05 min in analytical SFC) as a solid.

[000312] **41: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 8.66 (d, 1H), 7.84 - 7.75 (m, 2H), 7.68 (s, 1H), 7.47 - 7.37 (m, 2H), 5.39 (quin, 1H), 4.55 - 4.42 (m, 1H), 2.39 (s, 3H), 2.17 (s, 3H), 1.66 (d, 3H), 1.43 (d, 6H). **LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.0.

[000313] **40: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 8.66 (d, 1H), 7.85 - 7.76 (m, 2H), 7.68 (s, 1H), 7.48 - 7.37 (m, 2H), 5.39 (quin, 1H), 4.55 - 4.43 (m, 1H), 2.39 (s, 3H), 2.17 (s, 3H), 1.66 (d, 3H), 1.43 (d, 6H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.0.

Example 26. Synthesis of 42 and 43



[000314] **A-40:** A mixture of 2-(tert-butoxycarbonylamino)propanoic acid (296.58 mg, 1.57 mmol) and CDI (279.58 mg, 1.72 mmol) in DMF (30 mL) was stirred at 15 °C for 1 hour and then N'-hydroxy-3-(trifluoromethyl)benzamidine (320 mg, 1.57 mmol) was added. The reaction mixture was stirred at 110 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 10% to 30% to 50%) to give the product (500 mg, 1.24 mmol, 79% yield) as an oil. **LCMS** R_t = 0.95 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{11}F_3N_3O_3$ $[M\text{-}tBu+H]^+$ 302.1, found 302.1.

[000315] **A-41:** To a solution of tert-butyl N-[1-[3-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (240 mg, 0.67 mmol) in 1,4-dioxane (3 mL) was added 4M HCl/1,4-dioxane (20 mL) and the mixture was stirred at 20 °C for 8 hours. The mixture was concentrated to give the crude product (180 mg, 0.61 mmol, 79% yield) as a solid. **LCMS** R_t = 0.69 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{11}F_3N_3O$ $[M+H]^+$ 258.1, found 258.0.

[000316] **A-42:** A mixture of 5-isopropyl-2-methyl-pyrazole-3-carboxylic acid (54.41 mg, 0.32 mmol), HOBt (92.03 mg, 0.68 mmol), Et_3N (0.24 mL, 1.7 mmol), EDCI (97.92 mg, 0.51mmol) and 1-[3-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.34 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours under N_2. The reaction mixture was quenched with sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 20-80% B over 8 min) to give the product (85 mg, 0.21


mmol, 61% yield) as a an oil. **LCMS** R_t = 0.93 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_2$ $[M+H]^+$ 408.2, found 408.1.

[000317] **42 & 43:** Analytical SFC (Regis (R,R) Whelk-O1 (100 mm x 4.6 mm, 5 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.15 min and Rt = 3.68 min. The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH; 38 °C; 65 mL/min; 30% B; 7 min run; 6 injections, Rt of peak 1 = 4.43 min, Rt of peak 2 = 5.72 min) to give the enantiomer 1, randomly assigned as **42** (25.66 mg, 0.06 mmol, 31% yield) (Rt = 3.15 min in analytical SFC) as oil and enantiomer 2, randomly assigned as **43** (25.41 mg,0.06 mmol, 32% yield) (Rt = 3.68 min in analytical SFC) as an oil.

42: ^{1}H NMR (400MHz, CD_3CN) δ_H = 8.36 - 8.23 (m, 2H), 7.88 (d, 1H), 7.74 (t, 1H), 7.50 (d, 1H), 6.64 (s, 1H), 5.46 (quin, 1H), 3.99 (s, 3H), 3.00 - 2.85 (m, 1H), 1.71 (d, 3H), 1.23 (d, 6H). **LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_2$ $[M+H]^+$ 408.2, found 408.1.

43: ^{1}H NMR (400MHz, CD_3CN) δ_H = 8.34 - 8.25 (m, 2H), 7.88 (d, 1H), 7.74 (t, H), 7.50 (d1H), 6.65 (s, 1H), 5.46 (quin, 1H), 3.99 (s, 3H), 2.99 - 2.86 (m, 1H), 1.71 (d, 3H), 1.23 (d, 6H). **LCMS** R_t = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_2$ $[M+H]^+$ 408.2, found 408.1.


Example 27. Synthesis of 44



[000318] To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1R)-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (120.84 mg, 0.59 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05 % NH_4OH) and B = CH_3CN; 17-47 % B over 8 min) to give the product as an oil. Analytical SFC (Daicel CHIRALPAK AS-3 (150 mm × 4.6 mm, 3 μm), mobile phase: A: CO_2 B:ethanol (0.05% DEA), gradient:

from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C, ABPR: 1500 psi) showed 2.39 min (main peak, 91.6%) and 2.55 min (8.4%). Note: the condensation reaction leads to some racemization. Then the product was purified by SFC (Daicel CHIRALPAK AS-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 65 mL/min; 20% B; 8.60 min run; 50 injections, Rt of peak 1 = 5.57 min, Rt of peak 2 = 6.60 min) to give the product (66.75 mg, 0.19 mmol, 32% yield) as an oil. **^{1}H NMR** (400MHz, CD_3CN) δ = 8.00 (s, 1H), 7.89 (s, 1H), 7.86 (d, 1H), 7.48 - 7.38 (m, 2H), 7.06 (br d, 1H), 5.50 - 5.40 (m, 1H), 4.46 (quin, 1H), 2.44 (s, 3H), 2.39 (s, 3H), 1.69 (d, 3H), 1.48 (d, 6H). **LCMS** R_t = 1.13 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.2, found 354.0.

Example 28. Synthesis of 45



[000319] **A-10:** A mixture of 3-fluorobenzonitrile (2.2 g, 18.17 mmol), hydroxylamine hydrochloride (3.79 g, 54.5 mmol) and NaOH (2.18 g, 54.5 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours to give a mixture. After cooling to room temperature, the reaction mixture was concentrated to remove most of the EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (3200 mg) as a solid. **LCMS** R_t = 0.16 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8FN_2O$ $[M+1H]^+$ 155.05, found 155.1.


[000320] **A-45:** To a mixture of 2-(tert-butoxycarbonylamino)-3-methyl-butanoic acid (400 mg, 1.84 mmol) and CDI (328.39 mg, 2.03 mmol) in DMF (20 mL) was stirred at 20 °C for 1h before 3-fluoro-N-hydroxy-benzamidine (283.79 mg, 1.84 mmol) was added. The reaction mixture was stirred at 100 °C for 16 hours. After cooling to room temperature, the mixture was diluted with H_2O (10 mL) and extracted with EtOAc (20 mL x 2). The

combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (150 mg, 0.45 mmol, 24% yield) as an oil. **LCMS** R_t = 0.96 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{17}H_{23}FN_3O_3$ $[M+H\text{-}Boc]^+$ 280.1, found 280.1.

[000321] **A-46:** To a solution of of tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methyl-propyl]carbamate (200 mg, 0.60 mmol) in 1,4-dioxane (10 mL) was added 4M HCl/1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 20 °C for 2 hours. The mixture was concentrated, and the pH was adjusted with the addition of sat. $NaHCO_3$ to pH ~ 9. The mixture was extracted with EtOAc (40 mL x 2) and the combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (150 mg) as an oil. **LCMS** R_t = 0.696 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{14}FN_3O$ $[M+H]^+$ 236.11, found 236.1.

[000322] **45:** A mixture of 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methyl-propan-1-amine (150 mg, 0.64 mmol), HOBt (172.32 mg, 1.28 mmol), EDCI (183.34 mg, 0.96 mmol), TEA (0.44 mL, 3.19 mmol) and 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (105.96 mg, 0.64 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The mixture was diluted with H_2O (20mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 59-89% B over 8 min) to give the product (75.38 mg, 0.20 mmol, 31% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 (d, 1H), 7.79 (td, 1H), 7.52 - 7.42 (m, 1H), 7.23 (dt, 1H), 6.53 (br d, 1H), 6.33 (s, 1H), 5.44 (dd, 1H), 4.09 (s, 3H), 2.45 - 2.36 (m, 1H), 1.98 - 1.90 (m, 1H), 1.10 - 1.01 (m, 6H), 0.98 - 0.92 (m, 2H), 0.79 - 0.74 (m, 2H). **LCMS** R_t = 1.35 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. for $C_{20}H_{22}FN_5O_2$ $[M+H]^+$ 384.2, found 384.2.

Example 29. Synthesis of 46 and 47



33 46 47


[000323] Analytical SFC (Regis (S,S) Whelk-O1 (250 mm x 4.6 mm, 5 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold

40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 5.34 min and 6.12 min. The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = ethanol (0.1% DEA); 38 °C; 60 mL/min; 40% B; 8 min run; 5 injections, Rt of peak 1 = 4.6 min, Rt of peak 2 = 5.8 min to give the enantiomer 1, randomly assigned as **46** (25.02 mg, 69.2 µmol, 25% yield) (Rt = 5.34 min in analytical SFC) as an oil and the enantiomer 2, randomly assigned as **47** (28.95 mg, 80.1 µmol, 29% yield) (Rt = 6.12 min in analytical SFC) as an oil.

46: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.55 (d, 1H), 8.18 (t, 1H), 7.85 - 7.74 (m, 2H), 7.50 - 7.36 (m, 2H), 7.02 (s, 1H), 5.44 (quin, 1H), 2.39 (s, 3H), 2.28 (s, 3H), 1.66 (d, 3H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.14, found 362.1.

47: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.56 (d, 1H), 8.19 (t, 1H), 7.88 - 7.71 (m, 2H), 7.53 - 7.35 (m, 2H), 7.03 (s, 1H), 5.45 (quin, 1H), 2.40 (s, 3H), 2.29 (s, 3H), 1.67 (d, 3H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{18}F_2N_5O_2$ $[M+H]^+$ 362.14, found 362.0.

Example 30. Synthesis of 48




[000324] **A-10:** A mixture of 3-fluorobenzonitrile (2.2 g, 18.17 mmol), hydroxylamine hydrochloride (3.79 g, 54.5 mmol) and NaOH (2.18 g, 54.5 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours to give a mixture. After cooling to room temperature, the reaction mixture was concentrated to remove most of the EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (3200 mg) as a solid. **LCMS** R_t = 0.16 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8FN_2O$ $[M+1H]^+$ 155.05, found 155.1.

[000325] **A-47:** A mixture of 2-(tert-butoxycarbonylamino)butanoic acid (400 mg, 1.97 mmol) and CDI (351.04 mg, 2.16 mmol) in DMF (20 mL) was stirred at 20 °C for 1 hour before 3-fluoro-N-hydroxy-benzamidine (303.37 mg, 1.97 mmol) was added. The mixture was stirred at 100 °C for 16 hours. The mixture was cooled then diluted with H_2O (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (150 mg, 0.45 mmol, 23% yield) as an oil. **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{16}H_{21}FN_3O_3$ $[M+H\text{-}Boc]^+$ 266.1, found 266.1.

[000326] **A-48:** To a solution of tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propyl]carbamate (200 mg, 0.62 mmol) in 1,4-dioxane (10 mL) was added 4M HCl/1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 20 °C for 2 hours. The mixture was concentrated, and the pH was adjusted with the addition of sat. $NaHCO_3$ to pH ~ 9. The mixture was extracted with EtOAc (40 mL x 2) and the combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (150 mg) as an oil. **LCMS** R_t = 0.66 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{14}ClFN_3O$ $[M+H]^+$ 222.0, found 222.0.

[000327] **48:** A mixture of 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine (150 mg, 0.68 mmol), HOBt (183.24 mg, 1.36 mmol), EDCI (194.97 mg, 1.02 mmol), TEA (0.47 mL, 3.39 mmol) and 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (112.67 mg, 0.68 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The mixture was diluted with sat. NH_4Cl (10 mL) and extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 55-85% B over 8 min) to give the product (77.37 mg, 0.21 mmol, 31% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (dd, 1H), 7.82 - 7.76 (m, 1H), 7.51 - 7.44 (m, 1H), 7.26 - 7.19 (m, 1H), 6.50 (br d, 1H), 6.32 (s, 1H), 5.53 - 5.45 (m, 1H), 4.09 (s, 3H), 2.23 - 2.11 (m, 1H), 2.10 - 1.98 (m, 1H), 1.98 - 1.88 (m, 1H), 1.06 (t, 3H), 0.98 - 0.91 (m, 2H), 0.78 - 0.72 (m, 2H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$ 370.2, found 370.1.

Example 31. Synthesis of 49




[000328] **A-49:** A mixture of 2-(tert-butoxycarbonylamino)-3-phenyl-propanoic acid (400 mg, 1.51 mmol) and CDI (268.93 mg, 1.66 mmol) in DMF (20 mL) was stirred at 20°C for 1 hour before 3-fluoro-N-hydroxy-benzamidine (232.4 mg, 1.51 mmol) was added. The mixture was stirred at 100 °C for 16 hours. The mixture was cooled then diluted with H_2O (10 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (150 mg, 0.39 mmol, 26% yield,) as an oil. **LCMS** R_t = 0.98 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{21}H_{23}FN_3O_3$ $[M+2H-Boc]^+$ 328.1, found 328.1.

[000329] **A-50:** To tert-utyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-phenyl-ethyl]carbamate (200 mg, 0.52 mmol) in 1,4-dioxane (10 mL) was added 4M HCl/1,4-dioxane (10 mL, 40 mmol) add the reaction mixture was stirred at 20 °C for 2 hours. The mixture was concentrated, and the pH was adjusted with the addition of sat. $NaHCO_3$ to pH ~ 9. The mixture was extracted with EtOAc (40 mL x 2), and the combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (150 mg, 0.46 mmol, 88% yield) as oil. **LCMS** R_t = 0.73 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{16}H_{16}ClFN_3O$ $[M+H]^+$ 284.1, found 284.1.

[000330] **49:** A mixture of 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-phenyl-ethanamine (150 mg, 0.53 mmol), HOBt (143.1 mg, 1.06 mmol), EDCI (152.25 mg, 0.79 mmol), TEA (0.37 mL, 2.65 mmol) and 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (87.99 mg, 0.53 mmol) in DCM (20 mL) was stirred at 20 °C for 16 hours. The mixture was diluted with H_2O (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic

phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 62-92% B over 8 min) to give the product (42.55 mg, 0.10 mmol, 18% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.17 (d, 1H), 7.86 (d, 1H), 7.77 - 7.70 (m, 1H), 7.68 - 7.60 (m, 1H), 7.48 (dt, 1H), 7.35 - 7.26 (m, 4H), 7.24 - 7.18 (m, 1H), 6.60 (s, 1H), 5.62 - 5.50 (m, 1H), 3.84 (s, 3H), 3.49 - 3.42 (m, 1H), 3.38 - 3.34 (m, 1H), 1.90 - 1.81 (m, 1H), 0.90 - 0.84 (m, 2H), 0.64 - 0.58 (m, 2H). **LCMS** R_t = 1.37 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. for $C_{24}H_{23}FN_5O_2$ $[M+H]^+$ 432.2, found 432.1.

Example 32. Synthesis of 50




[000331] **A-51:** A mixture of 2-(tert-butoxycarbonylamino)-2-methyl-propanoic acid (580.16 mg, 2.85 mmol) and CDI (462.86 mg, 2.85 mmol) in DMF (6 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N-hydroxy-benzamidine (400 mg, 2.6 mmol) was added. The reaction mixture was then stirred at 100 °C for 2 hours. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 60%) to give the product (140 mg, 344.1 μmol, 13% yield) as a solid. **LCMS** Rt = 0.91 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{13}FN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 266.1, found 266.0.

[000332] **A-52:** To tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-1-methyl-ethyl]carbamate (140 mg, 0.44 mmol) in 1,4-dioxane (2 mL) was added 4M HCl in 1,4-dioxane (3 mL, 12 mmol) and the mixture was stirred at 15 °C for 1 hour. The mixture was concentrated to give the product of 2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-2-

amine hydrochloride (110 mg, 417. 8 μmol, 96% yield) as a solid. **LCMS** Rt = 0.64 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{15}H_{13}FN_3O$ $[M+H]^+$ 222.1, found 222.0.

[000333] **50:** To a mixture of 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (77.39 mg, 0.47 mmol), HATU (177.06 mg, 0.47 mmol) and 2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-2-amine hydrochloride (100 mg, 0.39 mmol) in DMF (6 mL) was added DIPEA (0.2 mL, 1.16 mmol) and the mixture was stirred at 20 °C for 2 hours. The mixture was diluted with water (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL x 2) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 52-82% B over 8 min) to give the product (116.91 mg, 0.32 mmol, 81% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 8.93 (s, 1H), 7.85 (d, 1H), 7.73 (d, 1H), 7.67 - 7.59 (m, 1H), 7.50 - 7.40 (m, 1H), 6.72 (s, 1H), 3.83 (s, 3H), 1.93 - 1.84 (m, 1H), 1.75 (s, 6H), 0.92 - 0.86 (m, 2H), 0.68 - 0.60 (m, 2H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$370.2, found 369.9.

Example 33. Synthesis of 51




[000334] **A-53:** A mixture of 2-(tert-butoxycarbonylamino)-2-phenyl-acetic acid (717.29 mg, 2.85 mmol) and CDI (462.86 mg, 2.85 mmol) in DMF (6 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N-hydroxy-benzamidine (400 mg, 2.6 mmol) was added. The reaction was then stirred at 100 °C for 2 hours to give a mixture. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered

and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 60%) to give the product (140 mg, 209.0 μmol, 8% yield) as a solid. **LCMS** Rt = 0.96 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{16}H_{13}FN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 314.1, found 314.1.

[000335] **A-54:** To tert-butyl N-[[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-phenyl-methyl]carbamate (140 mg, 0.38 mmol) in 1,4-dioxane (6 mL) was added 4M HCl in 1,4-dioxane (3 mL, 12 mmole) and the mixture was stirred at 15 °C for 1 hour. The mixture was concentrated to give the product (110 mg, 351.4 μmol, 93% yield) as a solid. **LCMS** Rt = 0.71min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{15}H_{13}FN_3O$ $[M+H]^+$ 270.1, found 270.0.

[000336] **51:** To a mixture of 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (91.32 mg, 0.55 mmol), HATU (208.94 mg, 0.55 mmol) and [3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-phenyl-methanamine hydrochloride (140 mg, 0.46 mmol) in DMF (3 mL) was added DIPEA (0.24 mL, 1.37 mmol) and the mixture was stirred at 20 °C for 2 hours. The mixture was diluted with water (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL x 2) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by rep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 60-90% B over 8 min) to give the product (88.39 mg, 0.21 mmol, 46% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.61 (d, 1H), 7.84 (d, 1H), 7.77 - 7.70 (m, 1H), 7.67 - 7.58 (m, 1H), 7.56 - 7.50 (m, 2H), 7.49 - 7.38 (m, 4H), 6.78 (s, 1H), 6.64 (d, 1H), 3.94 (s, 3H), 1.92 - 1.82 (m, 1H), 0.91 - 0.82 (m, 2H), 0.66 - 0.56 (m, 2H). **LCMS** R_t = 1.41 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{23}H_{21}FN_5O_2$ $[M+H]^+$418.2, found 418.0.

Example 34. Synthesis of 52




[000337] **A-57b:** To a mixture of Cs_2CO_3 (8.45 g, 25.95 mmol) and 2-bromopropane (3.19 g, 25.95 mmol) in DMF (30 mL) was added ethyl 3-methyl-1H-pyrazole-4-carboxylate (2 g, 12.97 mmol) and the reaction mixture was stirred at 100 °C for 2.5 hours. After cooling to room temperature, the reaction mixture was diluted with sat. NH_4Cl (50 mL), then extracted wtih EtOAc (50 mL x 2), the organic layer was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30%) to give the product (300 mg, 1.53 mmol, 12% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.85 (s, 1H), 4.51 - 4.34 (m, 1H), 4.28 (q, 2H), 2.46 (s, 3H), 1.50 (d, 6H), 1.34 (t, 3H). **LCMS** Rt = 3.25 min in 7.0 min chromatography, 0-60AB, MS ESI calcd. $C_{10}H_{17}N_2O_2$ $[M+H]^+$ 197.1, found 197.0.

[000338] **A-57:** To a solution of ethyl 1-isopropyl-3-methyl-pyrazole-4-carboxylate (300 mg, 1.53 mmol) in ethanol (5 mL) was slowly added a solution of NaOH (122.29 mg, 3.06 mmol) in water (5 mL). The resulting mixture was stirred at 25 °C for 2 hours. The mixture was concentrated under reduced pressure to remove the EtOH. To the aqueous phase was added 1N HCl (30 mL) to adjust the pH = 2 and the mixture was diluted with EtOAc (10 mL). The phases were separated, and the organic phase was washed with brine (10 mL), dried over Na_2SO_4 then concentrated to give the crude product (210 mg, 1.22 mmol, 80% yield) as a solid. **LCMS** Rt = 0.61 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_8H_{13}N_2O_2$ $[M+H]^+$ 169.1, found 168.8.

[000339] **52:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (120.84 mg, 0.59 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was concentrated, and the residue was diluted with H_2O (20 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 40-60% B over 9 min) to give the product. Analytical SFC (Column: Daicel CHIRALPAK AS-3 150 mm × 4.6 mm I.D., 3 μm, mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C, ABPR: 1500 psi) showed two peaks at 2.52 min (11.1%) and 2.73 min (main peak, 88.9%). Note: the condensation reaction leads to some racemization. The product was separated by SFC (Daicel CHIRALPAK AS-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38

°C; 60 mL/min; 20% B; 8 min run; 10 injections, Rt of peak 1 = 4.8 min, Rt of peak 2 = 6 min) to give the product (50.77 mg, 0.14 mmol, 24% yield) (Rt = 2.55 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, CD_3CN) δ_H = 7.96 (s, 1H), 7.86 (s, 1H), 7.83 (br d, 1H), 7.44 - 7.36 (m, 2H), 7.04 (br d, 1H), 5.46 - 5.37 (m, 1H), 4.47 - 4.36 (m, 1H), 2.41 (s, 3H), 2.36 (s, 3H), 1.66 (d, 3H), 1.44 (d, 6H). **LCMS** Rt = 1.20 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.2, found 354.1.


Example 35. Synthesis of 55



[000340] **A-59:** To a solution of ethyl 2-isopropyl-4-methyl-pyrazole-3-carboxylate (220 mg, 1.12 mmol) in ethanol (5 mL) was added a solution of NaOH (134.52 mg, 3.36 mmol) in water (5 mL). The mixture was stirred at 20 °C for 2 hours. The reaction mixture was concentrated to give a residue and the residue was diluted with H_2O (30 mL) and extracted with EtOAc (20 mL x 1). To the aqueous phase was added 1N HCl (20 mL) to adjust pH = 1 and it was diluted with H_2O (10 mL). The mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (70 mL), dried over Na_2SO_4, filtered and concentrated to give the product (130 mg) as a solid. **LCMS**R_t = 0.70 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_{13}N_2O_2$ $[M+H]^+$ 169.09, found 169.0.

[000341] **55:** To a mixture of 2-isopropyl-4-methyl-pyrazole-3-carboxylic acid (129.45 mg, 0.77 mmol), HATU (315.92 mg, 0.83 mmol), DIPEA (0.29 mL, 1.66 mmol) in DCM (8 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.42 mmol) and the reaction mixture was stirred at 20 °C for 12 hours. The mixture was concentrated and diluted with H_2O (10 mL) then extracted with EtOAc (20 mL x 2). The combined organic layer was washed with brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 44-74% B over 9 min) to give the product (75.42 mg, 213.4 μmol, 51% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ_H = 9.17 (d, 1H), 7.85 - 7.76 (m, 2H), 7.49 - 7.39 (m, 2H), 7.33 (s, 1H), 5.42 (quin, 1H), 4.80 (spt, 1H), 2.39 (s, 3H), 2.15 (s, 3H), 1.65 (d, 3H), 1.36 (d, 6H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.19, found 354.0.

Example 36. Synthesis of 56 and 57



[000342] **A-60b:** To a solution of ethyl 5-methyl-2,4-dioxo-hexanoate (2 g, 10.74 mmol) in ethanol (20 mL) at 5 °C was added methylhydrazine (1.36 g, 11.81 mmol, 40% in H_2O) in a dropwise manner. The reaction mixture was stirred at 5 °C for 3 hours and then was concentrated. The residue was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 50%) to give the product (700 mg, 3.57 mmol, 33% yield) as an oil. **^{1}H NMR**(400MHz, $CDCl_3$) δ_H = 6.70 - 6.59 (m, 1H), 4.43 - 4.24 (m, 2H), 4.12 (s, 3H), 3.07 - 2.77 (m, 1H), 1.38 (t, 3H), 1.26 (d, 6H).

[000343] **A-60:** A mixture of ethyl 5-isopropyl-2-methyl-pyrazole-3-carboxylate (700 mg, 3.57 mmol) and NaOH (428.03 mg, 10.70 mmol) in ethanol (10 mL) and water (10 mL) was stirred at 20 °C for 2 hours. The reaction mixture was concentrated under reduced pressure to remove the EtOH and then extracted with EtOAc (10 mL). The aqueous phase was acidified with HCl (1N) to pH ~ 2 and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (400 mg) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 13.17 (br s, 1H), 6.62 (s, 1H), 3.99 (s, 3H), 2.91 - 2.80 (m, 1H), 1.17 (d, 6H).

[000344] **A-61:** A mixture of 2-(tert-butoxycarbonylamino)propanoic acid (1227.52 mg, 6.49 mmol) and CDI (1157.16 mg, 7.14 mmol) in DMF (20 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N'-hydroxy-benzamidine (1000 mg, 6.49 mmol) was added. The reaction mixture was then stirred at 110 °C for 16 hours. The mixture was cooled then diluted with NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (600 mg, 1.95 mmol, 30% yield) as a solid.

^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.89 - 7.78 (m, 2H), 7.76 - 7.70 (m, 1H), 7.68 - 7.60 (m, 1H), 7.50 - 7.43 (m, 1H), 5.02 - 4.93 (m, 1H), 1.51 (d, 3H), 1.40 (s, 9H).

[000345] **A-62:** To a mixture of tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (200 mg, 0.65 mmol) in 1,4-dioxane (5 mL) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol). The reaction mixture was stirred at 20 °C for 16 hours. The reaction mixture was concentrated to give the crude product (200 mg) as a an oil. **LCMS** R_t = 0.61 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{11}FN_3O$ [M+H] 208.1, found 207.7.

[000346] **A-63:** A mixture of 5-isopropyl-2-methyl-pyrazole-3-carboxylic acid (120 mg, 0.71 mmol), DIPEA (0.62 mL, 3.57 mmol), HOBt (192.82 mg, 1.43 mmol) EDCI (205.16 mg, 1.07 mmol) and 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (173.85 mg, 0.71 mmol) in DCM (15 mL) was stirred at 20 °C for 16 hours. The mixture was diluted with NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 24-54% B over 8 min) to give the product (180 mg, 0.50 mmol, 70% yield) as an oil. **LCMS** R_t = 0.88 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{18}H_{21}FN_5O_2$ $[M+H]^+$ 358.2, found 358.1.


[000347] **56 & 57:** Analytical SFC (Daicel CHIRALPAK IC-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B:IPA (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 3.55 min and 4.32 min. The product was separated by SFC (Daicel CHIRALPAK IC (250 mm x 30 mm, 5 µm); A = CO_2 and B = *i*-PrOH; 38 °C; 65 mL/min; 30% B; 7 min run; 7 injections, Rt of peak 1 = 3.5 min, Rt of peak 2 = 4.7 min) to give the enantiomer 1, randomly assigned as **56** (75.09 mg, 0.21 mmol, 41% yield) (Rt = 3.55 min in analytical SFC) as an oil and enantiomer 2, randomly assigned as **57** (73.93 mg, 0.21 mmol, 41% yield) (Rt = 4.32 min in analytical SFC) as an oil.


56: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.91 - 7.86 (m, 1H), 7.82 - 7.76 (m, 1H), 7.51 - 7.44 (m, 1H), 7.26 - 7.19 (m, 1H), 6.56 (br d, 1H), 6.45 (s, 1H), 5.65 - 5.56 (m, 1H), 4.12 (s, 3H), 3.06 - 2.94 (m, 1H), 1.76 (d, 3H), 1.29 (d, 6H). **LCMS** R_t = 1.26 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{21}FN_5O_2$ $[M+H]^+$ 358.2, found 358.1.

57: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.91 - 7.86 (m, 1H), 7.82 - 7.75 (m, 1H), 7.52 - 7.44 (m, 1H), 7.26 - 7.19 (m, 1H), 6.56 (br d, 1H), 6.45 (s, 1H), 5.66 - 5.56 (m, 1H), 4.12 (s, 3H),

3.05 - 2.95 (m, 1H), 1.76 (d, 3H), 1.29 (d, 6H). **LCMS** R_t = 1.26 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{21}FN_5O_2$ $[M+H]^+$ 358.2, found 358.1.

Example 37. Synthesis of 58






[000348] **A-64:** A mixture of 3-benzyloxy-2-(tert-butoxycarbonylamino)propanoic acid (843.03 mg, 2.85 mmol) and CDI (462.86 mg, 2.85 mmol) in DMF (6 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N-hydroxy-benzamidine (400 mg, 2.6 mmol) was added. The reaction mixture was stirred at 110 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 60%) to give the product (260 mg, 0.58 mmol, 22% yield) as an oil. **LCMS** R_t = 1.45 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{22}H_{25}FN_3O_4$ $[M+H\text{-}tBu]^+$ 358.1, found 358.1.

[000349] **A-65:** A mixture of *tert*-butyl N-[2-benzyloxy-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl] ethyl]carbamate (260 mg, 0.63 mmol) and 4M HCl in 1,4-dioxane (5 mL, 20 mmol) was stirred at 25 °C for 16 hours. The mixture was concentrated to give the product (300 mg, 0.59 mmol, 94% yield) as a solid. **LCMS** R_t = 0.99 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}FN_3O_2$ $[M+H]^+$ 314.1, found 314.0.

[000350] **58:** To a mixture of 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (171.03 mg, 1.03 mmol), HATU (391.33 mg, 1.03 mmol) and DIPEA (0.45 mL, 2.57 mmol) in DMF (3 mL) was added 2-benzyloxy-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (300 mg, 0.86 mmol) and the mixture was stirred at 25 °C for 16 hours. The mixture was diluted with water (20 mL) and extracted with EtOAc (20 mL x

3). The combined organic phase was washed with water (20 mL x 2) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05 % NH_4OH) and B = CH_3CN; 55-85% B over 9 min) to give the product (49.27 mg, 0.11 mmol, 12% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ = 9.16 (d, 1H), 7.86 (d, 1H), 7.74 (m, 1H), 7.64 (m, 1H), 7.48 (m, 1H), 7.37 - 7.21 (m, 5H), 6.69 (s, 1H), 5.61 (q, 1H), 4.59 (s, 2H), 4.02 (m, 2H), 3.93 (s, 3H), 1.97 - 1.76 (m, 1H), 0.95 - 0.84 (m, 2H), 0.69 - 0.56 (m, 2H).**LCMS** R_t = 1.31 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{25}H_{25}FN_5O_3$ $[M+H]^+$ 462.1, found 462.1.


Example 38. Synthesis of 54



58 54

[000351] To a mixture of N-[2-benzyloxy-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-cyclopropyl-2-methyl-pyrazole-3-carboxamide (70 mg, 0.15 mmol) in DCM (3 mL) was added BBr_3 (0.03 mL, 0.30 mmol) and the mixture was stirred at 20 °C for 2 hours. The reaction mixture was diluted with H_2O (5 mL) and extracted with DCM (5 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH3CN; 39-69 % B over 9 min) to give the product (21.67 mg, 0.06 mmol, 38% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 9.01 (d, 1H), 7.87 (d, 1H), 7.80 - 7.73 (m, 1H), 7.65 (m, 1H), 7.48 (m, 1H), 6.73 (s, 1H), 5.26 - 5.44 (m, 2H), 3.88 - 4.05 (m, 5H), 1.81 - 1.99 (m, 1H), 0.79 - 0.99 (m, 2H), 0.53 - 0.74 (m, 2H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_3$ $[M+H]^+$ 372.1, found 372.1.

Example 39. Synthesis of 59




A-66 A-67 59

A-67: To a mixture of ethyl 1-isopropyl-5-methyl-pyrazole-4-carboxylate (650 mg, 3.31 mmol) in ethanol (5 mL) and water (5 mL) was added NaOH (264.97 mg, 6.62 mmol), and

the mixture was stirred at 50 °C for 3 hours. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the ethanol then the residue was diluted with H_2O (10 mL) and the mixture was washed with EtOAc (5 mL x 2). The pH of the aqueous phase was adjusted to pH~2 with 1N HCl, then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with H_2O (10 mL) and brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the product (520 mg, 3.09 mmol, 93% yield) as a solid. **LCMS** R_t = 0.63 min in 1.5 min chromatography, 10-80AB, MS ESI calcd. for $C_8H_{13}N_2O_2$ $[M+H]^+$ 169.1, found 169.0.

[000352] **59:** A mixture of 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (241.68 mg, 1.19 mmol), HOBt (321.37 mg, 2.38 mmol), DIPEA (0.62 mL, 3.57 mmol), EDCI (455.91 mg, 2.38 mmol) and 1-isopropyl-5-methyl-pyrazole-4-carboxylic acid (200 mg, 1.19 mmol) in DCM (10mL) was stirred at 20 °C for 16 hours The mixture was concentrated to give a residue. The residue was dissolved in EtOAc (20 mL) and washed with sat. NH_4Cl (10 mL), water (10 mL) and brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by flash chromatography on silica gel (EtOAc in PE =0% to 40% to 100%) to give the impure product. The impure product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 50-70% B over 9 min) to give the product (17.06 mg, 48.1 mol, 4% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.93 - 7.85 (m, 2H), 7.77 (s, 1H), 7.41 - 7.30 (m, 2H), 6.38 (br d, 1H), 5.64 (quin, 1H), 4.50 (spt, 1H), 2.59 (s, 3H), 2.43 (s, 3H), 1.74 (d, 3H), 1.50 (d, 6H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.2, found 354.2.

Example 40. Synthesis of 60 and 61



48 60 61


[000353] Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm × 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.00 min and Rt = 3.93 min). The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = ethanol (0.1% DEA); 38 °C; 60 mL/min; 20% B; 7 min run; 5 injections, Rt of peak 1 = 5.2 min, Rt of peak 2 = 5.4 min) to give the enantiomer 1, randomly assigned as **60** (16.59 mg, 44.9

μmol, 24% yield) (Rt = 3.00 min in analytical SFC) as oil and the enantiomer 2, randomly assigned as **61** (26.44 mg, 71.6 μmol, 38% yield) (Rt = 3.93 min in analytical SFC) as an oil.

60: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.84 - 7.75 (m, 1H), 7.50 - 7.42 (m, 1H), 7.26 - 7.20 (m, 1H), 6.50 (br d, 1H), 6.32 (s, 1H), 5.53 - 5.46 (m, 1H), 4.09 (s, 3H), 2.23 - 2.11 (m, 1H), 2.10 - 1.98 (m, 1H), 1.98 - 1.89 (m, 1H), 1.06 (t, 3H), 0.99 - 0.92 (m, 2H), 0.78 - 0.73 (m, 2H). **LCMS** R_t = 1.23 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$ 369.16, found 370.0.

61: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.83 - 7.74 (m, 1H), 7.51 - 7.41 (m, 1H), 7.27 - 7.20 (m, 1H), 6.50 (br d, 1H), 6.32 (s, 1H), 5.53 - 5.45 (m, 1H), 4.09 (s, 3H), 2.22 - 2.11 (m, 1H), 2.11 - 1.98 (m, 1H), 1.98 - 1.87 (m, 1H), 1.06 (t, 3H), 0.98 - 0.91 (m, 2H), 0.78 - 0.73 (m, 2H). **LCMS** R_t = 1.27 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$ 369.16, found 370.1.

Example 41. Synthesis of 62 and 63




[000354] Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA) gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C ABPR: 1500 psi) showed two peaks at 2.60 min and Rt = 3.27 min. The product was purified by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 15% B; 10 min run; 5 injections, Rt of peak 1 = 5.7 min, Rt of peak 2 = 8.5 min) to give the enantiomer 1, randomly assigned as **62** (22.57 mg, 58.9 μmol, 33% yield) (Rt = 2.60 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **63** (25.47 mg, 66.2 μmol, 37% yield) (Rt = 3.27 min in analytical SFC) as a solid.

62: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.02 (d, 1H), 7.86 (d, 1H), 7.74 (br d, 1H), 7.69 - 7.60 (m, 1H), 7.47 (dt, 1H), 6.75 (s, 1H), 5.13 (t, 1H), 3.91 (s, 3H), 2.46 - 2.36 (m, 1H), 1.92 - 1.84 (m, 1H), 1.05 (d, 3H), 0.95 - 0.85 (m, 5H), 0.66 - 0.60 (m, 2H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{23}FN_5O_2$ $[M+H]^+$ 384.2, found 384.2.

63: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.02 (d, 1H), 7.86 (d, 1H), 7.74 (br d, 1H), 7.70 - 7.60 (m, 1H), 7.47 (dt, 1H), 6.75 (s, 1H), 5.13 (t, 1H), 3.91 (s, 3H), 2.46 - 2.36 (m, 1H), 1.92 - 1.84 (m, 1H), 1.05 (d, 3H), 0.96 - 0.85 (m, 5H), 0.66 - 0.59 (m, 2H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{23}FN_5O_2$ $[M+H]^+$ 384.2, found 384.2.

Example 42. Synthesis of 64 and 65



[000355] **A-71b:** To a solution of ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (500 mg, 2.4 mmol), cyclopropylboronic acid (412.7 mg, 4.8 mmol) and Na_2CO_3 (509.22 mg, 4.8 mmol) in DCE (7 mL) was added a solution of $Cu(OAc)_2$ (436.32 mg, 2.4 mmol) and 2,2-bipyridine (450.25 mg, 2.88 mmol) in DCE (14 mL). The reaction mixture was stirred at 70 °C for 4 hours. After cooling to room temperature, the mixture was diluted with sat. NH_4Cl (20 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5% to 50%) to give the product (180 mg, 0.73 mmol, 30% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.07 (s, 1H), 4.43 - 4.36 (m, 3H), 1.41 (t, 3H), 1.36 - 1.30 (m, 2H), 1.14 - 1.07 (m, 2H).

[000356] **A-71:** To a solution of ethyl 2-cyclopropyl-5-(trifluoromethyl)pyrazole-3-carboxylate (180 mg, 0.73 mmol) in ethanol (3 mL) was added a solution of NaOH (87.03 mg, 2.18 mmol) in water (3 mL). The reaction mixture was stirred at 50 °C for 2 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (20 mL) and washed with EtOAc (20 mL x 1). The pH of the aqueous phase was adjusted to pH = 1 by adding 1N HCl (20 mL) and then diluted with H_2O (10 mL). The mixture was extracted with

EtOAc (50 mL x 2). The combined organic phase was washed with brine (70 mL), dried over Na_2SO_4, filtered and concentrated to give the product (160 mg, 661.7 μmol, 84% yield) as an oil. **LCMS** R_t = 0.79 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_8H_8F_3N_2O_2$ $[M+H]^+$ 221.05, found 221.1.

[000357] **A-12:** A mixture of 3-chlorobenzonitrile (1.2 g, 8.72 mmol), NH_2OH HCl (1818.45 mg, 26.17 mmol) and NaOH (1046.74 mg, 26.17 mmol) in ethanol (9 mL) and water (3 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the EtOH and then diluted with H_2O (100 mL). The mixture was extracted with EtOAc (150 mL x 2). The combined organic phase was washed with brine (100 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1760 mg, 6.62 mmol, 76% yield) as a solid. **LCMS** R_t = 0.21 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8ClN_2O$ $[M+H]^+$ 171.02, found 170.9.

[000358] **A-68:** A mixture of 2-(tert-butoxycarbonylamino)propanoic acid (1.24 g, 6.54 mmol) and CDI (1.17 g, 7.19 mmol) in DMF (60 mL) was stirred at 15 °C for 1 hour and then 3-chloro-N'-hydroxy-benzamidine (1.74g, 6.54mmol) was added. The reaction mixture was then stirred at 100 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (100 mL) and extracted with EtOAc (50 mL x 3). The combined organic phase was washed with brine (100 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (1480 mg, 4.29 mmol, 66% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.09 (s, 1H), 7.97 (d, 1H), 7.52 - 7.46 (m, 1H), 7.45 - 7.39 (m, 1H), 5.26 - 5.05 (m, 2H), 1.64 (br d, 3H), 1.47 (s, 9H). **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{15}H_{19}ClN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 268.1, found 268.1.

[000359] **A-69:** To tert-butyl N-[1-[3-(3-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (1480 mg, 4.57 mmol)was added 4 N HCl/1,4-dioxane (10 mL, 40 mmol) and the reaction micture was stirred at 20 °C for 0.5 hours. The mixture was concentrated to give the crude product (1160 mg, 4.30 mmol, 94% yield) as a solid. **LCMS** R_t = 0.66 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{11}ClN_3O$ $[M+H]^+$ 224.05, found 224.0.

[000360] **A-70:** To a mixture of 2-cyclopropyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (160 mg, 0.73 mmol), HATU (292.35 mg, 0.77 mmol), DIPEA (0.27 mL, 1.54 mmol) in DCM (8 mL) was added 1-[3-(3-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine


hydrochloride (100 mg, 0.38 mmol) and the mixture was stirred at 15 °C for 2 hours. The mixture was concentrated and diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The organic layer was washed brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 59-89% B over 8 min) to give the product (140 mg, 322.6 μmol, 84% yield) as a solid. **LCMS** R_t = 0.98 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{18}H_{16}ClF_3N_5O_2$ $[M+H]^+$ 426.19, found 426.1.


[000361] **64 & 65:** Analytical SFC (column: Regis (S,S) Whelk-O1 (100 mm × 4.6 mm I.D., 5.0 μm), mobile phase: A: CO_2 B: IPA (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column tem: 35 °C) showed two peaks at 3.78 min and 4.30 min. The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = IPA (0.1% DEA); 38 °C; 60 mL/min; 35% B; 9 min run; 8 injections, Rt of peak 1 = 4.7 min, Rt of peak 2 = 6.8 min) to give the enantiomer 1, randomly assigned as (54.51 mg, 128 μmol, 39% yield) (Rt = 3.78 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as (55.53 mg, 130.4 μmol, 39% yield) (Rt = 4.30 min in analytical SFC) as a solid.

65: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 8.08 (t, 1H), 7.97 (td, 1H), 7.54 - 7.48 (m, 1H), 7.48 - 7.41 (m, 1H), 6.91 (s, 1H), 6.69 (br d, 1H), 5.63 (quin, 1H), 4.39 (tt, 1H), 1.79 (d, 3H), 1.40 - 1.28 (m, 2H), 1.16 - 1.05 (m, 2H). **LCMS** Rt = 1.43 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}ClF_3N_5O_2$ $[M+H]^+$ 426.09, found 426.0.

64: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 8.08 (t, 1H), 7.97 (td, 1H), 7.54 - 7.49 (m, 1H), 7.47 - 7.41 (m, 1H), 6.91 (s, 1H), 6.70 (br d, 1H), 5.63 (quin, 1H), 4.39 (tt, 1H), 1.79 (d, 3H), 1.39 - 1.29 (m, 2H), 1.16 - 1.05 (m, 2H). **LCMS** Rt = 1.41 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}ClF_3N_5O_2$ $[M+H]^+$ 426.09, found 426.0.

Example 43. Synthesis of 66






[000362] **A-72:** A mixture of 1-(tert-butoxycarbonylamino)cyclohexanecarboxylic acid (0.79 g, 3.24 mmol) and CDI (0.58 g, 3.57 mmol) in DMF (10mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N-hydroxy-benzamidine (0.5 g, 3.24 mmol) was added. The reaction mixture was then stirred at 100 °C for 16 hours. After cooling to room temperature, the mixture was diluted with sat. NH_4Cl (30 mL) and the mixture was extracted with EtOAc (20 mL x 3). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (260 mg, 0.66 mmol, 20% yield) as a solid. **LCMS** R_t = 0.98 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{19}H_{25}FN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 306.2, found 306.2.

[000363] **A-73:** To tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]cyclohexyl]carbamate (260 mg, 0.72 mmol) was added 4 M HCl in 1,4-dioxane (3 mL, 12 mmol) and the reaction mixture was stirred at 20 °C for 0.5 hours. The mixture was concentrated to give the crude product (210 mg, 705.3 µmol, 98% yield) as a solid. **LCMS** R_t = 0.71 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{14}H_{17}FN_3O$ $[M+H]^+$ 262.13, found 262.2.

[000364] **66:** To a mixture of HATU (254.54 mg, 0.67 mmol), 5-cyclopropyl-2-methyl-pyrazole-3-carboxylic acid (55.62 mg, 0.33 mmol), 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]cyclohexanamine hydrochloride (100 mg, 0.33 mmol) in DCM (8 mL) was added DIPEA (0.23 mL, 1.34 mmol) and the mixture was stirred at 25 °C for 3.5 hours. The mixture was diluted with H_2O (20 mL) and extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (20 mL) and dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 x 25 mm, 5 µm), A = H_2O (0.04% NH_4OH + 10 mM NH_4HCO_3) and B = CH_3CN; 60-

90% B over 7.5 min) to give the product (26.87 mg, 65.6 µmol, 20% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 8.68 (s, 1H), 7.85 (d, 1H), 7.76 - 7.70 (m, 1H), 7.67 - 7.58 (m, 1H), 7.50 - 7.41 (m, 1H), 6.73 (s, 1H), 3.81 (s, 3H), 2.33 (br d, 2H), 2.15 - 2.01 (m, 2H), 1.94 - 1.80 (m, 1H), 1.74 - 1.51 (m, 5H), 1.49 - 1.35 (m, 1H), 0.93 - 0.83 (m, 2H), 0.71 - 0.58 (m, 2H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{22}H_{25}FN_5O_2$ $[M+H]^+$ 410.19, found 410.0.

Example 44. Synthesis of 67



A-74 A-75 A-76 A-77 67


[000365] **A-75:** A mixture of 3,4-difluorobenzonitrile (2 g, 14.38 mmol), hydroxylamine hydrochloride (3 g, 43.13 mmol) and NaOH (1.73 g, 43.13 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the ethanol and then it was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.5 g, 7.20 mmol, 50% yield) as a solid. **LCMS** R_t = 0.8 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_7F_2N_2O$ $[M+H]^+$ 173.0, found 173.0.

[000366] **A-76:** A mixture of 2-(tert-butoxycarbonylamino) propanoic acid (549.61 mg, 2.9 mmol) and CDI (518.11 mg, 3.2 mmol) in DMF (10 mL) was stirred at 15 °C for 1 hour and then 3,4-difluoro-N'-hydroxy-benzamidine (500 mg, 2.9 mmol) was added. The reaction mixture was then stirred at 110 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (10 mL) and extracted with EtOAc (10 mL x 3). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 % to 50%) to give the product (230 mg, 0.58 mmol, 20% yield) as an oil. **LCMS** R_t = 0.94 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{11}H_{10}F_2N_3O_3$ $[M+H\text{-}tBu]^+$ 270.1, found 270.0.

[000367] **A-77:** To tert-butyl N-[1-[3-(3,4-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (230 mg, 0.71 mmol) was added 4M HCl in 1,4-dioxane (6.91 mL, 27.63 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated, diluted with H_2O (20 mL) and the pH adjusted with addition of solid $NaHCO_3$ to pH~9. The solution was extracted with EtOAc (20 mL x 3). The combined organic phase was concentrated to give the crude product (300 mg, 0.99 mmol) as a solid.

[000368] **67:** A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (300 mg, 1.55 mmol) in DCM (15mL), DIPEA (0.86 mL, 6.18 mmol), HOBt (417.69 mg, 3.09 mmol) and EDCI (592.55 mg, 3.09 mmol) was stirred at 25 °C for 30 minutes and then 1-[3-(3,4-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (348.03 mg, 1.55 mmol) was added, and the mixture was stirred at 25 °C for 16 hours. The reaction mixture was washed with H_2O (20 mL), then extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Column (Boston Prime C18 (150 mm × 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 63-75% B over 9 min) to give the product (15.72 mg, 39.2 μmol, 3% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ = 9.45 (d, 1H), 8.12 - 7.94 (m, 1H), 7.88 (m, 1H), 7.72 - 7.60 (m, 1H), 7.44 (s, 1H), 5.45 (quin, 1H), 4.13 (s, 3H), 1.67 (d, 3H). **LCMS** R_t = 1.32 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{12}F_5N_5O_2$ $[M+H]^+$ 402.1, found 402.0.

Example 45. Synthesis of 68



[000369] **A-79:** A mixture of 5-methylpyridine-3-carbonitrile (1.9 g, 16.08 mmol), hydroxylamine hydrochloride (3.35 g, 48.25 mmol) and NaOH (1.93 g, 48.25 mmol) in ethanol (5 mL) and water (0.50 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the EtOH and then diluted with H_2O (40 mL). The mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (2.1 g, 12.79 mmol, 79% yield) as a solid.


LCMS R_t = 1.02 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_9N_3O$ $[M+H]^+$ 152.1, found 152.0.

[000370] **A-80:** A mixture of 2-(tert-butoxycarbonylamino) propanoic acid (1251.64 mg, 6.62 mmol) and CDI (1179.9 mg, 7.28 mmol) in DMF (10 mL) was stirred at 15 °C for 1 hour and then N'-hydroxy-5-methyl-pyridine-3-carboxamidine (1 g, 6.62 mmol) was added. The reaction mixture was stirred at 110 °C for 16 hours. The mixture was diluted with H_2O (50 mL) and extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 % to 50 %) to give the product (300 mg, 0.55 mmol, 8% yield) as a solid. **LCMS** R_t = 0.79 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{15}H_20N_4O_3$ $[M+H]^+$ 305.2, found 304.9.

[000371] **A-81:** A mixture of tert-butyl N-[1-[3-(5-methyl-3-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (300 mg, 0.99 mmol) and 4M HCl in 1,4-dioxane (9.63 mL, 38.52 mmol) was stirred at 25 °C for 16 hours. The mixture was concentrated and diluted with H_2O (20 mL) and the pH adjusted with addition of $NaHCO_3$ (solid) to pH~9. The mixture was extracted with EtOAc (20 mL x 3). The combined organic phase was concentrated to give the crude product (150 mg, 0.11 mmol, 11% yield) as an oil. **LCMS** R_t = 0.39 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{13}N_4O$ $[M+H]^+$ 205.1, found 204.9.

[000372] **68:** To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (140 mg, 0.72 mmol) in DCM (20 mL) was added DIPEA (0.4 mL, 2.88 mmol), HOBt (194.92 mg, 1.44 mmol) and EDCI (276.52 mg, 1.44 mmol) and the mixture was stirred at 25 °C for 30 minutes. Then 1-[3-(5-methyl-3-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (147.3 mg, 0.72 mmol) was added and the mixture was stirred at 25 °C for 16 hours. The reaction mixture was washed with H_2O (20 mL) and then the aqueous phase was extracted with DCM (20 mL x 2). The combined organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm × 30 mm, 5µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 17-47% B over 8 min) to give the product (91.9 mg, 0.24 mmol, 33% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ = 9.12 (d, 1H), 8.59 (d, 1H), 8.17 (s, 1H), 6.94 (s, 1H), 6.89 (br d, 1H), 5.63 (quin, 1H), 4.24 (s, 3H), 2.44 (s, 3H), 1.78 (d, 3H). **LCMS** R_t = 1.02 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{16}F_3N_6O_2$ $[M+H]^+$ 381.1, found 380.9.

Example 46. Synthesis of 69 and 70



[000373] **A-82b:** To a solution of Cs_2CO_3 (1.57 g, 4.8 mmol) and 2,2-difluoroethyl trifluoromethanesulfonate (1.03 g, 4.8 mmol) in CH_3CN (15 mL) was added ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (500 mg, 2.4 mmol). The reaction mixture was stirred at 20 °C for 3 hours. The mixture was concentrated to give a residue. The residue was diluted with sat. NH_4Cl (30 mL), and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with water (30 mL) and brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 15%) to give the product (400 mg, 1.44 mmol, 60% yield) as a solid. **LCMS** R_t = 4.52 min in 7.0 min chromatography, 0-60AB, MS ESI calcd. for $C_9H_{10}F_5N_2O_2$ $[M+H]^+$ 273.1, found 272.9.

[000374] **A-82:** To a mixture of ethyl 2-(2,2-difluoroethyl)-5-(trifluoromethyl)pyrazole-3-carboxylate (400 mg, 1.47 mmol) in ethanol (5 mL) and water (5 mL) was added NaOH (117.57 mg, 2.94 mmol), and the mixture was stirred at 50 °C for 3 hours. After cooling to room temperature, the reaction mixture was concentrated under reduced pressure to remove most of the ethanol and then diluted with H_2O (10 mL) and the mixture was washed with EtOAc (10 mL x 2, discarded). The pH of the aqueous phase was adjusted to pH ~ 2 with 1N HCl, then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with H_2O (10 mL) and brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (450 mg) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 7.36 (s, 1H), 6.62 - 6.31 (m, 1H), 5.10 (dt, 2H).

[000375] **A-83:** To a mixture of 2-(2,2-difluoroethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (224.06 mg, 0.92 mmol), HATU (634.51 mg, 1.67 mmol) and DIPEA (0.44 mL, 2.5 mmol) in DMF (10 mL) was added 1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (200 mg, 0.83 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with sat. NH_4Cl (20 mL) and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20


mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm 5 µm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 57-77% B over 9 min) to give the product (140 mg, 0.32 mmol, 39% yield) as a solid. **LCMS** R_t = 0.94 min in 1.5 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_5N_5O_2$ $[M+H]^+$ 430.1, found 430.2.

[000376] **69 & 70:** The product was analyzed by SFC (Daicel CHIRALCEL OJ-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C, ABPR: 1500 psi) showed two peaks 2.18 min and 2.59 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 15% B; 7 min run; 6 injections, Rt of peak 1 = 4.55 min, Rt of peak 2 = 5.55 min) to give the enantiomer 1, randomly assigned as **69** (37.48 mg, 87.3 mmol, 27% yield) (Rt = 2.18 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **70** (48.34 mg, 112.6 mmol, 34% yield) (Rt = 2.59 min in analytical SFC) as a solid.

69: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.59 (d, 1H), 7.85 - 7.75 (m, 2H), 7.54 (s, 1H), 7.48 - 7.38 (m, 2H), 6.57 - 6.25 (m, 1H), 5.46 (quin, 1H), 5.10 (dt, 2H), 2.39 (s, 3H), 1.67 (d, 3H). **LCMS** R_t = 1.38 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_5N_5O_2$ $[M+H]^+$ 430.1, found 430.0.

70: ^{1}H NMR (400MHz, DMSO-d_6) δ_H = 9.60 (d, 1H), 7.86 - 7.77 (m, 2H), 7.54 (s, 1H), 7.49 - 7.39 (m, 2H), 6.57 - 6.26 (m, 1H), 5.47 (quin, 1H), 5.10 (dt, 2H), 2.40 (s, 3H), 1.68 (d, 3H). **LCMS** R_t = 1.37 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_5N_5O_2$ $[M+H]^+$ 430.1, found 430.1.

Example 47. Synthesis of 71



[000377] **A-84:** A mixture of 6-methylpyridine-2-carbonitrile (10 g, 84.65 mmol), hydroxylamine hydrochloride (17.65 g, 253.94 mmol) in ethanol (24 mL) and NaOH (10.16

g, 253.94 mmol) in water (8 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature the mixture was concentrated to give a residue. The residue was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (8.2 g, 52.96 mmol, 62% yield) as a solid. **LCMS** Rt = 0.153 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_7H_{10}N_3O$ $[M+H]^+$ 152.07, found 152.0.


[000378] **A-85:** A mixture of 2-(tert-butoxycarbonylamino)propanoic acid (3754.91 mg, 19.85 mmol) and CDI (3539.69 mg, 21.83 mmol) in DMF (30 mL) was stirred at 15 °C for 1 hour. N-hydroxy-6-methyl-pyridine-2-carboxamidine (3 g, 19.85 mmol) was added and the reaction mixture was stirred at 110 °C for 16 hours. The mixture was diluted with NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (1800 mg, 3.82 mmol, 19% yield) as a solid. **LCMS** Rt = 0.805 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{15}H_{21}N_4O_3$ $[M+H]^+$ 305.15, found 305.2.

[000379] **A-86:** To tert-butyl N-[1-[3-(6-methyl-2-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (500 mg, 1.64 mmol) in 1,4-dioxane (10 mL) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 40 °C under N_2 for 5 hours. After cooling to room temperature, the mixture was concentrated to give a residue. The residue was poured into ice-water (20 mL) and the pH of mixture was adjusted with the addition of Na_2CO_3 (solid) to pH~ 9 and then extracted with EtOAc (20mL x 2). The organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (300 mg, 0.78 mmol, 69% yield) as a solid. **LCMS** Rt = 0.278 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{13}N_4O$ $[M+H]^+$ 205.1, found 205.1.

[000380] **71:** A mixture of 1-[3-(6-methyl-2-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (200 mg, 0.98 mmol), HOBt (264.66 mg, 1.96 mmol), EDCI (375.46 mg, 1.96 mmol), DIPEA (0.54 mL, 3.92 mmol) and 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (190.09 mg, 0.98 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was partially concentrated under reduced pressure to give a residue. The residue was diluted with H_2O (30 mL), and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC

(Boston Prime (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 30-60% B over 9 min) to give the product (78.77 mg, 0.20 mmol, 21% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.94 (d, 1H), 7.76 (t, 1H), 7.33 (d, 1H), 7.03 - 6.91 (m, 2H), 5.73 - 5.57 (m, 1H), 4.22 (s, 3H), 2.68 (s, 3H), 1.77 (d, 3H). **LCMS** R_t = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{16}F_3N_6O_2$ $[M+H]^+$ 381.12, found 381.1.

Example 48. Synthesis of 72



[000381] **A-90:** To a solution of tert-butoxycarbonyl tert-butyl carbonate (2.02 g, 9.23 mmol) in THF (8 mL) was added 2-amino-3-methoxy-propanoic acid (1 g, 8.39 mmol) in THF (18 mL) and NaOH (671.59 mg, 16.79 mmol) in water (18 mL) and the mixture was stirred at 20 °C for 16 hours. The mixture was partially concentrated to give a residue. The residue was diluted with H_2O (30 mL) and extracted with EtOAc (30 mL x 2), the organic phase was discarded. The aqueous phase was acidified with 10% aqueous acetic acid to pH ~ 4, and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the product (1650 mg, 7.53mmol, 89% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.05 - 7.75 (m, 1H), 5.41 (br s, 1H), 4.52 - 4.37 (m, 1H), 3.86 (br d, 1H), 3.63 (dd, 1H), 3.38 (d, 3H), 1.46 (d, 9H).

[000382] **A-3:** A mixture of 3-methylbenzonitrile (10 g, 85.36 mmol), hydroxylamine hydrochloride (17795.13 mg, 256.08 mmol) in ethanol (12 mL) and NaOH (10.24 g, 256.08 mmol) in water (4 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and

concentrated to give the crude product (12500 mg, 66.38 mmol, 77% yield) as a solid. **LCMS** Rt = 0.178 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_8H_{11}N_2O$ $[M+H]^+$ 151.1, found 151.1.


[000383] **A-88:** A mixture of N-hydroxy-3-methyl-benzamidine (650.78 mg, 4.33 mmol) and CDI (772.92 mg, 4.77 mmol) in DMF (20 mL) was stirred at 15 °C for 1 hour and then 2-(tert-butoxycarbonylamino)-3-methoxy-propanoic acid (950 mg, 4.33 mmol) was added. The reaction mixture was stirred at 110 °C for 16 hours. The mixture was diluted with sat. NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (290 mg, 0.86 mmol, 20% yield) as a solid. **LCMS** Rt = 0.92 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{17}H_{24}N_3O_4$ $[M+H\text{-}Boc]^+$ 278.1, found 334.2.

[000384] **A-89:** To tert-butyl N-[2-methoxy-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (290 mg, 0.87 mmol) in 1,4-dioxane (10 mL) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the mixture was stirred at 40 °C under N_2 for 5 hours. After cooling to room temperature, the mixture was concentrated under partial pressure to give a residue. The residue was poured into ice-water (20 mL) and the pH of the mixture was adjusted with Na_2CO_3 (solid) to pH~ 9 and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (180 mg, 0.63 mmol, 69% yield) as a solid. **LCMS** R_t = 0.678 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{16}N_3O_2$ $[M+H]^+$ 234.1, found 234.12.

[000385] **72:** A mixture of 2-methoxy-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (180 mg, 0.77 mmol), HOBt (208.54 mg, 1.54 mmol), EDCI (295.85 mg, 1.54 mmol), DIPEA (0.43 mL, 3.09 mmol) and 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (149.78 mg, 0.77 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was partially concentrated to give a residue. The residue was diluted with H_2O (20 mL), and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 50-80% B over 9 min) to give the product (77.3 mg, 0.19 mmol, 24% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.91 - 7.85 (m, 2H), 7.41 - 7.31 (m, 2H), 7.05 - 7.00 (m, 1H), 6.99 (s, 1H), 5.70 - 5.64 (m, 1H), 4.25 (s, 3H), 4.09

- 4.03 (m, 1H), 3.93 - 3.88 (m, 1H), 3.41 (s, 3H), 2.43 (s, 3H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}F_3N_5O_3$ $[M+H]^+$ 410.2, found 410.14.

Example 49. Synthesis of 73




[000386] **A-92:** To a solution of 2,2,2-trifluoroethanamine hydrochloride (4144.34 mg, 30.58 mmol) in toluene (50 mL) was added $NaNO_2$ (2109.94 mg, 30.58 mmol) in water (2.5 mL) and the reaction mixture was stirred at 0 °C for 0.5 hour. Then to the mixture was added ethyl prop-2-ynoate (1000 mg, 10.19 mmol) and the mixture was stirred at 20 °C for 1 hour. The mixture was diluted with H_2O (30 mL) and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (50 mL x 1) and brine (60 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (2000 mg, 9.61 mmol, 94% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 11.84 (br s, 1H), 7.11 (s, 1H), 4.45 (q, 2H), 1.42 (t, 3H).

[000387] **A-93:** To a solution of ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (1000 mg, 4.8 mmol) in MeCN (20 mL) was added Cs_2CO_3 (3130.59 mg, 9.61mmol) followed by sodium 2-chloro-2,2-difluoro-acetate (1464.98mg, 9.61mmol) and 18-crown-6 (253.98 mg, 0.96 mmol). The reaction mixture was stirred at 90 °C for 1.5 hours under N_2. After cooling to room temperature, the mixture was diluted with H_2O (30 mL) and then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5% to 10%) to give the product (100 mg, 387.4 µmol, 8.06% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.26 - 7.85 (m, 1H), 7.21 (s, 1H), 4.45 (q, 2H), 1.43 (t, 3H).

[000388] **A-94:** To a solution of ethyl 2-(difluoromethyl)-5-(trifluoromethyl)pyrazole-3-carboxylate (70 mg, 0.27 mmol) in ethanol (3 mL) was added a solution of NaOH (32.54 mg, 0.81 mmol) in water (3mL). The mixture was stirred at 20 °C for 2 hours. The reaction mixture was quenched by addition of 1N HCl (1 mL) and diluted with H_2O (10 mL) and extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the product (60 mg, 260.8 µmol, 96% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.21 - 7.81 (m, 1H), 7.30 (s, 1H).

[000389] **73:** To a mixture of 2-(difluoromethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (60 mg, 0.26 mmol), HATU (292.35 mg, 0.77 mmol) and DIPEA (0.27 mL, 1.54 mmol) in DCM (8 mL) was added 1-[3-(3-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (100 mg, 0.38 mmol) and the mixture was stirred at 25 °C for 2 hours. The mixture was diluted with H_2O (10 mL) and then extracted with EtOAc (20 mL x 2). The organic layer was washed brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 24%) to give the product (26.79 mg, 61.5 µmol, 16% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.31 - 7.99 (m, 2H), 7.99 - 7.95 (m, 1H), 7.55 - 7.49 (m, 1H), 7.48 - 7.42 (m, 1H), 7.10 (s, 1H), 6.95 (br d, 1H), 5.62 (quin, 1H), 1.80 (d, 3H). **LCMS** R_t = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{16}H_{12}ClF_5N_5O_2$ $[M+H]^+$ 436.1, found 435.9.

Example 50. Synthesis of 74 and 75



[000390] **A-96:** A mixture of 2,4-dichlorobenzonitrile (3 g, 17.44 mmol), hydroxylamine hydrochloride (3.64 g, 52.32 mmol) and NaOH (2.09 g, 52.32 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 16 hours to give a suspension. After cooling to room temperature, the reaction mixture was partially concentrated under vacuum to remove most of the EtOH and then diluted with H_2O (50 mL). The mixture was extracted with EtOAc (50 mL x 3). The combined organic phase was washed with brine (50

mL), dried over Na_2SO_4, filtered and concentrated to give the product (2 g, 7.72 mmol, 44% yield) as a solid. **LCMS** R_t = 0.232 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_7H_7Cl_2N_2O$ $[M+H]^+$ 205.1, found 204.8.

[000391] **A-97:** A mixture of 2-(tert-butoxycarbonylamino) propanoic acid (1.38 g, 7.32 mmol) and CDI (1.3 g, 8.05 mmol) in DMF (10 mL) was stirred at 15 °C for 1 hour and then 2,4-dichloro-N'-hydroxy-benzamidine (1.5 g, 7.32 mmol) was added. The reaction mixture was then stirred at 110 °C for 16 hours. After cooling to room temperature the mixture was diluted with H_2O (100 mL) and extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 % to 10%) to give the product (600 mg, 1.62 mmol, 22% yield) as a solid. **LCMS** R_t = 1.380 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{11}H_9Cl_2N_3O_3$ $[M+H\text{-}tBu]^+$ 302.0, found 301.9.

[000392] **A-98:** To tert-butyl N-[1-[3-(2,4-dichlorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (600 mg, 1.67 mmol) was added 4M HCl in 1,4-dioxane (16.36 mL, 65.45 mmol) and the mixture was stirred at 25 °C for 16 hours. The mixture was concentrated, diluted with H_2O (20 mL) and the pH was adjusted with $NaHCO_3$ (solid) to pH~9. The mixture was extracted with EtOAc (20 mL x 3). The combined organic phase was concentrated to give the crude product (500 mg, 0.92 mmol, 55% yield) as an oil. **LCMS** R_t = 0.906 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{10}Cl_2N_3O$ $[M+H\text{-}tBu]^+$ 258.0, found 257.8.


[000393] **A-99:** To a mixture of 2-methyl-5-(trifluoromethyl) pyrazole-3-carboxylic acid (280 mg, 1.44 mmol) in DCM (25 mL) was added DIPEA (0.8 mL, 5.77 mmol), HOBt (389.84 mg, 2.88 mmol), EDCI (553.05 mg, 2.88 mmol) and the mixture was stirred at 25 °C for 30 min. Then 1-[3-(2,4-dichlorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (372.3 mg, 1.44 mmol) was added and the reaction mixture was stirred at 25 °C for 16 hours. The reaction mixture was washed with H_2O (30 mL) and the mixture was extracted with DCM (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified from prep-HPLC (Boston Prime C18 (150 mm × 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 56-76% B over 9 min) to give the product (450 mg, 1.04 mmol, 72% yield) as oil. **LCMS** R_t = 1.318 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. For C16H13Cl2F3N5O2 $[M+H]^+$ 434.0, found 433.9.

[000394] **74 & 75:** Analytical SFC (Daicel CHIRALCEL OJ-H (150 mm x 4.6 mm, 5 μm), mobile phase: A: CO_2, B: ethanol (0.05% DEA), gradient: from 5 % to 40 % of B in 4.5 min and hold 40% for 5.5 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 2.94 min and 3.79 min. The product was purified by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 25% B; 7.76 min run; 45 injections, Rt of peak 1 = 4.39 min, Rt of peak 2 = 5.76 min) to give the enantiomer 1, randomly assigned as **74** (176.6 mg, 0.41 mmol, 44% yield) (Rt = 2.943 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **75** (211.15 mg, 0.49 mmol, 53 % yield) (Rt = 3.79 min in analytical SFC) as a solid.

74: ^{1}H NMR (400MHz, DMSO-d_6) δ = 9.47 (d, 1H), 7.96 (d, 1H), 7.91 (d, 1H), 7.66 (dd, 1H), 7.45 (s, 1H), 5.49 (quin, 1H), 4.14 (s, 3H), 1.68 (d, 3H). **LCMS** R_t = 1.38 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}Cl_2F_3N_5O_2$ $[M+H]^+$ 434.0, found 434.0.

75: **^{1}H NMR** (400MHz, DMSO-d_6) δ =9.46 (d, 1H), 7.96 (d, 1H), 7.91 (d, 1H), 7.66 (dd, 1H), 7.45 (s, 1H), 5.49 (quin, 1H), 4.14 (s, 3H), 1.68 (d, 3H). **LCMS** R_t = 1.38 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}Cl_2F_3N_5O_2$ $[M+H]^+$ 434.0, found 434.0.

Example 51. Synthesis of 76 and 77




67 76 77

[000395] Analytical SFC (Daicel CHIRALCEL OJ-H (150 mm x 4.6 mm, 5 μm), mobile phase: A: CO_2, B: ethanol (0.05% DEA), Gradient: from 5% to 40% of B in 4.5 min and hold 40% for 5.5 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 2.18 min and 2.40 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 10% B; 8.74 min run; 120 injections, Rt of peak 1 = 5.95 min, Rt of peak 2 = 6.74 min) to give the enantiomer 1, randomly assigned as **76** (178.71 mg, 0.44 mmol) (Rt = 2.18 min in analytical SFC) as a solid and the enantiomer 2 randomly assighned as 77 (138.05 mg, 0.34 mmol, 37% yield) (Rt = 2.40 min in analytical SFC) as a solid.

76: ^{1}H NMR (400MHz, DMSO-d_6) 400MHz δ = 9.46 (d, 1H), 8.05 - 7.96 (m, 1H), 7.94 - 7.83 (m, 1H), 7.74 - 7.59 (m, 1H), 7.45 (s, 1H), 5.46 (quin, 1H), 4.14 (s, 3H), 1.68 (d, 3H). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}F_5N_5O_2$ $[M+H]^+$ 402.1, found 402.0.

77: ^{1}H NMR (400MHz, DMSO-d_6) 400MHz δ = 9.46 (d, 1H), 8.10 - 7.95 (m, 1H), 7.93 - 7.82 (m, 1H), 7.67 (m, 1H), 7.45 (s, 1H), 5.46 (quin, 1H), 4.13 (s, 3H), 1.68 (d, 3H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}F_5N_5O_2$ $[M+H]^+$ 402.1, found 402.0.

Example 52. Synthesis of 78 and 79




68 78 79

[000396] Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm I.D., 5.0 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed 2 peaks at 3.78 min and 4.09 min. The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = *i*-PrOH(0.1% DEA) ; 38 °C; 60 mL/min; 35% B; 9 min run; 6 injections, Rt of peak 1 = 6.3 min, Rt of peak 2 = 7.3 min) to give the enantiomer 1, randomly assigned as **78** (35.11 mg, 92.3 µmol, 41% yield)(Rt = 3.78 min in analytical SFC) as an oil and enantiomer 2, randomly assigned as **79** (28.13 mg, 74 µmol, 33% yield) (Rt = 4.09 min in analytical SFC) as an oil.

78: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 9.13 (s, 1H), 8.60 (s, 1H), 8.22 (s, 1H), 6.96 (s, 1H), 6.90 (br d, 1H), 5.68 - 5.57 (m, 1H), 4.24 (s, 3H), 2.46 (s, 3H), 1.79 (d, 3H). **LCMS** R_t = 1.05 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{16}F_3N_6O_2$ $[M+H]^+$ 381.12, found 381.1.

79: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 9.13 (s, 1H), 8.64 - 8.54 (m, 1H), 8.23 - 8.16 (m, 1H), 6.95 (s, 1H), 6.85 - 6.75 (m, 1H), 5.69 - 5.55 (m, 1H), 4.24 (s, 3H), 2.46 (s, 3H), 1.79 (d, 3H). **LCMS** R_t = 1.05 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{16}F_3N_6O_2$ $[M+H]^+$ 381.12, found 381.1.

Example 53. Synthesis of 80 and 81




72 80 81

[000397] Analytical SFC (Daicel CHIRALPAK IC-3 (150 mm x 4.6 mm I.D., 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C, ABPR: 1500 psi) showed two peaks at 2.00 min and 2.57 min. The product was separated by SFC (Daicel CHIRALPAK IC (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 15% B; 7 min run; 6 injections, Rt of peak 1 = 4.4 min, Rt of peak 2 = 5.5 min) to give the enantiomer 1, randomly assigned as **80** (18.41 mg, 45.0 μmol, 26% yield) (Rt = 2.00 min in analytical SFC) as a oil and enantiomer 2, randomly assigned as **81** (19.73 mg, 48.2 μmol, 28% yield) (Rt = 2.57 min in analytical SFC) as an oil.

80: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.92 - 7.85 (m, 2H), 7.40 - 7.31 (m, 2H), 7.07 - 6.97 (m, 2H), 5.70 - 5.64 (m, 1H), 4.25 (s, 3H), 4.06 (dd, 1H), 3.90 (dd, 1H), 3.41 (s, 3H), 2.43 (s, 3H). **LCMS** R_t = 1.31 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}F_3N_5O_3$ $[M+H]^+$ 410.1, found 410.1.

81: ^{1}H NMR (400MHz, $CDCl_3$) δ_H = 7.92 - 7.85 (m, 2H), 7.40 - 7.32 (m, 2H), 7.06 - 6.98 (m, 2H), 5.70 - 5.65 (m, 1H), 4.25 (s, 3H), 4.06 (dd, 1H), 3.90 (dd, 1H), 3.41 (s, 3H), 2.43 (s, 3H). **LCMS** R_t = 1.31 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}F_3N_5O_3$ $[M+H]^+$ 410.1, found 410.1.

Example 54. Synthesis of 82 and 83



73 82 83


[000398] Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: methanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.16 min and 2.44 min. The product was separated by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = IPA (0.1% DEA); 38 °C; 60 mL/min; 35% B; 8 min run; 5 injections, Rt of peak 1 = 4.46 min, Rt of peak 2 = 5.3 min) to give the enantiomer 1, randomly assigned as **82** (3.68 mg, 8.4 μmol, 14% yield) (Rt = 2.16 min in

analytical SFC) as an oil and enantiomer 2, randomly assigned as **83** (7.25 mg, 16.6 μmol, 27 % yield) (Rt = 2.44 min in analytical SFC) as an oil.

[000399] **82:** ^{1}**H NMR** (400MHz, $CDCl_3$) δ_H = 8.31 - 7.99 (m, 2H), 7.99 - 7.94 (m, 1H), 7.54 - 7.50 (m, 1H), 7.46 (d, 1H), 7.10 (s, 1H), 6.99 - 6.91 (m, 1H), 5.68 - 5.57 (m, 1H), 1.80 (d, 3H). **LCMS** R_t = 1.39 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{12}ClF_5N_5O_2$ $[M+H]^+$ 436.05, found 436.0.

[000400] **83:** ^{1}**H NMR** (400MHz, $CDCl_3$) δ_H = 8.30 - 7.99 (m, 2H), 7.99 - 7.95 (m, 1H), 7.54 - 7.49 (m, 1H), 7.48 - 7.42 (m, 1H), 7.09 (s, 1H), 6.96 - 6.88 (m, 1H), 5.68 - 5.57 (m, 1H), 1.82 - 1.78 (m, 3H). **LCMS** R_t = 1.37 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{12}ClF_5N_5O_2$ $[M+H]^+$ 436.05, found 436.1.

Example 55. Synthesis of 84



[000401] **A-101:** A mixture of hydroxylamine hydrochloride (3.53 g, 50.79 mmol), 2-methylpyridine-4-carbonitrile (2 g, 16.93 mmol) and NaOH (2.03 g, 50.79 mmol) in ethanol (24 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was concentrated under vacuum to remove most of EtOH and then diluted with H_2O (30 mL). The mixture was extracted with EtOAc (30 mL x 2), and the combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filteredand concentrated to give the crude product (1.45 g, 9.59 mmol, 56% yield) as a soid. **LCMS** R_t = 0.11 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_{10}N_3O$ $[M+H]^+$ 152.1, found 152.0.


[000402] **A-102:** A mixture of 2-(tert-butoxycarbonylamino)propanoic acid (1.22 g, 6.48 mmol) and CDI (1.16 g, 7.13 mmol) in DMF (20 mL) was stirred at 25 °C for 1 hour and then N'-hydroxy-2-methyl-pyridine-4-carboxamidine (980 mg, 6.48 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with sat NaCl (30 mL), and the mixture was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (30 mL), dried over

Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (220 mg, 0.66 mmol, 10 % yield) as an oil. **LCMS** R_t = 0.70 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{15}H_{21}N_4O_3$ $[M+H]^+$ 305.2, found 305.7.

[000403] **A-103:** To a mixture of tert-butyl N-[1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (220 mg, 0.72 mmol) in ethyl acetate (3 mL) was added ethyl acetate/HCl (15 mL, 60 mmol), and the mixture was stirred at 25 °C for 3 hours. The reaction mixture was concentrated to give the crude product (180 mg, 0.74 mmol) as a solid. **LCMS** R_t =0.12 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{13}N_4O$ $[M+H]^+$ 205.1, found 205.2.

[000404] **84:** A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (77.58 mg, 0.4 mmol), HOBt (120.01 mg, 0.89 mmol), Et_3N (0.31 mL, 2.22 mmol) and EDCI (127.69 mg, 0.67 mmol) in DCM (7 mL) was stirred at 25 °C for 1 hour. Then to the mixture was added 1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethanamine hydrochloride (100 mg, 0.44 mmol) and it was stirred at 25 °C for 16 hours. The reaction mixture was quenched with sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 to 10% to 25% to 50%) to give the product (63.51 mg, 0.17 mmol, 38% yield) as a solid.**^{1}H NMR** (400MHz, CD_3CN) δ_H = 8.67 (d, 1H), 7.81 (s, 1H), 7.74 (d, 1H), 6.95 (s, 1H), 6.75 (d, 1H), 5.62 (q, 1H), 4.24 (s, 3H), 2.67 (s, 3H), 1.79 (d, 3H). **LCMS** R_t = 0.90 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{16}F_3N_6O_2$ $[M+H]^+$ 381.1, found 380.9.

Example 56. Synthesis of 85



[000405] **A-105:** A mixture of hydroxylamine hydrochloride (0.54 g, 7.8 mmol), benzene-1,3-dicarbonitrile (1 g, 7.8 mmol) and NaOH (0.31 g, 7.8 mmol) in ethanol (20 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of EtOH and then it was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.4 g, 6.06 mmol, 77% yield) as a solid. **LCMS** R_t = 0.14 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8N_3O$ $[M+H]^+$ 162.1, found 162.0.


[000406] **A-106:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (821.84 mg, 4.34 mmol) and CDI (774.73 mg, 4.78 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour and then 3-cyano-N'-hydroxy-benzamidine (700 mg, 4.34 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (60 mL) and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 30%) to give the product (400 mg, 0.77 mmol) as a solid. **LCMS** R_t = 1.15 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{10}N_4O_3$ $[M+H\text{-}tBu]^+$ 259.1, found 258.9.

[000407] **A-107:** A mixture of tert-butyl N-[(1S)-1-[3-(3-cyanophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (400 mg, 1.27 mmol) and 4M HCl in 1,4-dioxane (15 mL, 60 mmol) was stirred at 25 °C for 16 hours. The mixture was concentrated, diluted with H_2O (20 mL) and basified with $NaHCO_3$(solid) to pH~8. The mixture was extracted with EtOAc (30 mL x 3), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (410 mg, 1.60 mmol, 29% yield) as an oil.

[000408] **85:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160 mg, 1.19 mmol), EDCI (228 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and 3-[5-[(1S)-1-aminoethyl]-1,2,4-oxadiazol-3-yl]benzonitrile (152.84 mg, 0.71 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (66 mg, 0.18 mmol, 30% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ = 8.40 (s, 1H), 8.34 (d, 1H), 7.86

(s, 1H), 7.82 (d, 1H), 7.64 (t, 1H), 6.27 (br d, 1H), 5.70 - 5.57 (m, 1H), 4.53 - 4.40 (m, 1H), 2.56 (s, 3H), 1.76 (d, 3H), 1.54 (d, 6H). **LCMS** R_t = 1.10 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{20}N_6O_2$ $[M+H]^+$ 365.2, found 365.2.

Example 57. Synthesis of 86



[000409] **A-109:** A mixture of 4-chloro-3-fluoro-benzonitrile (1 g, 6.43 mmol), hydroxylamine hydrochloride (1.34 g, 19.29 mmol), NaOH (0.77 g, 19.29 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the mixture was partially concentrated under reduced pressure to give a residue. The residue was diluted with H_2O (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1 g, 2.74 mmol, 43% yield) as a solid. **LCMS** Rt = 0.33 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_7H_7ClFN_2O$ $[M+H]^+$ 189.02, found 188.7.

[000410] **A-110:** To a mixture of 4-chloro-3-fluoro-N-hydroxy-benzamidine (1 g, 5.3 mmol) and CDI (945.78 mg, 5.83 mmol) in DMF (30 mL) was stirred at 25 °C for 1 hour before (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1 g, 5.3 mmol) was added. The reaction mixture was then stirred at 80 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (350 mg, 0.99 mmol, 19% yield) as a solid. **LCMS** Rt = 0.96 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{15}H_{18}ClFN_3O_3$ $[M+H\text{-}BOC]^+$ 286.09, found 286.1.


[000411] **A-111:** To tert-butyl N-[(1S)-1-[3-(4-chloro-3-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (350 mg, 1.02 mmol) in 1,4-dioxane (10 mL) was added 4M HCl in 1,4-

dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 30 °C under N_2 for 5 hours. After cooling to room temperature, the mixture was concentrated under reduced pressure to give a residue. The residue was poured into ice-water (20 mL) and the mixture was basified with Na_2CO_3 (solid) to pH~ 9, and extracted with EtOAc (20 mL x 2), the organic phase was washed with brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (250 mg, 0.99 mmol, 69% yield) as an oil. **LCMS** Rt = 0.68 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{10}ClFN_3O$ $[M+H]^+$ 242.04, found 241.8.

[000412] **86:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(4-chloro-3-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethanamine (172.41 mg, 0.71 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was concentrated to give a residue. The residue was diluted with H_2O (30 mL), and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50%) to give the product.

[000413] Analytical SFC (Regis (S,S) Whelk-01 (100 mm × 4.6 mm I.D., 5.0 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 4.41 (12.4%) min and 5.48 min (87.6%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 40% B; 10 min run; 5 injections, Rt of peak 1 = 5.6 min, Rt of peak 2 = 8.2 min) to give the product (100.73 mg, 0.26 mmol, 43% yield) (Rt = 5.48 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.91 - 7.80 (m, 3H), 7.53 (t, 1H), 6.26 (br d, 1H), 5.74 - 5.53 (m, 1H), 4.52 - 4.38 (m, 1H), 2.54 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H). **LCMS** Rt = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{20}ClFN_5O_2$ $[M+H]^+$ 392.12, found 391.9.

Example 58. Synthesis of 87



A-112 A-113 A-114 A-115 87

[000414] **A-113:** To a mixture of 3,4-difluorobenzonitrile (2 g, 14.38 mmol) and hydroxylamine hydrochloride (3 g, 43.13 mmol) in ethanol (24 mL) was added NaOH (1.73 g, 43.13 mmol). The mixture was stirred at 40 °C for 16 hours. The mixture was diluted with H_2O (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (3 g, 16.11 mmol) as a solid. **LCMS** R_t = 0.13 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_7F_2N_2O$ $[M+H]^+$ 173.0, found 173.1.


[000415] **A-114:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1099.23 mg, 5.81 mmol) and CDI (1036.22 mg, 6.39 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour and then 3,4-difluoro-N'-hydroxy-benzamidine (1000 mg, 5.81 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with bine (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (700 mg, 2.15 mmol, 37% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H =7.96 - 7.88 (m, 1H), 7.88 - 7.83 (m, 1H), 7.32 - 7.28 (m, 1H) , 5.16 (s, 2H), 1.63 (d, 3H), 1.47 (s, 9H).

[000416] **A-115:** A corlorless mixture of tert-butyl N-[(1S)-1-[3-(3,5-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (700 mg, 2.15 mmol) and HCl/EtOAc (10 mL, 4M) was stirred at 25 °C for 6 hours. The mixture was concentrated to give the crude product as a solid, which was used directly without any further purification. **LCMS** R_t = 0.61 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{10}F_2N_3O$ $[M+H]^+$ 226.1, found 225.8.

[000417] **87:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (15 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg,

1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(3,5-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (171.13 mg, 0.65 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction mixture was quenched with saturated NH_4Cl (15 mL), then extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the impure product, the impure product was purified by prep-TLC (silica gel, PE:EtOAc = 5:2) to give the product. The product was triturated from DCM (1 mL) and n-hexane (5 mL) to give the product (160.64 mg, 0.43 mmol, 72% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ_H = 8.55 (d, 1H), 8.27 (s, 1H), 8.02 - 7.94 (m, 1H), 7.90 - 7.83 (m, 1H), 7.70 - 7.60 (m, 1H), 5.41 - 5.30 (m, 1H), 4.48 - 4.33 (m, 1H), 2.29 (s, 3H), 1.61 (d, 3H), 1.40 (d, 6H). **LCMS** R_t = 1.13 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{20}F_2N_5O_2$ $[M+H]^+$ 376.2, found 376.0.

Example 59. Synthesis of 88



[000418] **A-117:** A mixture of 3-chloro-4-fluoro-benzonitrile (1 g, 6.43 mmol), hydroxylamine hydrochloride (1.34 g, 19.29 mmol), NaOH (0.77 g, 19.29 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours. After cooling to room temperature, the mixture was partially concentrated under reduced pressure to give the residue and the residue was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1 g, 3.35 mmol, 52% yield) as a solid. **LCMS** Rt = 0.29 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_7H_7ClFN_2O$ $[M+H]^+$ 189.0, found 188.8.

[000419] **A-118:** A mixture of 3-chloro-4-fluoro-N-hydroxy-benzamidine (1 g, 5.3 mmol) and CDI (945.78 mg, 5.83 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour before (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1 g, 5.3 mmol) was added. The

reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (350 mg, 1.00 mmol, 19% yield) as a solid. **LCMS** Rt = 0.96 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{15}H_{18}ClFN_3O_3$ $[M+H\text{-}Boc]^+$ 286.1, found 286.0.


[000420] **A-119:** To tert-butyl N-[(1S)-1-[3-(3-chloro-4-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (350 mg, 1.02 mmol) in 1,4-dioxane (10 mL) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the mixture was stirred at 30 °C under N_2 for 5 hours. After cooling to room temperature, the mixture was concentrated to give a residue. The residue was poured into ice-water (20 mL) and the pH of the mixture was basified with Na_2CO_3 (solid) to pH ~ 9, and then extracted with EtOAc (20 mL x 2). The organic phase was washed with brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (250 mg, 0.99 mmol, 69% yield) as an oil. **LCMS** Rt = 0.67 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{10}ClFN_3O$ $[M+H]^+$ 242.0, found 241.8.

[000421] **88:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(3-chloro-4-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethanamine (172.41 mg, 0.71 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was concentrated to give a residue. The residue was diluted with H_2O (30 mL), and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50%) to give the product.

[000422] Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm, 5.0 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min.) showed two peaks at 4.45 (17.1%) and 5.55 min (main peak, 82.9%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 50% B; 8 min run; 4 injections, Rt of peak 1 = 4.4 min, Rt of peak 2 = 6.5 min) to give the product (80.08 mg, 0.20 mmol, 34 % yield) (Rt = 5.55 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, CD_3CN) δ_H = 8.14 (dd, 1H), 8.03 - 7.97 (m, 1H), 7.96 (s, 1H), 7.40 (t, 1H), 7.02 (br d, 1H), 5.54 - 5.30 (m, 1H), 4.53 - 4.31 (m, 1H), 2.35 (s, 3H), 1.66 (d,

3H), 1.44 (d, 6H). **LCMS** Rt = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{20}ClFN_5O_2$ $[M+H]^+$ 392.12, found 391.9.

Example 60. Synthesis of 89



A-120 A-121 A-122 A-123 89

[000423] **A-121:** A mixture of 3-(trifluoromethyl)benzonitrile (2 g, 11.69 mmol), $NH_2OH.HCl$ (2.44 g, 35.06 mmol) and NaOH (1.4 g, 35.06 mmol) in ethanol (30 mL) was stirred at 30 °C for 16 hours. The reaction mixture was partially concentrated under reduced pressure to remove the EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (2.2 g, 9.36 mmol, 80% yield) as a solid. **LCMS** R_t = 0.33 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8F_3N_2O$ $[M+H]^+$ 205.1, found 205.1.

[000424] **A-122:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (648.78 mg, 3.43 mmol) and CDI (611.59 mg, 3.77 mmol) in DMF (10 mL) was stirred at 25 °C for 1 hour and then N'-hydroxy-3-(trifluoromethyl)benzamidine (700 mg, 3.43 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (30 mL), and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (250 mg, 0.65 mmol, 19% yield) as a solid. **LCMS** R_t = 0.96 min in 1.5 min chromatography, 5-95AB, , MS ESI calcd. for $C_{12}H_{11}F_3N_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 302.1, found 302.1.


[000425] **A-123:** To tert-butyl N-[(1S)-1-[3-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (250 mg, 0.70 mmol) was added 4N HCl in 1,4-dioxane (20

mL, 0.70mmol) and the mixture was stirred at 25 °C for 2 hours. The mixture was concentrated, diluted with water (20 mL) and basified with Na_2CO_3 (solid) to pH ~9. The mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the product (160 mg, 0.57 mmol, 82% yield) as a solid. **LCMS** R_t = 0.71 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{11}F_3N_3O$ $[M+H]^+$ 258.1, found 258.1.

[000426] **89:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-[3-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine (152.93 mg, 0.59 mmol) in CH_2Cl_2 (10 mL) was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (20 mL) and extracted with CH_2Cl_2 (20 mL x 2). The combined organic phase was washed with brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 40%), then triturated from *n*-hexane/DCM (2:1, 3 mL) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm, 5.0μμm), mobile phase: A: CO_2 B: ethanol (0.05% DEA) gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 3.79 (38.1%) min and 4.80 min (main peak, 61.9%).

[000427] The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 35% B; 10 min run; 7 injections, Rt of peak 1 = 5 min, Rt of peak 2= 8 min) to give the product (152.93 mg, 0.59 mmol) (Rt = 4.80 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.36 (s, 1H), 8.27 (d, 1H), 7.85 (s, 1H), 7.78 (d, 1H), 7.66 - 7.60 (m, 1H), 6.31 (br d, 1H), 5.66 (quin, 1H), 4.50 - 4.39 (m, 1H), 2.55 (s, 3H), 1.75 (d, 3H), 1.52 (d, 6H). **LCMS** R_t = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_2$ $[M+H]^+$ 408.2, found 408.0.

Example 61. Synthesis of 90



A-124 A-125 A-126 A-127 90

[000428] **A-125:** A mixture of 4-fluorobenzonitrile (2 g, 16.51 mmol), $NH_2OH.HCl$ (3.44 g, 49.54 mmol) and NaOH (1.98 g, 49.54 mmol) in ethanol (30 mL) and was stirred at 30 °C for 16 hours. The reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH, then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (2.2 g, 11.03 mmol, 67% yield) as a solid. **LCMS** R_t = 0.15 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8FN_2O$ $[M+H]^+$ 155.1, found 155.0.


[000429] **A-126:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (736.52 mg, 3.89 mmol) and CDI (694.3 mg, 4.28 mmol) in DMF (10 mL) was stirred at 25 °C for 1 hour and then 4-fluoro-N'-hydroxy-benzamidine (600 mg, 3.89 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 40%) to give the product (257 mg, 0.80 mmol, 21% yield) as a solid. **LCMS** R_t = 0.90 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{11}FN_3O_3$ $[M+H-t\text{-}Bu]^+$ 252.1, found 252.1.

[000430] **A-127:** To tert-butyl N-[(1S)-1-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (257 mg, 0.84 mmol) was added 4M HCl in 1,4-dioxane (30 mL, 0.84 mmol) and the reaction mixture was stirred at 25 °C for 2 hours. The mixture was concentrated, diluted with water (30 mL), and basified with Na_2CO_3 (solid) to pH ~ 9. and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the product (180 mg, 0.77 mmol, 92% yield) as a solid. **LCMS** R_t = 0.61 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{11}FN_3O$ $[M+H]^+$ 208.1, found 208.1.

[000431] **90:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (147.83 mg, 0.71 mmol) in CH_2Cl_2 (6 mL) was stirred at 25 °C for 16 hours. The reaction mixture was quenched with H_2O (10 mL) and then extracted with CH_2Cl_2 (10 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 40%), then triturated from *n*-hexane/DCM (2:1, 3 mL) to give the product.

[000432] analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm, 5.0 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA) gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 4.28 (7.2%) min and 5.36 min (main peak, 92.8%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 40% B; 10 min run; 5 injections, Rt of peak 1 = 5.4 min, Rt of peak 2= 8.1 min) to give the product (84.82 mg, 0.24 mmol, 40% yield) (Rt = 5.36 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.12 - 8.04 (m, 2H), 7.85 (s, 1H), 7.21 - 7.15 (m, 2H), 6.34 (br d, 1H), 5.64 (quin, 1H), 4.44 (spt, 1H), 2.55 (s, 3H), 1.73 (d, 3H), 1.51 (d, 6H). **LCMS** R_t = 1.10 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{21}FN_5O_2$ $[M+H]^+$ 358.2, found 358.0.

Example 62. Synthesis of 91



[000433] **A-129:** A mixture of 3-chloro-5-fluoro-benzonitrile (1 g, 6.43 mmol), hydroxylamine hydrochloride (1.34 g, 19.29 mmol), and NaOH (0.77 g, 19.29 mmol) in ethanol (24 mL) and water (8 mL) was stirred at 40 °C for 12 hours. After cooling to room

temperature, the mixture was partially concentrated under reduced pressure to give a residue. The residue was diluted with H_2O (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1 g, 5.27 mmol, 82% yield) as a solid. **LCMS** Rt = 0.313 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_7H_7ClFN_2O$ $[M+H]^+$ 189.0, found 188.8.

[000434] **A-130:** To a mixture of 3-chloro-5-fluoro-N-hydroxy-benzamidine (500 mg, 2.65 mmol) and CDI (472.89 mg, 2.92 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour before (2S)-2-(tert-butoxycarbonylamino)propanoic acid (501.64 mg, 2.65 mmol) was added. The reaction was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product (250 mg, 0.72 mmol, 27% yield) as a solid. **LCMS** Rt = 0.97 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{15}H_{18}ClFN_3O_3$ $[M+H\text{-}Boc]^+$ 286.1, found 286.0.


[000435] **A-131:** To tert-butyl N-[(1S)-1-[3-(3-chloro-5-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (250 mg, 0.73 mmol) in 1,4-dioxane (10mL) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 40 °C under N_2 for 5 hours. After cooling to room temperature, the mixture was concentrated to give a residue. The residue was poured into ice-water (20 mL) and the pH of the mixture was basified with Na_2CO_3 (solid) to pH~ 9, and then extracted with EtOAc (20 mL x 2). The organic phase was washed with brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (180 mg, 0.69 mmol, 69% yield) as an oil. **LCMS** Rt = 0.69 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{10}ClFN_3O$ $[M+H]^+$ 242.0, found 242.0.

[000436] **91:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(3-chloro-5-fluoro-phenyl)-1,2,4-oxadiazol-5-yl]ethanamine (172.41 mg, 0.71 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced to give a residue. The residue was diluted with H_2O (30 mL) and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude

product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50%) to give the product.

[000437] Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm I.D., 5.0 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 4.23 (12.8%) min and 5.29 min (main peak, 87.2%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 50% B; 8 min run; 4 injections, Rt of peak 1 = 4.2 min, Rt of peak 2 = 6.0 min) to give the product (63.77 mg, 0.16 mmol, 27% yield) (Rt = 5.29 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, CD_3CN) δ_H = 7.96 (s, 1H), 7.88 (s, 1H), 7.73 (br d, 1H), 7.42 (br d, 1H), 7.04 (br s, 1H), 5.46 - 5.35 (m, 1H), 4.47 - 4.36 (m, 1H), 2.35 (s, 3H), 1.67 (d, 3H), 1.45 (d, 6H). **LCMS** Rt = 1.19 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{20}ClFN_5O_2$ $[M+H]^+$ 392.12, found 391.9.

Example 63. Synthesis of 92



[000438] **A-133:** A mixture of hydroxylamine hydrochloride (1.5 g, 21.57 mmol), 3,5-difluorobenzonitrile (1 g, 7.19 mmol) and NaOH (0.86 g, 21.57 mmol) in ethanol (24 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH, diluted with H_2O (20 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.28 g, 4.19 mmol, 58% yield) as a solid. **LCMS** R_t = 0.18 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_7F_2N_2O$ $[M+H]^+$ 173.0, found 173.0.


[000439] **A-134:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.41 g, 7.44 mmol) and CDI (1.33 g, 8.18 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour

and then 3,5-difluoro-N'-hydroxy-benzamidine (1.28 g, 7.44 mmol) was added. The reaction mixture was stirred at 70 °C for 16 hours then cooled to room temperature. The residue was diluted with H_2O (20 mL), and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with and brine (20 mL x 2), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 15%) to give the product (660 mg, 2.03 mmol, 27% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.68 - 7.54 (m, 2H), 7.02 - 6.91 (m, 1H), 5.17 (s, 2H), 1.64 (d, 3H), 1.47 (s, 9H).

[000440] **A-135:** A mixture of tert-butyl N-[(1S)-1-[3-(3,5-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (660 mg, 2.03 mmol) and HCl/EtOAc (10.mL, 4M) was stirred at 25 °C for 6 hours. The mixture was concentrated under reduced pressure to give the crude product as a solid. The mixture was used directly without any further purification. **LCMS** R_t = 0.63 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{10}F_2N_3O$ $[M+H]^+$ 226.1, found 226.1.

[000441] **92:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (15 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38mmol) and (1S)-1-[3-(3,5-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (186.69 mg, 0.71 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction mixture was quenched with saturated NH_4Cl (15 mL), then extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by prep-TLC (silica gel, PE:EtOAc = 5:2) to give the impure product. The impure product was triturated from DCM (1 mL) and n-hexane (5 mL) to give the product (120.53 mg, 0.32 mmol, 53% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.84 (s, 1H), 7.66 - 7.58 (m, 2H), 7.01 - 6.92 (m, 1H), 6.28 (d, 1H), 5.70 - 5.57 (m, 1H), 4.50 - 4.40 (m, 1H), 2.54 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H). **LCMS** R_t = 1.15 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{20}F_2N_5O_2$ $[M+H]^+$ 376.2, found 375.9.

Example 64. Synthesis of 93 and 94





[000442] Analytical SFC (Daicel CHIRALCEL OJ-3 (100 x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA). gradient: from 5% to 40% of B in 4 min and hold 40% for 2.5 min, then 5% of B for 1.5 min, flow rate: 2.8 mL/min, column temp: 35 °C, ABPR: 1500 psi) showed two peaks at 2.29 min and 2.50 min. The product was purified by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 15% B; 13 min run; 6 injections, Rt of peak 1 = 8.7 min, Rt of Peak 2 = 11.2 min) to give the enantiomer 1, randomly assigned as **93** (4.23 mg, 12 µmol, 22% yield) (Rt = 2.29 min in analytical SFC) as a solid and enantiomer 2, randomly assigned as **94** (5.91 mg, 17 µmol, 30% yield) (Rt = 2.50 min in analytical SFC) as a solid.

[000443] **93: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.79 (td, 1H), 7.47 (dt, 1H), 7.25 - 7.15 (m, 1H), 6.53 (br d, 1H), 6.31 (s, 1H), 5.59 (quin, 1H), 4.10 (s, 3H), 1.98 - 1.88 (m, 1H), 1.75 (d, 3H), 0.99 - 0.91 (m, 2H), 0.78 - 0.71 (m, 2H). **LCMS** R_t = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_2$ $[M+H]^+$ 356.1, found 355.9.

[000444] **94: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.79 (td, 1H), 7.47 (dt, 1H), 7.26 - 7.15 (m, 1H), 6.55 (br d, 1H), 6.31 (s, 1H), 5.59 (quin, 1H), 4.10 (s, 3H), 1.99 - 1.87 (m, 1H), 1.75 (d, 3H), 0.99 - 0.91 (m, 2H), 0.78 - 0.71 (m, 2H). **LCMS** R_t = 1.22 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_2$ $[M+H]^+$ 356.1, found 356.1.

Example 65. Synthesis of 281 & 95





95

[000445] **281:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (0.74 g, 3.89 mmol) and CDI (0.69 g, 4.28 mmol) in DMF (18 mL) was stirred at 15 °C for 1 hour and then 3-fluoro-N'-hydroxy-benzamidine (0.6 g, 3.89 mmol) was added. The reaction mixture


was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with H_2O (30 mL) and extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 15% to 40%) to give the product (420 mg, 1.37 mmol, 35% yield) as an oil. **LCMS** R_t = 0.91 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{11}FN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 252.07, found 252.1.

[000446] **A-138:** To tert-butyl N-[(1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (420 mg, 1.37 mmol) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 20 °C for 12 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with $NaHCO_3$ (solid) to a pH~8. The mixture was extracted with EtOAc (20 mL x 2), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (270 mg, 1.30 mmol, 95% yield) as an oil. **LCMS** R_t = 0.41 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{11}FN_3O$ $[M+H]^+$ 208.08, found 207.9.

[000447] **95:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (147.83 mg, 0.71 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (30 mL) and brine (5 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 15% to 40%) to give the product (150 mg) as a solid. Analytical SFC analysis (Regis (S,S) Whelk-O1 (100 × 4.6 mm, 5 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B in 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed 2 peaks at 4.27 min (8.1%), and 5.32 min (main peak, 91.9%) The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = ethanol (0.1% NH_4OH) ; 38 °C; 60 mL/min; 50% B; 8 min run; 8 injections, Rt of peak 1 = 4.1 min, Rt of peak 2 = 6 min) to give the product (105.13 mg, 294.2 µmol, 70% yield) (Rt = 5.32 min in analysis analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.85 (s, 1H), 7.81 - 7.76 (m, 1H), 7.51 - 7.43 (m, 1H), 7.25 - 7.19 (m, 1H), 6.43 - 6.21 (m, 1H), 5.76 - 5.55 (m, 1H), 4.50 - 4.87 (m, 1H), 2.55 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H).

LCMS R_t = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{21}FN_5O_2$ $[M+H]^+$ 358.16, found 358.1.

Example 66. Synthesis of 96



[000448] **A-12:** A mixture of 3-chlorobenzonitrile (2 g, 14.54 mmol), $NH_2OH.HCl$ (3.03 g, 43.61 mmol) and NaOH (1.74 g, 43.61 mmol) in ethanol (18 mL) and water (6 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of EtOH and then diluted with H_2O (30 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product as a solid. **LCMS** R_t = 0.2 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8ClN_2O$ $[M+H]^+$ 171.0, found 171.0.

[000449] **A-139:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (0.43 g, 2.26 mmol) and CDI (0.4 g, 2.48 mmol) in DMF (18 mL) was stirred at 15 °C for 1 hour and then 3-chloro-N'-hydroxy-benzamidine (0.6 g, 2.26 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated NH_4Cl (30 mL), and the mixture was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 20% to 50%) to give the product (370 mg, 1.14 mmol, 50% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.09 (t, 1H), 7.97 (td, 1H), 7.52 - 7.46 (m, 1H), 7.45 - 7.40 (m, 1H), 5.18 (br s, 2H), 1.66 - 1.62 (m, 3H), 1.47 (s, 9H). **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{11}ClN_3O_3$ $[M+H\text{-}t\text{-}Bu]^+$ 268.04, found 268.0.


[000450] **A-140:** To tert-butyl N-[(1S)-1-[3-(3-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (370 mg, 1.14 mmol) was added 4M HCl in 1,4-dioxane (5 mL, 20 mmol)

and the reaction mixture was stirred at 20 °C for 1.5 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with $NaHCO_3$ (solid) to pH~8. The mixture was extracted with EtOAc (20 mL x 2), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (250 mg, 1.07 mmol, 94% yield) as an oil. **LCMS** R_t = 0.67 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{11}ClN_3O$ $[M+H]^+$ 224.05, found 224.0.

[000451] **A-141:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(3-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (159.58 mg, 0.71 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with sat. NH_4Cl (30 mL) and brine (5 mL), extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 15% to 40%) to give the product (150 mg) as a solid.

[000452] **96:** Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 5 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B in 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed 2 peaks at 4.74 min (6.9%), and 5.89 min (93.1%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = ethanol (0.1% NH_4OH); 38 °C; 60 mL/min; 50% B; 9 min run; 6 injections, Rt of peak 1 = 4.8 min, Rt of peak 2 = 7 min) to give the product (84.2 mg, 225.2 µmol, 56% yield) (Rt = 5.89 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.08 (t, 1H), 7.97 (td, 1H), 7.85 (s, 1H), 7.52 - 7.47 (m, 1H), 7.45 - 7.39 (m, 1H), 6.37 - 6.23 (m, 1H), 5.71 - 5.59 (m, 1H), 4.51 - 4.39 (m, 1H), 2.55 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H). **LCMS** R_t = 1.22 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{21}ClN_5O_2$ $[M+H]^+$ 374.1, found 374.1.

Example 67. Synthesis of 97



A-19 A-20 A-142 A-143 97

[000453] **A-20:** A mixture of 4-chlorobenzonitrile (2 g, 14.54 mmol), $NH_2OH.HCl$ (3.03 g, 43.61 mmol) and NaOH (1.74 g, 43.61 mmol) in ethanol (18 mL) and water (6 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH and then diluted with H_2O (30 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (2.1 g, 11.81 mmol, 81% yield) as a solid. **LCMS** R_t = 0.18 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_7H_8ClN_2O$ $[M+H]^+$ 171.02, found 171.0.


[000454] **A-142:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (0.67 g, 3.52 mmol) and CDI (0.63 g, 3.87 mmol) in DMF (18 mL) was stirred at 15 °C for 1 hour and then 4-chloro-N'-hydroxy-benzamidine (0.6 g, 3.52 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated NH_4Cl (30 mL), and the mixture was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 15% to 40%) to give the product (580 mg, 1.79 mmol, 51% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.05 - 8.03 (m, 1H), 8.02 - 8.01 (m, 1H), 7.49 - 7.46 (m, 1H), 7.46 - 7.44 (m, 1H), 5.17 (br s, 2H), 1.65 - 1.62 (m, 3H), 1.47 (s, 9H).

[000455] **A-143:** To tert-butyl N-[(1S)-1-[3-(4-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (580 mg, 1.79 mmol) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 20 °C for 1.5 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with $NaHCO_3$ (solid) to pH~8. The mixture was extracted with EtOAc (20 mL x 2), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and

concentrated to give the crude product (320 mg, 1.42 mmol, 79% yield) as an oil. **LCMS** R_t = 0.85 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{10}H_{11}ClN_3O$ $[M+H]^+$ 224.05, found 223.9.

[000456] **97:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(4-chlorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (159.58 mg, 0.71 mmol) and it was stirred at 25 °C for 16 hours. The reaction mixture was quenched with sat. NH_4Cl (30 mL) and brine (5 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 15% to 40%) to give the product (100 mg, 267.5 µmol, 45% yield) as a solid. Analytical SFC (Regis (S,S) Whelk-O1 (150 mm × 4.6 mm, 5 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B in 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed 2 peaks at 4.70 min (10.2%) and 5.82 min (89.8%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = ethanol (0.1% NH_4OH); 38 °C; 60 mL/min; 50% B; 9 min run; 6 injections, Rt of peak 1 = 4.8 min, Rt of peak 2 = 7 min) to give the product (91.16 mg, 243.8 µmol, 61% yield) (Rt= 5.82 in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.05 - 7.99 (m, 2H), 7.84 (s, 1H), 7.50 - 7.44 (m, 2H), 6.39 - 6.23 (m, 1H), 5.72 - 5.53 (m, 1H), 4.51 - 4.38 (m, 1H), 2.55 (s, 3H), 1.73 (d, 3H), 1.52 (d, 6H). **LCMS** R_t = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{21}ClN_5O_2$ $[M+H]^+$ 374.1, found 374.0.

Example 68. Synthesis of 98



[000457] **A-145:** A mixture of 4-(trifluoromethyl)benzonitrile (1 g, 5.84 mmol), hydroxylamine hydrochloride (1.22 g, 17.53 mmol) and NaOH (0.7 g, 17.53 mmol) in ethanol (20mL) as stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduce pressure to remove most of the EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.46 g, 6.96 mmol) as a solid.**LCMS** R_t = 0.41 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8F_3N_2O$ $[M+H]^+$ 205.1, found 205.0.

[000458] **A-146:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (463.41 mg, 2.45 mmol) and CDI (436.85 mg, 2.69 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour and then N'-hydroxy-4-(trifluoromethyl)benzamidine (500 mg, 2.45 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (60 mL), and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 30%) to give the product (550 mg, 1.54 mmol, 63 % yield) as a solid. **LCMS** R_t = 1.34 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{10}F_3N_3O_3$ $[M+H-tBu]^+$ 302.1, found 301.9.


[000459] **A-147:** To tert-butyl N-[(1S)-1-[3-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (550 mg, 1.54 mmol) was added 4M HCl in1,4-dioxane (15 mL) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with $NaHCO_3$(solid) to pH~8. The mixture was extracted with EtOAc (30 mL x 3), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (440 mg, 1.71 mmol) as an oil.

[000460] **98:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-[4-(trifluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine (183.51 mg, 0.71 mmol) and the resultant mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified

with flash chromatography on silica gel (EA in PE = 0/1 to 1/5) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 5.0 µm), mobile phase: A: CO2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40%, for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 3.78 min (6.7%) and 4.74 min (main peak, 93.3%).

[000461] Then the product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 40% B; 7.80 min run; 4 injections, Rt of peak 1 = 4.00 min, Rt of peak 2 = 5.80 min) to give the product (58.15 mg, 0.14 mmol) (Rt = 4.74 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ = 8.21 (d, 2H), 7.85 (s, 1H), 7.76 (d, 2H), 6.30 (br d, 1H), 5.66 (quin, 1H), 4.50 - 4.36 (m, 1H), 2.55 (s, 3H), 1.75 (d, 3H), 1.52 (d, 6H). **LCMS** R_t = 1.24 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_2$ $[M+H]^+$ 408.2, found 408.1.

Example 69. Synthesis of 99



A-148 A-149 A-150 A-151 99

[000462] **A-149:** A mixture of hydroxylamine hydrochloride (1.11 g, 16.03 mmol), 4-(trifluoromethoxy)benzonitrile (1 g, 5.34 mmol) and NaOH (0.64 g, 16.03 mmol) in ethanol (24 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH and then it was diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.48 g, 6.24 mmol) as a solid.**LCMS** R_t = 0.59 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8F_3N_2O_2$ $[M+H]^+$ 221.0, found 221.0.

[000463] **A-150:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (429.73 mg, 2.27 mmol) and CDI (405.1 mg, 2.5 mmol) in DMF (15mL) was stirred at 25 °C for 1 hour and then N'-hydroxy-4-(trifluoromethoxy)benzamidine (500 mg, 2.27 mmol) was


added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (60 mL) and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 50%) to give the product (610 mg, 1.63 mmol, 72 % yield) as a solid.**LCMS** R_t = 1.38 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{10}F_3N_3O_4$ $[M+H\text{-}tBu]^+$ 318.1, found 318.0.

[000464] **A-151:** To tert-butyl N-[(1S)-1-[3-[4-(trifluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (610 mg, 1.63 mmol) was added 4M HCl in 1,4-dioxane (15 mL) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with $NaHCO_3$(s) to pH~8. The mixture was extracted with EtOAc (30 mL x 2), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (510 mg, 1.87 mmol) as an oil.

[000465] **99:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-[4-(trifluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine (194.93 mg, 0.71 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with flash chromatography on silica gel (0% to 20% EtOAc in PE) to give the product. Analytical SFC (Regis (S,S) Whelk-01 (100 × 4.6 mm, 5.0 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40%, for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 3.75 min (18.7%) and 4.71 min (main peak, 81.3%).

[000466] The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 45% B; 7.50 min run; 4 injections, Rt of peak 1 = 4.00 min, Rt of peak 2 = 5.50 min) to give the product (53.39 mg, 0.13 mmol, 21% yield) (Rt = 4.71 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) 400MHz δ = 8.20 - 8.07 (m, 2H), 7.85 (s, 1H), 7.34 (d, 2H), 6.32 (br d, 1H), 5.70 - 5.57 (m, 1H), 4.50 - 4.40 (m, 1H), 2.55 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H).**LCMS** R_t = 1.26 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{20}F_3N_5O_3$ $[M+H]^+$ 424.2, found 424.1.

Example 70. Synthesis of 100



[000467] **A-153:** A mixture of hydroxylamine hydrochloride (1.11 g, 16.03 mmol), 3-(trifluoromethoxy)benzonitrile (1 g, 5.34 mmol) and NaOH (0.64 g, 16.03 mmol) in ethanol (24 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of the EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.2 g, 3.07 mmol, 57% yield) as an oil. **LCMS** R_t = 0.59 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8F_3N_2O_2$ $[M+H]^+$ 221.0, found 221.0.


[000468] **A-154:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (429.73 mg, 2.27 mmol) and CDI (405.1 mg, 2.5 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour and then N'-hydroxy-3-(trifluoromethoxy)benzamidine (500 mg, 2.27 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (60 mL) and extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30%) to give the product (640 mg, 1.71 mmol, 75% yield) as a solid. **LCMS** R_t = 1.38 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{10}F_3N_3O_4$ $[M+H\text{-}tBu]^+$ 318.1, found 317.9.

[000469] **A-155:** To tert-butyl N-[(1S)-1-[3-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (640 mg, 1.71 mmol) was added 4M HCl in 1,4-dioxane (15 mL) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced pressure and diluted with H_2O (20 mL) and basified with

$NaHCO_3$ (solid) to pH~8. The mixture was extracted with EtOAc (30 mL x 2), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (500 mg, 1.83 mmol) as an oil.

[000470] **100:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-[3-(trifluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine (194.93 mg, 0.71 mmol) and the resultant reaction mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product.Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 5.0 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40%, for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 3.71 min (12.9%) and 4.67 min (87.1%).The product as purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 50% B; 7.50 min run; 4 injections, Rt of peak 1 = 3.71 min, Rt of peak 2 = 4.67 min) to give the product (64.7 mg, 0.16 mmol, 25% yield) (Rt = 4.67 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ = 8.12 (d, 2H), 7.85 (s, 1H), 7.33 (d, 2H), 6.33 (br d, 1H), 5.65 (quin, 1H), 4.55 - 4.33 (m, 1H), 2.55 (s, 3H), 1.74 (d, 3H), 1.52 (d, 6H).**LCMS** R_t = 1.26 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}F_3N_5O_3$ $[M+H]^+$ 424.2, found 424.1.

Example 71. Synthesis of 101



[000471] **A-157:** A mixture of hydroxylamine hydrochloride (0.49 g, 7.1 mmol), 3-(difluoromethoxy)benzonitrile (0.4 g, 2.37 mmol) and NaOH (0.28 g, 7.1 mmol) in ethanol (24 mL) was stirred at 40 °C for 16 hours. After cooling to room temperature, the reaction mixture was partially concentrated under reduced pressure to remove most of EtOH and then diluted with H_2O (20 mL). The mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (0.52 g, 2.43 mmol) as a solid. **LCMS** R_t = 0.23 min in in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_9F_2N_2O_2$ $[M+H]^+$ 203.1, found 203.1.


[000472] **A-158:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (486.69 mg, 2.57 mmol) and CDI (458.79 mg, 2.83 mmol) in DMF (15 mL) was stirred at 25 °C for 1 hour and then 3-(difluoromethoxy)-N'-hydroxy-benzamidine (520 mg, 2.57 mmol) was added. The reaction mixture was then stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with water (60 mL) and extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 30%) to give the product (440 mg, 1.23 mmol, 48% yield) as a solid. **LCMS** R_t = 1.38 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{12}F_2N_3O_4$ $[M+H\text{-}tBu]^+$ 300.1, found 299.9.

[000473] **A-160:** To tert-butyl N-[(1S)-1-[3-[3-(difluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (440 mg, 1.24 mmol) was added 4M HCl in 1,4-dioxane (15 mL, 60 mmol) was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced pressure, diluted with H_2O (20 mL) and basified with $NaHCO_3$ (solid) to pH~8. The mixture was extracted with EtOAc (30 mLx 3), and the combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (220 mg, 0.86 mmol) as an oil.

[000474] **101:** To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-[3-(difluoromethoxy)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine (182.09 mg, 0.71 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was

purified with flash chromatography on silica gel (EtOAc in PE =0% to 20%) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 5.0 μm); mobile phase: A: CO_2 B:ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40%, for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min) showed two peaks at 4.29 min (13.4%) and 5.43 min (main peak, 86.6%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 50% B; 8.00 min run; 5 injections, Rt of peak 1 = 4.00 min, Rt of peak 2 = 6.00 min) to give the product (51.8 mg, 0.13 mmol, 21% yield) (Rt = 5.43 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ = 7.96 (s, 1H), 7.92 (d, 1H), 7.78 (s, 1H), 7.58 (t, 1H), 7.34 (dd, 1H), 7.08 - 6.61 (m, 2H), 5.42 (quin, 1H), 4.48 - 4.36 (m, 1H), 2.35 (s, 3H), 1.67 (d, 3H), 1.45 (d, 6H). **LCMS** R_t = 1.13 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{22}F_2N_5O_3$ $[M+H]^+$ 406.2, found406.0.

Example 72. Synthesis of 102




A-138 102

[000475] A mixture of 1-(difluoromethyl)-3-methyl-pyrazole-4-carboxylic acid (91.8 mg, 0.52 mmol), HOBt (156.52 mg, 1.16 mmol), Et_3N (0.4 mL, 2.9 mmol) and EDCI (166.53 mg, 0.87 mmol) in DMF (20 mL) was stirred at 25 °C for 1 hour and then (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (120 mg, 0.58 mmol) was added and the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 to 10% to 25% to 50%) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 3 μm); mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.93 min (9.1%) and 3.38 min (main peak, 90.9%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 25% B; 9 min run; 4 injections, Rt of peak 1 = 6 min, Rt of peak 2 = 7.5 min) to give the product (68.18 mg, 0.18 mmol, 32% yield) (Rt = 3.38 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz,

$CDCl_3$) δ_H = 7.88 (d, 1H), 7.82-7.75 (m, 1H), 7.52 - 7.44 (m, 1H), 7.26 - 7.19 (m, 1H), 6.89 (s, 1H), 6.86 - 6.55 (m, 2H), 5.65 - 5.50 (m, 1H), 4.20 (s, 3H), 1.77 (d, 3H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{15}F_3N_5O_2$ $[M+H]^+$ 366.1, found 366.1.


Example 73. Synthesis of 103



A-135 103

[000476] A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (77.58 mg, 0.4 mmol), HOBt (120.01 mg, 0.89 mmol), Et_3N (0.31 mL, 2.22 mmol) and EDCI (127.69 mg, 0.67 mmol) in DCM (15 mL) was stirred at 25 °C for 1 hours and to the mixture was added (1S)-1-[3-(3,5-difluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine (100 mg, 0.44 mmol) with additional stirring at 25 °C for 16 hours to give a solution. The reaction was quenched with sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 to 10% to 25% to 50%) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.37 min (18.1%) and 2.66 min (main peak, 81.9%). The product was purified by SFC (Regis (S,S) Whelk -O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 20% B; 8 min run; 5 injections, Rt of peak 1 = 5.5 min, Rt of peak 2 = 6.5 min) to give the product (41.14 mg, 0.1 mmol, 23% yield) (Rt = 2.66 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.68 - 7.56 (m, 2H), 7.04 - 6.95 (m, 1H), 6.93 (s, 1H), 6.67 (d, 1H), 5.61 (q, 1H), 4.24 (s, 3H), 1.78 (d, 3H). **LCMS** R_t = 1.31 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}F_5N_5O_2$ $[M+H]^+$ 402.1, found 402.0.


Example 74. Synthesis of 104



[000477] A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (97.86 mg, 0.5 mmol), HOBt (151.39 mg, 1.12 mmol), Et_3N (0.39 mL, 2.8 mmol) and EDCI (161.08 mg, 0.84 mmol) in DCM (15 mL) was stirred at 25 °C for 1 hour and then to the mixture was added 3-[5-[(1S)-1-aminoethyl]-1,2,4-oxadiazol-3-yl]benzonitrile (120 mg, 0.56 mmol) and the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with sat. NH_4Cl (20 mL), and the mixture was extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0 to 10% to 25% to 50%) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm × 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.38 min (5.6%) and 3.76 min (main peak, 94.4%). Note: the condensation reaction might lead to some racemization. The analogue have the risk of racemization under basic condition.

[000478] The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 30% B; 8 min run; 5 injections, Rt of peak 1 = 5.2 min, Rt of peak 2 = 6.5 min) to give the product (50.78 mg, 0.13 mmol, 23% yield) (Rt = 3.76 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.39 (s, 1H), 8.32 (d, , 1H), 7.82 (d, 1H), 7.64 (t, 1H), 6.94 (s, 1H), 6.69 (d, 1H), 5.62 (quin, 1H), 4.24 (s, 3H), 1.79 (d, 3H). **LCMS** R_t = 1.23 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{14}F_3N_6O_2$ $[M+H]^+$ 391.1, found 391.3.

Example 75. Synthesis of 105



[000479] **A-109a:** A mixture of (2S)-2-(tert-butoxycarbonylamino)butanoic acid (263.71 mg, 1.3 mmol), DCC (534.58 mg, 2.6 mmol), 3-fluoro-N-hydroxy-benzamidine (200 mg, 1.3 mmol) in 1,4-dioxane (6 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated NH_4Cl (30 mL), and the mixture was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10%) to give the product (290 mg, 0.89 mmol, 68% yield) as a solid. **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, , MS ESI calcd. for $C_{12}H_{13}FN_3O_3$ $[MH\text{-}tBu+H]^+$ 266.09, found 266.1.


[000480] **A-110a:** To tert-butyl (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propyl]carbamate (290.mg, 0.9000mmol) was added 4M HCl in 1,4-dioxane (10 mL, 40 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated to give the crude product as a solid. **LCMS** R_t = 0.68 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{13}FN_3O$ $[M+H]^+$ 222.10, found 222.1.

[000481] **105:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (105.46 mg, 0.54 mmol), EDCI (104.15 mg, 0.54 mmol) in MeCN (4 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (140 mg, 0.54 mmol) and the reaction mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the crude product. The crude product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the product (25.53 mg, 64.3 µmol, 12% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.91 - 7.86 (m, 1H), 7.81 - 7.76 (m,

1H), 7.52 - 7.44 (m, 1H), 7.26 - 7.20 (m, 1H), 6.94 (s, 1H), 6.66 (br d, 1H), 5.56 - 5.38 (m, 1H), 4.23 (s, 3H), 2.24 - 2.14 (m, 1H), 2.13 - 2.03 (m, 1H), 1.08 (t, 3H). **LCMS** R_t = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{17}H_{16}F_4N_5O_2$ $[M+H]^+$ 398.12, found 398.1.

Example 76. Synthesis of 106 and 107

a) Synthesis of 107



Synthesis of tert-butyl N-[(1S)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-337)

[000482] To a stirred solution of compound **A-336** (300 mg, 1.98 mmol) in 1,4-dioxane (20 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (375 mg, 1.98 mmol) and DCC (449 mg, 2.18 mmol) at room temperature. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 20% EtOAc/PE to afford compound **A-337** (390 mg, 1.27 mmol, 64% yield). **LCMS**: 305.1 (M+H), Rt 1.66 min (Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min)

Synthesis of (1S)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (A-338)


[000483] To a stirred solution of compound **A-337** (390 mg, 1.27 mmol) in DCM (10.0 mL) was added TFA (1.47 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (5

mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-338** (230 mg). The compound was used for the next step without further purification.

Synthesis of 2-methyl-N-[(1S)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (107)

[000484] To a stirred solution of compound **A-338** (230 mg, 1.13 mmol) in THF (10.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (218 mg, 1.13 mmol) followed by T3P (2.01 mL, 3.38 mmol) and Et_3N (0.47mL, 3.38mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **107** (140 mg, 0.36 mmol, 32% yield) as a solid. Prep. HPLC method: Rt 8.86; Column: Atlantis C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.1 min, 99.4% Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 381.2 (M+H), Rt 1.82 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.87 min, SFC column: (R,R)-Whelk-01(250 mm x 4.6 mm, 5 µm); mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.61 (d, 1H), 7.95 (s, 1H), 7.86 (d, 1H), 7.26 (s, 1H), 5.54 (q, 1H), 4.19 (s, 3H), 2.64 (s, 3H), 1.77 (d, 3H).

b) Synthesis of 106



Synthesis of tert-butyl N-[(1R)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-339)

[000485] To a stirred solution of compound **A-336** (400 mg, 2.65 mmol) in 1,4-dioxane (10.0 mL) at room temperaturewas added Boc-D-alanine (600 mg, 3.18 mmol) and DCC (817 mg, 3.97 mmol)and the reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 24% EtOAc/PE to afford compound **A-339** (600 mg, 1.9 mmol, 73% yield). **LCMS**: 305.2 (M+H), Rt 1.65 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min


Synthesis of (1R)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (A-340)

[000486] To a stirred solution of compound **A-339** (600.mg, 1.97 mmol)in DCM (7.0 mL)was added TFA (1.51 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-340** (280 mg). The compound was used for the next step without further purification.

Synthesis of 2-methyl-N-[(1R)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (106)

[000487] To a stirred solution of compound **A-340** (280 mg, 1.37 mmol)in THF (10.0 mL)was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (266 mg, 1.37 mmol)followed by T3P (2.44 mL, 4.11 mmol) and Et_3N (0.57 mL, 4.11 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **106** (272 mg, 0.70 mmol, 51% yield) as a solid. Prep. HPLC method: Rt 11.1; Column: X-Bridge C8 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.09 min, 99.2% Column: XBridge C8 (50 x 4.6 mm, 3.5 µm); Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 381.1 (M+H), Rt 1.79 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm, 3.5 µm); Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.14 min, SFC column: Regis (R,R)-Whelk-O1 (250 mm x 4.6 mm, 5 µm); mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.47 (d, 1H), 8.66 (d, 1H), 7.81 (s, 1H), 7.74-7.73 (dd, 1H), 7.45 (s, 1H), 5.50-5.46 (m, 1H), 4.13 (s, 3H), 2.58 (s, 3H), 1.68 (d, 3H).

Example 77. Synthesis of 108 and 109



[000488] **A-110b:** A mixture of 3-fluoro-N-hydroxy-benzamidine (500 mg, 3.24 mmol) 2-(tert-butoxycarbonylamino)-2-cyclopropyl-acetic acid (698.23 mg, 3.24 mmol) and DCC (1336.45 mg, 6.49 mmol) in 1,4-dioxane (10 mL) was stirred at 100 °C for 16 hours. After

cooling to room temperature, the mixture was concentrated under reduced pressure to give a residue. The residue was diluted with H_2O (20 mL) and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30%) to give the product (800 mg, 2.39 mmol, 74% yield) as an oil. **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{13}FN_3O$ [M+H- *t*Bu] 278.1, found 278.1.

[000489] **A-111c:** To tert-butyl N-[cyclopropyl-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]methyl]carbamate (800 mg, 2.4 mmol) was added 4M HCl in 1,4-dioxane (8 mL, 32 mmol) an the reaction mixture was stirred 40 °C at for 16 hours. The reaction mixture was concentrated under reduced pressure and diluted with EtOAc (20 mL) and H_2O (20 mL). The EtOAc phase was discarded and the pH of the aqueous phase was basified with NaOH (solid) to pH = 9 and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over anhydrous Na_2SO_4 to give the crude product (500 mg, 1.90 mmol, 79% yield) as an oil. **LCMS** R_t = 0.64 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{13}FN_3O$ [M+H] 234.1, found 233.8.

[000490] **A-112c:** A mixture of 1-cyclopropyl-3-methyl-pyrazole-4-carboxylic acid (150 mg, 0.90 mmol), EDCI (346.07 mg, 1.81 mmol), DIPEA (0.47 mL, 2.71 mmol), HOBt (243.95 mg, 1.81 mmol) and cyclopropyl-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]methanamine (210.53 mg, 0.90 mmol) in DCM (2 mL) was stirred at 20 °C for 16 hours. The residue was diluted with sat. NH_4Cl (20 mL) and extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5% to 60%) to give the product (400 mg, 0.95 mmol) as a solid. **LCMS** R_t = 0.88 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{20}H_{21}FN_5O_2$ $[M+H]^+$ 382.2, found 382.0.


[000491] **108 & 109:** The product was analyzed by SFC (Diacel CHIRALPAK AD-3 (50 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 2 min and hold 40% for 1.2 min, then 5% of B for 0.8 mim, flow rate: 4 mL/min, column temp: 35 °C) to show two peaks at 0.98 min and 1.31 min. The product was separated by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 35 °C; 30 mL/min; 30% B; 10 min run; 9 injections, Rt of peak 1 = 7.5 min, Rt of peak 2 = 8.44 min) to give the enantiomer 1, randomly assigned as **108** (104.77 mg, 0.3 mmol, 26% yield) (Rt = 0.98 min in analytica SFC) as an oil, and the enantiomer 2,

randomly assigned as **109** (103.84 mg, 0.3 mmol, 25% yield) (Rt = 1.30 min in analytical SFC) as an oil.

[000492] **108: ^{1}H NMR** (400MHz, $CDCl_3$)δ_H = 7.92 - 7.85 (m, 1H), 7.82 - 7.75 (m, 1H), 7.51 - 7.42 (m, 1H), 7.26 - 7.18 (m, 1H), 6.64 (d, 1H), 6.35 (s, 1H), 4.96 - 4.86 (m, 1H), 4.07 (s, 3H), 1.98 - 1.89 (m, 1H), 1.42 - 1.32 (m, 1H), 0.99 - 0.92 (m, 2H), 0.82 - 0.58 (m, 6H). **LCMS** R_t = 1.26 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{21}FN_5O_2$ $[M+H]^+$ 382.2, found 382.1.

[000493] **109: ^{1}H NMR** (400MHz, $CDCl_3$)δ_H = 7.92 - 7.85 (m, 1H), 7.82 - 7.76 (m, 1H), 7.52 - 7.42 (m, 1H), 7.26 - 7.18 (m, 1H), 6.65 (d, 1H), 6.35 (s, 1H), 4.96 - 4.85 (m, 1H), 4.07 (s, 3H), 1.99 - 1.88 (m, 1H), 1.43 - 1.32 (m, 1H), 0.99 - 0.91 (m, 2H), 0.79 - 0.58 (m, 6H). **LCMS** R_t = 1.25 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{21}FN_5O_2$ $[M+H]^+$ 382.2, found 382.1.

Example 78. Synthesis of 110



[000494] **A-110d:** DAST (8 mL, 60.05 mmol) was added dropwise to a solution of 3-formylbenzonitrile (3 g, 22.88 mmol) in DCM (30 mL). The resulting mixture was stirred at 20 °C for 16 hours to give a solution. The reaction mixture was added dropwise into saturated $NaHCO_3$ aqueous (200 mL) and then extracted with DCM (100 mL x 2). The organic phase was washed with brine (100 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (DCM in PE = 0% to 2%) to give the product (2.4 g, 15.67 mmol, 68% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.84 - 7.73 (m, 3H), 7.65 - 7.57 (m, 1H), 6.85 - 6.51 (m, 1H).

[000495] **A-111d:** A mixture of 3-fluorobenzonitrile (2.4 g, 19.82 mmol), hydroxylamine hydrochloride (4.13 g, 59.45 mmol) and NaOH (2.38 g, 59.45 mmol) in

ethanol (45 mL) and water (15 mL) was stirred at 40 °C for 16 hours to give a mixture. After cooling to room temperature, it was concentrated under reduced pressure and then water (50 mL) and EtOAc (50 mL) were added. The phases were separated, and the organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.79 (s, 1H), 7.76 (d, 1H), 7.61 - 7.55 (m, 1H), 7.54 - 7.47 (m, 1H), 6.83 - 6.50 (m, 1H), 4.95 (s, 2H).


[000496] **A-112d:** A mixture of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (3.05 g, 16.12 mmol) and CDI (2.87 g, 17.73 mmol) in DMF (30 mL) was stirred at 25 °C for 1 hour. To the mixture was added 3-(difluoromethyl)-N-hydroxy-benzamidine (3 g, 16.1 2mmol) and the resulting mixture was stirred at 70 °C for 16 hours. After cooling to room temperature, water (50 mL) was added and the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the product (1.8 g, 5.30 mmol, 33% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.26 - 8.18 (m, 2H), 7.71 - 7.65 (m, 1H), 7.63 - 7.57 (m, 1H), 6.87 - 6.56 (m, 1H), 5.19 (s, 2H), 1.68 - 1.63 (m, 3H), 1.48 (s, 9H).

[000497] **A-113d:** To tert-butyl N-[(1S)-1-[3-[3-(difluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (1.8 g, 5.3 mmol) was added 4M HCl in 1,4-dioxane (20 mL, 5.3 mmol) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated under reduced pressure to give the crude product (1.4 g) as a solid. **LCMS** R_t = 0.39 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{11}H_{12}F_2N_3O$ $[M\text{-}NH2]^+$ 240.1, found 239.9.

[000498] **110:** A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (152.52 mg, 0.91 mmol), HATU (689.62 mg, 1.81 mmol), DIPEA (0.38 mL, 2.72 mmol) and then (1S)-1-[3-[3-(difluoromethyl)phenyl]-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (250 mg, 0.91 mmol) in MeCN (15 mL) was stirred at 25 °C for 16 hours. The reaction mixture was concentrated under reduced pressure and water (20 mL) was added to the residue and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm, 5.0 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min , then 5% of B, for 1.5 min, flow rate: 2.5 mL/min,

column temp: 40 °C, ABPR: 100 bar) showed two peaks at 4.36 min (8.8%) and 5.50 min (91.2%). The impure product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 38 °C; 50 mL/min; 45% B; 8 min run; 10 injections, Rt of peak 2 = 7.2 min) to give the product (93.5 mg, 240.1µmol, 26% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) 400MHz δ_H = 8.58 (d, 1H), 8.29 (s, 1H), 8.21 - 8.14 (m, 2H), 7.84 - 7.78 (m, 1H), 7.77 - 7.70 (m, 1H), 7.18 (t, 1H), 5.43 - 5.33 (m, 1H), 4.47 - 4.36 (m, 1H), 2.30 (s, 3H), 1.63 (d, 3H), 1.41 (d, 6H). **LCMS** R_t = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{22}F_2N_5O_2$ $[M+H]^+$ 390.2, found 390.1.


Example 79. Synthesis of 111




A-110e 111

[000499] To a mixture of 2-(2,2-difluoroethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (210 mg, 0.86 mmol), EDCI (164.91 mg, 0.86 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (221.67 mg, 0.86 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with 1N HCl (10 mL) and then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with water (15 mL) and brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by prep-TLC (silica gel, PE:EtOAc = 5:1) to give the impure product. The impure product was triturated from *n*-hexane/DCM (10:1, 10 mL) to give the product (98.1 mg, 0.22 mmol, 25% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) 400MHz δ_H = 9.54 (d, 1H), 7.86 (d, 1H), 7.74 (d, 1H), 7.68 - 7.60 (m, 1H), 7.56 (s, 1H), 7.47 (dt, 1H), 6.40 (tt, 1H), 5.35 - 5.25 (m, 1H), 5.09 (dt, 2H), 2.19 - 1.98 (m, 2H), 1.02 (t, 3H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_6N_5O_2$ $[M+H]^+$ 448.1, found 448.1.

Example 80. Synthesis of 112



A-71a A-111e A-94



112


[000500] **A-111e:** To a solution of ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (1000 mg, 4.8 mmol) in MeCN (20 mL) was added Cs_2CO_3 (3130.59 mg, 9.61 mmol), sodium 2-chloro-2,2-difluoro-acetate (1464.98 mg, 9.61 mmol) and 18-crown-6 (253.98 mg, 0.96 mmol). The reaction mixtire was stirred at 90 °C for 1.5 hours. After cooling to room temperature, the mixture was diluted with H_2O (30 mL), and then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 15%) to give the product (140 mg, 437.4 µmol, 10% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.29 - 7.89 (m, 1H), 7.21 (s, 1H), 4.44 (q, H), 1.43 (t, 3H). **LCMS** R_t = 4.45 min in 7.0 min chromatography, 0-60AB, MS ESI calcd. for $C_8H_8F_5N_2O_2$ $[M+H]^+$ 259.04, found 258.9.

[000501] **A-94:** To a solution of ethyl 2-(difluoromethyl)-5-(trifluoromethyl)pyrazole-3-carboxylate (140 mg, 0.54 mmol) in ethanol (4 mL) was added a solution of NaOH (65.08 mg, 1.63 mmol) in water (4mL). The mixture was stirred at 20 °C for 2 hours. The reaction mixture was quenched by addition of 1N HCl (15 mL) and diluted with H_2O (15 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the product (100 mg) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.39 (t, 1H), 7.45 (s, 1H).

[000502] **112:** To a mixture of 2-(difluoromethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (100 mg, 0.43 mmol), EDCI (104.15 mg, 0.54 mmol) in MeCN (4 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (167.99 mg, 0.65 mmol) and the mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated

to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the product (4.5 mg, 10 μmol, 2% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.13 (t, 1H), 7.88 (d, 1H), 7.78 (d, 1H), 7.54 - 7.44 (m, 1H), 7.25 - 7.19 (m, 1H), 7.09 (s, 1H), 6.86 - 6.75 (m, 1H), 5.63 - 5.41 (m, 1H), 2.28 - 2.16 (m, 1H), 2.15 - 2.03 (m, 1H), 1.09 (t, 3H). **LCMS** R_t = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{17}H_{14}F_6N_5O_2$ $[M+H]^+$ 434.10, found 433.9.

Example 81. Synthesis of 113




[000503] **A-113e:** To a solution of ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (600 mg, 2.88 mmol), cyclopropylboronic acid (495.24 mg, 5.77 mmol) and Na_2CO_3 (611.07 mg, 5.77 mmol) in DCE (7 mL) was added a solution of $Cu(OAc)_2$ (523.58 mg, 2.88 mmol) and 2,2-bipyridine (540.29 mg, 3.46 mmol) in DCE (14 mL). The reaction mixture was stirred at 70 °C for 4 hours. After cooling to room temperature, the mixture was diluted with sat. NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (40 mL) and brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5% to 50%) to give the product (180 mg, 0.73 mmol, 25% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.07 (s, 1H), 4.40 (q, 3H), 1.41 (t, 3H), 1.36 - 1.30 (m, 2H), 1.14 - 1.07 (m, 2H). **LCMS** R_t = 0.94 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{12}F_3N_2O_2$ $[M+H]^+$ 249.1, found 249.1.

[000504] **A-114a:** To a solution of ethyl 2-cyclopropyl-5-(trifluoromethyl)pyrazole-3-carboxylate (180 mg, 0.73 mmol) in ethanol (3 mL) was added a solution of NaOH (87.03 mg, 2.18 mmol) in water (3 mL). The reaction mixture was stirred at 50 °C for 2 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (20 mL) and

extracted with EtOAc (20 mL). The pH of the aqueous phase was adjusted with the addition of 1 N HCl to pH ~ 2, and it was then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with water (50 mL) and brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the product (140 mg) as a solid. **LCMS** R_t = 0.81 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_8F_3N_2O_2$ $[M+H]^+$ 221.0, found 221.1.

[000505] **113:** To a mixture of 2-cyclopropyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (140 mg, 0.64 mmol), EDCI (121.91 mg, 0.64 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (163.87 mg, 0.64 mmol), the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with 1 N HCl (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic phase was washed with water (15 mL) and brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by prep-TLC (silica gel, PE:EtOAc = 5:1) to give the impure product. The impure product was triturated from *n*-hexane:DCM (10:1, 6 mL) to give the product (34 mg, 80.3 μmol, 13% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.43 (d, 1H), 7.87 (d, 1H), 7.78 - 7.71 (m, 1H), 7.64 (dt, 1H), 7.48 (dt, 1H), 7.41 (s, 1H), 5.34 - 5.26 (m, 1H), 4.45 - 4.34(m, 1H), 2.19 - 1.98 (m, 2H), 1.16 - 0.97 (m, 7H). **LCMS** R_t = 1.40 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_2$ $[M+H]^+$ 424.1, found 424.1.

Example 82. Synthesis of 116



A-117a A-118a

A-119a 116


[000506] **A-118a:** A mixture of 3-fluoro-N-hydroxy-benzamidine (5 g, 32.44 mmol), 2-(tert-butoxycarbonylamino)-4-methoxy-4-oxo-butanoic acid (8.02g, 32.44mmol) and DCC (13.36 g, 64.88 mmol) in 1,4-dioxane (150 mL) was stirred at 70 °C for 16 hours. The

mixture was cooled to room temperature and concentrated under reduced pressure and then EtOAc (250 mL) and water (150 mL) were added to the residue and the mixture was filtered. After the phases were separated, the organic phase was washed with brine (150 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the product (9 g, 24.63 mmol, 75% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.86 (d, 1H), 7.79 - 7.72 (m, 1H), 7.48 - 7.40 (m, 1H), 7.23 - 7.16 (m, 1H), 5.82 (d, 1H), 5.45 - 5.36 (m, 1H), 3.70 (s, 3H), 3.29 - 3.18 (m, 1H), 3.12 - 3.03 (m, 1H), 1.48 (s, 9H).

[000507] **A-119a:** To methyl 3-(tert-butoxycarbonylamino)-3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propanoate (9 g, 24.63 mmol) was added 4M HCl in 1,4-dioxane (50 mL, 24.63 mmol) and the reaction mixture was stirred at 20 °C for 2 hour to give a solution. The solution was concentrated under reduced pressure to give the crude product (7.4 g) as a solid. **LCMS** R_t = 0.65 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{13}FN_3O_3$ $[M+H]^+$ 332.2, found 266.1.

[000508] **116:** A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (4.76 g, 24.53 mmol), HATU (13.99 g, 36.79 mmol), DIPEA (6.82 mL, 49.06 mmol) and methyl 3-amino-3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propanoate hydrochloride (7.4 g, 24.53 mmol) in DMF (30 mL) was stirred at 20 °C for 3 hours to give a solution. To the mixture was added water (100 mL) and the mixture was extracted with EtOAc (80 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 30%) to give the product (7.7 g, 17.44 mmol) as a solid. The crude product (300 mg) was triturated from DCM (10 mL) and n-hexane (10 mL) at 65 °C to give the product (238.05 mg, 534.6 µmol, 78% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (d, 1H), 7.78 - 7.75 (m, 1H), 7.55 (d, 1H), 7.49 - 7.44 (m, 1H), 7.25 - 7.20 (m, 1H), 6.94 (s, 1H), 5.86 - 5.81 (m, 1H), 4.26 (s, 3H), 3.76 (s, 3H), 3.40 (dd, 1H), 3.14 (dd, , 1H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_4N_5O_4$ $[M+H]^+$ 442.1, found 442.1.

Example 83. Synthesis of 114 and 115



[000509] **A-115:** A mixture of methyl 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoate (3 g, 6.8 mmol) and LiOH.H_2O (570.44 mg, 13.59 mmol) in THF (10 mL) and water (5 mL) was stirred at 20 °C for 16 hours. To the mixture was added 1N HCl (50 mL) to adjust the pH = 2 and then extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 12.59 (br s, 1H), 9.51 (d, 1H), 7.86 (d, 1H), 7.78 - 7.71 (m, 1H), 7.64 (dt, 1H), 7.53 - 7.43 (m, 1H), 7.40 (s, 1H), 5.67 (q, 1H), 4.13 (s, 3H), 3.23 (dd, 1H), 3.11 (dd, 1H).

[000510] **A-116:** To a solution of 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoic acid (500 mg, 1.17 mmol) and HATU (667.37 mg, 1.7 6mmol) in DMF (10 mL) was added $MeNH_2$/THF (2M, 0.59 mL, 1.17 mmol) and DIPEA (0.49 mL, 3.51 mmol). The mixture was stirred at 20 °C for 16 hours. The reaction was quenched with H_2O (20 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20% to 60%) to give the product (200 mg, 425.7 µmol, 36% yield) as a solid. **LCMS** R_t = 0.85 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{18}H_{17}F_4N_6O_3$ $[M+H]^+$ 441.12, found 441.2.


[000511] **114 & 115:** Analytical SFC (Daicel CHIRALPAK AD-3 (150 mm x 4.6 mm I.D., 3 µm), mobile phase: A: CO_2 B: methanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C,ABPR: 1500 psi) showed two peaks at 3.45 min and 4.34 min. The product was separated by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% DEA); 38 °C; 50 mL/min; 30% B; 8 min run; 11 injections, Rt of

peak 1 = 4.1 min, Rt of peak 2 = 5.3 min) to give the enantiomer 1, randomly assigned as **114** (50.1 mg, 113.8 µmol, 25% yield) (Rt = 3.45 min in analytical SFC) as a solid, and the enantiomer 2, randomly assigned as **115** (51.5 mg, 117 µmol, 25% yield) (Rt = 4.34 min in analytical SFC) as a solid.

[000512] **114: ^{1}H NMR** (400MHz $CDCl_3$) δ_H = 8.46 (br d, 1H), 7.90 - 7.82 (m, 1H), 7.78 - 7.72 (m, 1H), 7.50 - 7.41 (m, 1H), 7.25 - 7.16 (m, 1H), 7.00 (s, 1H), 5.87 - 5.77 (m, 1H), 5.71 (br d, 1H), 4.26 (s, 3H), 3.19 (dd, 1H), 3.03 - 2.95 (m, 1H), 2.83 (d, 3H). **LCMS** Rt = 1.15 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_4N_6O_3$ $[M+H]^+$ 441.1, found 441.1.

[000513] **115: ^{1}H NMR** (400MHz $CDCl_3$) δ_H = 8.46 (br d, 1H), 7.90 - 7.81 (m, 1H), 7.77 - 7.72 (m, 1H), 7.49 - 7.41 (m, 1H), 7.24 - 7.16 (m, 1H), 7.00 (s, 1H), 5.87 - 5.77 (m, 1H), 5.71 (br d, 1H), 4.26 (s, 3H), 3.19 (dd, 1H), 3.03 - 2.95 (m, 1H), 2.83 (d, 3H). **LCMS** Rt = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_4N_6O_3$ $[M+H]^+$ 441.1, found 441.1.

Example 84. Synthesis of 117




[000514] To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (159.32 mg, 0.82 mmol), EDCI (157.34 mg, 0.82 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (200 mg, 0.82 mmol) and the reaction mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic phase was washed with water (15 mL) and brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) after flash chromatography on silica gel (EtOAc in PE = 0% to 40%) to give the impure product.

[000515] Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm x 4.6 mm I.D., 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.25 min (96.2%) and 2.70 min (3.8%). The impure product was purified by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5

μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 15% B; 8 min run; 6 injections, Rt of peak 1 = 4.2 min, Rt of peak 2 = 6.5 min) to give the product (59.08 mg, 0.15 mmol, 67% yield) (Rt = 2.25 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.46 (d, 1H), 7.86 (d, 1H), 7.79 - 7.72 (m, 1H), 7.67 - 7.58 (m, 1H), 7.51 - 7.43 (m, 2H), 5.46 (quin, 1H), 4.13 (s, 3H), 1.67 (d, 3H). **LCMS** R_t = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{14}F_4N_5O_2$ $[M+H]^+$ 384.1, found 383.9.

Example 85. Synthesis of 118




[000516] **A-140a:** A mixture of 3-fluoro-N-hydroxy-benzamidine (251.38 mg, 1.63 mmol) and DCC (671.91 mg, 3.26 mmol), 4-(tert-butoxycarbonylamino)tetrahydropyran-4-carboxylic acid (400 mg, 1.63 mmol) in 1,4-dioxane (5 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was filtered and diluted with EtOAc (10 mL). To the combined organic layer was added saturatred NH_4Cl (30 mL) and it was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10%) to give the product (400 mg, 1.09 mmol, 66% yield) as a solid. **LCMS** R_t = 0.9 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{18}H_{23}FN_3O_4$ $[M+H-56]^+$364.16, found 308.1.

[000517] **A-141a:** To tert-butyl N-[4-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]tetrahydropyran-4-yl]carbamate (200 mg, 0.54 mmol) was added 4M HCl in 1,4-dioxane (10 mL, 0.54 mmol) and the reaction mixture was stirred at 25 °C for 2 hours. The mixture was concerned under reduced pressure to give the crude product (130 mg, 0.49 mmol,) as a solid. **LCMS** R_t = 0.65 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{13}H_{15}FN_3O_2$ $[M+H]^+$264.1, found 264.1.

[000518] **118:** To a solution of 4-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]tetrahydropyran-4-amine hydrochloride (130 mg, 0.49 mmol), 2-methyl-5-

(trifluoromethyl)pyrazole-3-carboxylic acid (105.43 mg, 0.54 mmol) and HATU (375.51 mg, 0.99 mmol) in DMF (2 mL) was added TEA (0.21 mL, 1.48 mmol). The mixture was stirred at 25 °C for 16 hours. The mixture was diluted with H_2O (10 mL) and extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% $NH_3.H_2O$) and B = CH_3CN; 50-80% B over 9 min) to give the product (159.97 mg, 0.36 mmol, 73% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 (d, 1H), 7.80 (td, 1H), 7.47 (dt, 1H), 7.22 (dt, 1H), 6.92 (s, 1H), 6.38 (s, 1H), 4.12 (s, 3H), 4.01 - 3.81 (m, 4H), 2.72 - 2.53 (m, 2H), 2.43 - 2.25 (m, 2H). **LCMS** R_t = 1.3 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$440.1, found 440.2.

Example 86. Synthesis of 119




A-142a A-143a A-144a 119

[000519] **A-143a:** CDI (298.59 mg, 1.84 mmol) was added to a stirred solution of 3-(tert-butoxycarbonylamino)oxetane-3-carboxylic acid (200 mg, 0.92 mmol) in DMF (10 mL). After 40 min, 3-fluoro-N-hydroxy-benzamidine (283.84 mg, 1.84 mmol) was added in one portion and the resulting solution was stirred for 30 min before it was heated to 100 °C and stirred for 16 hours. After cooling to room temperature, the reaction was diluted with H_2O (40 mL), then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with flash chromatography on silica gel (EtOAc in PE = 0% to 5%) to give the product (180 mg, 0.54 mmol, 58% yield) as an oil. **LCMS** R_t = 0..88 min in 1.5 min chromatography, 5-95% AB, MS ESI calcd. for $C_{16}H_{19}FN_3O_4$ $[M+H\text{-}tBu]^+$ 280.1, found 279.7.

[000520] **A-144a:** To the solution of tert-butyl N-[3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]oxetan-3-yl]carbamate (120 mg, 0.36 mmol) in DCM (2 mL) was added TFA (0.5 mL, 6.83 mmol) and the resulting solution was stirred at 25 °C for 30 min. The reaction mixture was concentrated under reduced pressure to give the crude product (150 mg, 0.40 mmol) as an oil. **LCMS** R_t = 0.87 min in 2.0 min chromatography, 10-80% AB, MS ESI calcd. for $C_{13}H_{12}F_4N_3O_4$ $[M+H]^+$ 236.1, found 235.9.

[000521] **119:** A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (100 mg, 0.52 mmol), HOBt (139.23 mg, 1.03 mmol), EDCI (197.52 mg, 1.03 mmol), Et_3N (0.29 mL, 2.06 mmol), 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]oxetan-3-amine 2,2,2-trifluoroacetic acid (150 mg, 0.43 mmol) in DCM (10 mL) was stirred at 25 °C for 16 hours. The reaction mixture was concentrated and then diluted with H_2O (20 mL) and extracted with DCM (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with prep-HPLC (Boston Prime C18 (150 mm × 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 45-75% B over 9 min) to give the product (62.95 mg, 0.15 mmol, 30% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ = 10.25 (s, 1H), 7.90 (d, 1H), 7.84 - 7.75 (m, 1H), 7.66 (dt, 1H), 7.58 - 7.44 (m, 2H), 5.17 (d, 2H), 5.00 (d, 2H), 4.09 (s, 3H). **LCMS:** R_t = 1.22 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{14}F_4N_5O_3$ $[M+H]^+$ 421.1, found 411.6.

Example 87. Synthesis of 120




A-145a **120**

[000522] To a mixture of 5-(difluoromethyl)-2-methyl-pyrazole-3-carboxylic acid (150.65 mg, 0.86 mmol), EDCI (148.78 mg, 0.78 mmol) in MeCN (4 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (200 mg, 0.78 mmol), and the mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give an impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the product (32.2 mg, 84.9 μmol,

10% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 (d, 1H), 7.82 - 7.77 (m, 1H), 7.52 - 7.43 (m, 1H), 7.26 - 7.20 (m, 1H), 6.90 (s, 1H), 6.86 - 6.55 (m, 2H), 5.58 - 5.41 (m, 1H), 4.20 (s, 3H), 2.25 - 2.13 (m, 1H), 2.09 (s, 1H), 1.08 (t, 3H). **LCMS** R_t = 1.19 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{17}H_{17}F_3N_5O_2$ $[M+H]^+$ 380.13, found 379.9.

Example 88. Synthesis of 121 and 122



[000523] **A-146a:** A mixture of 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoic acid (300 mg, 0.70 mmol), HATU (400.42 mg, 1.05 mmol), DIPEA (0.29 mL, 2.11 mmol), and 2-aminoethanol (57.25 mg, 0.94 mmol) in DMF (10 mL) was stirred at 20 °C for 16 hours. The residue was diluted with sat. NH_4Cl (20 mL) and extracted with EtOAc (20 mL). The combined organic phase was washed with brine (15 mL x 2), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50% to 90%) to give the product (120 mg, 0.25 mmol, 36% yield) as a solid. **LCMS** R_t = 0.81 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{19}H_{19}F_4N_6O_4$ $[M+H]$ 471.1, found 471.0.


[000524] **121 & 122:** Analytical SFC (Daicel CHIRALPAK AS-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.49 min and 2.93 min. The product was separated by SFC (Daicel CHIRALPAK AS-H(250 mm x 30 mm, 5 µm); A = CO_2 and B = ETOH (0.1% NH_3H_2O) ; 38 °C; 70 mL/min; 30% B; 5 min run; 14 injections, Rt of peak 1 = 2.4 min, Rt of peak 2 = 3.0 min) to give the enantiomer 1, randomly assigned as **120** (10.72 mg, 0.02 mmol, 8% yield) (Rt = 2.49 min in analytical SFC) as a solid, and the enantiomer 2,

randomly assigned as **121** (11.6 mg, 0.02 mmol, 9% yield) (Rt = 2.93 min in analytical SFC) as a solid.

[000525] **121: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.49 (d, 1H), 8.20 - 8.12 (m, 1H), 7.85 (d, 1H), 7.77 - 7.70 (m, 1H), 7.68 - 7.60 (m, 1H), 7.52 - 7.43 (m, 1H), 7.41 (s, 1H), 5.77 - 5.61 (m, 1H), 4.66 (t, 1H), 4.12 (s, 3H), 3.42 - 3.35 (m, 2H), 3.17 - 3.05 (m, 3H), 2.98 - 2.88 (m, 1H). **LCMS** R_t = 1.08 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{19}F_4N_6O_4$ $[M+H]^+$ 471.1, found 471.0.

[000526] **122: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.48 (d, 1H), 8.19 - 8.11 (m, 1H), 7.85 (d, H), 7.76 - 7.70 (m, 1H), 7.68 - 7.59 (m, 1H), 7.51 - 7.43 (m, 1H), 7.40 (s, 1H), 5.72 - 5.64 (m, 1H), 4.65 (t, 1H), 4.12 (s, 3H), 3.40 - 3.34 (m, 2H), 3.15 - 3.05 (m, 3H), 2.97 - 2.89 (m, 1H). **LCMS** R_t = 1.07 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{19}F_4N_6O_4$ $[M+H]^+$ 471.1, found 471.0.

Example 89. Synthesis of 123 and 124



[000527] **A-147a:** A mixture of 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoic acid (600 mg, 1.4 mmol), HATU (800.84 mg, 2.11 mmol), DIPEA (0.59 mL, 4.21 mmol) and azetidin-3-ol hydrochloride (461.47 mg, 4.21 mmol) in DMF (5 mL) was stirred at 25 °C for 4 hours. The reaction was quenched with sat. NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (30 mL) and brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give an impure product. The impure product was triturated from *n*-hexane/DCM (10: 1, 11 mL) to give the product (210 mg, 0.42 mmol,


30% yield) as a solid. **LCMS** R_t = 0.82 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{20}H_{19}F_4N_6O_4$ $[M+H]^+$ 483.1, found 483.0.

[000528] **123 & 124:** Analytical SFC (Daicel CHIRALPAK AS-3 (150 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA) gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.42 min and 3.82 min. The product was separated by SFC (Daicel CHIRALPAK AS-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38 °C; 60 mL/min; 35% B; 8 min run; 4 injections, Rt of peak 1 = 2.52 min, Rt of peak 2= 6.45 min) to give the enantiomer 1, randomly assigned as **123** (57.28 mg, 118.7 μmol, 27% yield) (Rt = 2.42 min in analytical SFC) as a solid, and the enantiomer 2, randomly assigned as **124** (81.28 mg, 164.8 μmol, 37% yield) (Rt = 3.82 min in analytical SFC) as a solid.

[000529] **123: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.47 (br t, 1H), 7.91 - 7.84 (m, 1H), 7.83 - 7.72 (m, 1H), 7.50 - 7.43 (m, 1H), 7.25 - 7.18 (m, 1H), 6.97 (d, 1H), 5.82 - 5.75 (m, 1H), 4.78 - 4.70 (m, 1H), 4.52 - 4.40 (m, 1H), 4.30 - 4.24 (m, 4H), 4.22 - 4.07 (m, 1H), 3.95 - 3.84 (m, 1H), 3.27 - 3.18 (m, 1H), 2.86 - 2.74 (m, 1H), 2.34 (m, 1H). **LCMS** R_t = 1.15 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{19}F_4N_6O_4$ $[M+H]^+$ 483.1.1, found 483.1.

[000530] **124: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.47 (br t, 1H), 7.91 - 7.84 (m, 1H), 7.83 - 7.73 (m, 1H), 7.50 - 7.42 (m, 1H), 7.25 - 7.19 (m, 1H), 6.97 (d, 1H), 5.82 - 5.76 (m, 1H), 4.78 - 4.70 (m, 1H), 4.52 - 4.41 (m, 1H), 4.30 - 4.24 (m, 4H), 4.22 - 4.07 (m, 1H), 3.94 - 3.83 (m, 1H), 3.26 - 3.18 (m, 1H), 2.86 - 2.74 (m, 1H), 2.31 - 2.25 (m, 1H). **LCMS** R_t = 1.13 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{19}F_4N_6O_4$ $[M+H]^+$ 483.1.1, found 483.1.

Example 90. Synthesis of 125 and 126




[000531] **A-119c:** A mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (315.26 mg, 1.62 mmol), HATU (926.34 mg, 2.44 mmol), DIPEA (0.45 mL, 3.25 mmol) and methyl 3-amino-3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]propanoate hydrochloride (490 mg, 1.62 mmol) in DMF (30 mL) was stirred at 20 °C for 3 hours. To the mixture was added

water (50 mL) and then it was extracted with EtOAc (80 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10% to 30%) to give the product. Analytical SFC (Daicel CHIRALCEL OJ-H (150 mm x 4.6 mm, 5 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 2.65 min and 2.78 min. The product was separated by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 15% B; 8 min run; 41 injections, Rt of peak 1 = 5.21 min, Rt of peak 2 = 5.69 min) to give the enantiomer 1 (Rt = 2.65 min in analytical SFC, 87% ee) and the enantiomer 2 (Rt = 2.78 min in analytical SFC, 65% ee). The impure enantiomer 1 was purified again by SFC (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 15% B; 8 min run; 22 injections, Rt of Peak 1 = 5.38 min) to give the enantiomer 1, randomly assigned as **125** (15.5 mg, 34.6 µmol, 2% yield) (Rt = 2.65 min in analytical SFC) as a solid. The impure enantiomer 2 was purified again by SFC two times (Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 15% B; 8 min run; 20 injections, Rt of peak 2 = 5.71 min. Daicel CHIRALCEL OJ-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 15% B; 8 min run; 10 injections, Rt of peak 2 = 6.14 min) to give the enantiomer 2, randomly assigned as **126** (31.41 mg, 68.8 µmol, 4% yield) (Rt = 2.78 min in analytical SFC) as a solid.

[000532] **125: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (d, 1H), 7.81 - 7.74 (m, 1H), 7.55 (d, 1H), 7.50 - 7.44 (m, 1H), 7.26 - 7.20 (m, 1H), 6.94 (s, 1H), 5.86 - 5.81 (m, 1H), 4.27 (s, 3H), 3.76 (s, 3H), 3.40 (dd, 1H), 3.14 (dd, 1H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_4N_5O_4$ $[M+H]^+$ 442.1, found 442.2.

[000533] **126: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (d, 1H), 7.79 - 7.75 (m, 1H), 7.55 (d, 1H), 7.49 - 7.44 (m, 1H), 7.26 - 7.20 (m, 1H), 6.94 (s, 1H), 5.86 - 5.81 (m, 1H), 4.26 (s, 3H), 3.76 (s, 3H), 3.40 (dd, 1H), 3.14 (dd, 1H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_4N_5O_4$ $[M+H]^+$ 442.1, found 442.1.


Example 91. Synthesis of 127



A-94 127

[000534] To a mixture of 2-(difluoromethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (120 mg, 0.52 mmol), EDCI (99.98 mg, 0.52 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (134.39 mg, 0.55 mmol) and the mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with water (15 mL) and brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30%) to give an impure product. The impure product was triturated with *n*-hexane/DCM (10:1, 5 mL) to give the product (44.1 mg, 0.1 mmol, 20% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.81 (d, 1H), 8.51 - 8.17 (m, 1H), 7.87 (d, 1H), 7.78 - 7.72 (m, 1H), 7.70 - 7.61 (m, 2H), 7.48 (dt, 1H), 5.54 - 5.45 (m, 1H), 1.69 (d, 3H). **LCMS** R_t = 1.29 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{12}F_6N_5O_2$ $[M+H]^+$ 420.1, found 420.1.

Example 92. Synthesis of 128 and 129



116 128 129

[000535] A solution of methyl 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoate (500 mg, 1.13 mmol) and $NH_3.H_2O$ (5 mL, 1.13 mmol) in methanol (5 mL) was stirred at 20 °C for 16 hours. To the mixture was added water (20 mL) and it was then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product.


[000536] Analytical SFC (Daicel CHIRALPAK AS-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.66 min and 3.38 min. The product was separated by SFC (Daicel CHIRALPAK AS-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 35 °C; 60 mL/min; 35% B; 8 min run; 6 injections, Rt of peak 1 = 2.88 min, Rt of peak 2 = 4.66 min) to give the enantiomer 1, randomly assigned as **128** (74.06 mg, 171.7 µmol, 15% yield) (Rt = 2.66 min in analytical SFC) as a solid, and the enantiomer 2 (Rt = 3.38 min in analytical SFC, 93.1% ee).

[000537] The impure enantiomer 2 was purified again by SFC (Daicel CHIRALPAK AS-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH; 35 °C; 60 mL/min; 25% B; 8 min run; 6 injections, Rt of Peak 2 = 4.6 min) to give the enantiomer 2, randomly assigned as **129** (62.93 mg, 147.6 µmol, 13% yield) (Rt = 3.38 min in analytical SFC) as a solid.

[000538] **128: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.49 (d, 1H), 7.85 (d, 1H), 7.74 (d, 1H), 7.68 - 7.57 (m, 2H), 7.51 - 7.44 (m, 1H), 7.41 (s, 1H), 7.09 (s, 1H), 5.67 (q, 1H), 4.13 (s, 3H), 3.11 - 3.05 (m, 1H), 2.97 - 2.90 (m, 1H). **LCMS** R_t = 1.11 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{15}F_4N_6O_3$ $[M+H]^+$ 427.1, found 427.1.

[000539] **129: ^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.49 (d, 1H), 7.85 (d, 1H), 7.74 (d, 1H), 7.68 - 7.57 (m, 2H), 7.51 - 7.44 (m, 1H), 7.41 (s, 1H), 7.09 (s, 1H), 5.67 (q, 1H), 4.13 (s, 3H), 3.13 - 3.03 (m, 1H), 2.99 - 2.88 (m, 1H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{15}F_4N_6O_3$ $[M+H]^+$ 427.1, found 427.1.

Example 93. Synthesis of 130



[000540] **A-149a:** To a solution of 2,2-difluoroethanamine (1.65 g, 20.39 mmol) in chloroform (300 mL) was added tert-butyl nitrite (2.5 g, 24.46 mmol) and AcOH (0.23 mL, 4.08 mmol). The reaction mixture was stirred at 78 °C for 15 mins then cooled to room

temperature. Ethyl prop-2-ynoate (1 g, 10.19 mmol) was then added to the reaction mixture and the mixture was stirred at 20 °C for 22 hours. The mixture was concentrated under reduced pressure. The residue was diluted with H_2O (50 mL), and the mixture was extracted with DCM (40 mL x 2). The combined organic phase was washed with water (60 mL) and brine (60 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10% to 15%) to give the product (1.7 g, 8.94 mmol, 62% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 11.63 - 10.72 (m, 1H), 7.11 - 7.03 (m, 1H), 6.75 (s, 1H), 4.48 - 4.35 (m, 2H), 1.41 (t, 3H).


[000541] **A-150a:** To a solution of ethyl 3-(difluoromethyl)-1H-pyrazole-5-carboxylate (900 mg, 4.73 mmol) in MeCN (10 mL) was added Cs_2CO_3 (3.08 g, 9.47 mmol) and sodium 2-chloro-2,2-difluoro-acetate (1.44 g, 9.47 mmol) and 18-crown-6 (250.21 mg, 0.95 mmol). The reaction mixture was stirred at 90 °C for 1.5 hours. After cooling to room temperature, the mixture was diluted with H_2O (50 mL), and then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (60 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 7%) to give the product (100 mg, 416.4 µmol, 8% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.27 - 7.85 (m, 1H), 7.20 (s, 1H), 6.94 - 6.47 (m, 1H), 4.43 (q, 2H), 1.42 (t, 3H).

[000542] **A-151a:** To a solution of ethyl 2,5-bis(difluoromethyl)pyrazole-3-carboxylate (100 mg, 0.42 mmol) in ethanol (8 mL) was added a solution of NaOH (49.97 mg, 1.25 mmol) in water (8 mL) and the mixture was stirred at 20 °C for 12 hours. The reaction mixture was quenched by addition of 1N HCl (15 mL) and diluted with H_2O (15 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the product (50 mg) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.46 - 8.04 (m, 1H), 7.31 - 6.97 (m, 2H).

[000543] **130:** To a mixture of 2,5-bis(difluoromethyl)pyrazole-3-carboxylic acid (69.72 mg, 0.33 mmol), EDCI (70.02 mg, 0.37 mmol) in MeCN (2 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (89 mg, 0.37 mmol) and the reaction mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give an impure product. The impure product was purified by prep-

TLC (silica gel, PE: EtOAc = 3:1) to give the product (17.27 mg, 43 μmol, 11% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.31 - 7.99 (m, 1H), 7.89 (d, 1H), 7.82 - 7.76 (m, 1H), 7.49 (d, 1H), 7.27 - 7.20 (m, 1H), 7.06 (s, 1H), 6.93 - 6.62 (m, 2H), 5.66 - 5.56 (m, 1H), 1.80 (d, 3H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{13}F_5N_5O_2$ $[M+H]^+$ 402.0, found 402.09.

Example 94. Synthesis of 131




[000544] **A-153a:** A mixture of (2S,3R)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoic acid (1 g, 4.29 mmol) and DCC (1.77 g, 8.57 mmol), 3-fluoro-N-hydroxy-benzamidine (660.81 mg, 4.29 mmol) in 1,4-dioxane (15 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated aqueous NH_4Cl (30 mL) and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with water (50 mL) and brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 30%) to give the product (780 mg, 2.1 mmol, 48% yield) as an oil. **LCMS** R_t = 0.93 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{13}H_{15}FN_3O_4$ $[M+H-t\text{-}Bu]^+$ 296.1, found 296.1.

[000545] **A-154a:** To tert-butyl N-[(1S,2R)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propyl]carbamate (780 mg, 2.22 mmol) was added 4M HCl in 1,4-dioxane (20 mL, 2.22 mmol) and the reaction mixture was stirred at 25 °C for 3 hours. The mixture was concentrated under reduced pressure to give the product (700 mg) as a solid. **LCMS** R_t = 0.68 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{15}FN_3O_2$ $[M+H]^+$ 252.1, found 252.1.

[000546] **131:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (202.39 mg, 1.04 mmol), EDCI (199.88 mg, 1.04 mmol) in CH_3CN (4 mL) was added

(1S,2R)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propan-1-amine hydrochloride (300 mg, 1.04 mmol) and the reaction mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (20 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (30 mL) and brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 40%) to give an impure product. The impure product was purified by prep-TLC (silica gel, PE: EtOAc = 3: 1) to give the product (91.9 mg, 0.2 mmol, 20% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.45 (d, 1H), 7.87 (d, 1H), 7.80 - 7.72 (m, 1H), 7.69 - 7.61 (m, 2H), 7.53 - 7.43 (m, 1H), 5.58 (dd, 1H), 4.15 - 4.06 (m, 4H), 3.31 (s, 3H), 1.22 (d, 3H). **LCMS** R_t = 1.30 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_3$ $[M+H]^+$ 428.1, found 428.0.

Example 95. Synthesis of 132



[000547] **A-149a:** To a solution of 2,2-difluoroethanamine (1.65 g, 20.39 mmol) in chloroform (300 mL) was added tert-butyl nitrite (2.52 g, 24.46 mmol) and AcOH (0.23 mL, 4.08 mmol and the reaction mixture was stirred at 78 °C for 15 minutes. The mixture was cooled to room temperature and then ethyl prop-2-ynoate (1 g, 10.19 mmol) was added. The mixture was stirred at 20 °C for 22 hours. The mixture was concentrated under reduced pressure to give the crude product (1.6 g) as an oil. A second batch was obtained of the crude product (600 mg) from 500 mg of 1, and the combined crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 15%) to give the product (1.7 g, 8.9 mmol) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 11.63 - 10.72 (m, 1H), 7.11 - 7.03 (m, 1H), 6.75 (s, 1H), 4.48 - 4.35 (m, 2H), 1.41 (t, 3H).


[000548] **A-155a:** To a solution of ethyl 3-(difluoromethyl)-1H-pyrazole-5-carboxylate (300 mg, 1.44 mmol), cyclopropylboronic acid (247.62 mg, 2.88 mmol) and Na_2CO_3 (305.53 mg, 2.88 mmol) in DCE (3 mL) was added a solution of $Cu(OAc)_2$ (261.79 mg, 1.44 mmol)

and 2,2-bipyridine (270.15 mg, 1.73 mmol) in DCE (6 mL). The reaction mixture was stirred at 70 °C for 4 hours. After cooling to room temperature, the mixture was diluted with sat. NH_4Cl (20 mL) and then the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (40 mL) and brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 5% to 50%) to give the product (210 mg, 0.9 mmol, 63% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.04 (s, 1H), 6.79 - 6.48 (m, 1H), 4.42 - 4.30 (m, 3H), 1.44 - 1.37 (m, 3H), 1.31 - 1.26 (m, 2H), 1.12 - 1.05 (m, 2H). **LCMS** R_t = 0.88 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{13}F_2N_2O_2$ $[M+H]^+$ 231.1, found 231.1.

[000549] **A-156a:** A mixture of ethyl 2-cyclopropyl-5-(difluoromethyl)pyrazole-3-carboxylate (210 mg, 0.91 mmol) and NaOH (109.47 mg, 2.74 mmol) in ethanol (4 mL) and water (4 mL) was stirred at 25 °C for 2 hours. The reaction mixture was quenched by addition of 1N HCl (15 mL) and diluted with H_2O (15 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the product (150 mg, 0.7 mmol, 79% yield) as a solid. **LCMS** R_t = 0.73 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_8H_9F_2N_2O_2$ $[M+H]^+$ 203.1, found 203.1.

[000550] **132:** To a mixture of 2-cyclopropyl-5-(difluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.74 mmol) and EDCI (142.24 mg, 0.74 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (180.8 mg, 0.74 mmol) and the mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL) and extracted with EtOAc (10 mL x 2). The combined organic phase was washed with water (15 mL) and brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30%) to give the impure product. The impure product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 45-75% B over 8 min) to give the product (35.8 mg, 91.6 µmol, 12% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.41 (d, 1H), 7.87 (td, 1H), 7.78 - 7.72 (m, 1H), 7.68 - 7.61 (m, 1H), 7.51 - 7.44 (m, 1H), 7.20 (s, 1H), 7.17 - 6.87 (m, 1H), 5.50 - 5.41 (m, 1H), 4.47 - 4.39 (m, 1H), 1.68 (d, 3H), 1.13 - 1.07 (m, 2H), 1.02 - 0.94 (m, 2H). **LCMS** R_t = 1.24 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_3N_5O_2$ $[M+H]^+$ 392.1, found 392.0.


Example 96. Synthesis of 133



A-62 133

[000551] A mixture of 2-methyl-4,5,6,7-tetrahydroindazole-3-carboxylic acid (100 mg, 0.55 mmol), EDCI (212.76 mg, 1.11 mmol), DIPEA (0.29 mL, 1.66 mmol), HOBt (149.98 mg, 1.11 mmol) and 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (148.74 mg, 0.61 mmol) in DCM (5 mL) was stirred at 20 °C for 16 hours. The reaction was concentratred and the residue was diluted with water (20 mL) and extracted with DCM (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 μm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-70% B over 10 min) to give the impure product. Further purification by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 μm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-70% B over 10 min gave the product (26.88 mg, 0.07 mmol, 13% yield) as a solid. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 7.87 (d, 1H), 7.82 - 7.74 (m, 1H), 7.52 - 7.42 (m, 1H), 7.26 - 7.19 (m, 1H), 6.46 (br d, 1H), 5.68 - 5.58 (m, 1H), 4.12 (s, 3H), 2.90 - 2.74 (m, 2H), 2.71 (t, 2H), 1.94 - 1.82 (m, 4H), 1.76 (d, 3H). **LCMS** R_t = 1.15 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$ 370.2, found 370.2.

Example 97. Synthesis of 134



A-157a A-158a A-159a A-160a 134

[000552] **A-158a:** A mixture of (2S)-3-amino-2-(tert-butoxycarbonylamino)propanoic acid (15 g, 73.45 mmol), pyridine (6.52 mL, 80.8 mmol) and Ac_2O (9 g, 88.14 mmol) in DCM (30 mL) was stirred at 25 °C for 16 under N_2. The mixture was acidified with 10%

aqueous acetic acid to pH ~4, extracted with EtOAc (50 mL x 2). The combined organic phase was washed with water (20 mL x 2) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (10.5 g, 13.51 mmol, 18% yield) as an oil. **LCMS** Rt = 0.56 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{10}H_{19}N_2O_5$ $[M+H\text{-}BOC]^+$ 147.1, found 147.0.


[000553] **A-159a:** A mixture of rac-(2S)-3-acetamido-2-(tert-butoxycarbonylamino)propanoic acid (10 g, 40.61 mmol), CDI (7.24 g, 44.67 mmol), 3-fluoro-N-hydroxy-benzamidine (6.26 g, 40.61 mmol) in 1,4-dioxane (100 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was diluted with saturated aqueous NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50% to 100%) to give the product (1.8 g, 1.42 mmol, 3% yield) as a solid. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 7.92 - 7.82 (m, 1H), 7.81 - 7.70 (m, 1H), 7.50 - 7.41 (m, 1H), 7.27 (s, 1H), 7.26 - 7.17 (m, 1H), 6.25 - 6.05 (m, 1H), 5.84 (br s, 1H), 5.24 - 5.11 (m, 1H), 3.98 - 3.70 (m, 2H), 2.02 (s, 3H), 1.47 (s, 9H). **LCMS** Rt = 0.83 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{17}H_{22}FN_4O_4$ $[M+H\text{-}BOC]^+$ 265.15, found 265.2.

[000554] **A-160a:** To a mixture of tert-butyl N-[(1S)-2-acetamido-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (800 mg, 2.2 mmol) in 1,4-dioxane (5 mL) and 4M HCl in 1,4-dioxane (10 mL, 40 mmol) was stirred at 25 °C for 16 hours. The mixture was filtered and concentrated to give the crude product (500 mg) as a solid. **LCMS** Rt = 0.56 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{12}H_{15}ClFN_4O_2$ $[M+H\text{-}HCl]^+$ 265.08, found 265.1.

[000555] **134:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (322.74 mg, 1.66 mmol), EDCI (318.74 mg, 1.66 mmol) in MeCN (10 mL) was added N-[(2S)-2-amino-2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]acetamide hydrochloride (500 mg, 1.66 mmol) and the mixture was stirred at 25 °C for 2 hours. The reaction was quenched with 1N HCl (10 mL), then extracted with EtOAc (10 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the product as a solid. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm I.D., 5.0 μm), mobile phase: A: CO_2 B: IPA

(0.05% DEA), gradient: from 5% to 40% of B in 5.5 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C, ABPR: 100 bar.) showed two peaks at 3.57 min (8%) and 4.13 min (92%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% DEA) ; 35 °C; 60 mL/min; 30% B; 9 min run; 7 injections, Rt of peak 1 = 4.4 min, Rt of peak 2 = 7.9 min) to give the product (36.7 mg, 0.8 mmol) (Rt = 4.13 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.48 (d, 1H), 8.23 (br t, 1H), 7.88 (d, 1H), 7.77 (d, 1H), 7.69 - 7.61 (m, 1H), 7.48 (dt, 1H), 7.43 (s, 1H), 5.41 (d, 1H), 4.12 (s, 3H), 3.94 - 3.81 (m, 1H), 3.66 - 3.54 (m, 1H), 1.82 (s, 3H). **LCMS** Rt = 1.19 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{17}F_4N_6O_3$ $[M+H]^+$ 341.12, found 441.1.

Example 98. Synthesis of 135 and 136



[000556] **A-161:** A mixture of methyl 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoate (3 g, 6.8 mmol) and LiOH.H_2O (570.44 mg, 13.59 mmol) in THF (10 mL) and water (5 mL) was stirred at 20 °C for 16 hours. To the mixture was added 1N HCl aqueous (50 mL) to adjust the pH = 2 and then the mixture was extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (50 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (2.5 g, 5.85 mmol, 86% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 12.59 (br s, 1H), 9.51 (d, 1H), 7.86 (d, 1H), 7.78 - 7.71 (m, 1H), 7.66 - 7.61 (m, 1H), 7.52 - 7.44 (m, 1H), 7.40 (s, 1H), 5.67 (q, 1H), 4.13 (s, 3H), 3.15 - 3.05 (m, 2H).


[000557] **135 & 136:** A mixture of 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoic acid (300 mg, 0.70 mmol), HATU (400.42 mg, 1.05 mmol), DIPEA (0.29 mL, 2.11 mmol) and N-dimethylamine

hydrochloride (57.25 mg, 0.70 mmol) in DMF (5 mL) was stirred at 20 °C for 16 hours. To the mixture was added water (20 mL) and it was then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the product. Analytical SFC (Daicel CHIRALPAK AD-3 (150 mm x 4.6 mm, 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% of B for 2.5 min, then hold 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 40 °C) showed two peaks at 2.59 min and 2.92 min. The product was separated by SFC (Phenomenex-Amylose-1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 38 °C; 55 mL/min; 15% B; 8 min run; 8 injections, Rt of peak 1 = 5.5 min, Rt of peak 2 = 7.5 min) to give the enantiomer 1, randomly assigned as **135** (23.42 mg, 51.5 µmol, 7% yield) (Rt = 2.59 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **136** (29.42 mg, 64.1 µmol, 9% yield) (Rt = 2.92 min in analytical SFC) as a solid.

[000558] **135: ^{1}H NMR** ($CDCl_3$ 400MHz) δ_H = 8.20 (d, 1H), 7.86 (d, 1H), 7.80 - 7.73 (m, 1H), 7.47 - 7.42 (m, 1H), 7.23 - 7.18 (m, 1H), 6.96 (s, 1H), 5.86 - 5.77 (m, 1H), 4.27 (s, 3H), 3.56 (dd, 1H), 3.10 (s, 3H), 3.03 (dd, 1H), 2.95 (s, 3H). **LCMS** R_t = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{19}F_4N_6O_3$ $[M+H]^+$ 455.1, found 455.1.

[000559] **136: ^{1}H NMR** ($CDCl_3$ 400MHz) δ_H = 8.20 (d, 1H), 7.86 (d, 1H), 7.77 - 7.74 (m, 1H), 7.47 - 7.42 (m, 1H), 7.23 - 7.18 (m, 1H), 6.96 (s, 1H), 5.87 - 5.78 (m, 1H), 4.27 (s, 3H), 3.56 (dd, 1H), 3.10 (s, 3H), 3.03 (dd, 1H), 2.95 (s, 3H). **LCMS** R_t = 1.20 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{19}F_4N_6O_3$ $[M+H]^+$ 455.1, found 455.1.

Example 99. Synthesis of 137



[000560] **A-163:** A mixture of (2R)-3-amino-2-(tert-butoxycarbonylamino)propanoic acid (10.00 g, 48.97 mmol), pyridine (4.35 mL, 53.86 mmol) and Ac_2O (6 g, 58.76 mmol) in DCM (30.00 mL) was stirred at 25 °C for 24 under N_2. The mixture was concentrated and diluted with sat. citric acid (40 mL) and then the mixture was extracted with EtOAc (40 mL x 2). The combined organic phase was washed with brine (40 mL), dried over Na_2SO_4, filtered and concentrated to give the product (5.00 g) as a solid.


[000561] **A-164:** A mixture of (2R)-3-acetamido-2-(tert-butoxycarbonylamino)propanoic acid (4.35 g, 17.66 mmol), CDI (3.15 g, 19.43 mmol) and 3-fluoro-N-hydroxy-benzamidine (2.72 g, 17.66 mmol) in DMF (30.00 mL) was stirred at 70 °C for 16 hours. The mixture was cooled to room temperature, filtered and solid washed with EtOAc (30 mL). The filtrate was diluted with saturated aqueous NH_4Cl (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by chromatography flash column on silica gel (EtOAc in PE = 0% to 10% to 30% to 50% to 100%) to give the product (800 mg, 2.20 mmol, 12% yield) as a solid. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 7.92 - 7.83 (m, 1H), 7.81 - 7.71 (m, 1H), 7.51-7.42 (m, 1H), 7.27 (s, 1H), 7.26 - 7.17 (m, 1H), 6.09 (br s, 1H), 5.84 (br s, 1H), 5.25 - 5.07 (s, 1H), 3.98 - 3.70 (m, 2H), 2.02 (s, 3H), 1.47 (s, 9H).

[000562] **A-165:** To a mixture of tert-butyl N-[(1R)-2-acetamido-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (300.0 mg, 0.82 mmol) in 1,4-dioxane (1.00 mL) was added 4M HCl in 1,4-dioxane (3.00 mL) and the reaction mixture was stirred at 25 °C for 16 hours. The mixture was concentrated to give the crude product (240.0 mg) as a solid. **LCMS** R_t = 0.35 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{12}H_{14}FN_4O_2$ $[M+H]^+$ 265.1, found 264.9.

[000563] **137:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (154.9 mg, 0.80 mmol), EDCI (153.0 mg, 0.80 mmol) in MeCN (10.00 mL) was added N-[(2R)-2-amino-2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]acetamide hydrochloride (240.0 mg, 0.80 mmol) and the reaction mixture was stirred at 25 °C for 3 hours. The reaction was quenched with the addition of 1 N HCl (15 mL) and then extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50% to 100%) to give the product (70.0 mg, 0.16 mmol) as a solid. Analytical SFC (Regis (S,S) Whelk-01 (150 mm x 4.6 mm, 5 µm), mobile phase: A: CO_2 B: methanol (0.05% DEA), gradient: from 5%

to 40% of B in 5 min and hold 40% for 2.5 min, then 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 3.57 min (70.5%) and 4.13 min (29.5%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = IPA (0.1% NH_4OH); 38 °C; 60 mL/min; 30% B; 10 min run; 3 injections, Rt of peak 1 = 5.7 min, Rt of peak 2 = 8.0 min) to give the product (22.47 mg, 51 μmol, 45% yield) (Rt = 3.57 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 8.78 (d, 1H), 7.87 (d, 1H), 7.82 - 7.73 (m, 1H), 7.52 - 7.41 (m, 1H), 7.26 - 7.19 (m, 1H), 7.04 (s, 1H), 6.12 - 5.96 (m, 1H), 5.52 - 5.38 (m, 1H), 4.23 (s, 3H), 4.08 - 3.82 (m, 2H), 2.09 (s, 3H). **LCMS** Rt = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_4N_6O_3$ $[M+H]^+$ 441.1, found 441.1.

Example 100 Synthesis of 138



[000564] **A-167:** To a solution of (2R,3S)-2-amino-3-hydroxy-butanoic acid (1 g, 8.39 mmol) and $NaHCO_3$ (1.0 g, 12.59 mmol) in water (18 mL) and methanol (18 mL) was added Boc_2O (2.75 g, 12.59 mmol). The reaction mixture was stirred at 20 °C for 16 hours. The mixture was concentrated, and the residue was partitioned between 1% HCl (1%, 40 mL) and EtOAc (30 mL). The aqueous layer was extracted with EtOAc (30 mL x 2). The combined organic phase was dried over Na_2SO_4, filtered and concentrated to give the crude product (1.8 g, 8.2 mmol, 97% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ_H = 12.48 (br s, 1H), 6.31 (d, 1H), 4.67 (br s, 1H), 3.91 - 3.83 (m, 1H), 1.38 (s, 9H), 1.07 (d, 3H).

[000565] **A-168:** To a solution of (2R,3S)-2-(tert-butoxycarbonylamino)-3-hydroxy-butanoic acid (1.6 g, 7.3 mmol) in MeCN (140 mL) was added Ag_2O (8.45 g, 36.49 mmol) followed by CH_3I (6.8 mL, 109.2 mmol) at 0 °C. The reaction mixture was warmed and stirred at 20 °C for 48 hours. The reaction mixture was filtered through Celite, eluted with EtOAc (30 mL x 2) and the filtrate was concentrated to give the product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 12% to 30%) to give the product (490 mg, 1.9 mmol, 27% yield) as an oil. **^{1}H NMR** (400MHz $CDCl_3$) δ_H

=5.22 (br d, 1H), 4.32 - 4.24 (m, 1H), 3.97 - 3.86 (m, 1H), 3.77 (s, 3H), 3.29 (s, 3H), 1.46 (s, 9H), 1.21 (d, 3H).

[000566] **A-169:** To a solution of methyl (2R,3S)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoate (390 mg, 1.58 mmol) in THF (8 mL) was added $LiOH{\cdot}H_2O$ (397.05 mg, 9.46 mmol) in water (4 mL). The mixture was stirred at 25 °C for 16 hours. The reaction mixture was quenched by the addition of 1N HCl (20 mL), diluted with H_2O (15 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL) and dried over Na_2SO_4, filtered and concentrated to give the product (300 mg, 1.2 mmol, 81% yield) as an oil. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 5.30 (br d, 1H), 4.35 (br d, 1H), 4.04 - 3.94 (m, 1H), 3.39 (s, 3H), 1.46 (s, 9H), 1.22 (d, 3H).


[000567] **A-170:** A mixture of 3-fluoro-N-hydroxy-benzamidine (198 mg, 1.28 mmol) and (2R,3S)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoic acid (299.63 mg, 1.28 mmol), DCC (529.23 mg, 2.57 mmol) in 1,4-dioxane (5 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was filtered and the solid washed with EtOAc (10 mL). The combined organic layer was diluted with saturated aqueous NH_4Cl (30 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 15%) to give the product (400 mg, 1.1 mmol, 88% yield) as an oil. **LCMS** R_t = 1.0 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{13}H_{15}FN_3O_4$ $[M+H\text{-}t\text{-}Bu]^+$ 296.10, found 295.8.

[000568] **A-171:** To tert-butyl N-[(1R,2S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propyl]carbamate (400 mg, 1.14 mmol) was added 4M HCl in 1,4-dioxane (20 mL, 80 mmol) and the reaction mixture was stirred at 20 °C for 12 hours. The mixture was concentrated to give the crude product (370 mg, 371.3 μmol, 32% yield) as an oil. **LCMS** R_t = 0.69 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{12}H_{15}FN_3O_2$ $[M+H]^+$ 252.11, found 252.1.

[000569] **138:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (249.62 mg, 1.29 mmol), EDCI (246.52 mg, 1.29 mmol) in MeCN (8 mL) was added (1R,2S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propan-1-amine hydrochloride (370 mg, 1.29 mmol) and the mixture was stirred at 0 °C for 2 hours. The reaction was quenched with the addition of 1N HCl (20 mL) and then extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified

by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the impure product (60 mg) as a solid. Analytical SFC (Regis (S,S) Whelk-O1 (100 mm x 4.6 mm I.D., 5 µm), mobile phase: A: CO_2 B: methanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C, ABPR: 100 bar) showed two peaks at 2.52 min (main peak, 88.7%) and 2.93 min (11.3%). The product was purified by SFC (Regis (S,S) Whelk-O1 (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O) ; 38 °C; 60 mL/min; 25% B; 8 min run; 7 injections, Rt of peak 1 = 4.7 min, Rt of peak 2 = 5.1 min) to give the product (37.97 mg, 88.2 µmol, 47% yield) (Rt = 2.52 min in analytical SFC) as an oil. **^{1}H NMR** (400MHz DMSO-d_6) δ_H = 9.49 - 9.35 (m, 1H), 7.87 (d, 1H), 7.79 - 7.73 (m, 1H), 7.68 - 7.61 (m, 2H), 7.48 (dt, 1H), 5.58 (dd, 1H), 4.13 (s, 3H), 4.12 - 4.07 (m, 1H), 3.31 (s, 3H), 1.22 (d, 3H). **LCMS** Rt = 1.40 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_3$ $[M+H]^+$ 428.13, found 428.1.

Example 101. Synthesis of 139



[000570] **A-173:** To a mixture of (2R,3R)-2-amino-3-hydroxy-butanoic acid (2 g, 16.79 mmol) in methanol (18 mL) and water (18 mL) at 0 °C was added $NaHCO_3$ (2.12 g, 25.18 mmol) and Boc_2O (5.5 g, 25.18 mmol) and it was stirred at 20 °C for 16 hours. The mixture was partially concentrated under reduced pressure to remove the MeOH. The residue was diluted with H_2O (20 mL), and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (1.8 g, 6.93 mmol, 41% yield) as an oil. **LCMS** R_t = 0.60 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_9H_{18}NO_5$ $[M+H-t\text{Bu}]$ 164.1, found 163.8.

[000571] **A-174:** To a mixture of (2R,3R)-2-(tert-butoxycarbonylamino)-3-hydroxy-butanoic acid (4 g, 18.25 mmol) in MeCN (100 mL) at 0 °C was added Ag_2O (21.14 g, 91.23 mmol) and iodomethane (38.85 g, 273.69 mmol) and the mixture was stirred at 20 °C for 2 days. The mixture was filtered through Celite, eluted with EtOAc (20 mL) and the filtrate was concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 30% to 50%) to give the product of (900 mg, 3.64 mmol, 19% yield) as an oil. **^{1}H NMR** $CDCl_3$ δ_H = 5.34 - 5.19 (m, 1H), 4.50 - 4.39 (m, 1H), 3.77 (s, 3H), 3.68 - 3.59 (m, 1H), 3.37 (s, 3H), 1.46 (s, 9H), 1.21 (d, 3H).


[000572] **A-175:** To a solution of methyl (2R,3R)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoate (500 mg, 2.02 mmol) in THF (5 mL) was added $LiOH.H_2O$ (509.04 mg, 12.13 mmol) in water (5 mL) and the mixture was stirred at 25 °C for 2 hours. The mixture was partially concentrated under reduced pressure to give remove the THF. The mixture was washed with EtOAc (5 mL), and the water phase was acidified with 1N HCl solution to pH = 4. The mixture was extracted with EtOAc (15 mL x 2). The combined organic phase was washed brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product (400 mg, 0.94 mmol, 46% yield) as an oil. **LCMS** R_t = 0.83 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{10}H_{20}NO_5$ [M+H-Boc] 134.1, found 133.8.

[000573] **A-176:** A mixture of 3-fluoro-N-hydroxy-benzamidine (264 mg, 1.71 mmol), (2R,3R)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoic acid (399.51 mg, 1.71 mmol) and DCC (705.64 mg, 3.43 mmol) in 1,4-dioxane (10 mL) was stirred at 70 °C for 16 hours. After cooling to room temperature, the mixture was filtered and the solid washed with EtOAc (10 mL). The filtrate was diluted with saturated aqueous NH_4Cl (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 10%) to give the product (200 mg, 0.33 mmol, 19% yield) as a solid. **LCMS** R_t = 0.92 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{17}H_{23}FN_3O_4$ [M+H-*t*Bu] 296.2, found 296.1.

[000574] **A-177:** A mixture of tert-butyl N-[(1R,2R)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propyl]carbamate (230 mg, 0.65 mmol) and 4M HCl in 1,4-dioxane (10 mL, 40 mmol) was stirred at 20 °C for 16 hours. The mixture was concentrated under reduced pressure to give the crude product (200 mg) as an oil. **LCMS** R_t = 0.68 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{15}FN_3O_2$ [M+H] 252.1, found 252.0.

[000575] **139:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (134.93 mg, 0.70 mmol), EDCI (133.25 mg, 0.70 mmol) in MeCN (4 mL) was added (1R,2R)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propan-1-amine hydrochloride (200 mg, 0.70 mmol) and the reaction mixture was stirred at 0 °C for 2 hours. The reaction was quenched with the addition of 1N HCl (15 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by flash chromatography on silica gel (EtOAc in PE = 0% to 40%) to give the impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) (twice) to give the product. The product was triturated from DCM/n-hexane (1:20, 3 mL) to give the product (18.8 mg, 0.04 mmol, 6% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 9.46 (d, 1H), 7.87 (d, 1H), 7.79 - 7.73 (m, 1H), 7.69 - 7.61 (m, 1H), 7.53 (s, 1H), 7.51 - 7.44 (m, 1H), 5.40 (t, 1H), 4.11 (s, 3H), 4.05 - 3.94 (m, 1H), 3.27 (s, 3H), 1.29 (d, 3H). **LCMS** R_t = 1.37 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_3$ $[M+H]^+$ 428.1, found 428.1.

Example 102. Synthesis of 140 and 141




[000576] A mixture of N-[3-amino-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-oxo-propyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (180 mg, 0.42 mmol), pyridine (0.14 mL, 1.69 mmol) and TFAA (0.12 mL, 0.84 mmol) in THF (5 mL) was stirred at 0 °C for 16 hours. The reaction mixture was warmed to room temperature and the pH of the mixture was adjusted pH = 2 with 1N HCl aqueous (20 mL) and then extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL), saturated $NaHCO_3$ aqueous (20 mL) and brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 μm) A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 45-73% B over 8 min) to give the product. Analytical SFC (Daicel CHIRALPAK AD-3 (150 mm x 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and hold 40% of B for 2.5 min, then hold 5% of B for 2.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at

2.40 min and 2.80 min. The product was separated by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH; 38 °C; 50 mL/min; 20% B; 8 min run; 22 injections, Rt of peak 1 = 4.25 min, Rt of peak 2 = 5.25 min) to give the enantiomer 1 (Rt = 2.40 min in analytical SFC, 65% ee) and the enantiomer 2 (Rt = 2.80 min in analytical SFC, 71% ee). The impure enantiomer 1 was purified by SFC (Phenomenex-Amylose-1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 38 °C; 50 mL/min; 20% B; 8 min run; 16 injections, Rt of peak 1 = 4.4 min) to give the enantiomer 1, randomly assigned as **140** (21.11 mg, 51.7 μmol, 12% yield) (Rt = 2.40 min in analytical SFC) as a solid. The impure enantiomer 2 was purified by SFC (Phenomenex-Amylose-1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = 0.1% $NH_3.H_2O$-EtOH; 38 °C; 50 mL/min; 20% B; 8 min run; 11 injections, Rt of peak 2 = 5.4 min) to give the the enantiomer 2, randomly assigned as **141** (19.51 mg, 47.8 μmol, 11% yield) (Rt = 2.80 min in analytical SFC) as a solid.

[000577] **140: ^{1}H NMR** (DMSO-d_6 400MHz) δ_H = (d, 1H), 7.88 (d, 1H), 7.77 (d, 1H), 7.70 - 7.62 (m, 1H), 7.54 - 7.46 (m, 1H), 7.43 (s, 1H), 5.89 - 5.77 (m, 1H), 4.15 (s, 3H), 3.52 - 3.38 (m, 2H). **LCMS** R_t = 1.07 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{13}F_4N_6O_2$ $[M+H]^+$ 409.1, found 409.1.

[000578] **141: ^{1}H NMR** (DMSO-d_6 400MHz) δ_H = 9.82 (d, 1H), 7.88 (d, 1H), 7.77 (d, 1H), 7.70 - 7.62 (m, 1H), 7.54 - 7.46 (m, 1H), 7.43 (s, 1H), 5.88 - 5.79 (m, 1H), 4.15 (s, 3H), 3.52 - 3.37 (m, 2H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{13}F_4N_6O_2$ $[M+H]^+$ 409.1, found 409.1.

Example 103. Synthesis of 142



[000579] **A-180:** To a solution of (2S,3S)-2-amino-3-hydroxy-butanoic acid (3 g, 25.18 mmol) and $NaHCO_3$ (3.17 g, 37.78 mmol) dissolved in water (18 mL) and methanol

(25 mL) was added Boc_2O (8.24 g, 37.78 mmol) and the reaction mixture was stirred at 20 °C for 16 hours. The mixture was partially concentrated, and the residue was partitioned between 1% HCl (80 mL) and EtOAc (80 mL). The aqueous layer was extracted with EtOAc (50 mL). The combined organic phase was dried over Na_2SO_4, filtered and concentrated to give the crude product (2 g, 9.12 mmol, 36% yield) as an oil. **^{1}H NMR** (400MHz, DMSO-*d*$_6$) δ_H = 6.80 (d, 1H), 4.10 (s, 1H), 3.85 (d, 1H), 3.16 (s, 2H), 1.38 (s, 9H), 1.08 (d, 3H).

[000580] **A-181:** To a mixture of (2S,3S)-2-(tert-butoxycarbonylamino)-3-hydroxy-butanoic acid (4 g, 18.25 mmol) in MeCN (20 mL) was added Ag_2O (21.14 g, 91.23 mmol) and iodomethane (41.12 g, 289.7 mmol) at 0 °C. The resulting mixture was then stirred at 20 °C for 48 hours. The reaction mixture was filtered through Celite, eluted with EtOAc (30 mL x 2) and the filtrate was concentrated to give the product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 12% to 30%) to give the product (1.6 g, 6.47 mmol, 35% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 5.28 (br d, 1H), 4.43 (dd, 1H), 3.76 (s, 3H), 3.67 - 3.59 (m, 1H), 3.36 (s, 3H), 1.45 (s, 9H), 1.20 (d, 3H).


[000581] **A-182:** To a solution of methyl (2S,3S)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoate (1.6 g, 6.47 mmol) in THF (12 mL) and water (6 mL) was added $LiOH.H_2O$ (1.63 g, 38.82 mmol) and the resulting mixture was stirred at 20 °C for 4 hours. The mixture was partially concentrated under reduced pressure. The residue was partitioned between 1% HCl (40 mL) and EtOAc (30 mL). The aqueous layer was extracted with EtOAc (30 mL x 2). The combined organic phase was dried over Na_2SO_4, filtered and concentrated to give the crude product (1.5 g, 6.43 mmol, 99% yield) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 5.40 - 5.21 (m, 1H), 4.50 - 4.40 (m, 1 H), 3.75 - 3.67 (m, 1H), 3.38 (s, 3H), 1.45 (s, 9H), 1.27 - 1.25 (m, 3H).

[000582] **A-183:** To a mixture of (2S,3S)-2-(tert-butoxycarbonylamino)-3-methoxy-butanoic acid (1.5 g, 6.43 mmol) and 3-fluoro-N-hydroxy-benzamidine (991.21 mg, 6.43 mmol) in 1,4-dioxane (20 mL) was added DCC (2.65 g, 12.86 mmol) and the mixture was stirred at 70 °C for 16 hours. The mixture was cooled to room temperature and filtered through Celite. The filtrate was concentrated to give the crude product. The crude product was purified by flash column chromatography on silica gel (EtOAc in PE = 0% to 15% to 30%) to give the product (1.4 g, 3.72 mmol, 57% yield) as an oil. **LCMS** R_t = 0.93 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{13}H_{15}FN_3O_4$ $[M+H\text{-}t\text{-}Bu]^+$ 296.1, found 296.1.

[000583] **A-184:** A solution of tert-butyl N-[(1S,2S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propyl]carbamate (300 mg, 0.85 mmol) in 4M HCl in 1,4-dioxane (10 mL, 0.85 mmol) was stirred at 20 °C for 2 hours. The solution was concentrated to give the crude product (245 mg) as an oil. **LCMS** R_t = 0.72 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{15}FN_3O_2$ $[M+H]^+$ 252.1, found 251.6.

[000584] **142:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (165.29 mg, 0.85 mmol), EDCI (163.24 mg, 0.85 mmol) in MeCN (8 mL) was added (1S,2S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methoxy-propan-1-amine hydrochloride (245 mg, 0.85 mmol), the resulting mixture was stirred at 0 °C for 2 hours. The reaction was quenched with 1N HCl (20 mL), then extracted with EtOAc (15 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was partially purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 20%) to give the impure product. The impure product was purified by prep-TLC (silica gel, PE:EtOAc = 3:1) to give the product (112.34 mg, 262.9 μmol, 30% yield) as a an oil. **^{1}H NMR** (DMSO-d_6 400MHz) δ_H = 9.46 (d, 1H), 7.87 (d, 1H), 7.79 - 7.73 (m, 1H), 7.67-7.61 (m, 1H), 7.52 (s, 1H), 7.50 - 7.45 (m, 1H), 5.40 (t, 1H), 4.11 (s, 3H), 4.05 - 3.94 (m, 1H), 3.27 (s, 3H), 1.29 (d, 3H). **LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_3$ $[M+H]^+$ 428.1, found 428.2.

Example 104. Synthesis of 143 and 144



[000585] **A-186:** To a solution of 2-amino-4-methylsulfonyl-butanoic acid (2 g, 11.04 mmol) and Boc_2O (3.61 g, 16.56 mmol) in methanol (20 mL) and water (20 mL) was added $NaHCO_3$ (1.39 g, 16.56 mmol) and the mixture was stirred at 20 °C for 16 hours. The mixture was partially concentrated, and the residue was partitioned between sat. citric acid (20 mL) and EtOAc (15 mL). The organic layer was separated, and the aqueous layer was

extracted with EtOAc (15 mL x 2). The combined organic phase was dried over Na_2SO_4, filtered and concentrated to give the crude product (2.9 g) as a solid.

[000586] **A-187:** A mixture of 2-(tert-butoxycarbonylamino)-4-methylsulfonyl-butanoic acid (1.5 g, 5.33 mmol), 3-fluoro-N-hydroxy-benzamidine (0.82 g, 5.33 mmol) and DCC (2.2 g, 10.66 mmol) in 1,4-dioxane (15 mL) was stirred at 70 °C for 16 hours. The mixture was diluted with H_2O (100 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 50%) to give the product (1.5 g, 2.87 mmol, 54% yield) as a solid. **LCMS** R_t = 0.86 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{17}H_{23}FN_3O_5S$ $[M+H\text{-}tBu]^+$344.13, found 344.1.

[000587] **A-188:** To tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-methylsulfonyl-propyl]carbamate (1.5 g, 3.76 mmol) was added 4M HCl in 1,4-dioxane (25 mL, 3.76 mmol) and the mixture was stirred at 25 °C for 10 mins. The mixture was concentrated to give the crude product (1.5 g) as a solid.

[000588] **A-189:** To a solution of 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-methylsulfonyl-propan-1-amine hydrochloride (400 mg, 1.34 mmol), 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (285.34 mg, 1.47 mmol) and HATU (1.02 g, 2.67 mmol) in DMF (5 mL) was added Et_3N (0.92 mL, 6.68 mmol) and mixture was stirred at 25 °C for 16 hours. The mixture was diluted with H_2O (20 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Prime C18 (150 mm x 30 mm, 5 μm), A = H_2O (0.05% NH_4OH) and B = CH_3CN; 46-66% B over 9 min) to give the product (280 mg, 0.58 mmol, 44% yield) as a solid. **LCMS** R_t = 1.19 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_4S$ $[M+H]^+$476.1, found 476.1.


[000589] **143 & 144:** Analytical SFC (Daicel CHIRALPAK AD-3 (150 mm × 4.6 mm, 3 μm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5.5 min and hold 40% for 3 min, then 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 40 °C ABPR: 100 bar) showed two peaks at 2.97 min and 3.71 min. The product was separated by SFC (Phenomenex-Amylose-1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O ETOH); 38 °C; 50 mL/min; 30% B; 8 min run; 15 injections, Rt of peak 1 = 5.2 min, Rt of peak 2 = 8.3 min) to give enantiomer 1 (100 mg, Rt = 2.97 min in analytical SFC, ee% = 92.34%) and enantiomer 2 (100 mg, Rt = 3.71 min in analytical SFC,

ee% = 93.37%). The impure enantiomer 1 was purified by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O) ; 38 °C; 50 mL/min; 30% B; 10 min run; 7 injections, Rt of peak 1 = 5.2 min, Rt of peak 2 = 8.3 min) to give the enantiomer 1, randomly assigned as **143** (70.77 mg, 0.15 mmol, 71% yield) (Rt = 2.97 min in analytical SFC) as a solid. The impure enantiomer 2 was purified by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O) ; 38 °C; 50 mL/min; 30% B; 10 min run; 6 injections, Rt of peak 1 = 5.2 min, Rt of peak 2 = 8.3 min) to give the enantiomer 2, randomly assigned as **144**(40.42 mg, 85 µmol, 40% yield) (Rt = 3.71 min in analytical SFC) as a solid.

[000590] **143: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.82 - 7.75 (m, 1H), 7.65 (br d, 1H), 7.53 - 7.44 (m, 1H), 7.26 - 7.21 (m, 1H), 7.05 (s, 1H), 5.69 (dt, 1H), 4.24 (s, 3H), 3.43 - 3.34 (m, 1H), 3.32 - 3.22 (m, 1H), 3.05 (s, 3H), 2.86 - 2.63 (m, 2H). **LCMS** R_t = 1.25 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_4S$ $[M+H]^+$476.1, found 476.0.

[000591] **144: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.81 - 7.75 (m, 1H), 7.65 (br d, 1H), 7.54 - 7.42 (m, 1H), 7.25 - 7.20 (m, 1H), 7.04 (s, 1H), 5.69 (dt, 1H), 4.24 (s, 3H), 3.44 - 3.34 (m, 1H), 3.32 - 3.22 (m, 1H), 3.06 (s, 3H), 2.88 - 2.63 (m, 2H). **LCMS** R_t = 1.26 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}F_4N_5O_4S$ $[M+H]^+$476.1, found 476.0.

Example 105. Synthesis of 147



A-191 A-192 A-193

147


[000592] **A-192:** To a solution of 2-(tert-butoxycarbonylamino)-2-(oxetan-3-yl)acetic acid (500 mg, 2.16 mmol) in DMF (5 mL) was added CDI (525.89 mg, 3.24 mmol) at 25 °C and the mixture was stirred at 25 °C for 30 min. Then 3-fluoro-N-hydroxy-benzamidine (399.93 mg, 2.59 mmol) was added to the solution and the mixture was stirred at 25 °C for 2

hours then heated to 110 °C and stirred for 16 hours. The reaction was cooled and was poured into water (30 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4 filtered and concentrated. The crude was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (360 mg, 1.03 mmol, 48% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.86 (d, 1H), 7.76 (d, 1H), 7.50 - 7.44 (m, 1H), 7.26 - 7.19 (m, 1H), 5.52 - 5.26 (m, 2H), 4.91 - 4.80 (m, 2H), 4.72 - 4.57 (m, 2H), 3.63 - 3.48 (m, 1H), 1.49 (s, 9H).

[000593] **A-193:** To a solution of tert-butyl N-[[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-(oxetan-3-yl) methyl] carbamate (200 mg, 0.57 mmol) in DCM (4 mL) was added TFA (1.26 mL, 17.17 mmol) at 0 °C. The mixture was then warmed to 25 °C and stirred for 30 mins. The reaction was concentrated to give the crude product (207 mg) as a solid. **LCMS** Rt = 0.68 min in 2.0 min chromatography, 5-95AB, MS ESI calcd. For $C_{12}H_{12}FN_3O_2$ $[M+H]^+$ 249.1, found 249.6.

[000594] **147:** To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (165.92 mg, 0.85 mmol) and Et_3N (0.28 mL, 1.99 mmol) in DCM (5 mL) was added HATU (325.01 mg, 0.85 mmol) at 25 °C and the mixture was stirred at 25 °C for 15 mins. Then [3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-(oxetan-3-yl)methanamine 2,2,2-trifluoroacetic acid (207 mg, 0.57 mmol) was added to the solution and the mixture was stirred at 25 °C for 16 hours. The mixture was poured in to water (20 mL) and extracted with DCM (10 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4 filtered and concentrated. The crude was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm x 5 µm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-70% over 10 min) to give the product (11.84 mg, 0.03 mmol, 5% yield) as a solid. **^{1}H NMR** (400MHz, DMSO-*d6*) δ_H = 7.86 (d, 1H), 7.76 (td, 1H), 7.48 (dt, 1H), 7.26 - 7.19 (m, 1H), 6.96 (s, 1H), 6.84 (d, 1H), 5.93 (t, 1H), 4.97 - 4.87 (m, 2H), 4.70 - 4.64 (m, 2H), 4.25 (s, 3H), 3.70 - 3.58 (m, 1H). **LCMS** Rt = 1.15 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. For $C_{18}H_{15}F_4N_5O_3$ $[M+H]^+$ 425.1, found 426.1.

Example 106. Synthesis of 148 and 149



A-194 A-195 A-196

A-197 148 149

[000595] **A-195:** A mixture of 3-fluoro-N-hydroxy-benzamidine (200 mg, 1.3 mmol), 4-(tert-butoxycarbonylamino)tetrahydrofuran-3-carboxylic acid (302.66 mg, 1.31 mmol) and DCC (534.58 mg, 2.6 mmol) in 1,4-dioxane (15 mL) was stirred at 100 °C for 16 hours. After cooling to room temperature, the mixture was filtered and the solid washed with EtOAc (10 mL). The combined organic layer was diluted with NH_4Cl (30 mL) and extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by flash column chromatography flash column on silica gel (EtOAc in PE = 0% to 10% to 30%) to give the product (370 mg, 1.06 mmol, 81% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.92 - 7.86 (m, 1H), 7.83 - 7.75 (m, 1H), 7.50 - 7.41 (m, 1H), 7.25 - 7.16 (m, 1H), 5.46 - 5.14 (m, 1H), 4.40 - 4.28 (m, 1H), 4.22 - 4.09 (m, 3H), 2.83 - 2.70 (m, 1H), 2.50 (s, 1H), 1.52 - 1.35 (m, 9H).

[000596] **A-196:** A mixture of tert-butyl N-[3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]tetrahydrofuran-3-yl]carbamate (370 mg, 1.06 mmol) in 4M HCl in 1,4-dioxane (15 mL, 60 mmol) was stirred at 20 °C for 16 hours. The mixture was concentrated to give the crude product (350 mg) as oil. **LCMS** R_t = 0.65 min in 2 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{13}FN_3O_2$ $[M+H]^+$ 250.1, found 249.8.

[000597] **A-197:** To a mixture of 3-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]tetrahydrofuran-3-amine hydrochloride (302 mg, 1.06 mmol), 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (205.18 mg, 1.06 mmol) in DCM (10 mL) was added EDCI (405.27mg, 2.11 mmol) and the reaction mixture was stirred at 20 °C for 2 hours. The reaction was quenched with the addition of sat. NH_4Cl (20 mL), then extracted with DCM (15 mL). The combined organic phase was washed brine (15 mL x 2), dried over Na2SO4, filtered and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 40%) to give the give the


product (100 mg, 0.23 mmol, 22% yield) as oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.90 - 7.84 (m, 1H), 7.81 - 7.75 (m, 1H), 7.50 - 7.42 (m, 1H), 7.25 - 7.18 (m, 1H), 7.06 - 7.00 (m, 1H), 6.99 (s, 1H), 4.33 (s, 2H), 4.27 - 4.17 (m, 2H), 4.15 (s, 3H), 3.03 - 2.90 (m, 1H), 2.74 - 2.63 (m, 1H).

[000598] **148 & 149:** Analytical SFC (Daicel CHIRALCEL OJ-3 (150 mm × 4.6 mm I.D., 3 µm), mobile phase: A: CO_2 B: ethanol (0.05% DEA), gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min, flow rate: 2.5 mL/min, column temp: 35 °C) showed two peaks at 2.71 min and 3.02 min. The product was separated by SFC (Daicel CHIRALPAK AD-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH ; 38 °C; 60 mL/min; 15% B; 9 min run; 10 injections, Rt of peak 1 = 6.16 min, Rt of peak 2 = 7.95 min) to give the enantiomer 1, randomly assigned as **148** (10.45 mg, 24.6 µmol, 10% yield) (Rt = 2.71 min in analytical SFC) as a solid and the enantiomer 2, randomly assigned as **149** (3.18 mg, 7.50 µmol, 3% yield) (Rt = 3.02 min in analytical SFC) as a solid.

[000599] **148: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.91 - 7.85 (m, 1H), 7.82 - 7.75 (m, 1H), 7.51 - 7.43 (m, 1H), 7.26 - 7.19 (m, 1H), 6.95 (s, 1H), 6.70 (s, 1H), 4.38 - 4.29 (m, 2H), 4.25 - 4.18 (m, 2H), 4.16 (s, 3H), 3.06 - 2.89 (m, 1H), 2.74 - 2.61 (m, 1H). **LCMS** R_t = 1.32 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_4N_5O_3$ $[M+H]^+$ 426.1, found 426.2.

[000600] **149: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.91 - 7.84 (m, 1H), 7.83 - 7.75 (m, 1H), 7.51 - 7.43 (m, 1H), 7.25 - 7.18 (m, 1H), 6.95 (s, 1H), 6.73 (s, 1H), 4.39 - 4.30 (m, 2H), 4.23 - 4.18 (m, 2H), 4.16 (s, 3H), 3.03 - 2.93 (m, 1H), 2.73 - 2.64 (m, 1H). **LCMS** R_t = 1.31 min in 2 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{16}F_4N_5O_3$ $[M+H]^+$ 426.1, found 426.1.

Example 107. Synthesis of 150




A-198 150

[000601] To a mixture of 1-tert-butyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.55 mmol) and EDCI (105.2 mg, 0.55 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (133.72 mg, 0.55 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of


1N HCl (5 mL), extracted with EtOAc (5 mL) and concentrated under reduced pressure to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 μm) A = H_2O (0.075% TFA) and B = CH_3CN; 45-70% B over 7.5 min) to give the product (24.1 mg, 65.0 μmol, 12% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.93 (s, 1H), 7.88 (d, 1H), 7.81 - 7.76 (m, 1H), 7.47 (dt, 1H), 7.25 - 7.19 (m, 1H), 6.32 (br d, 1H), 5.70 - 5.61 (m, 1H), 2.56 (s, 3H), 1.74 (d, 3H), 1.59 (s, 9H). **LCMS** R_t = 1.23 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{23}FN_5O_2$ $[M+H]^+$ 372.2, found 372.1.

Example 108. Synthesis of 151




[000602] To a mixture of 3-cyclopropyl-1-(2,2,2-trifluoroethyl)pyrazole-4-carboxylic acid (100 mg, 0.43 mmol), EDCI (81.86 mg, 0.43 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (104.05 mg, 0.43 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched by the addition of 1N HCl (5 mL), then extracted with EtOAc (5 mL), and concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 μm) A = H_2O (0.075% TFA) and B = CH_3CN; 49-79% B over 9 min) to give the product (4.3 mg, 10.1 mmol, 2% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.94 (s, 1H), 7.87 (d, 1H), 7.77 (br d, 1H), 7.51 - 7.44 (m, 1H), 7.25 - 7.20 (m, 1H), 6.70 (br d, 1H), 5.71 - 5.63 (m, 1H), 4.88 (q, 2H), 1.95 - 1.87 (m, 1H), 1.76 (d, 3H), 1.33 - 1.26 (m, 2H), 1.00 - 0.86 (m, 2H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_2$ $[M+H]^+$ 424.1, found 424.1.

Example 109. Synthesis of 152




[000603] To a mixture of methyl 3-methyl-1-tetrahydropyran-4-yl-pyrazole-4-carboxylate (100 mg, 0.45 mmol), EDCI (85.48 mg, 0.45 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (108.66 mg, 0.45 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched by the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and the organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm) A = H_2O (0.075% TFA) and B = CH_3CN; 43-68% B over 7.5 min) to give the product (10.1 mg, 25.3 mmol, 5% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 - 7.86 (m, 2H), 7.78 (td, 1H), 7.47 (dt, 1H), 7.25 - 7.20 (m, 1H), 6.33 (br d, 1H), 5.69 - 5.61 (m, 1H), 4.34 - 4.25 (m, 1H), 4.15 - 4.10 (m, 2H), 3.54 (dt, 2H), 2.55 (s, 3H), 2.16 - 1.99 (m, 4H), 1.74 (d, 3H). **LCMS** R_t = 1.13 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{23}FN_5O_3$ $[M+H]^+$ 400.2, found 400.1.

Example 110. Synthesis of 153



[000604] To a mixture of 3-cyclopropyl-1-isopropyl-pyrazole-4-carboxylic acid (100 mg, 0.51 mmol), EDCI (98.7 mg, 0.51 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (125.45 mg, 0.51 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with by the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and the organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm) A = H_2O (0.075% TFA) and B = CH_3CN; 53-78% B over 7.5 min) to give the product (15.4 mg, 39.9 µmol, 8% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.93 (s, 1H), 7.87 (d, 1H), 7.77 (br d, 1H), 7.50 - 7.43 (m, 1H), 7.25 - 7.19 (m, 2H), 5.73 - 5.65 (m, 1H), 4.48 - 4.38 (m, 1H), 2.17 - 2.09 (m, 1H), 1.75 (d, 3H), 1.48 (d, 6H), 1.12 - 1.00 (m, 4H). **LCMS** R_t = 1.28 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{23}FN_5O_2$ $[M+H]^+$ 384.2, found 384.1.


Example 111. Synthesis of 154



A-202 154

[000605] To a mixture of 1-(4-fluorophenyl)-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.45 mmol), EDCI (87.06 mg, 0.45 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (110.66 mg, 0.45 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL), and the organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm), A = H_2O (0.075% TFA) and B = CH_3CN; 55-80% B over 7.5 min) to give the product (8.1 mg, 19.7 µmol, 4% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.26 (s, 1H), 7.88 (td, 1H), 7.81 - 7.76 (m, 1H), 7.68 - 7.63 (m, 2H), 7.51 - 7.44 (m, 1H), 7.26 - 7.15 (m, 3H), 6.44 (br d, 1H), 5.72 - 5.63 (m, 1H), 2.64 (s, 3H), 1.77 (d, 3H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{21}H_{18}F_2N_5O_2$ $[M+H]^+$ 410.1, found 410.1.

Example 112. Synthesis of 155




A-203 155

[000606] To a mixture of 3-cyclobutyl-1-methyl-pyrazole-4-carboxylic acid (100 mg, 0.55 mmol), EDCI (106.38 mg, 0.55 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (135.22 mg, 0.55 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched by the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and the organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm), A = H_2O (0.075% TFA) and B = CH_3CN; 45-70% B over 7.5 min) to give the product (13.5 mg, 6.4 µmol, 6% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 (td, 1H), 7.82 - 7.77 (m, 2H), 7.48 (dt, 1H), 7.26 - 7.20 (m, 1H), 6.33 (br d, 1H), 5.66 - 5.58 (m, 1H), 3.94 - 3.85 (m, 4H), 2.50 - 2.40 (m, 4H), 2.17 - 2.04 (m, 1H),


2.01 - 1.92 (m, 1H), 1.72 (d, 3H). **LCMS** R_t = 1.22 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{21}FN_5O_2$ $[M+H]^+$ 370.2, found 370.1.

Example 113. Synthesis of 156



[000607] To a mixture of 3-(methoxymethyl)-1-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol), EDCI (112.65 mg, 0.59 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (143.19 mg, 0.59 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and the organic phase was concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm), A = H_2O (0.075% TFA) and B = CH_3CN; 39-69% B over 9 min) to give the product (5.3 mg, 14.7 µmol, 2% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.90 - 7.84 (m, 2H), 7.78 (br d, 1H), 7.50 - 7.43 (m, 1H), 7.25 - 7.18 (m, 2H), 5.68 - 5.59 (m, 1H), 4.85 - 4.76 (m, 2H), 3.95 (s, 3H), 3.45 (s, 3H), 1.73 (d, 3H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{19}FN_5O_3$ $[M+H]^+$ 360.1, found 360.1.

Example 114. Synthesis of 157



[000608] **A-205:** To the solution of 3-amino-2-(tert-butoxycarbonylamino)propanoic acid (1 g, 4.9 mmol), K_2CO_3 (1.35 g, 9.79 mmol), KOH (0.27 g, 4.9 mmol) in THF (15 mL) and water (5 mL) was added benzyl carbonochloridate (1.25 g, 7.35 mmol) in a dropwise manner at -5 °C. The suspension was stirred at 25 °C for 16 hours. The mixture was partially


concentrated to remove THF. The residue was acidified with solid citric acid to pH~4 and extracted with DCM (20 mL x 2). The combined organic phase was dried over Na_2SO_4, filtered and concentrated to give product (1 g) as an oil. **LCMS** R_t = 0.78 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{16}H_{22}N_2O_6$ $[M+Na]^+$ 361.2 , found 360.9.

[000609] **A-206:** A mixture of 3-(benzyloxycarbonylamino)-2-(tert-butoxycarbonylamino)propanoic acid (900 mg, 2.66 mmol) and CDI (862.6 mg, 5.32 mmol) in DMF (25 mL) was sirred at 25 °C for 30 minutes, then 3-fluoro-N-hydroxy-benzamidine (819.99 mg, 5.32 mmol) was added and the resulting mixture was stirred at 100 °C for 16 hours. After cooling to room temperature, the reaction mixture was diluted with H_2O (50 mL) and then extracted with EtOAc (50 mL x 2). The combined organic phase was washed with brine (50 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified with flash chromatography on silic gel (EtOAc in PE = 0/1 to 1/5) to give the product (800 mg, 1.70 mmol, 64% yield). **LCMS** R_t = 1.33 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{23}H_{25}FN_4O_5$ $[M+Na]^+$ 479.2, found 479.2.

[000610] **A-207:** A solution of tert-butyl N-[2-(benzyloxycarbonylamino)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (400 mg, 0.88 mmol) and 4M HCl in 1,4-dioxane (5 mL, 20 mmol) in DCM (10 mL) was stirred at 25 °C for 2 hours. The reaction mixture was concentrated to give a crude product (350 mg, 0.89 mmol) as a solid. **LCMS** R_t = 0.74 min in 1.5 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{18}FN_4O_3$ $[M+H]^+$ 357.1, found 357.0.


[000611] **157:** To a solution of benzyl N-[2-amino-2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate hydrochloride (336 mg, 0.86 mmol), 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (182.64 mg, 0.94 mmol) and HATU (650.5 mg, 1.71 mmol) in DMF (4 mL) was added Et_3N (0.59 mL, 4.28 mmol). The mixture was stirred at 25 °C for 16 hours. The reaction was quenched with the addition of sat. NH_4Cl (20 mL) and the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Phenomenex Gemini-NX (150 mm x 30 mm, 5 μm), A = H_2O (0.04% NH_3H_2O+10 mM NH_4HCO_3) and B = CH_3CN; 38-68% B over 8.5 mins) to give the product (216.9 mg, 0.40 mmol, 47% yield) as a solid. **^{1}H NMR** (400M Hz $CDCl_3$) δ_H = 9.46 (d, 1H), 7.87 (d, 1H), 7.76 (br d, 1H), 7.71 - 7.59 (m, 2H), 7.48 (dt, 1H), 7.42 (s, 1H), 7.28 (s, 5H), 5.44 (q, 1H), 5.06 - 4.94 (m, 2H), 4.11 (s, 3H), 3.82 (td, 1H), 3.65 (td, 1H). **LCMS** R_t = 1.39 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{24}H_{21}F_4N_6O_4$ $[M+H]^+$ 533.2, found 533.1.

Example 116. Synthesis of 160



[000612] To a mixture of 1-methyl-3-(trifluoromethyl)pyrazole-4-carboxylic acid (100 mg, 0.52 mmol), EDCI (98.76 mg, 0.52 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (125.53 mg, 0.52 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL), and concentrated to give the crude product. The crude product was partially purified by prep-HPLC (Boston Green ODS (150 mm x 30 mm, 5 μm) A = H_2O (0.075% TFA) and B = CH_3CN; 55-75% B over 7 min) to give the impure product. The impure product was triturated with *n*-hexane:DCM (10:1, 2 mL) to give the product (5.0 mg, 13.0μmol, 2% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.00 (s, 1H), 7.88 (br d, 1H), 7.78 (br d, 1H), 7.50 - 7.44 (m, 1H), 7.25 - 7.19 (m, 1H), 6.75 (br s, 1H), 5.64 - 5.55 (m, 1H), 4.00 (s, 3H), 1.74 (d, 3H). **LCMS** R_t = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{14}F_4N_5O_2$ $[M+H]^+$ 384.1, found 384.1.


Example 117. Synthesis of 161



[000613] To a mixture of 1-cyclopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.60 mmol) and EDCI (115.36 mg, 0.60 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (124.68 mg, 0.51 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 μm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 35-65% B over 10 min) to give the product (18.6 mg, 52.3 μmol, 8% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 - 7.84 (m, 2H), 7.81 - 7.75 (m, 1H), 7.47 (dt, 1H), 7.25 - 7.19 (m, 1H), 6.30 (br d,

1H), 5.68 - 5.59 (m, 1H), 3.60 - 3.53 (m, 1H), 2.53 (s, 3H), 1.73 (d, 3H), 1.15 - 1.09 (m, 2H), 1.08 - 1.02 (m, 2H). **LCMS** R_t = 1.16 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{19}FN_5O_2$ $[M+H]^+$ 356.1, found 356.1.


Example 118. Synthesis of 162



A-124 162

[000614] To a mixture of 3-tert-butyl-1-methyl-pyrazole-4-carboxylic acid (100 mg, 0.55 mmol) and EDCI (105.2 mg, 0.55 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (113.71 mg, 0.47 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL) and concentrated to give the crude product. The crude product was purified by prep-HPLC (Phenomenex Gemini-NX (150 mm x 30 mm, 5 µm), A = H_2O (0.04% NH_3H_2O + 10 mM NH_4HCO_3) and B = CH_3CN; 43-73% B over 8.5 mins) to give the product (28.5 mg, 76.6µmol, 14% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (td, 1H), 7.80 - 7.75 (m, 1H), 7.64 (s, 1H), 7.47 (dt, 1H), 7.25 - 7.19 (m, 1H), 6.26 (br d, 1H), 5.62 - 5.54 (m, 1H), 3.87 (s, 3H), 1.72 (d, 3H), 1.42 (s, 9H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{23}FN_5O_2$ $[M+H]^+$ 372.2, found 372.1.

Example 119. Synthesis of 163




A-215 163

[000615] To a mixture of 1-(2,2-difluoroethyl)-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.53 mmol) and EDCI (100.82 mg, 0.53 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (128.15 mg, 0.53 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with the addition of 1N HCl (5 mL), extracted with EtOAc (5 mL), and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm), A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 30-60% B over


10 min) to give the product (16.9 mg, 44.7 µmol, 8% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.89 - 7.86 (m, 2H), 7.80 - 7.76 (m, 1H), 7.47 (dt, 1H), 7.25 - 7.19 (m, 1H), 6.34 (br d, 1H), 6.26 - 5.96 (m, 1H), 5.68 - 5.59 (m, 1H), 4.42 (dt, 2H), 2.54 (s, 3H), 1.74 (d, 3H). **LCMS** R_t = 1.18 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}F_3N_5O_2$ $[M+H]^+$ 380.1, found 380.0.

Example 120. Synthesis of 164



[000616] To a mixture of 3-methyl-1-(2,2,2-trifluoroethyl)pyrazole-4-carboxylic acid (100 mg, 0.48 mmol), EDCI (92.1 mg, 0.48 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (117.07 mg, 0.48 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with 1N HCl (5 mL) and then extracted with EtOAc (5 mL) and concentrated to give the crude product. The crude product was purified by prep-HPLC (Boston Green ODS (150 mm x 30 mm, 5 µm); A = H_2O (0.075% TFA) and B = CH_3CN; 53-73% B over 8 min) to give the impure product. The impure product was purified by prep-TLC (PE:EtOAc = 3:1) to give the product (8.25 mg, 20.8 µmol, 4% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.93 (s, 1H), 7.90 - 7.86 (m, 1H), 7.80 - 7.76 (m, 1H), 7.48 (dt, 1H), 7.26 - 7.20 (m, 1H), 6.37 (br d, 1H), 5.68 - 5.60 (m, 1H), 4.70 - 4.63 (m, 2H), 2.57 - 2.53 (m, 3H), 1.75 (d, 3H). **LCMS** R_t = 1.21 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{16}F_4N_5O_2$ $[M+H]^+$ 398.1, found 398.0.


Example 121. Synthesis of 165



[000617] To a mixture of 1-ethyl-3-(trifluoromethyl)pyrazole-4-carboxylic acid (100 mg, 0.48 mmol), EDCI (92.1 mg, 0.48 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (117.07 mg, 0.48 mmol) and


the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with 1N HCl (5 mL), extracted with EtOAc (5 mL) and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm); A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-70% B over 10 min) to give the impure product. The impure product was purified by prep-TLC (PE:EtOAc = 3:1) to give the product (4.7 mg, 11.9µmol, 2% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.04 (s, 1H), 7.88 (d, 1H), 7.78 (br d, 1H), 7.47 (dt, 1H), 7.25 - 7.18 (m, 1H), 6.76 (br s, 1H), 5.65 - 5.55 (m, 1H), 4.25 (q, 2H), 1.74 (d, 3H), 1.55 (t, 3H). **LCMS** R_t = 1.25 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{16}F_4N_5O_2$ $[M+H]^+$ 398.1, found 398.0.

Example 122. Synthesis of 166



[000618] To a mixture of 1-methyl-3-tetrahydropyran-4-yl-pyrazole-4-carboxylic acid (100 mg, 0.48 mmol), EDCI (91.19 mg, 0.48 mmol) in CH_3CN (3 mL) was added (1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (115.91 mg, 0.48 mmol) and the mixture was stirred at 0 °C for 3 hours. The reaction was quenched with 1N HCl (5 mL), extracted with EtOAc (5 mL) and concentrated to give the crude product. The crude product was purified by prep-HPLC (Welch Xtimate C18 (150 mm x 25 mm, 5 µm); A = H_2O (0.075% TFA) and B = CH_3CN; 37-62% B over 7.5 min) then the product was re-dissolved in CH_2Cl_2 (5 mL), washed with water (5 mL), brine (5 mL), dried over Na_2SO_4, filtered and concentrated to give the product (11.4 mg, 28.5 µmol, 6% yield) as a solid. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (td, 1H), 7.81 - 7.75 (m, 1H), 7.71 (s, 1H), 7.47 (dt, 1H), 7.25 - 7.19 (m, 1H), 6.24 (br d, 1H), 5.66 - 5.57 (m, 1H), 4.08 - 4.01 (m, 2H), 3.91 (s, 3H), 3.60 - 3.51 (m, 2H), 3.49 - 3.40 (m, 1H), 1.99 - 1.87 (m, 4H), 1.72 (d, 3H). **LCMS** R_t = 1.12 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{20}H_{23}FN_5O_3$ $[M+H]^+$ 400.2, found 400.1.


Example 123. Synthesis of 167



157 167

[000619] To a mixture of benzyl N-[2-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]ethyl]carbamate (500 mg, 0.94 mmol) in HBr/AcOH (0.5 mL,33%), and the mixture was stirred at 25 °C for 1 hour. The mixture was concentrated to give the crude product. The crude product was purified by purified by prep-HPLC (Boston Green ODS (150 mm x 30 mm, 5 μm); A = H_2O (0.075% TFA) and B = CH_3CN; 33-63% B over 7 min) to give the product (123.9 mg, 0.31 mmol, 33% yield) as a solid. **LCMS** R_t = 1.03 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{16}H_{15}F_4N_6O_2$ $[M+H]^+$399.1, found 399.0. **^{1}H NMR** (400MHz, DMSO-d_6) δ_H = 7.87 (d, 1H), 7.81 - 7.71 (m, 1H), 7.64 (dt, 1H), 7.55 - 7.42 (m, 2H), 5.29 (t, 1H), 4.12 (s, 3H), 3.26 - 3.08 (m, 2H).


Example 124. Synthesis of 168



167 168


[000620] To a solution of N-[2-amino-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (180 mg, 0.45 mmol) and Et_3N (0.19 mL, 1.36 mmol) in DCM (2 mL) was added methyl carbonochloridate (42.71 mg, 0.45 mmol) and the reaction mixture was stirred at 25 °C for 2 hours. The reaction was quenched with sat. NH_4Cl (10 mL) and then the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by purified by prep-HPLC (Boston Green ODS (150 mm x 30 mm, 5 μm); A = H_2O (0.075% TFA) and B = CH_3CN; 53-83% B over 9 min) to give the product (10.82 mg, 23.7μmol, 5% yield) as a solid. **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{18}H_{17}F_4N_6O_4$ $[M+H]^+$457.1, found 457.0. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 8.36 (br d, 1H), 7.87 (d, 1H), 7.78 (td, 1H), 7.47 (dt, 1H), 7.23 (dt, 1H), 7.02 (s, 1H), 5.49 (br d, 1H), 5.24 (br s, 1H), 4.23 (s, 3H), 3.98 - 3.83 (m, 2H), 3.75 (s, 3H).

Example 125. Synthesis of 169



[000621] To a solution of N-[2-amino-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (60 mg, 0.15 mmol) and Et_3N (0.06 mL, 0.45 mmol) in DCM (2 mL) was added methanesulfonyl chloride (34.51 mg, 0.30 mmol) and the reaction mixture was stirred at 25 °C for 2 hours. The reaction was quenched with sat. NH_4Cl (10 mL) and then the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5 µm); A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 40-67% B over 9 min) to give the product (9.54 mg, 19.7µmol, 13% yield) as a solid. **LCMS** R_t = 1.23 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{17}F_4N_6O_4S$ $[M+H]^+$477.1, found 477.0. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.87 (d, 1H), 7.77 (td, 1H), 7.62 (br d, 1H), 7.54 - 7.44 (m, 1H), 7.26 - 7.21 (m, 1H), 7.04 (s, 1H), 5.61 - 5.55 (m, 1H), 5.13 (br t, 1H), 4.24 (s, 3H), 4.00 - 3.90 (m, 1H), 3.90 - 3.80 (m, 1H), 3.09 (s, 3H).

Example 126. Synthesis of 170



[000622] **A-221:** To a solution of benzyl carbonochloridate (8.93 g, 52.34 mmol) in 1,4-dioxane (50 mL) at 0 °C was added 2-amino-4-hydroxy-butanoic acid (5 g, 41.97 mmol) in water (200 mL). The mixture was stirred for 0.5 hour then $NaHCO_3$ (9.99 g, 118.95 mmol) was added portionwise and the reaction mixture was stirred at 20 °C for 16 hours. The mixture was washed with EtOAc (50 mL x 2). The aqueous layer was acidified to pH 2 with HCl (1N) and extracted with EtOAc (50 mL). The combined organic layers were washed

with water (30 mL) and brine (30 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (9.68 g) as a an oil. **LCMS** Rt = 0.64 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{12}H_{16}NO_5$ $[M+H-44]^+$ 210.1, found 210.1.

[000623] **A-222:** To a solution of 2-(benzyloxycarbonylamino)-4-hydroxy-butanoic acid (3 g, 10.36 mmol) in MeCN (60 mL) at 0 °C was added Ag_2O (12.01 g, 51.82 mmol) followed by CH_3I (9.68 mL, 155.45 mmol). The reaction mixture was warmed and stirred at 20 °C for 48 hours. The reaction mixture was filtered through Celite, eluted with EtOAc (50 mL x 2) and the filtrate was concentrated to give the product. The crude product was purified by flash chromatography on silica gel (DCM in PE = 0% to 50%) to give the product (1.25 g, 2 mmol, 22% yield) as an oil. **LCMS** Rt = 2.1 min in 4 min chromatography, 10-80AB, MS ESI calcd. $C_{14}H_{20}NO_5$ $[M+H]^+$ 282.13, found 281.9.

[000624] **A-223:** To a solution of methyl 2-(benzyloxycarbonylamino)-4-methoxy-butanoate (2.78 g, 9.88 mmol) in THF (28 mL) and water (28 mL) was added $LiOH{\cdot}H_2O$ (1.24 g, 29.65 mmol). The mixture was stirred at 25 °C for 16 hours. The mixture was washed with EtOAc (10 mL x 2). The aqueous layer was acidified to pH ~2 with HCl (1N) and then extracted with EtOAc (50 mL). The combined organic layers were washed with water (30 mL) and brine (20 mL), dried over anhydrous Na_2SO_4, filtered and concentrated to give the crude product (780 mg) as a an oil. **LCMS** Rt = 1.76 min in 4 min chromatography, 10-80AB, MS ESI calcd. $C_{13}H_{18}NO_5$ $[M+H]^+$ 268.11, found 267.9.


[000625] **A-224:** A mixture of 2-(benzyloxycarbonylamino)-4-methoxy-butanoic acid (780 mg, 2.92 mmol), 3-fluoro-N-hydroxy-benzamidine (494.81 mg, 3.21 mmol) and DCC (601.17 mg, 2.92 mmol) in 1,4-dioxane (10 mL) was stirred at 100 °C for 16 hours. After cooling to room temperature, the mixture was concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (DCM in PE = 0% to 40%) to give the product (720 mg, 1.13 mmol, 39% yield) as an oil. **LCMS** Rt = 0.91 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{20}H_{21}FN_3O_4$ $[M+H]^+$ 386.14, found 386.2.

[000626] **A-225:** To a solution of benzyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-methoxy-propyl]carbamate (230 mg, 0.60 mmol) in acetic acid (6 mL) was added HBr/AcOH (0.5 mL, 33%). Then the mixture was stirred at 25 °C for 1 hours. The mixture was concentrated to give the crude product (200 mg, 0.62 mmol) as an oil. **LCMS** Rt = 0.67 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{12}H_{15}FN_3O_2$ $[M+H]^+$ 252.1, found 252.0.

[000627] **170:** To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (265 mg, 0.14 mmol), Et_3N (0.06 mL, 0.46 mmol) and HATU (69.21 mg, 0.18 mmol) in

DCM (10 mL) was added 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-3-methoxy-propan-1-amine (200 mg, 0.09 mmol). Then the mixture was stirred at 25 °C for 16 hours. The mixture was diluted with H_2O (10 mL) and extracted with DCM (10 mL x 2). The combined organic phase was washed with brine (10 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified by prep-HPLC (Waters XBridge (150 mm x 25 mm, 5μμm); A = H_2O (10 mM NH_4HCO_3) and B = CH_3CN; 43-73% B over 10 min) to give the product (22.83 mg, 0.05 mmol) as a an oil. **^{1}H NMR** (400MHz $CDCl_3$) δ_H = 7.95 - 7.85 (m, 2H), 7.82 - 7.75 (m, 1H), 7.47 (dt, 1H), 7.26 - 7.19 (m, 1H), 6.82 (s, 1H), 5.67 - 5.58 (m, 1H), 4.23 (s, 3H), 3.71 - 3.62 (m, 1H), 3.61 - 3.52 (m, 1H), 3.43 (s, 3H), 2.51 - 2.39 (m, 1H), 2.38 - 2.29 (m, 1H). **LCMS** Rt = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{18}H_{18}F_4N_5O_3$ $[M+H]^+$ 428.13, found 428.0.

Example 127. Synthesis of 171



[000628] **A-227:** To a mixture of 2-amino-3-sulfanyl-propanoic acid hydrochloride (3 g, 19.03 mmol, 1.00 eq) in methanol (20 mL) was added NaOMe (4.11 g, 76.13 mmol, 4.00 eq), and the mixture was stirred at 25 °C for 1 h and then MeI (4.05 g, 28.55 mmol, 1.5 eq) was added. The mixture was stirred at 25 °C for 16 hours under N_2. Isopropyl ether (100 mL) was added and the precipitated solid was collected by filtration, washed with cold ethanol and dried to give the product (3 g) as a solid.**^{1}HNMR** (400 M Hz, D_2O) 3.93 - 3.85 (m, 1 H), 3.10 - 2.40 (m, 2 H), δ_H = 2.13 (s, 3 H).

[000629] **A-228:** To a mixture of 2-amino-3-methylsulfanyl-propanoic acid (2 g, 14.8 mmol, 1 eq) and $NaHCO_3$ (4.97 g, 59.18 mmol, 4 eq) in 1,4-dioxane (20 mL) and H_2O (5 mL) was added di-tert butyl dicarbonate (6.46 g, 29.59 mmol, 1 eq), and the mixture was stirred at 25 °C for 5 hours. The reaction mixture quenched with sat. $NaHCO_3$ and extracted with EtOAc (50 mL x 3). The combined organic phrase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the product (2.7 g). **^{1}HNMR** (400MHz, $CDCl_3$) δ_H = 10.42 - 11.05 (m, 1 H) 5.30 - 5.61 (m, 1 H) 4.13 - 4.64 (m, 1 H) 2.69 - 3.08 (m, 2 H) 2.09 (s, 3 H) 1.32 - 1.43 (m, 9 H).

[000630] **A-229:** To a mixture of 2-(tert-butoxycarbonylamino)-3-methylsulfanyl-propanoic acid (500 mg, 2.12 mmol, 1 eq) in 1,4-dioxane (5 mL) was added CDI (1.37 g, 8.5 mmol, 4 eq) and the mixture was stirred at 25 °C for 30 min and then 3-fluoro-N-hydroxy-benzamidine (655.08 mg, 4.25 mmol) was added and the mixture was stirred at 25 °C for 2 hours under N2. The mixture was warmed to 90 °C and stirred for 2 hours under N_2. After cooling to 25 °C, the mixture was extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the product (290 mg).**LCMS** R_t = 1.01 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{12}H_{12}FN_3O_3S$ $[M-56+H]^+$ 297.9, found 297.9.


[000631] **A-230:** To a mixture of tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methylsulfanyl-ethyl]carbamate (290 mg, 0.82 mmol) in 1,4-dioxane (5 mL) and water (2 mL) was added oxone (1 g, 1.64 mmol), and the mixture was stirred at 25 °C for 2 hours. Then the mixture was concentrated and the residue was extracted with EtOAc (30 mL x 3). The combined organic phase was washed with brine (50 mL), dried over $NaSO_4$, filtered and concentrated to give the product (330 mg). **LCMS** R_t = 0.92 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. for $C_{16}H_{20}FN_3O_5S$ Na $[M+Na]^+$ 408.0, found 408.0.

[000632] **A-231:** A mixture of tert-butyl N-[1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methylsulfonyl-ethyl]carbamate (330 mg, 0.86 mmol) in 4M HCl/1,4-dioxane (3 M, 6 mL) was stirred at 25 °C under N_2 for 16 hours to give a mixture. The reaction was quenched with sat. $NaHCO_3$ (20 mL) and then extracted with EtOAc (30 mL x 2). The combined organic phase was washed with brine (15 mL), dried over Na_2SO_4, filtered and concentrated to give the product (300 mg). ^{1}**HNMR** (400MHz, $CDCl_3$) δ_H = 9.24 - 9.68 (m, 2 H) 7.86 - 7.92 (m, 1 H) 7.75 - 7.81 (m, 1 H) 7.64 - 7.73 (m, 1 H) 7.47 - 7.55 (m, 1 H) 5.31 - 5.60 (m, 1 H) 4.08 - 4.18 (m, 2 H) 3.18 - 3.27 (m, 3 H).

[000633] **171:** To a mixture of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.53 mmol, 1 eq) in MeCN (5 mL) was added EDCI (302.38 mg, 1.58 mmol, 3 eq) and the mixture was stirred at 25 °C for 5 min and then 1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-2-methylsulfonyl-ethanamine (153 mg, 0.79 mmol, 1.5 eq) was added. The mixture was stirred at 25 °C for 16 hours under N_2. The reaction mixture was extracted with EtOAc (30 mL x 3). The combined organic phrase was washed with brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product, the crude product was purified by flash chromatography on silica gel (EtOAc in PE, 0% to 10%) to give the product (110 mg). The impure product was triturated from n-hexane (5 mL) to give the product (86.28 mg, 0.19 mmol).^{1}**HNMR** (400MHz, $CDCl_3$) δ_H = 7.84 - 7.95 (m, 1 H) 7.72 - 7.82 (m,

1 H) 7.43 - 7.66 (m, 2 H) 7.23 - 7.27 (m, 1 H) 6.90 - 7.06 (m, 1 H) 5.83 - 6.09 (m, 1 H) 4.19 - 4.28 (m, 3 H) 4.05 - 4.15 (m, 1 H) 3.84 - 3.96 (m, 1 H) 2.98 - 3.09 (m, 3 H). **LCMS** R_t = 1.26 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{17}H_{16}F_4N_5O_4S$ $[M+H]^+$ 462.0, found 462.0.

Example 128. Synthesis of 172, 173, 174, and 175



[000634] **A-233:** To a solution of 2-amino-2-tetrahydrofuran-2-yl-acetic acid (300 mg, 2.07 mmol) and $NaHCO_3$ (434.06 mg, 5.17 mmol) in 1,4-dioxane (4 mL) and water (1 mL) was added Boc_2O (676.58 mg, 3.1 mmol) at 25 °C. The mixture was stirred at 30 °C for 2 hours. The mixture was poured into water (30 mL) and acidified with 30% citric acid to pH ~2 and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4 filtered and concentrated to give crude product (500 mg) as an oil. **^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 5.24 (d, 1H), 4.47 - 4.26 (m, 1H), 4.16 - 4.07 (m, 1H), 4.02 - 3.91 (m, 1H), 3.89 - 3.76 (m, 1H), 2.15 - 1.88 (m, 4H), 1.46 (d, 9H).

[000635] **A-234:** To a solution of 2-(tert-butoxycarbonylamino)-2-tetrahydrofuran-2-yl-acetic acid (400 mg, 1.63 mmol) in 1,4-dioxane (6 mL) was added CDI (528.89 mg, 3.26 mmol) and the mixture was stirred at 25°C for 30 mins. Then 3-fluoro-N-hydroxy-benzamidine (301.66 mg, 1.96 mmol) was added to the solution and the mixture was stirred at 25 °C for 3 hours and then stirred at 100 °C for 16 hours. The mixture was cooled to room temperature and poured into water (30 mL) and extracted with EtOAc (20 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4 filtered and concentrated to give the crude product. The crude was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product (320 mg, 0.80 mmol, 49% yield) as an oil. **LCMS** Rt = 0.92 min in 1.5 min chromatography, 5-95AB MS ESI calcd. For $C_{14}H_{15}FN_3O_4$ $[M\text{-}tBu+H]^+$ 308.16, found 308.2.

[000636] **A-235:** To tert-butyl N-[[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-tetrahydrofuran-2-yl-methyl] carbamate (320 mg, 0.88 mmol) was added 4M HCl/1,4 dixoane (5 mL, 20 mmol) at 25 °C. The mixture was stirred at 25 °C for 16 hours. The

reaction mixture was concentrated to give the crude product (263 mg) as a solid.**LCMS** Rt = 0.75 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. For $C_{13}H_{15}FN_3O_2$ $[M+H]^+$ 264.11, found 263.9.

[000637] **A-236:** To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (204.39 mg, 1.05 mmol) and TEA (0.43 mL, 3.07 mmol) in DCM (5 mL) was added HATU (500.47 mg, 1.32 mmol) at 25°C. The mixture was stirred at 25 °C for 15 mins and then [3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]-tetrahydrofuran-2-yl-methanamine hydrochloride (263 mg, 0.88 mmol) was added to the solution. The mixture was stirred at 25 °C for 16 hrs. The mixture was poured into water (30 mL) and extracted with DCM (10 mL x 2). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4 filtered and concentrated to give the crude product. The crude was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 20%) to give the product, a mixture of 4 diastereomers (260 mg, 0.58 mmol, 66% yield) as an oil. **LCMS** Rt = 1.02 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. For $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$ 440.1, found 440.0.

[000638] **172, 173, 174, and 175:** Analytical SFC: Analysis by SFC (Phenomenex Cellulose-2 (100 mm × 4.6 mm, 3 µm); mobile phase: A: CO_2 B: methanol (0.05% DEA); gradient: from 5% to 40% of B in 4 min and hold 40% for 2.5 min, then 5% of B for 1.5 min; flow rate: 2.8 mL/min, column temp: 35 °C; ABPR: 1500 psi). Showed four peaks at = 1.56 min, 1.64 min, 2.04 min and 2.27 min). The isomers were partially separated by SFC (Phenomenex Cellulose-2 (250 mm x 30 mm, 5µµm); A = CO_2 and B = MeOH (0.1% NH_3H_2O); 35 °C; 60 mL/min; 15% B; 11 min run; 150 injections, Rt of peak 1 = 4.4 min, peak 2 = 5.3 min, peak 3 = 7.4 min and peak 4 = 9.2 min) Stereoisomers 1 and 2 were randomly assigned as **172** and **173** (65 mg) (Rt = 1.56 and 1.64 min in analytical SFC) as a solid, and the stereoisomer isomer 3, randomly assigned as **174** (32.25 mg, 0.074 mmol) (Rt = 2.04 min in analytical SFC) as a solid and the stereoisomer 4, randomly assigned as **175** (50 mg, impure) as a solid. **175** (50 mg, impure) was further purified by SFC (Phenomenex Cellulose-2 (250 mm x 30 mm, 5 µm); A = CO_2 and B = MeOH (0.1% NH_3H_2O); 35 °C; 60 mL/min; 15% B; 11 min run; 80 injections, Rt = 9.2 min) to give the isomer 4 (26.86 mg, 0.06 mmol) (Rt = 2.27 min in analytical SFC) as a solid. Separation for Stereoisomers 1 and 2 (which were randomly assigned as **172**and **173**)

[000639] Analysis by SFC (Daicel CHIRALPAK AD-3 (150 mm × 4.6 mm, 3 µm) mobile phase: A: CO_2 B: ethanol (0.05% DEA); gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5 min, hold 5% of B for 1.5 min; flow rate: 2.5 mL/min, column temp: 35 °C). Showed two peaks at = 1.94 min and 2.39 min). The mixture of

isomers 1 and 2 was separated by SFC (Phenomenex Amylose-1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = 0.1% NH_3H_2O EtOH; 35 °C; 50 mL/min; 15% B; 9 min run; 80 injections, Rt of peak 1 = 4.7 min and peak 2 = 6.6 min) to give the isomer 1, randomly assigned as **173** (28.78 mg,0.065 mmol) (Rt = 1.94 min in analytical SFC) as a solid, and isomer 2, randomly assigned as **172** (28.63 mg,0.065 mmol) (Rt = 2.39 min in analytical SFC) as a solid.


[000640] **172: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.90 (d, 1H), 7.80 (d, 1H), 7.51 - 7.42 (m, 1H), 7.25 - 7.21 (m, 1H), 7.09 (d, 1H), 6.98 (s, 1H), 5.58 (dd, 1H), 4.44 – 4.39 (m, 1H), 4.21 (s, 3H), 3.78 (t, 2H), 2.25 - 2.13 (m, 1H), 2.12 - 2.01 (m, 1H), 1.97 - 1.87 (m, 1H), 1.73 - 1.62 (m, 1H). **LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$ 440.1, found 439.9.

[000641] **173: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.79 (d, 1H), 7.49 – 7.44 (m, 1H), 7.25 - 7.18 (m, 1H), 6.95 (s, 1H), 6.88 (d, 1H), 5.58 (dd, 1H), 4.56 - 4.52 (m, 1H), 4.24 (s, 3H), 4.04 - 3.96 (m, 1H), 3.90 - 3.81 (m, 1H), 2.24 - 2.14 (m, 1H), 2.07 - 1.96 (m, 2H), 1.92 - 1.80 (m, 1H).**LCMS** R_t = 1.37 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$ 440.1, found 439.9.

[000642] **174: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.88 (d, 1H), 7.80 (d, 1H), 7.49 - 7.44 (m, 1H), 7.25 - 7.18 (m, 1H), 6.96 (s, 1H), 6.89 (d, 1H), 5.59 (dd, 1H), 4.56 - 4.52 (m, 1H), 4.24 (s, 3H), 4.03 - 3.96 (m, 1H), 3.90 - 3.81 (m, 1H), 2.24 - 2.13 (m, 1H), 2.06 - 1.95 (m, 2H), 1.91 – 1.82 (m, 1H).**LCMS** R_t = 1.36 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$ 440.1, found 439.9.

[000643] **175: ^{1}H NMR** (400MHz, $CDCl_3$) δ_H = 7.90 (d, 1H), 7.80 (d, 1H), 7.51 - 7.45 (m, 1H), 7.25 - 7.21 (m, 1H), 7.10 (d, 1H), 6.98 (s, 1H), 5.59 (dd, 1H), 4.44 - 4.39 (m, 1H), 4.21 (s, 3H), 3.78 (t, 2H), 2.23 - 2.15 (m, 1H), 2.10 - 2.03 (m, 1H), 1.97 - 1.87 (m, 1H), 1.72 - 1.63 (m, 1H).**LCMS** R_t = 1.35 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{18}F_4N_5O_3$ $[M+H]^+$ 440.1, found 440.0.

Example 129. Synthesis of 176



Synthesis of N'-hydroxy-3-(hydroxymethyl)benzimidamide (A-238)

[000644] To a stirred solution of 3-(hydroxymethyl)benzonitrile (2.0 g, 14.72 mmol) in ethanol (25 mL) was added hydroxylamine hydrochloride (2.56 g, 36.8 mmol) and DIPEA (6.14 mL, 44.16 mmol) at room temperature under nitrogen. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The residue was treated with water (25 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-238** (1.47 g). The crude compound was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(3-(hydroxymethyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-239)

[000645] To a stirred solution of compound **A-238** (1.47 g, 8.86 mmol) in DMF (25 mL) was added DL-Boc-alanine (1.68 g, 8.86 mmol), DIPEA (1.85 mL, 13.29 mmol) and HATU (5.05 g, 13.29 mmol) at 10 °C under nitrogen. The reaction mixture was slowly warmed to room temperature, stirred for 2 h and then heated at 110 °C for 6 h. The reaction mixture was cooled to room temperature, diluted with water (50 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (2 x 20 mL), dried over Na_2SO_4 and concentrated. The crude product was purified by column chromatography on silica gel with 25% EtOAc/PE to afford compound **A-239** (1.61 g, 5.05 mmol, 57% yield).**^{1}H NMR (400 MHz, DMSO-d_6):** δ 7.99 (s, 1H), 7.86 (m, 1H), 7.79 (d, 1H), 7.51 (m, 2H), 5.39 (t, 1H), 4.99-4.96 (m, 1H), 4.59 (d, 2H), 1.52 (d, 3H), 1.41 (s, 9H).

Synthesis of (3-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)phenyl)methanol (A-240)


[000646] To a stirred solution of compound **A-239** (1.61 g, 5.05 mmol) in DCM (30 mL) was added TFA (1.85 mL) at 0 °C under nitrogen. The reaction mixture was slowly warmed to room temperature and stirred for 12 h. The mixture was concentrated under

reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-240** (0.71 g). The crude compound was used for the next step without further purification.

Synthesis of N-(1-(3-(3-(hydroxymethyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (176)

[000647] To a solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (500 mg, 2.52 mmol) in DCM (10 mL) was added compound **A-240** (663 mg, 3.03 mmol) followed by TEA (1.05 mL, 7.57 mmol) and T3P (50% in ethyl acetate, 3.0 mL, 5.05 mmol) at 10 °C under nitrogen. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was diluted with water (25 mL) and extracted with dichloromethane (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude product was purified by flash column chromatography on silica get with 40% EtOAc/pet ether to afford **176** (430 mg, 1.08 mmol, 43% yield) as a liquid. **HPLC**: Rt 4.19 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm); Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 396.2 (M+H), Rt 2.03 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.52 (brs, 1H), 8.05 (s, 1H), 7.94-7.90 (m, 1H), 7.57-7.54 (m, 2H), 7.51 (s, 1H), 5.54-5.49 (m, 1H), 5.44 (brs, 1H), 4.64 (s, 2H), 4.19 (s, 3H), 1.73 (d, 3H).

Example 130. Synthesis of 177




176 177

Synthesis of N-(1-(3-(3-(hydroxymethyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)-N,1-dimethyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (177)

[000648] To a stirred solution of compound **176** (150 mg, 0.38 mmol) in THF (9.0 mL) was added dimethyl sulphate (95 mg, 0.76 mmol) and KOH (64 mg, 1.14 mmol) under nitrogen atmosphere. The reaction mixture was heated at 65 °C for 3 h. The reaction mixture was cooled to room temperature, treated with water (25 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by flash column chromatography on silica get with

50% EtOAc/PE to afford **177** (38 mg, 0.092 mmol, 24% yield) as a liquid. **HPLC**: Rt 4.31 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 410.2 (M+H), Rt 1.96 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 8.01 (s, 1H), 7.89 (m, 1H), 7.56-7.53 (m, 2H), 7.22 and 7.09 (s, 1H), 5.90 and 5.59 (m, 1H), 5.39 (t, 1H), 4.61-4.57 (m, 2H), 3.98 (s, 3H), 3.08 and 2.93 (s, 3H), 1.76 (d, 3H).

Example 131. Synthesis of 178



Synthesis of 3-bromo-N'-hydroxybenzimidamide (A-242)

[000649] To a solution of 3-bromobenzonitrile (10 g, 55.25 mmol) in ethanol (200 mL) was added hydroxylamine hydrochloride (11.45 g, 164.82 mmol) followed by DIPEA (29.25 mL, 164.82 mmol). The reaction mixture was heated at 70 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (100 mL) and extracted with ethyl acetate (2 x 150 mL). The organic layer was washed with brine (40 mL), dried over Na_2SO_4 and concentrated to afford compound **A-242** (10 g). It was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(3-bromophenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-243)

[000650] To a solution of compound **A-242** (2.0 g, 9.3 mmol) in 1,4-dioxane (40 mL) was added 2-(tert-butoxycarbonylamino)propanoic acid (1.76 g, 9.3 mmol) followed by DCC (2.1 g, 10.23 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The residue was treated with water (30 mL) and extracted with EtOAc (2 x 50 mL). The organic layer was washed with brine (40 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 25% EtOAc/PE to afford compound **A-243** (1.3 g, 3.54 mmol, 38% yield). **LCMS**: 366.0 (M-H) and 368.1 (M+2-H). Rt 2.73 min; Column:

ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of tert-butyl (1-(3-(3-(cyanomethyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-244)

[000651] To a PTFE screw-capped vial containing 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)isoxazole (0.83 g, 4.24 mmol) and compound **A-243** (1.3 g, 3.54 mmol) in DMSO (20 mL) was added KF (0.62 g, 10.59 mmol) and water (0.19 mL, 10.62 mmol). The reaction mixture was degassed and $Pd(dppf)Cl_2$.DCM (0.29 g, 0.35 mmol) was added under nitrogen. The reaction mixture was heated at 130 °C for 16 h. The reaction mixture was cooled to room temperature and filtered over celite. The filtrate was treated with saturated NaCl solution (25 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was taken up in methanol (10 mL) and KF (0.62 g, 10.59 mmol) was added. The reaction mixture was stirred at room temperature for 3 h. The mixture was concentrated, the residue was treated with water (20 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 20% EtOAc/PE to afford compound **A-244** (242 mg, 0.74 mmol, 20% yield). **LCMS**: 327.1 (M-H), Rt 2.20 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm. Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of 2-(3-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)phenyl)acetonitrile (A-245)


[000652] To a stirred solution of compound **A-244** (120 mg, 0.37 mmol) in DCM (4 mL) was added TFA (0.5 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 8 h. The mixture was concentrated under reduced pressure and treated with saturated $NaHCO_3$ solution (10.0 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (10 mL), dried over Na_2SO_4 and concentrated to afford compound **A-245** (66 mg). It was used for the next step without further purification.

Synthesis of N-(1-(3-(3-(cyanomethyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (178)

[000653] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (62 mg, 0.32 mmol) in THF (10 mL) was added T3P (0.35 mL, 0.58 mmol) and TEA (0.12 mL, 0.87 mmol). The reaction mixture was stirred at room temperature for 10 min and compound **A-245** (66 mg, 0.27 mmol) was added. The reaction mixture was stirred for 16 h at room temperature. The reaction mixture was concentrated, treated with water (30

mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica get with 45% EtOAc/PE to afford **178** (26 mg, 0.06 mmol, 21% yield) as a solid. **HPLC**: Rt 4.78 min, 98.4%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 405.1 (M+H), Rt 2.41 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.09-8.07 (m, 2H), 7.56-7.54 (m, 2H), 6.96 (s, 1H), 6.73 (d, 1H), 5.67-5.59 (m, 1H), 4.25 (s, 3H), 3.86 (s, 2H), 1.79 (d, 3H).

Example 132. Synthesis of 179



Synthesis of N'-hydroxy-3-(methylsulfonyl)benzimidamide (A-247)

[000654] To a solution of 3-methylsulfonylbenzonitrile (**A-246**, 1.0 g, 5.52 mmol)n ethanol (20 mL)was added hydroxylamine hydrochloride (1.15 g, 16.56 mmol)and DIPEA (2.94 mL, 16.56 mmol). The reaction mixture was then heated at 70 ºCfor 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (25 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-247** (0.6 g). The compound was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(3-(methylsulfonyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-248)

[000655] To a solution of compound **A-247** (0.6 g, 2.8 mmol) in 1,4-dioxane (20 mL) was added 2-(tert-butoxycarbonylamino)propanoic acid (0.77 g, 4.05 mmol) and DCC (846.15 mg, 4.11 mmol). The reaction mixture was heated at 100 ºC for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (20 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by

column chromatography on silica gel with 20% EtOAc/PE to afford compound **A-248** (0.5 g, 1.35 mmol, 48% yield). **LCMS**: 366.1 (M-H), Rt 2.05 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of 1-(3-(3-(methylsulfonyl)phenyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-249)


[000656] To a stirred solution of compound **A-248** (300 mg, 0.81 mmol) in DCM (3 mL) was added TFA (0.5 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 8 h. The reaction mixture was concentrated and treated with saturated $NaHCO_3$ solution (20 mL). The mixture was extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-249** (200 mg). The compound was used for the next step without further purification.

Synthesis of 1-methyl-N-(1-(3-(3-(methylsulfonyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (179)

[000657] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (160 mg, 0.82 mmol) in THF was added T3P (1.33 mL, 2.24 mmol) and triethylamine (0.52 mL, 3.74 mmol) at 0 °C. The reaction mixture was stirred for 10 min and compound **A-249** (200 mg, 0.75 mmol) was added. The reaction mixture was slowly warmed to room temperature and stirred for 16 h. The reaction mixture was concentrated, and the residue was treated with water (20 mL). The mixture was extracted with EtOAc (2 x 25 mL) and washed with brine (20 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica get with 35% EtOAc/PE to afford **179** (120 mg, 0.27 mmol, 36% yield) as a solid.**HPLC**: Rt 4.43 min, 99.8%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm

[000658] Mobile phase: A: 0.1% TFA in water, B: ACN; Flow Rate: 2.0 mL/min; **LCMS**: 444.0 (M+H), Rt 2.12 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min; **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.67 (t, 1H), 8.39-8.36 (m, 1H), 8.13-8.10 (m, 1H), 7.74 (t, 1H), 6.99 (s, 1H), 6.78 (d, 1H), 5.67-5.60 (m, 1H), 4.25 (s, 3H), 3.13 (s, 3H), 1.80 (d, 3H).

Example 133. Synthesis of 180



Synthesis of N'-hydroxy-3-sulfamoylbenzimidamide (A-251)

[000659] To a stirred solution of 3-cyanobenzenesulfonamide (1.0 g, 5.49 mmol) in ethanol (15 mL) at room temperature was added hydroxylamine hydrochloride (572 mg, 8.23 mmol) and N,N-diisopropylethylamine (2.87 mL, 16.47 mmol). The reaction mixture was heated at 77 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (20 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-251** (800 mg). The compound was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(3-sulfamoylphenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-252)

[000660] To a stirred solution of compound **A-251** (800 mg, 3.72 mmol) in 1,4-dioxane (30 mL) was added 2-(tert-butoxycarbonylamino)propanoic acid (703 mg, 3.72 mmol) and DCC (843 mg, 4.09 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica get with 60% EtOAc/PE to afford compound **A-252** (685 mg, 1.86 mmol, 50% yield) as a solid. **LCMS**: 367.2 (M-H), Rt 1.91 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of 3-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)benzenesulfonamide (A-253)


[000661] To a stirred solution of compound **A-252** (500 mg, 1.35 mmol) in DCM (12 mL) was added TFA (2.5 mL). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was concentrated and treated with saturated sodium bicarbonate

solution (10 mL). The mixture was extracted with EtOAc (2 x 20 mL) and washed with brine (10 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-253** (260 mg). The compound was used for the next step without further purification.

Synthesis of 1-methyl-N-(1-(3-(3-sulfamoylphenyl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (180)

[000662] To a stirred solution of compound **A-253** (260 mg, 0.97 mmol) and 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (225 mg, 1.16 mmol) in THF (10 mL) was added T3P (1.73 mL, 2.91 mmol) and DIPEA (0.4 mL, 2.91 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude product was purified by preparative HPLC to afford **180** (80 mg, 0.17 mmol, 18% yield) as a solid. prep-HPLC method: Rt 10.59; Column: XBridge C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.19 min, 99.4%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 445.1 (M+H), Rt 2.03 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.48 (d, 1H), 8.45 (t, 1H), 8.24-8.22 (m, 1H), 8.05-8.03 (m, 1H), 7.79 (t, 1H), 7.57 (brs, 2H), 7.46 (s, 1H), 5.51-5.47 (m, 1H), 4.14 (s, 3H), 1.70 (d, 3H).

Example 134. Synthesis of 181



Synthesis of methyl 3-(N'-hydroxycarbamimidoyl)benzoate (A-255)

[000663] To a solution of methyl 3-cyanobenzoate (10 g, 62.05 mmol) in ethanol (200 mL) was added hydroxylamine hydrochloride (12.94 g, 186.15 mmol) followed by DIPEA (6.28 mL, 62.05 mmol). The reaction mixture was stirred at 70 °C for 16 h. The reaction

mixture was cooled to room temperature and concentrated. The mixture was treated with water (100 mL) and extracted with ethyl acetate (2 x 80 mL). The organic layer was washed with brine (40 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-255** (8.47 g). It was used for the next step without further purification.

Synthesis of methyl 3-(5-(1-((tert-butoxycarbonyl)amino)ethyl)-1,2,4-oxadiazol-3-yl)benzoate (A-256)

[000664] To a solution of compound **A-255** (1.68 g, 8.7 mmol) in 1,4-dioxane (60 mL) was added 2-(tert-butoxycarbonylamino)propanoic acid (2.11 g, 11.17 mmol) and DCC (2.33 g, 11.33 mmol). The reaction mixture heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The residue was treated with water (30 mL) and extracted with EtOAc (2 x 40 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica get with 25% EtOAc/PE to afford compound **A-256** (1.2 g, 3.4 mmol, 33% yield). **LCMS**: 346.1 (M-H), Rt 2.45 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of methyl 3-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)benzoate (A-257)


[000665] To a stirred solution of compound **A-256** (1.2 g, 3.45 mmol) in DCM (3 mL) was added TFA (2.1 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and concentrated. The mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (1 x 10 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-257** (524 mg). The compound was used for the next step without further purification.

Synthesis of methyl 3-(5-(1-(1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamido)ethyl)-1,2,4-oxadiazol-3-yl)benzoate (181)

[000666] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (0.5 mL, 2.22 mmol) in THF (10 mL) was added T3P (6.0 mL, 10.11 mmol) and TEA (0.84 mL, 6.07 mmol). The reaction mixture was stirred at room temperature for 10 min and compound **A-257** (500 mg, 2.02 mmol) was added. The reaction mixture was stirred at room temperature for 16 h and concentrated. The mixture was treated water (20 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (1 x 10 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 25% EtOAc/PE to afford **181** (350 mg, 0.81 mmol, 40% yield). **HPLC**: Rt 5.04 min, 98.3%Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile

phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 424.1 (M+H), Rt 2.43 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.73 (s, 1H), 8.26 (d, 1H), 8.19 (d, 1H), 7.59 (t, 1H), 6.96 (s, 1H), 6.79 (d, 1H), 5.65-5.60 (m, 1H), 4.24 (s, 3H), 3.97 (s, 3H), 1.78 (d, 3H).

Example 135. Synthesis of 182




181 182

Synthesis of 3-(5-(1-(1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamido)ethyl)-1,2,4-oxadiazol-3-yl)benzoic acid (182)

[000667] To a stirred solution of compound **181** (330 mg, 0.78 mmol) in THF (8 mL), methanol (2 mL) and water (1 mL) was added $LiOH.H_2O$ (170.62 mg, 4.07 mmol) at room temperature. The reaction mixture was stirred at room temperature for 8 h and concentrated. The mixture was treated with 1 N HCl (8 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (1 x 20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 75% EtOAc/PE to afford **182** (250 mg, 0.60 mmol, 78% yield). **HPLC**: Rt 4.37 min, 98.9%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 410.1 (M+H), Rt 2.02 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min; **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.48 (d, 1H), 8.56 (s, 1H), 8.22 (d, 1H), 8.14 (d, 1H), 7.70 (t, 1H), 7.46 (s, 1H), 5.51-5.47 (m, 1H), 4.14 (s, 3H), 1.69 (d, 3H).

Example 136. Synthesis of 183




182 183

Synthesis of N-(1-(3-(3-(dimethylcarbamoyl)phenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (183)

[000668] To a stirred solution of compound **182** (150 mg, 0.37 mmol) in THF (15 mL) was added T3P (0.43 mL, 0.73 mmol) and TEA (0.15 mL, 1.1 mmol). The reaction mixture was stirred at room temperature for 10 min and dimethylamine (2.0 M in THF, 0.92 mL, 1.83 mmol) was added. The reaction mixture was stirred at room temperature for 16 h and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (1 x 15 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 75% EtOAc/PE to afford **183** (65 mg, 0.14 mmol, 40% yield). **HPLC**: Rt 4.19 min, 98.7%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 437.2 (M+H), Rt 2.03 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.10-8.07 (m, 2H), 7.54-7.52 (m, 2H), 7.45 (d, 1H), 7.06 (s, 1H), 5.62-5.55 (m, 1H), 4.23 (s, 3H), 3.15 (s, 3H), 2.99 (s, 3H), 1.74 (d, 3H).

Example 137. Synthesis of 184



Synthesis of 2-methoxyisonicotinonitrile (A-259)

[000669] To a solution of 2-chloropyridine-4-carbonitrile (10 g, 72.18 mmol) in 1,4-dioxane (20 mL) was added sodium methoxide (25% in methanol, 16.4 mL, 72.18 mmol) at room temperature. The reaction mixture was heated at 100 °C for 5 h. The reaction mixture was cooled to room temperature and left in the refrigerator at 0 °C for 16 h. The precipitate was filtered and washed with cold methanol (60 mL). The filtrate was concentrated and ice water (50 mL) was added. The precipitate was filtered, washed with water (100 mL) and

dried to afford compound **A-259** (4.43 g) as a solid. It was used for the next step without further purification.

Synthesis of N'-hydroxy-2-methoxyisonicotinimidamide (A-260)

[000670] To a stirred solution of compound **A-259** (4.43 g, 33.15 mmol) in ethanol (50 mL) was added DIPEA (9.22 mL, 66.3 mmol) and hydroxylamine hydrochloride (4.6 g, 66.3 mmol). The reaction mixture was heated at 100 °C for 16 h and concentrated. The mixture was treated with water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (1 x 20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-260** (3.75 g) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(2-methoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-261)

[000671] To a solution of compound **A-260** (3.75 g, 22.43 mmol) in 1,4-dioxane (40 mL) was added DL-Boc-alanine (4.67 g, 24.68 mmol) and DCC (5.08 g, 24.68 mmol) at room temperature. The reaction mixture was then heated at 100 °Cfor 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (20 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentration. The crude was purified by column chromatography on silica gel with 30% EtOAc/PE to afford compound **A-261** (5.3 g, 16.3 mmol, 72% yield) as a solid. **LCMS**: 321.1 (M+H), Rt 2.33 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of 1-(3-(2-methoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-262)

[000672] To a stirred solution of compound **A-261** (5.3 g, 16.3 mmol) in DCM (50 mL) was added TFA (10.02 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 1 h. The reaction mixture was concentrated, treated with 10% $NaHCO_3$ solution (30 mL) and extracted with DCM (2 x 40 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-262** (3 g) as a solid. It was used for the next step without further purification.


Synthesis of N-(1-(3-(2-methoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (184)


[000673] To a solution of compound **A-262** (250 mg, 1.13 mmol) in DMF (10 mL) was added DIPEA (0.47 mL, 3.4 mmol) and 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (241.7 mg, 1.25 mmol) followed by HATU (861 mg, 2.26 mmol). The

reaction mixture was stirred at room temperature for 12 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude product was purified by column chromatography on silica gel with 45% EtOAc/PE to afford **184** (122 mg, 0.30 mmol, 27% yield) as a solid. **HPLC**: Rt 4.65 min, 99.3%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 397.1 (M+H), Rt 2.35 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min; **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.47 (d, 1H), 8.39 (d, 1H), 7.54-7.52 (m, 1H), 7.45 (s, 1H), 7.31 (s, 1H), 5.50-5.46 (m, 1H), 4.13 (s, 3H), 3.93 (s, 3H), 1.68 (d, 3H).

Example 138. Synthesis of 145 and 146

a) Synthesis of 145





Synthesis of tert-butyl N-[(1R)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-342)

[000674] To a stirred solution of compound **A-341** (320 mg, 1.91 mmol) in 1,4-dioxane (10.0 mL) was added Boc-D-alanine (434 mg, 2.3 mmol) and DCC (591 mg, 2.87 mmol) at room temperature. The reaction mixture was then stirred at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **A-342** (592 mg, 1.85 mmol, 96% yield). **LCMS**: 321.3 (M+H), Rt 2.32 min, Column: ZORBAX XDB C-

18 (50 x 4.6 mm), 3.5 µm m Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min


Synthesis of (1R)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (A-343)

[000675] To a stirred solution of compound **A-342** (592 mg, 1.85 mmol) in DCM (7.0 mL) was added TFA (1.43 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-343** (340 mg). The compound was used for the next step without further purification.

Synthesis of 2-methyl-N-[(1R)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (145)

[000676] To a stirred solution of compound **A-343** (340 mg, 1.54 mmol)in THF (10.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (299 mg, 1.54 mmol) followed by T3P (2.76 mL, 4.63 mmol) and Et_3N (0.64 mL, 4.63 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **145** (346 mg, 0.86 mmol, 55% yield) as a solid. Prep. HPLC method: Rt 13.0; Column: YMC C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.63 min, 99.3%, Column: XBridge C8 (50 x 4.6) mm, 3.5 µm, Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 397.1 (M+H), Rt 2.34 min, Column: ZORBAX XDB C-18 (50 x 4.6) mm, 3.5 µm, Mobile phase: A: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min **Chiral method:** Rt 2.54 min, SFC column: CHIRALCEL OJ-H; mobile phase: 80:20 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.47 (d, 1H), 8.39 (d, 1H), 7.54-7.52 (dd, 1H), 7.45 (s, 1H), 7.31 (s, 1H), 5.50-5.45 (m, 1H), 4.13 (s, 3H), 3.92 (s, 3H), 1.67 (d, 3H).

b) Synthesis of 146



Synthesis of tert-butyl N-[(1S)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-344)

[000677] To a stirred solution of compound **A-341** (320.mg, 1.9 mmol) in 1,4-dioxane (10.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (359 mg, 1.9 mmol) and DCC (430 mg, 2.09 mmol) at room temperature. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (30 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **A-344** (530 mg, 1.65 mmol, 87% yield). **LCMS**: 321.1 (M+H), Rt 2.32 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm , Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min


Synthesis of (1S)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine (A-345)

[000678] To a stirred solution of compound **A-344** (530 mg, 1.65 mmol) in DCM (10.0 mL) was added TFA (1.94 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-345** (310 mg). The compound was used for the next step without further purification.

Synthesis of 2-methyl-N-[(1S)-1-[3-(2-methoxy-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (146)

[000679] To a stirred solution of compound **A-345** (310 mg, 1.4 mmol) in THF (10.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (273 mg, 1.41 mmol) followed by T3P (2.51mL, 4.22mmol) and Et_3N (0.59mL, 4.22mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **146** (133 mg, 0.32 mmol, 23% yield) as a solid. Prep. HPLC method: Rt 11.21; Column: Atlantis (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.64 min, 96.8%, Column: XBridge C8 (50 x 4.6) mm, 3.5 µm, Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 397.1 (M+H), Rt 2.20 min, Column: XBridge C8 (50 x 4.6) mm, 3.5 µm, Mobile phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 1.99 min, SFC column: Chiralcel OJ-H; mobile phase: 80:20 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.32-8.30 (dd, 1H), 7.57-7.55 (dd, 1H), 7.415-7.41 (dd, 1H), 7.25 (s, 1H), 5.53 (q, 1H), 4.19 (s, 3H), 3.98 (s, 3H), 1.77 (d, 3H).

Example 139. Synthesis of 185



Synthesis of N',3-dihydroxybenzimidamide (A-264)

[000680] To a stirred solution of 3-hydroxybenzonitrile (5.0 g, 41.97 mmol) in ethanol (50 mL) was added DIPEA (11.67 mL, 83.95 mmol) and hydroxylamine hydrochloride (5.83 g, 83.95 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The reaction mixture was treated with water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-264** (4.4 g). The compound was used for the next step without further purification.

Synthesis of tert-butyl (1-(3-(3-hydroxyphenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-265)

[000681] To a solution of compound **A-264** (2.5 g, 16.44 mmol) in 1,4-dioxane (100 mL) was added DL-Boc-alanine (3.11 g, 16.44 mmol) and DCC (3.72 g, 18.08 mmol) and the reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (40 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 15% EtOAc/PE to afford compound **A-265** (4.2 g, 13.6 mmol, 82% yield) as a solid. **LCMS**: 304.2 (M-H), Rt 2.00 min Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of 3-(5-(1-((tert-butoxycarbonyl)amino)ethyl)-1,2,4-oxadiazol-3-yl)phenyl acetate (A-266)

[000682] To a solution of compound **A-265** (500 mg, 1.62 mmol) in DCM (15 mL) was added pyridine (0.39 mL, 4.83 mmol), DMAP (19.81 mg, 0.16 mmol) and acetic anhydride (0.17 mL, 1.78 mmol). The reaction mixture was stirred at room temperature for 3 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **A-266** (500 mg) as a solid. **LCMS**: 346.1 (M-H), Rt 2.32 min Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of 3-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)phenyl acetate (A-267)


[000683] To a solution of compound **A-266** (500 mg, 1.42 mmol) in DCM (10 mL) was added TFA (0.55 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The reaction mixture was concentrated, treated with 10% $NaHCO_3$ solution (10 mL) and extracted with DCM (2 x 20 mL). The organic layer was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-267** (265 mg). The compound was used for the next step without further purification.

Synthesis of 3-(5-(1-(1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamido)ethyl)-1,2,4-oxadiazol-3-yl)phenyl acetate (185)

[000684] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (200 mg, 1.03 mmol) in DMF (10 mL) was added HATU (1.17 g, 3.09 mmol), DIPEA (0.55 mL, 3.09 mmol) and compound **A-267** (247 mg, 0.99 mmol). The

reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 35% EtOAc/PE to afford **185** (68 mg, 0.16 mmol, 15% yield) as a solid. **HPLC**: Rt 4.93 min, 99.9%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: ACN; Flow Rate: 2.0 mL/min. **LCMS**: 424.1 (M+H), Rt 2.42 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.46 (d, 1H), 7.90 (d, 1H), 7.77 (s, 1H), 7.63 (t, 1H), 7.46 (s, 1H), 7.38 (d, 1H), 5.50-5.43 (m, 1H), 4.14 (s, 3H), 2.31 (s, 3H), 1.68 (d, 3H).

Example 140. Synthesis of 186




185 186

Synthesis of N-(1-(3-(3-hydroxyphenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (186)

[000685] To a stirred solution of compound **185** (60 mg, 0.14 mmol) in methanol (10 mL) and water (10 mL) was added $LiOH.H_2O$ (18 mg, 0.43 mmol) at room temperature. The reaction mixture was heated at 50 °C for 1 h. The reaction mixture was cooled to room temperature, treated with 1.5 N hydrochloric acid (2.0 mL) and extracted with EtOAc (2 x 15 mL). The organic layer was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 5% MeOH/DCM to afford **186** (48 mg, 0.12 mmol, 87% yield) as a solid. **HPLC**: Rt 4.40 min, 98.0%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 382.1 (M+H), Rt 2.11 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.86 (s, 1H), 9.44 (d, 1H), 7.45-7.34 (m, 4H), 6.99-6.97 (m, 1H), 5.47-5.43 (m, 1H), 4.14 (s, 3H), 1.66 (d, 3H).


Example 141. Synthesis of 282 (tert-butyl (R)-(1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate)



A-10 282

[000686] To a solution of compound **A-10** (0.25 g, 1.62 mmol) in 1,4-dioxane (10.0 mL) was added (2R)-2-(tert-butoxycarbonylamino)propanoic acid (0.33 g, 1.76 mmol) and DCC (0.37 g, 1.78 mmol the reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated. The mixture was treated with water (15 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 8% EtOAc/PE to afford **282** (270 mg, 0.86 mmol, 53% yield) as a solid. **HPLC**: Rt 5.02 min, 99.3%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 306.1 (M-H), Rt 2.51 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.94 min, SFC column: YMC Amylose-C; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 7.89 (d, 1H), 7.82-7.78 (m, 1H), 7.50-7.45 (m, 1H), 7.25-7.20 (m, 1H), 5.19 (m, 2H), 1.65 (d, 3H), 1.49 (s, 9H).

Example 143. Synthesis of 187



282 A-210 187


Synthesis of (R)-1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-210)

[000687] To a solution of compound **282** (270 mg, 0.88 mmol) in DCM (5 mL) was added TFA (1.3 mL) and the mixture stirred at room temperature for 3 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with DCM (2 x 20 mL). The organic layer was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-210 (**170 mg) as a liquid. The compound was used for the next step without further purification.

Synthesis of (R)-3-(difluoromethyl)-N-(1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-1H-pyrazole-4-carboxamide (187)

[000688] To a solution of compound **A-210** (170 mg, 0.82 mmol) in THF (5 mL) was added 3-(difluoromethyl)-1-methyl-pyrazole-4-carboxylic acid (158.95 mg, 0.90 mmol) followed by T3P (50% in ethyl acetate, 1.47 mL, 2.46 mmol) and Et_3N (0.34 mL, 2.46 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (15 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (15 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 60% EtOAc/PE to afford **187** (85 mg, 0.23 mmol, 28% yield) as a solid. **HPLC**: Rt 4.22 min, 98.6%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 366.1 (M+H), Rt 2.02 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 3.27 min, SFC column: YMC Cellulose-SC; mobile phase: 60:40 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.00 (s, 1H), 7.91-7.89 (m, 1H), 7.82-7.79 (m, 1H), 7.50-7.45 (m, 1H), 7.25-7.20 (m, 1H), 7.03 (brs, 1H), 6.90 (t, 1H), 5.65-5.58 (m, 1H), 3.97 (s, 3H), 1.75 (d, 3H).

Example 144. Synthesis of 188




Synthesis of (S)-1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-138)

[000689] To a solution of compound **281** (300 mg, 0.98 mmol) in DCM (5mL) was added TFA (1.45 mL) and the mixture was stirred at room temperature for 3 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with DCM (2 x 20 mL). The organic layer was washed with brine (10 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-138** (200 mg). The compound was used for the next step without further purification.

Synthesis of (S)-3-(difluoromethyl)-N-(1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-1H-pyrazole-4-carboxamide (188)

[000690] To a solution of compound **A-138** (200 mg, 0.97 mmol) in THF (5.0 mL) was added 3-(difluoromethyl)-1-methyl-pyrazole-4-carboxylic acid (187 mg, 1.06 mmol) followed by T3P (50% in ethyl acetate, 1.73 mL, 2.91 mmol) and Et_3N (0.4 mL, 2.9 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (15 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (15 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 60% EtOAc/PE to afford **188** (140 mg, 0.38 mmol, 39% yield) as a solid. **HPLC**: Rt 4.22 min, 97.9%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 366.1 (M+H), Rt 2.10 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**; Chiral method:** Rt 1.76 min, SFC column: YMC Cellulose-SC; mobile phase: 60:40 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 7.99 (s, 1H), 7.90-7.88 (m, 1H), 7.81-7.78 (m, 1H), 7.49-7.44 (m, 1H), 7.24-7.19 (m, 1H), 7.02 (brs, 1H), 6.89 (t, 1H), 5.64-5.56 (m, 1H), 3.96 (s, 3H), 1.74 (d, 3H).

Example 145. Synthesis of 189




282 **189**

Synthesis of (R)-N-(1-(3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (189)

[000691] To a solution of compound **282** (145 mg, 0.70 mmol) in THF (5.0 mL) was added 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (135 mg, 0.70 mmol) followed by T3P (1.25 mL, 2.1 mmol) and Et_3N (0.29 mL, 2.1 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (20 mL) and extracted with EtOAc (2 x 20 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 60% EtOAc/PE to afford **189** (130 mg, 0.33 mmol, 47%

yield) as a solid. **HPLC**: Rt 5.17 min, 99.0%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 384.1 (M+H), Rt 2.50 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**; Chiral method:** Rt 2.76 min, SFC column: YMC Cellulose-SJ; mobile phase: 85:15 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 7.91-7.88 (m, 1H), 7.81-7.78 (m, 1H), 7.52-7.47 (m, 1H), 7.27-7.22 (m, 1H), 6.94 (s, 1H), 6.69 (d, 1H), 5.64-5.60 (m, 1H), 4.25 (s, 3H), 1.79 (d, 3H).

Example 150. Synthesis of 193




182 193

Synthesis of N-(1-(3-(3-carbamoylphenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (193)

[000692] To a stirred solution of compound **182** (100 mg, 0.24 mmol) in THF (10.0 mL) was added pentafluorophenol (66 mg, 0.36 mmol) and DCC (74 mg, 0.36 mmol). The reaction mixture was stirred at room temperature for 16 h and NH_3 (2.0 M in methanol, 2 mL) was added. The mixture was stirred for 1 h at room temperature and concentrated under reduced pressure. The residue was treated with water (25 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (10 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 45% EtOAc/PE to afford **193** (13 mg, 0.03 mmol, 13% yield) as a solid. **HPLC**: Rt 3.84 min, 99.6%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 409.1 (M+H), Rt 1.88 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, DMSO-d_6):** δ 9.47 (d, 1H), 8.51 (s, 1H), 8.21 (s, 1H), 8.14 (d, 1H), 8.09 (d, 1H), 7.66 (t, 1H), 7.55 (s, 1H), 7.46 (s, 1H), 5.50-5.46 (m, 1H), 4.14 (s, 3H), 1.69 (d, 3H).

Example 151. Synthesis of 194 and 195



Synthesis of (Z)-2-chloro-N'-hydroxyisonicotinimidamide (A-273)

[000693] To a solution of 2-chloropyridine-4-carbonitrile (20 g, 144 mmol) in ethanol (200 mL) was added hydroxylamine hydrochloride (15.05 g, 216 mmol) followed by DIPEA (59.6 mL, 360 mmol). The reaction mixture was heated at 75 °C for 12 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (100 mL) followed by saturated sodium bicarbonate solution (50 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (50 mL), dried over Na_2SO_4 and concentrated to afford compound **A-273** (23 g) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-chloropyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-274)

[000694] To a solution of compound **A-273** (4.0 g, 23.3 mmol) in 1,4-dioxane (50.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (4.78 g, 25.29 mmol) followed by DCC (5.28 g, 25.64 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 7% ethyl acetate/PE to afford compound **A-274** (3.2 g, 9.8 mmol, 42% yield) as a solid. **LCMS**: 325.1 (M+H), Rt 2.36 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate:1.5 mL/min.

Synthesis of tert-butyl (1-(3-(2-(pyrrolidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-275)

[000695] To a solution of compound **A-274** (1.0 g, 3.0 mmol) in NMP (8.0 mL) was added pyrrolidine (429 mg, 6.04 mmol) and irradiated in microwave at 120 °C for 1 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The

mixture was extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-275** (720 mg, 2.0 mmol, 66% yield) as a solid. **LCMS**: 360.2 (M+H), Rt 1.41 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate:1.5 mL/min.

Synthesis of 1-(3-(2-(pyrrolidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-276)

[000696] To a solution of compound **A-275** (400 mg, 1.11 mmol) in DCM (5.0 mL) was added TFA (1.22 mL) at 0 ºC. The reaction mixture was slowly warmed to room temperature and stirred for 3 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-276** (200 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(2-(pyrrolidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (194) and (R)-1-methyl-N-(1-(3-(2-(pyrrolidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (195)


[000697] To a solution of compound **A-276** (200 mg, 0.77 mmol) in THF (10.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (164 mg, 0.85 mmol) followed by T3P (50% in EtOAc, 1.38 mL, 2.31 mmol) and Et_3N (0.32 mL, 2.31 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 30% EtOAc/PE to afford 100 mg of racemic compound. The racemic mixture was separated by SFC purification to afford **194** (35 mg, 0.08 mmol, 10% yield) and **195** (38 mg, 0.09 mmol, 11% yield) as solids. **Chiral method:** SFC column: CHIRALCEL OX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. The stereochemistry of **194** and **195** was randomly assigned.

[000698] **194: HPLC**: Rt 3.46 min, 99.1%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min.

LCMS: 436.1 (M+H), Rt 1.52 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 6.55 min, SFC column: CHIRALCELOX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.47 (d, 1H), 8.24 (d, 1H), 7.45 (s, 1H), 7.06-7.04 (m, 1H), 6.91 (s, 1H), 5.51-5.41 (m, 1H), 4.13 (s, 3H), 3.44 (t, 4H), 1.97 (t, 4H), 1.67 (d, 3H).

[000699] **195: HPLC**: Rt 3.46 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 436.2 (M+H), Rt 1.54 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 7.16 min, SFC column: Chiralcel OX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.47 (s, 1H), 8.24 (d, 1H), 7.45 (s, 1H), 7.05 (d, 1H), 6.91 (s, 1H), 5.47 (m, 1H), 4.13 (s, 3H), 3.44 (m, 4H), 1.97 (m, 4H), 1.67 (d, 3H).

Example 152. Synthesis of 283 & 196



Synthesis of (Z)-N'-hydroxy-3-methoxybenzimidamide (A-278)

[000700] To a stirred solution of 3-methoxybenzonitrile (5.0 g, 37.5 mmol) in ethanol (50.0 mL) was added hydroxylamine hydrochloride (3.91 g, 56.3 mmol) followed by DIPEA (19.6 mL, 112.6 mmol). The reaction mixture was heated at 80 ºC for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (100 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (100 mL), dried over Na_2SO_4 and concentrated to afford compound **A-278** (6.0 g) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(3-methoxyphenyl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (283)

[000701] To a stirred solution of compound **A-278** (2.7 g, 16.2 mmol) in 1,4-dioxane (30 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (3.06 g, 16.2 mmol) and DCC (3.67 g, 17.8 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (100 mL) and extracted with ethyl acetate (2 x 70 mL). The organic layer was washed with brine (100 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 9% EtOAc/PE to afford compound **283** (4.6 g, 14.4 mmol, 88% yield) as a solid. **LCMS**: 318.1 (M-H), Rt 2.45 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(3-methoxyphenyl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-280)


[000702] To a stirred solution of compound **283** (0.5 g, 1.5 mmol) in DCM (10 mL) was added TFA (1.1 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5.0 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-280** (320 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(3-methoxyphenyl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (196)

[000703] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-280** (170 mg, 0.77 mmol). To the reaction mixture TEA (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) were added and the mixture was stirred at RT for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 30% EtOAc/PE to afford **196** (95 mg, 0.24 mmol, 30% yield) as a colourless liquid. **HPLC**: Rt 5.02 min, 99%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 394.1 (M-H), Rt 2.48 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.41 min, SFC column: YMC

Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.45 (d, 1H), 7.59 (d, 1H), 7.52-7.45 (m, 3H), 7.19-7.17 (m, 1H), 5.49-5.42 (m, 1H), 4.14 (s, 3H), 3.84 (s, 3H), 1.67 (d, 3H).

Example 153. Synthesis of 197 and 198



Synthesis of tert-butyl (1-(3-(2-(dimethylamino)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-281)

[000704] To a solution of compound **A-274** (0.5 g, 1.51 mmol) in NMP (2.0 mL) was added dimethylamine solution (2.0 M in THF, 3.77 mL, 7.54 mmol) and irradiated in microwave at 120 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The mixture was extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-281** (220 mg, 0.66 mmol, 43% yield) as a solid. **LCMS**: 334.3 (M+H), Rt 1.43 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of 4-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)-N,N-dimethylpyridin-2-amine (A-282)

[000705] To a solution of compound **A-281** (220 mg, 0.66 mmol) in DCM (5.0 mL) was added TFA (0.72 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 3 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-282** (140 mg). The compound was used for the next step without further purification.


Synthesis of (S)-N-(1-(3-(2-(dimethylamino)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (197) and (R)-N-(1-(3-(2-(dimethylamino)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (198)

[000706] To a solution of compound **A-282** (140 mg, 0.60 mmol) in THF (8.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (116 mg, 0.60 mmol) followed by Et_3N (0.25 mL, 1.8 mmol) and T3P (50% in EtOAc, 1.07 mL, 1.8 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 32% EtOAc/PE to afford 110 mg of racemic compound. The racemic mixture was separated by SFC purification to afford **1977** (24 mg, 0.06 mmol, 9% yield) and **198** (34 mg, 0.08 mmol, 13% yield) as solids. **Chiral method:** SFC column: LUX C3; mobile phase: 80:20 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 220 nm. The stereochemistry of **197** and **198** was randomly assigned.

[000707] **197: HPLC**: Rt 3.23 min, 97.1%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 410.1 (M+H), Rt 1.49 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.88 min, SFC column: LUX C3; mobile phase: 80:20 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.21 (d, 1H), 7.27 (s, 1H), 7.26 (s, 1H), 7.18 (d, 1H), 5.53 (q, 1H), 4.19 (s, 3H), 3.16 (s, 6H), 1.77 (d, 3H).

[000708] **198: HPLC**: Rt 3.21 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 410.1 (M+H), Rt 1.49 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.41 min, SFC column: LUX C3; mobile phase: 80:20 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.21-8.20 (m, 1H), 7.27 (s, 1H), 7.26 (s, 1H), 7.19-7.17 (m, 1H), 5.53 (q, 1H), 4.19 (s, 3H), 3.15 (s, 6H), 1.77 (d, 3H).

Example 154. Synthesis of 199



Synthesis of (Z)-N'-hydroxy-2-methylisonicotinimidamide (A-101)

[000709] To a stirred solution of 2-methylpyridine-4-carbonitrile (1.0 g, 8.46 mmol) in ethanol (20 mL) was added hydroxylamine hydrochloride (0.88 g, 12.7 mmol) followed by DIPEA (4.41 mL, 25.4 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (20 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over Na_2SO_4 and concentrated to afford compound **A-101** (1.1 g) as a a liquid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)carbamate (284)

[000710] To a stirred solution of compound **A-101** (1.1 g, 7.3 mmol) in 1,4-dioxane (20 mL) was added (2S)-2-(tert-butoxycarbonylamino)butanoic acid (1.41 g, 6.9 mmol) followed by DCC (1.57 g, 7.6 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (60 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **284** (1.7 g, 5.3 mmol, 76% yield) as a a liquid. **LCMS**: 319.2 (M+H), Rt 1.89 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.


Synthesis of (S)-1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propan-1-amine (A-284)

[000711] To a stirred solution of compound **284** (700 mg, 2.2 mmol) in DCM (14 mL) was added TFA (2.7 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (30 mL). The mixture was treated with 10% aqueous Na_2CO_3 solution (5.0 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-284** (270 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (199)

[000712] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6 mL) was added compound **A-284** (170 mg, 0.77 mmol). To the reaction mixture TEA (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) were added and the mixture was stirred at RT for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to **199** (51 mg, 0.13 mmol, 16% yield) as a colourless liquid. Prep. HPLC method: Rt 9.30; Column: XBridge (150 x 19 mm), 5.0 µm; Mobile phase: 10 mM NH_4OAc in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.94 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: ACN; Flow Rate: 2.0 mL/min**LCMS**: 395.2 (M+H), Rt 2.01 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.31 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.61 (d, 1H), 7.96 (s, 1H), 7.86 (d, 1H), 7.28 (s, 1H), 5.40-5.36 (m, 1H), 4.19 (s, 3H), 2.64 (s, 3H), 2.28-2.09 (m, 2H), 1.13 (t, 3H).

Example 155. Synthesis of 200



Synthesis of (Z)-N'-hydroxy-4-methylpicolinimidamide (A-286)

[000713] To a stirred solution of 4-methylpyridine-2-carbonitrile (5.0 g, 42.3 mmol) in ethanol (50.0 mL) was added DIPEA (20.9 mL, 126.7 mmol) followed by hydroxylamine hydrochloride (2.94 g, 42.3 mmol). The reaction heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (80 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (100 mL), dried over Na_2SO_4 and concentrated to afford compound **A-286** (5.6 g) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(4-methylpyridin-2-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (285)

[000714] To a stirred solution of compound **A-286** (3.0 g, 19.8 mmol) in1,4-dioxane (30 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (3.74 g, 19.8 mmol) followed by DCC (4.48 g, 21.7 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (80 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (100 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 18% EtOAc/PE to afford compound **285** (3 g, 9.8 mmol, 49% yield) as a solid. **LCMS**: 305.2 (M+H), Rt 2.02 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.


Synthesis of (S)-1-(3-(4-methylpyridin-2-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-288)

[000715] To a stirred solution of compound **285** (700 mg, 2.3 mmol) in DCM (14 mL) was added TFA (1.8 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (30 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5.0 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-288** (230 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(4-methylpyridin-2-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (200)

[000716] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-288** (169 mg, 0.83 mmol). To the reaction mixture TEA (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) were added and the mixture was stirred at RT for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to obtain **200** (58 mg, 0.15 mmol, 19% yield) as a solid. Prep. HPLC method: Rt 11.04; Column: Sunfire C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.83 min, 99.6%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 381.2 (M+H), Rt 2.11 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.38 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.56 (d, 1H), 8.05 (s, 1H), 7.45 (d, 1H), 7.28 (s, 1H), 5.59-5.53 (m, 1H), 4.19 (s, 3H), 2.50 (s, 3H), 1.78 (d, 3H).

Example 156. Synthesis of 286 & 201



Synthesis of 2-isopropoxyisonicotinonitrile (A-290)

[000717] To the isopropyl alcohol (45.0 mL) at 0 °C was added NaH (60% in mineral oil, 952 mg, 23.8 mmol) in small portions. The resulting suspension was stirred for 5 min and 2-chloropyridine-4-carbonitrile (3.0 g, 21.65 mmol) was added in small portions. The reaction mixture was heated at 80 °C for 1 h. The reaction mixture was cooled to 10 °C and treated with ice water (50 mL). The mixture was extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford compound **A-290** (980 mg, 6.0 mmol, 27% yield).

LCMS: 163.1 (M+H), Rt 2.32 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-N'-hydroxy-2-isopropoxyisonicotinimidamide (A-291)

[000718] To a stirred solution of compound **A-290** (0.98 g, 6.0 mmol) in ethanol (20.0 mL) was added hydroxylamine hydrochloride (0.63 g, 9.0 mmol) followed by DIPEA (3.16 mL, 18.13 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The mixture was treated with 10% sodium carbonate solution (10 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-291** (1.1 g). It was used for next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-isopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (286)

[000719] To a stirred solution of compound **A-291** (1.1 g, 5.6 mmol) in 1,4-dioxane (20.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.07 g, 5.6 mmol) and DCC (1.28 g, 6.2 mmol). The reaction mixture was heated to 100 °C for 16 h. The

reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 14% EtOAc/PE to afford compound **286** (1.5 g, 4.3 mmol, 76% yield) as a solid. **LCMS**: 349.1 (M+H), Rt 2.64 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(2-isopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-293)


[000720] To a stirred solution of compound **286** (700 mg, 2.01 mmol) in DCM (14.0 mL) was added TFA (0.77 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (10 mL). The mixture was treated with 10% Na_2CO_3 solution (5.0 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (25 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-293** (280 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-isopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (201)

[000721] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-293** (191 mg, 0.77 mmol) followed by TEA (0.32 mL, 2.32 mmol). To the reaction mixture T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) was added and the mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **201** (280 mg, 0.3 mmol, 38% yield) as a solid. Prep. HPLC method: Rt 9.82; Column: XBridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.27 min, 99.7%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 423.1 (M-H), Rt 2.71 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.29 min, SFC column: Chiralcel OX-H; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz,**

CD_3OD): δ 8.28 (dd, 1H), 7.52-7.50 (m, 1H), 7.33-7.32 (m, 1H), 7.25 (s, 1H), 5.53 (q, 1H), 5.40-5.31 (m, 1H), 4.19 (s, 3H), 1.77 (d, 3H), 1.37 (d, 6H).

Example 157. Synthesis of 287 & 202



Synthesis of 2-methoxy-6-methylisonicotinonitrile (A-295)

[000722] To a stirred solution of 2-chloro-6-methyl-pyridine-4-carbonitrile (1.0 g, 6.55 mmol) in 1,4-dioxane (10.0 mL) was added NaOMe (0.39 g, 7.21 mmol) at room temperature. The reaction mixture was heated at 60 °C for 4 h. The reaction mixture was cooled, treated with ice water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford compound **A-295** (650 mg, 4.3 mmol, 66% yield) as a solid. **LCMS**: 149.2 (M+H), Rt 2.05 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-N'-hydroxy-2-methoxy-6-methylisonicotinimidamide (A-296)

[000723] To a stirred solution of compound **A-295** (650 mg, 4.3 mmol) in ethanol (20.0 mL) was added hydroxylamine hydrochloride (452 mg, 6.51 mmol) followed by DIPEA (2.26 mL, 13.03 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (20 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-296** (680 mg) as a solid. The compound was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-methoxy-6-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (287)

[000724] To a stirred solution of compound **A-296** (680 mg, 3.7 mmol) in 1,4-dioxane (15.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (701 mg, 3.7 mmol) and DCC (840 mg, 4.1 mmol). The reaction mixture was heated at 100 ℃ for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **287** (1.0 g, 3.0 mmol, 80% yield) as a solid. **LCMS**: 335.2 (M+H), Rt 2.55 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(2-methoxy-6-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-298)


[000725] To a stirred solution of compound **287** (400 mg, 1.2 mmol) in DCM (10 mL) was added TFA (0.88 mL) at 0 ℃. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5.0 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-298** (260 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-methoxy-6-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (202)

[000726] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-298** (181 mg, 0.77 mmol) followed by TEA (0.32mL, 2.32mmol). To the reaction mixture T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) was added and the mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (25 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **202** (135 mg, 0.33 mmol, 42% yield) as a solid. Prep. HPLC method: Rt 10.49; Column: Sunfire C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.82 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 411.1 (M+H), Rt 2.54 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.97

min, SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 7.42 (d, 1H), 7.25 (s, 1H), 7.19 (d, 1H), 5.52 (q, 1H), 4.19 (s, 3H), 3.96 (s, 3H), 2.51 (s, 3H), 1.76 (d, 3H).

Example 158. Synthesis of 288 & 203



[000727] Synthesis of 2-cyclopropylisonicotinonitrile (A-300)

[000728] To a solution of 2-chloropyridine-4-carbonitrile (2.0 g, 14.4 mmol) in 1,4-dioxane (25 mL) was added potassium cyclopropyltrifluoroborate (6.41 g, 43.3 mmol) followed by K_2CO_3 (7.98 g, 57.7 mmol) and RuPhos (1.35 g, 2.89 mmol). The resulting mixture was degassed with N_2 gas for 10 min and $Pd(OAc)_2$ (324 mg, 1.44 mmol) was added. The mixture was stirred at 100 °C for 1 h. The reaction mixture was cooled to room temperature and filtered through celite. The filtrate was concentrated under reduced pressure and the crude was purified by column chromatography on silica gel with 15% EtOAc/PE to afford compound **A-300** (1.1 g, 7.6 mmol, 50% yield) as a solid.**LCMS**: 145.1 (M+H), Rt 1.87 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-2-cyclopropyl-N'-hydroxyisonicotinimidamide (A-301)

[000729] To a solution of compound **A-300** (450 mg, 3.1 mmol) in ethanol (15.0 mL) was added hydroxylamine hydrochloride (312 mg, 4.4 mmol) followed by DIPEA (1.49 mL, 8.99 mmol) at room temperature. The reaction mixture was heated at 80 °C for 5 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (30 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (10 mL), dried over Na_2SO_4 and concentrated to afford compound **A-301** (420 mg) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (288)

[000730] To a solution of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (0.44 g, 2.31 mmol) in 1,4-dioxane (10.0 mL) was added compound **A-301** (0.41 g, 2.31 mmol) followed by DCC (0.52 g, 2.55 mmol). The resulting mixture was stirred at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 15% ethyl acetate/PE to afford compound **288** (570 mg, 1.72 mmol, 74% yield) as a solid.**LCMS**: 331.3 (M+H), Rt 2.22 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-303)


[000731] To a solution of compound **288** (410 mg, 1.24 mmol) in DCM (5.0 mL) was added TFA (1.36 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 3 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-303** (240 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (203)

[000732] To a solution of compound **A-303** (240 mg, 1.04 mmol) in THF (8.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (202 mg, 1.04 mmol) followed by Et_3N (0.43 mL, 3.13 mmol) and T3P (50% in EtOAc, 1.86 mL, 3.13 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica get with 60% EtOAc/PE to afford **203** (216 mg, 0.53 mmol, 51% yield) as a solid. **HPLC**: Rt 3.48 min, 97.3%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 407.1 (M+H), Rt 1.77 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5

mL/min. **Chiral method:** Rt 1.42 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, $CDCl_3$):** δ 8.61 (d, \ 1H), 7.79 (s, 1H), 7.68 (dd, 1H), 6.95 (s, 1H), 6.70 (d, 1H), 5.67-5.60 (m, 1H), 4.25 (s, 3H), 2.20-2.13 (m, 1H), 1.80 (d, 3H), 1.14-1.10 (m, 4H).

Example 159. Synthesis of 289 & 204



Synthesis of 2-(methoxymethyl)isonicotinonitrile (A-305)

[000733] To a stirred solution of 2-(hydroxymethyl)pyridine-4-carbonitrile (500 mg, 3.73 mmol) in THF (8.0 mL) was added NaH (60% in mineral oil, 164 mg, 4.1 mmol) at 0 °C in small portions followed by iodomethane (0.23 mL, 3.73 mmol). The reaction temperature was slowly raised to room temperature and stirred for 2 h. The reaction mixture was cooled to 10 °C, treated with ice water (20 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford compound **A-305** (470 mg, 3.17 mmol, 85% yield). **LCMS**: 149.1 (M+H), Rt 1.31 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-N'-hydroxy-2-(methoxymethyl)isonicotinimidamide (A-306)

[000734] To a stirred solution of compound **A-305** (470 mg, 3.17 mmol) in ethanol (10.0 mL) was added hydroxylamine hydrochloride (330 mg, 4.76 mmol) followed by DIPEA (1.57 mL, 9.52 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (30 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20

mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-306** (420 mg) as a colourless liquid. The compound was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(methoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (289)

[000735] To a stirred solution of compound **A-306** (420 mg, 2.32 mmol) in 1,4-dioxane (10.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (438 mg, 2.32 mmol) and DCC (525 mg, 2.55 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated to under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 8% ethyl acetate/PE to afford compound **289** (560 mg,1.6 mmol, 72% yield) as a solid. **LCMS**: 335.3 (M+H), Rt 2.23 min; Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 10 mM NH_4HCO_3 in H_2O, B: ACN; Flow Rate: 1.5 mL/min.

Synthesis of (S)-1-(3-(2-(methoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-308)


[000736] To a stirred solution of compound **289** (420 mg, 1.26 mmol) in DCM (8.0 mL) was added TFA (0.96 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (30 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-308** (210 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(methoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (204)

[000737] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-308** (168 mg, 0.72 mmol). To the reaction mixture TEA (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) were added and the mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **204** (96 mg, 0.23 mmol, 30% yield) as a solid. Prep. HPLC method: Rt 10.75; Column: Atlantis C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt

3.58 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 411.0 (M+H), Rt 2.90 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 10 mM NH_4HCO_3 in H_2O, B: ACN; Flow Rate:0.8 mL/min. **Chiral method:** Rt 1.81 min, SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.67 (d, 1H), 8.12 (s, 1H), 7.95-7.93 (m, 1H), 7.25 (s, 1H), 5.54 (q, 1H), 4.64 (s, 2H), 4.18 (s, 3H), 3.51 (s, 3H), 1.77 (d, 3H).

Example 160. Synthesis of 290 & 205



Synthesis of 2-ethoxyisonicotinonitrile (A-309)

[000738] To a stirred solution of 2-chloropyridine-4-carbonitrile (5.0 g, 36.1 mmol) in 1,4-dioxane (50.0 mL) was added NaOEt (2.46 g, 36.1 mmol) at one portion. The reaction mixture was heated at 60 °C for 4 h. The reaction mixture was cooled to room temperature, treated with ice cold water (50 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica get with 6% EtOAc/PE to afford compound **A-309** (3.5 g, 23.5 mmol, 65% yield). **LCMS**: 149.1 (M+H), Rt 2.06 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-2-ethoxy-N'-hydroxyisonicotinimidamide (A-310)

[000739] To a stirred solution of 2-ethoxypyridine-4-carbonitrile (1.5 g, 10.12 mmol) in ethanol (30.0 mL) was added hydroxylamine hydrochloride (1.06 g, 15.19 mmol) followed by DIPEA (5.29 mL, 30.37 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (30 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with

brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-310** (1.7 g). The compound was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-ethoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (290)

[000740] To a stirred solution of (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.78 g, 9.38 mmol) in 1,4-dioxane (34.0 mL) was added compound **A-310** (1.7 g, 9.38 mmol) and DCC (2.13 g, 10.32 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (100 mL) and extracted with ethyl acetate (2 x 100 mL). The organic layer was washed with brine (100 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 14% EtOAc/PE to afford compound **290** (2.4 g, 7.07 mmol, 75% yield) as a solid. **LCMS**: 335.1 (M+H), Rt 3.22 min; Column: XBridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 10 mM NH_4HCO_3 in H_2O, B: ACN; Flow Rate:0.8 mL/min.

Synthesis of (S)-1-(3-(2-ethoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-312)


[000741] To a stirred solution of compound **290** (400 mg, 1.2 mmol) in DCM (10.0 mL) was added TFA (0.9 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-312** (240 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-ethoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (205)

[000742] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added compound **A-312** (181 mg, 0.77 mmol) followed by TEA (0.32 mL, 2.32 mmol). To the reaction mixture T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) was added and the mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **205** (186 mg, 0.45 mmol, 58% yield) as a solid. Prep. HPLC method: Rt 9.81; Column: Sunfire (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.93 min, 99.1%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA

in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 411.2 (M+H), Rt 3.22 min, Column: XBridge C8 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 10 mM NH_4HCO_3 in H_2O, B: ACN; Flow Rate:0.8 mL/min. **Chiral method:** Rt 2.01 min, SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm.**^{1}H NMR (400 MHz, CD_3OD):** δ 8.27 (d, 1H), 7.54-7.53 (m, 1H), 7.37 (s, 1H), 7.25 (s, 1H), 5.53 (q, 1H), 4.40 (q, 2H), 4.19 (s, 3H), 1.77 (d, 3H), 1.42 (t, 3H).

Example 161. Synthesis of 291 & 206



Synthesis of (Z)-2-(difluoromethyl)-N'-hydroxyisonicotinimidamide (A-313)

[000743] To a stirred solution of 2-(difluoromethyl)pyridine-4-carbonitrile (500 mg, 3.24 mmol) in ethanol (20.0 mL) was added hydroxylamine hydrochloride (338 mg, 4.87 mmol) followed by DIPEA (1.69 mL, 9.73 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over Na_2SO_4 and concentrated to afford compound **A-313** (580 mg) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(difluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (291)

[000744] To a stirred solution of compound **A-313** (580 mg, 3.05 mmol) in 1,4-dioxane (15 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic) acid (586 mg, 3.1 mmol) followed by DCC (702 mg, 3.41 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature, concentrated and was diluted with ethyl acetate (30 mL). The organics washed with water (2 x 20 mL) and then with brine (20 mL). The organic layer was dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 8% EtOAc/PE to afford compound **291** (1.0 g,

2.93 mmol, 94% yield) as a solid. **LCMS**: 341.1 (M+H), Rt 2.24 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.


Synthesis of (S)-1-(3-(2-(difluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-315)

[000745] To a stirred solution of compound **291** (400 mg, 1.18 mmol) in DCM (8.0 mL) was added TFA (0.9 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (30 mL). The mixture was treated with 10% aqueous $NaHCO_3$ solution (5 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-315** (260 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(difluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (206)

[000746] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6 mL) was added compound **A-315** (200 mg, 0.77 mmol). To the reaction mixture TEA (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) were added and the mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **206** (97 mg, 0.23 mmol, 29% yield) as a solid. Prep. HPLC method: Rt 9.60; Column: Sunfire C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 4.68 min, 99%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 417.2 (M+H), Rt 2.29 min,Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.74 min; SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm.**^{1}H NMR (400 MHz, CD_3OD):** δ 8.85 (d, 1H), 8.29 (s, 1H), 8.17 (d, 1H), 7.26 (s, 1H), 6.84 (t, 1H), 5.56 (q, 1H), 4.19 (s, 3H), 1.79 (d, 3H).

Example 162. Synthesis of 292 & 207



A-272 A-316 A-317 292

A-319 207

Synthesis of 2-(methylamino)isonicotinonitrile (A-316)

[000747] To a stirred solution of NaH (0.32 g, 7.94 mmol) in THF (20.0 mL) was added cyclopropanol (0.5 g, 8.66 mmol) and 2-chloropyridine-4-carbonitrile (1.0 g, 7.22 mmol) at 0 °C under a nitrogen atmosphere. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was cooled to 10 °C and treated with ice water (10 mL). The mixture was extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 15% ethyl acetate/PE to afford compound **A-316** (550 mg, 3.43 mmol, 43% yield).

LCMS: 161.1 (M+H), Rt 2.37 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (Z)-2-cyclopropoxy-N'-hydroxyisonicotinimidamide (A-317)

[000748] To a stirred solution of compound **A-316** (550 mg, 3.43 mmol) in ethanol (15 mL) was added DIPEA (1.7 mL, 10.31 mmol) and hydroxylamine hydrochloride (357 mg, 5.15 mmol) at room temperature under nitrogen atmosphere. The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (15 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-317** (646 mg). It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-cyclopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (292)

[000749] To a stirred solution of compound **A-317** (646 mg, 3.35 mmol) in 1,4-dioxane (10 mL) was added DCC (937 mg, 4.55 mmol) and (2S)-2-(tert-

butoxycarbonylamino)propanoic acid (783 mg, 4.14 mmol) at room temperature. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 40% ethyl acetate/PE to afford compound **292** (550 g, 1.58 mmol, 47% yield). **LCMS**: 347.2 (M+H), Rt 2.38 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(2-cyclopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-319)


[000750] To a stirred solution of compound **292** (550 mg, 1.58 mmol) in DCM (10 mL) was added TFA (1.1 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-319** (250 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-cyclopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (207)

[000751] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (10.0 mL) was added compound **A-319** (190 mg, 0.77 mmol) followed by Et_3N (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **207** (130 mg, 0.30 mmol, 39% yield) as a solid. Prep. HPLC method: Rt 13.48; Column: XBridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min.**HPLC**: Rt 4.80 min, 99.2%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 423.0 (M+H), Rt 3.14 min, Column: XBridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 10 mM NH_4HCO_3 in H_2O, B: ACN; Flow Rate: 0.8 mL/min; **Chiral method:** Rt 1.17 min, SFC column: YMC Cellulose-SJ; mobile phase: 60:40 (A: B), A =

liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.34-8.32 (m, 1H), 7.63-7.62 (m, 1H), 7.52 (s, 1H), 7.24 (s, 1H), 5.53 (q, 1H), 4.24-4.20 (m, 1H), 4.18 (s, 3H), 1.76 (d, 3H), 0.86-0.83 (m, 2H), 0.79-0.77 (m, 2H).

Example 163. Synthesis of 293 & 208



Synthesis of 2-(3,3-difluorocyclobutoxy)isonicotinonitrile (A-320)

[000752] To a stirred solution of NaH (158 mg, 3.97 mmol) in THF (10.0 mL) was added 3,3-difluorocyclobutanol (468 mg, 4.33 mmol) and 2-chloropyridine-4-carbonitrile (500 mg, 3.61 mmol) at 0 °C under a nitrogen atmosphere. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was cooled to 10 °C and treated with ice water (10 mL). The mixture was extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 20% ethyl acetate/PE to afford compound **A-320** (320 mg, 1.5 mmol, 42% yield). **^{1}H NMR (400 MHz, CD_3OD):** δ 8.30 (dd, 1H), 7.13 (dd, 1H), 7.03 (m, 1H), 5.20-5.15 (m, 1H), 3.20-3.10 (m, 2H), 2.81-2.69 (m, 2H).

Synthesis of (Z)-2-(3,3-difluorocyclobutoxy)-N'-hydroxyisonicotinimidamide (A-321)

[000753] To a stirred solution of compound **A-320** (320 mg, 1.52 mmol) in ethanol (10.0 mL) was added DIPEA (589 mg, 4.57 mmol) and hydroxylamine hydrochloride (95.27mg, 3.97mmol) at room temperature under nitrogen. The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 40 mL). The organic layer was washed with brine (30 mL), dried over Na_2SO_4 and concentrated to afford compound **A-321** (326 mg). It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(3,3-difluorocyclobutoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (293)

[000754] To a stirred solution of compound **A-321** (320 mg, 1.32 mmol) in 1,4-dioxane (10.0 mL) was added DCC (294 mg, 1.43 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (248 mg, 1.32 mmol) at room temperature under nitrogen. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 40% ethyl acetate/PE to afford compound **293** (404 mg, 1.02 mmol, 77% yield).**LCMS**: 397.2 (M+H), Rt 2.64 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min.

Synthesis of (S)-1-(3-(2-(3,3-difluorocyclobutoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-323)


[000755] To a stirred solution of compound **293** (350 mg, 0.88 mmol) in DCM (10.0 mL) was added TFA (1.5 mL) at 0 °C under nitrogen. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-323** (245 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(3,3-difluorocyclobutoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (208)

[000756] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (10.0 mL) was added compound **A-323** (230.mg, 0.77 mmol) followed by Et_3N (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 40% ethyl acetate/PE to afford **208** (160 mg, 0.33 mmol, 44% yield) as a solid. **HPLC**: Rt 5.46 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min.

LCMS: 473.1 (M+H), Rt 2.49 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 1.15 min, SFC column: YMC Cellulose-SJ; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.30 (d, 1H), 7.58 (dd, 1H), 7.43 (s, 1H), 7.24 (s, 1H), 5.52 (q, 1H), 5.21-5.17 (m, 1H), 4.18 (s, 3H), 3.19-3.09 (m, 2H), 2.80-2.68 (m, 2H), 1.76 (d, 3H).

Example 164. Synthesis of 209 and 210



Synthesis of tert-butyl (1-(3-(2-(piperidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-324)

[000757] To a solution of compound **A-274** (1.2 g, 3.7 mmol) in NMP (10.0 mL) was added piperidine (629 mg, 7.39 mmol), and the mixture was irradiation in microwave at 120 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The mixture was extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-324** (962 mg, 2.58 mmol, 69% yield) as a solid.**LCMS**: 374.3 (M+H), Rt 2.00 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of 1-(3-(2-(piperidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-325)

[000758] To a stirred solution of compound **A-324** (887 mg, 2.38 mmol) in DCM (10.0 mL) was added TFA (1.0 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and

treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-325** (316 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(2-(piperidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (209) and (R)-1-methyl-N-(1-(3-(2-(piperidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (210)


[000759] To a solution of A-325 (210 mg, 0.77 mmol) in THF (5.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) followed by Et_3N (0.5 mL, 3.58 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The racemic mixture was separated by SFC purification to afford **209** (19.0 mg, 0.04 mmol, 5% yield) and **210** (20.0 mg, 0.05 mmol, 5% yield) as solids. Chiral method: SFC column: CHIRALCEL OX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. The stereochemistry of **209** and **210** was randomly assigned.

[000760] **209: HPLC**: Rt 3.68 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 450.3 (M+H), Rt 2.05 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 4.8 min, SFC column: CHIRALCEL OX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.21 (d, 1H), 7.40 (s, 1H), 7.26 (s, 1H), 7.19-7.18 (m, 1H), 5.53 (q, 1H), 4.19 (s, 3H), 3.64-3.61 (m, 4H), 1.77 (d, 3H), 1.73-1.67 (m, 6H).

[000761] **210: HPLC**: Rt 3.67 min, 99.1%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 450.2 (M+H), Rt 2.05 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 5.05 min, SFC column: CHIRALCEL OX-H; mobile phase: 90:10 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.21-8.20 (m, 1H), 7.39 (s, 1H), 7.25 (s,

1H), 7.18-7.17 (m, 1H), 5.52 (q, 1H), 4.18 (s, 3H), 3.63-3.60 (m, 4H), 1.76 (d, 3H), 1.72-1.67 (m, 6H).

Example 165. Synthesis of 294 & 211



Synthesis of 2-(2,2-difluoroethoxy)isonicotinonitrile (A-324a)

[000762] NaH (60% in mineral oil, 866 mg, 21.65 mmol) was added in small portions to a solution of 2,2-difluoroethanol (2.13 g, 25.98 mmol) in THF (35 mL) at 0 °C . The resulting suspension was stirred for 5 min and 2-chloropyridine-4-carbonitrile (3.0 g, 21.65 mmol) was added to the mixture in small portions. The mixture was stirred at room temperature for 3 h. The reaction mixture was cooled to 10 °C and treated with ice water (50 mL). The mixture was extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford compound **A-324a** (2.9 g, 15.7 mmol, 72% yield). **LCMS**: 185.1 (M+H), Rt 2.46 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of (Z)-2-(2,2-difluoroethoxy)-N'-hydroxyisonicotinimidamide (A-325a)

[000763] To a stirred solution of compound **A-324a** (2.9 g, 15.7 mmol) in ethanol (20 mL) was added hydroxylamine hydrochloride (1.63 g, 23.44 mmol) and DIPEA (8.16 mL, 46.87 mmol). The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (30 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-325a** (3.4 g) as a solid. It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(2,2-difluoroethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (294)

[000764] To a solution of compound **A-325a** (3.3 g, 15.3 mmol) in 1,4-dioxane (30 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (2.9 g, 15.3 mmol) followed by DCC (3.48 g, 16.88 mmol). The reaction mixture was heated at 100 ºC for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (50 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 7% ethyl acetate/PE to afford compound **294** (5.2 g, 14.0 mmol, 91% yield) as a solid. **LCMS**: 371.1 (M+H), Rt 2.48 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-1-(3-(2-(2,2-difluoroethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-327)


[000765] To a stirred solution of compound **294** (0.9 g, 2.43 mmol) in DCM (15.0 mL) was added TFA (2.0 mL) at 0 ºC. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-327** (297 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(2,2-difluoroethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (211)

[000766] To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (8.0 mL) was added compound **A-327** (240 mg, 0.88 mmol) followed by Et_3N (0.32 mL, 2.32 mmol) and T3P (50% in EtOAc, 1.38 mL, 2.32 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **211** (146 mg, 0.32 mmol, 42% yield) as a solid. Prep. HPLC method: Rt 9.18; Column: XBridge (150 x 19 mm), 5.0 µm; Mobile phase: 10 mM NH_4HCO_3 in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.16 min, 99.2%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0

mL/min. **LCMS**: 447.1 (M+H), Rt 2.48 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.35 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.34-8.33 (m, 1H), 7.64-7.63 (m, 1H), 7.48 (s, 1H), 7.25 (s, 1H), 6.24 (tt, 1H), 5.54 (q, 1H), 4.60 (dt, 2H), 4.19 (s, 3H), 1.77 (d, 3H).

Example 166. Synthesis of 212



Synthesis of 2-(methylamino)isonicotinonitrile (A-328)

[000767] To a stirred solution of 2-chloropyridine-4-carbonitrile (800 mg, 5.77 mmol) in NMP (8.0 mL) at room temperature was added methylamine (2.0 M in THF, 3.4 mL, 6.8 mmol). The reaction mixture was irradiated in microwave at 120 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The mixture was extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 12% EtOAc/PE to afford compound **A-328** (720 mg, 5.4 mmol, 93% yield). **LCMS**: 134.2 (M+H), Rt 1.71 min; Column: ZORBAX Extend C-18 (50 x 4.6 mm), 5 µm; Mobile Phase: A: 10 mM Ammonium acetate in water, B: ACN; Flow Rate:1.2 mL/min

Synthesis of (Z)-N'-hydroxy-2-(methylamino)isonicotinimidamide (A-329)

[000768] To a stirred solution of compound **A-328** (700 mg, 5.26 mmol) in ethanol (10.0 mL) was added DIPEA (2.75 mL, 15.77 mmol) and hydroxylamine hydrochloride (547 mg, 7.89 mmol) at room temperature. The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with

brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-329** (860 mg). It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(methylamino)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-330)

[000769] To a stirred solution of compound **A-329** (860 g, 5.2 mmol) in 1,4-dioxane (20 mL) was added DCC (1.64 g, 7.94 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.37 g, 7.22 mmol) at room temperature. The reaction heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (40 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 40% ethyl acetate/PE to afford compound **A-330** (600 mg, 1.87 mmol, 36% yield).**LCMS**: 320.2 (M+H), Rt 1.24 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-4-(5-(1-aminoethyl)-1,2,4-oxadiazol-3-yl)-N-methylpyridin-2-amine (A-331)


[000770] To a stirred solution of compound **A-330** (600 mg, 1.87 mmol) in DCM (15 mL) was added TFA (1.5 mL) at 0 °C under nitrogen. The reaction mixture was slowly warmed to room temperature and stirred for 3 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-331** (270 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(2-(methylamino)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (212)

[000771] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (239 mg, 1.23 mmol) in THF (10 mL) was added compound **A-331** (270 mg, 1.23 mmol) followed by TEA (0.51 mL, 3.69 mmol) and T3P (50% in EtOAc, 2.2 mL, 3.69 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **212** (178 mg, 0.44 mmol, 36% yield) as a solid. Prep. HPLC method: Rt 11.39; Column: Sunfire (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile;

Flow Rate: 15.0 mL/min.**HPLC**: Rt 3.16 min, 99.8%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 396.3 (M+H), Rt 1.45 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.4 min, SFC column: YMC Amylose-SA; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.10-8.09 (m, 1H), 7.26 (s, 1H), 7.17-7.16 (m, 1H), 7.14-7.12 (m, 1H), 5.52 (q, 1H), 4.19 (s, 3H), 2.92 (s, 3H), 1.76 (d, 3H).

Example 167. Synthesis of 213



Synthesis of 2-(3-methoxyazetidin-1-yl)isonicotinonitrile (A-332)

[000772] To a stirred solution of 2-chloropyridine-4-carbonitrile (500 mg, 3.61 mmol) in NMP (5.0 mL) was added DIPEA (1.89 mL, 10.83 mmol) and 3-methoxyazetidine hydrochloride (0.54 g, 4.33 mmol) at room temperature under nitrogen and the reaction mixture was irradiated in microwave at 120 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (30 mL). The mixture was extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% EtOAc/PE to afford compound **A-332** (650 mg, 3.43 mmol, 95% yield). **LCMS**: 190.2 (M+H), Rt 1.29 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min

Synthesis of (Z)-N'-hydroxy-2-(3-methoxyazetidin-1-yl)isonicotinimidamide (A-333)

[000773] To a stirred solution of compound **A-332** (650 mg, 3.43 mmol) in ethanol (10.0 mL) was added DIPEA (1.73 mL, 10.46 mmol) and hydroxylamine hydrochloride (363 mg, 5.23 mmol) at room temperature. The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution

(10 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-333** (760 mg). It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(3-methoxyazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-334)

[000774] To a stirred solution of compound **A-333** (760 mg, 3.43 mmol) in 1,4-dioxane (20 mL) was added DCC (866 mg, 4.21 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (723 mg, 3.82 mmol) at room temperature. The reaction heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (40 mL) and extracted with ethyl acetate (2 x 40 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 60% ethyl acetate/PE to afford compound **A-334** (671 mg, 1.78 mmol, 51% yield). **LCMS**: 376.2 (M+H), Rt 1.49 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 μm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-1-(3-(2-(3-methoxyazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-335)


[000775] To a stirred solution of compound **A-334** (671 mg, 1.78 mmol) in DCM (20 mL) was added TFA (1.5 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 6 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-335** (290 mg). The compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(3-methoxyazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (213)

[000776] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (183 mg, 0.94 mmol) in THF (10.0 mL) was added compound **A-335** (260 mg, 0.94 mmol) followed by TEA (0.39 mL, 2.83 mmol) and T3P (50% in EtOAc, 1.69 mL, 2.83 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **213** (186 mg, 0.41 mmol, 43% yield) as a solid. Prep. HPLC method: Rt 9.32;

Column: Sunfire C18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min.**HPLC**: Rt 3.38 min, 99.5%; Column: XBridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 452.2 (M+H), Rt 1.67 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.45 min, SFC column: YMC Amylose-SA; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.18-8.16 (m, 1H), 7.26-7.24 (m, 2H), 7.02 (s, 1H), 5.52 (q, 1H), 4.42-4.38 (m, 1H), 4.30-4.26 (m, 2H), 4.18 (s, 3H), 3.92-3.89 (m, 2H), 3.36 (s, 3H), 1.76 (d, 3H).

Example 168. Synthesis of 214



Synthesis of 2-(cyclopropylmethoxy)isonicotinonitrile (A-346)

[000777] To a solution of cyclopropylmethanol (0.58 g, 8.0 mmol) in THF (15.0 mL) was added NaH (60% in mineral oil, 290 mg, 7.26 mmol) and stirred at room temperature for 15 min. 2-chloropyridine-4-carbonitrile (1.0 g, 7.22 mmol) was added to the reaction mixture and stirred for 3 h at room temperature. The reaction mixture was cooled to 10 °C and treated with ice water (30 mL). The mixture was extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford A-346 (900 mg, 5.17 mmol, 71% yield). **LCMS**: 175.2 (M+H), Rt 2.28 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of 2-(cyclopropylmethoxy)-N'-hydroxyisonicotinimidamide (A-347)

[000778] To a stirred solution of compound **A-346** (900 mg, 5.17 mmol) in ethanol (20 mL) was added hydroxylamine hydrochloride (0.54 g, 7.74 mmol) and DIPEA (2.7 mL, 15.5 mmol). The reaction mixture was heated at 80 °C for 3 h. The reaction mixture was cooled to room temperature and treated with water (20 mL). The mixture was treated with 10% sodium carbonate solution (10 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-347** (1.05 g). It was used for next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(cyclopropylmethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-348)

[000779] To a solution of compound **A-347** (1.05 g, 5.1 mmol) in 1,4-dioxane (30 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.22 g, 6.47 mmol) and DCC (1.47 g, 7.11 mmol). The reaction mixture was heated to 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 14% EtOAc/PE to afford compound **A-348** (1.47 g, 4.1 mmol, 80% yield) as a solid.**LCMS**: 361.2 (M+H), Rt 2.71 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-1-(3-(2-(cyclopropylmethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-349)


[000780] To a stirred solution of compound **A-348** (1.2 g, 3.33 mmol) in DCM (12 mL) was added TFA (3.9 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and THF (20 mL) was added. The mixture was treated with silicon carbonate (3.0 g), stirred for 30 min. and filtered over celite. The filtrate was concentrated to afford compound **A-349** (720 mg). It was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(cyclopropylmethoxy)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (214)

[000781] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (6.0 mL) was added A-349 (201 mg, 0.77 mmol) followed by Et_3N (0.32 mL, 2.32 mmol) and T3P (1.38 mL, 2.32 mmol). The reaction mixture stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated

sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **214** (111 mg, 0.25 mmol, 32% yield) as a solid. Prep. HPLC method: Rt 11.55; Column: X-Select C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.45 min, 99.1%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 437.2 (M+H), Rt 2.67 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.44 min, SFC column: Lux C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.27 (d, 1H), 7.53 (dd, 1H), 7.39 (d, 1H), 7.26 (s, 1H), 5.53 (q, 1H), 4.20-4.18 (m, 5H), 1.77 (d, 3H), 1.35-1.29 (m, 1H), 0.65-0.60 (m, 2H), 0.40-0.36 (m, 2H).

Example 169. Synthesis of 215




A-284 215

Synthesis of (S)-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-phenyl-1H-pyrazole-5-carboxamide (215)

[000782] To a stirred solution of compound **A-284** (120 mg, 0.55 mmol) in THF (8.0 mL) was added 2-phenylpyrazole-3-carboxylic acid (106 mg, 0.57 mmol) followed by Et_3N (0.23 mL, 1.65 mmol) and T3P (50% in EtOAc, 0.98 mL, 1.65 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **215** (56 mg, 0.14 mmol, 25% yield) as a solid. Prep. HPLC method: Rt 5.85; Column: YMC C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.80 min, 97.6%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 389.3 (M+H), Rt 1.63 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1%

HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.76 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.62 (d, 1H), 7.96 (s, 1H), 7.88 (d, 1H), 7.77 (d, 1H), 7.48-7.38 (m, 5H), 6.97 (d, 1H), 7.26 (dd, 1H), 2.65 (s, 3H), 2.23-2.03 (m, 2H), 1.09 (t, 3H).

Example 169. Synthesis of 216




A-284 216

Synthesis of (S)-1-cyclopentyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1H-pyrazole-5-carboxamide (216)

[000783] To a stirred solution of compound **A-284** (120 mg, 0.55 mmol) in THF (8.0 mL) was added 2-cyclopentylpyrazole-3-carboxylic acid (106 mg, 0.59 mmol) followed by Et_3N (0.23 mL, 1.65 mmol)and T3P (50% in EtOAc, 0.98 mL, 1.65 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (20 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **216** (72 mg, 0.18 mmol, 34% yield) as a solid. Prep. HPLC method: Rt 8.05; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.30 min, 99.7%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 381.3 (M+H), Rt 2.00 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**Chiral method:** Rt 1.6 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.60 (d, 1H), 7.96 (s, 1H), 7.86 (d, 1H), 7.53 (d, 1H), 6.86 (d, 1H), 5.58-5.50 (m, 1H), 5.35 (dd, 1H), 2.64 (s, 3H), 2.28-1.88 (m, 8H), 1.71-1.65 (m, 2H), 1.13 (t, 3H).

Example 170. Synthesis of 217




A-284 217

Synthesis of (S)-3-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-(tetrahydro-2H-pyran-4-yl)-1H-pyrazole-5-carboxamide (217)

[000784] To a stirred solution of compound **A-284** (123 mg, 0.57 mmol) in THF (6.0 mL) was added 5-methyl-2-tetrahydropyran-4-yl-pyrazole-3-carboxylic acid (120 mg, 0.57 mmol) followed by Et_3N (0.24 mL, 1.7 mmol) and T3P (50% in EtOAc, 1.01 mL, 1.7 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **217** (68 mg, 0.16 mmol, 29% yield) as a solid. Prep. HPLC method: Rt 12.65; Column: X-Select (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% HCOOH in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.68 min, 98.3%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 411.2 (M+H), Rt 1.47 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**; Chiral method:** Rt 1.53 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.61 (d, 1H), 7.95 (s, 1H), 7.86 (d, 1H), 6.66 (s, 1H), 5.34 (dd, 1H), 5.23-5.19 (m, 1H), 4.05-4.00 (m, 2H), 3.55-3.46 (m, 2H), 2.64 (s, 3H), 2.29 (s, 3H), 2.26-2.09 (m, 4H), 1.90-1.86 (m, 2H), 1.13 (t, 3H).

Example 171. Synthesis of 218




A-284 218

Synthesis of (S)-3-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-phenyl-1H-pyrazole-5-carboxamide (218)

[000785] To a stirred solution of compound **A-284** (120 mg, 0.55 mmol) in THF (5.0 mL) was added 5-methyl-2-phenyl-pyrazole-3-carboxylic acid (122 mg, 0.60 mmol) followed by Et_3N (0.23 mL, 1.65 mmol) and T3P (50% in EtOAc, 1.5 mL, 2.52 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **218** (20 mg, 0.05 mmol, 8% yield) as a solid. Prep. HPLC method: Rt 6.95; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 10 mM NH_4OAc in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.95 min, 97.1%Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 403.1 (M+H), Rt 1.54 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.62 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 9.44 (d, 1H), 8.68 (d, 1H), 7.82 (s, 1H), 7.73 (d, 1H), 7.38-7.30 (m, 5H), 6.78 (s, 1H), 5.19-5.13 (m, 1H), 2.58 (s, 3H), 2.28 (s, 3H), 2.07-2.00 (m, 2H), 0.99 (t, 3H).

Example 172. Synthesis of 219




A-284 219

Synthesis of (S)-1-benzyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1H-pyrazole-5-carboxamide (219)

[000786] To a stirred solution of compound **A-284** (120 mg, 0.55 mmol) in THF (5.0 mL) was added 2-benzylpyrazole-3-carboxylic acid (122 mg, 0.60 mmol) followed by Et_3N (0.23 mL, 1.65 mmol) and T3P (50% in EtOAc, 1.5 mL, 2.52 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **219** (95 mg, 0.23 mmol, 42% yield) as a solid. Prep. HPLC method: Rt 7.46; Column: YMC C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.29 min, 98.3%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 403.1 (M+H), Rt 1.66 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.81 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.60 (d, 1H), 7.93 (s, 1H), 7.84 (d, 1H), 7.59 (d, 1H), 7.26-7.15 (m, 5H), 6.96 (d, 1H), 5.74 (d, 2H), 5.32 (q, 1H), 2.64 (s, 3H), 2.24-2.17 (m, 1H), 2.11-2.03 (m, 1H), 1.06 (t, 3H).

Example 173. Synthesis of 220



Synthesis of 6-methylpyrimidine-4-carbonitrile (A-351)

[000787] To a stirred solution of 4-chloro-6-methyl-pyrimidine (**A-350**, 2.0 g, 15.56 mmol) in DMSO (25.0 mL) and water (7.0 mL) was added DABCO (0.87 g, 7.78 mmol) and NaCN (1.68 g, 34.23 mol) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 7 h. The reaction mixture was treated with water (80 mL) and extracted with EtOAc (2 x 80 mL). The organic layer was washed with saturated sodium bicarbonate solution (60 mL), washed with brine (60 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 10% ethyl acetate/PE to afford A-351 (850 mg, 7.13 mmol, 45% yield). **LCMS**: 120.2 (M+H), Rt 1.01 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of N'-hydroxy-6-methylpyrimidine-4-carboximidamide (A-352)

[000788] To a stirred solution of compound **A-351** (830 mg, 6.97 mmol) in ethanol (10.0 mL) was added hydroxylamine hydrochloride (726 mg, 10.45 mmol) and DIPEA (3.45 mL, 20.9 mmol). The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford A-352 (850 mg). The compound was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(6-methylpyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-353)

[000789] To a stirred solution of compound **A-352** (530 mg, 3.47 mmol) in 1, 4-dioxane (20 mL) was added DCC (859 mg, 4.16 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (984 mg, 5.21 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 29% EtOAc/PE to afford A-353 (520 mg, 1.7 mmol, 49% yield). **LCMS**: 306.1 (M+H), Rt 2.04 min; Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 10 mM NH_4HCO_3 in water, B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-1-(3-(6-methylpyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-354)


[000790] To a stirred solution of compound **A-353** (520 mg, 1.71 mmol) in DCM (10.0 mL) was added TFA (1.25 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% $NaHCO_3$ solution (10 mL) and extracted with DCM (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-354** (225 mg). The crude compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-N-(1-(3-(6-methylpyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (220)

[000791] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (213 mg, 1.1 mmol) and compound **A-354** (280 mg, 1.1 mmol) in THF (10.0 mL) was added Et_3N (0.46 mL, 3.3 mmol) followed by T3P (50% in EtOAc, 0.98 mL, 3.3 mmol) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by preparative HPLC to afford **220** (220 mg, 0.57 mmol, 52% yield) as a solid. Prep. HPLC method: Rt 11.40; Column: X-Bridge (150 x

19 mm), 5.0 µm; Mobile phase: 10 mM NH_4OAc in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.75 min, 99.2%Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min; **LCMS**: 382.2 (M+H), Rt 1.89 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.21 min, SFC column: YMC Cellulose-C; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 9.17 (s, 1H), 8.10 (s, 1H), 7.26 (s, 1H), 5.55 (q, 1H), 4.18 (s, 3H), 2.66 (s, 3H), 1.77 (d, 3H).

Example 174. Synthesis of 221




A-284 221

Synthesis of (S)-1-(difluoromethyl)-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (221)

[000792] To a stirred solution of 2-(difluoromethyl)-5-trifluoromethyl-pyrazole-3-carboxylic acid (100 mg, 0.57 mmol) and compound **A-284** (123 mg, 0.57 mmol) in THF (6.0 mL) was added Et_3N (0.24 mL, 1.7 mmol) and T3P (50% in EtOAc, 1.01 mL, 1.7 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **221** (125 mg, 0.33 mmol, 58% yield) as a solid. Prep. HPLC method: Rt 12.23; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 10 mM NH_4OAc in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.84 min, 99.6%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 377.2 (M+H), Rt 1.67 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.4 min, SFC column: YMC

Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.60 (d, 1H), 8.13 (t, 1H), 7.95 (s, 1H), 7.86 (d, 1H), 6.94 (s, 1H), 5.38 (dd, 1H), 2.64 (s, 3H), 2.36 (s, 3H), 2.28-2.20 (m, 1H), 2.16-2.05 (m, 1H), 1.12 (t, 3H).

Example 175. Synthesis of 222



Synthesis of 2-(3,3-difluoroazetidin-1-yl)isonicotinonitrile (A-355)

[000793] To a stirred solution of 3,3-difluoroazetidine (**A-272**, 1.4 g, 10.86 mmol) in NMP (8.0 mL) was added DIPEA (1.9 mL, 10.86 mmol) and 2-chloropyridine-4-carbonitrile (750 mg, 5.4 mmol). The reaction mixture was heated in a microwave at 120 °C for 2 h. The reaction mixture was cooled to room temperature and treated with water (50 mL). The mixture was extracted with ethyl acetate (2 x 40 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 20% EtOAc/PE to afford A-355 (750 mg, 3.84 mmol, 71% yield).**LCMS**: 196.1 (M+H), Rt 1.86 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of 2-(3,3-difluoroazetidin-1-yl)-N'-hydroxyisonicotinimidamide (A-356)

[000794] To a stirred solution of compound **A-355** (750 mg, 3.84 mmol) in ethanol (10.0 mL) was added hydroxylamine hydrochloride (0.4 g, 5.81 mmol) and DIPEA (2.0 mL, 11.52 mmol). The reaction mixture was heated at 80 °C for 4 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (20 mL) followed by saturated sodium bicarbonate solution (20 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over

anhydrous Na_2SO_4 and concentrated to afford compound **A-356** (840 mg). The compound was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(3,3-difluoroazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-357)

[000795] To a stirred solution of compound **A-356** (840 mg, 3.68 mmol) in 1,4-dioxane (30mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (0.7 g, 3.68 mmol) followed by DCC (0.83 g, 4.05 mmol). The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude compound was purified by column chromatography on silica gel with 40% EtOAc/PE to afford compound **A-357** (1.25 g, 3.3 mmol, 90% yield) as a solid.**LCMS**: 382.3 (M+H), Rt 2.29 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of (S)-1-(3-(2-(3,3-difluoroazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-358)


[000796] To a stirred solution of compound **A-357** (680 mg, 1.78 mmol) in DCM (8.0 mL) was added TFA (2.0 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with 10% $NaHCO_3$ solution (10.0 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-358** (465 mg). The crude compound was used for the next step without further purification.

Synthesis of (S)-N-(1-(3-(2-(3,3-difluoroazetidin-1-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (222)

[000797] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in THF (8.0 mL) was added compound **A-358** (260 mg, 0.92 mmol) followed by T3P (50% in EtOAc, 1.38 mL, 2.32 mmol) and Et_3N (0.32 mL, 2.32 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20

mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **222** (23 mg, 0.05 mmol, 6% yield) as a solid. Prep. HPLC method: Rt 10.23; Column: Atlantis C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.87 min, 99.9%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 458.2 (M+H), Rt 2.33 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.06 min, SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm.**^{1}H NMR (400 MHz, CD_3OD):** δ 8.28 (dd, 1H), 7.40 (dd, 1H), 7.25 (s, 1H), 7.18-7.17 (m, 1H), 5.53 (q, 1H), 4.45 (t, 4H), 4.19 (s, 3H), 1.77 (d, 3H).

Example 176. Synthesis of 223




A-293 223

Synthesis of (S)-3-(4-fluorophenyl)-N-(1-(3-(2-isopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-1H-pyrazole-5-carboxamide (223)

[000798] To a stirred solution of compound **A-293** (120 mg, 0.48 mmol) in THF (6.0 mL) was added 5-(4-fluorophenyl)-2-methyl-pyrazole-3-carboxylic acid (106 mg, 0.48 mmol) followed by Et_3N (0.2 mL, 1.45 mmol) and T3P (50% in EtOAc, 0.86 mL, 1.45 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **223** (80 mg, 0.17 mmol, 36% yield) as a solid. Prep. HPLC method: Rt 12.41; Column: YMC C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.38 min, 98.5%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 451.1 (M+H), Rt 2.44 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm, Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in

ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.94 min, SFC column: YMC Cellulose-SC; mobile phase: 70:30 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.28 (d, 1H), 7.85-7.82 (m, 2H), 7.52 (d, 1H), 7.33 (s, 1H), 7.23 (s, 1H), 7.17 (t, 2H), 5.55 (q, 1H), 5.36-5.29 (m, 1H), 4.16 (s, 3H), 1.79 (d, 3H), 1.37 (d, 6H).

Example 177. Synthesis of 224




A-293 224

Synthesis of (S)-1-(2-(dimethylamino)-2-oxoethyl)-N-(1-(3-(2-isopropoxypyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (224)

[000799] To a stirred solution of 2-[2-(dimethylamino)-2-oxo-ethyl]-5-(trifluoromethyl)pyrazole-3-carboxylic acid (128 mg, 0.48 mmol) and compound **A-293** (120 mg, 0.48 mmol) in THF (6.0 mL) was added Et_3N (0.2 mL, 1.45mmol) and T3P (50% in EtOAc, 0.86 mL, 1.45 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **224** (143 mg, 0.28 mmol, 59% yield) as a solid. Prep. HPLC method: Rt 15.01; Column: X-Select (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min.**HPLC**: Rt 4.82 min, 99.3%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µmMobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 496.1 (M+H), Rt 2.23 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.52 min, SFC column: YMC Cellulose-SB; mobile phase: 70:30 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.27 (d, 1H), 7.50 (d, 1H), 7.35-7.33 (m, 2H), 5.61-5.46 (m, 3H), 5.35-5.31 (m, 1H), 3.12 (s, 3H), 2.94 (s, 3H), 1.75 (d, 3H), 1.37 (d, 6H).

Example 178. Synthesis of 295 & 225



Synthesis of 2-(trifluoromethyl)isonicotinamide (A-360)

[000800] To a stirred solution of 2-(trifluoromethyl)pyridine-4-carboxylic acid (**A-359**, 2.0 g, 10.47 mmol) in DCM (20.0 mL) was added oxalyl chloride (1.39 g, 10.99 mmol) and catalytic amount of DMF (0.05 mL) at 0 °C. The reaction mixture was warmed to room temperature and stirred for 2 h. The reaction mixture was concentrated under reduced pressure and aqueous NH_4OH (2.0 mL) was added at 0 °C dropwise followed by MeCN (10.0 mL). The reaction mixture was stirred at room temperature for 30 min and diluted with ethylacetate (100 mL). The organic layer was washed with water (2 x 50 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-360** (1.3 g). The compound was used for the next step without further purification.

Synthesis of 2-(trifluoromethyl)pyridine-4-carbonitrile (A-361)

[000801] $POCl_3$ (3.04 mL, 32.61 mmol) was added dropwise to the compound **A-360** (1.3 g, 6.84 mmol) at 0 °C. The reaction mixture heated at 100 °C for 3 h. The reaction mixture was cooled to room temperature and treated with 50% NaOH solution (10 mL). The reaction mixture was diluted with water (50 mL) and extracted with ethylacetate (2 x 70 mL). The organic layer was washed with brine (40 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 8% ethyl acetate/PE to afford compound **A-361** (520 mg, 3.0 mmol, 44 % yield). **LCMS**: 173.1 (M+H), Rt 1.84 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min

Synthesis of N'-hydroxy-2-(trifluoromethyl)pyridine-4-carboxamidine (A-362)

[000802] To a stirred solution of compound **A-361** (520 mg, 3.02 mmol) in ethanol (10.0 mL) was added hydroxylamine hydrochloride (314 mg, 4.53 mmol) and DIPEA (1.58 mL, 9.05 mmol) at room temperature under nitrogen atmosphere. The reaction mixture was heated at 80 °C for 2 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The crude was treated with water (15 mL) followed by saturated sodium bicarbonate solution (10 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-362** (580 mg). It was used for the next step without further purification.

Synthesis of tert-butyl (S)-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (295)

[000803] To a stirred solution of compound **A-362** (580 mg, 2.83 mmol) in 1,4-dioxane (20.0 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (534 mg, 2.83 mmol) and DCC (640 mg, 3.11 mmol). The reaction mixture was stirred at 100 °C for 16 h. The reaction mixture was cooled to room temperature and concentrated under reduced pressure. The mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica gel with 12% ethyl acetate/PE to afford compound **295** (840 mg, 2.34 mmol, 82% yield). **LCMS**: 359.2 (M+H), Rt 2.42 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min


Synthesis of (1S)-1-[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine (A-364)

[000804] To a stirred solution of compound **295** (400 mg, 1.12 mmol) in DCM (8.0 mL) was added TFA (1.5 mL) at 0 °C. The reaction mixture was slowly warmed to room temperature and stirred for 2 h. The mixture was concentrated under reduced pressure and treated with ice water (20 mL). The mixture was treated with saturated $NaHCO_3$ solution (10 mL) and extracted with EtOAc (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated to afford compound **A-364** (260 mg). The compound was used for the next step without further purification.

Synthesis of (S)-1-methyl-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (225)

[000805] To a stirred solution of compound **A-364** (260 mg, 1.01 mmol) in THF (10.0 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (195 mg, 1.01 mmol) followed by Et_3N (0.42 mL, 3.02 mmol) and T3P (50% in EtOAc, 1.8 mL, 3.02 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **225** (130 mg, 0.3 mmol, 29% yield) as a solid. Prep. HPLC method: Rt 7.15; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 5.10 min, 99.5%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 435.0 (M+H), Rt 2.56 min, Column: Atlantis dc-18 (50 x 4.6 mm), 5.0 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.73 min, SFC column: LUX C3; mobile phase: 85:15 (A: B), A = liquid CO_2, B = methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.92 (d, 1H), 8.38 (s, 1H), 8.28 (dd, 1H), 7.25 (s, 1H), 5.55 (q, 1H), 4.18 (s, 3H), 1.78 (d, 3H).

Example 179. Synthesis of 226




A-284 **226**

Synthesis of (S)-1-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-3-phenyl-1H-pyrazole-5-carboxamide (226)

[000806] To a stirred solution of compound **A-284** (120 mg, 0.55 mmol) in THF (5.0 mL) was added 2-methyl-5-phenyl-pyrazole-3-carboxylic acid (122 mg, 0.60 mmol) followed by Et_3N (0.23 mL, 1.65 mmol) and T3P (50% in EtOAc, 1.5 mL, 2.52 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with

water (40 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **226** (85 mg, 0.21 mmol, 38% yield) as a solid. Prep. HPLC method: Rt 7.65; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.53 min, 99.8%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 403.1 (M+H), Rt 1.75 min, Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.47 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.61 (d, 1H), 7.96 (s, 1H), 7.88-7.87 (m, 1H), 7.84-7.81 (m, 2H), 7.45-7.42 (m, 2H), 7.36-7.31 (m, 1H), 7.30 (s, 1H), 5.40 (dd, 1H), 4.17 (s, 3H), 2.64 (s, 3H), 2.31-2.12 (m, 2H), 1.15 (t, 3H).

Example 180. Synthesis of 227




A-303 **227**

Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1,3-dimethyl-1H-pyrazole-5-carboxamide (227)

[000807] To a solution of compound **A-303** (100 mg, 0.43 mmol) in THF (8.0 mL) was added 2,5-dimethylpyrazole-3-carboxylic acid (84 mg, 0.60 mmol) followed by Et_3N (0.18 mL, 1.3 mmol) and T3P (50% in EtOAc, 0.78 mL, 1.3 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 50 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **227** (75 mg, 0.21 mmol, 48% yield) as a solid. Prep. HPLC method: Rt 10.61; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 10 mM NH_4OAc in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.42 min, 99.8%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1%

TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 353.2 (M+H), Rt 1.66 min, Column: Zorbax XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.48 min, SFC column: YMC Amylose-C; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.53 (d, 1H), 7.85 (s, 1H), 7.76 (dd, 1H), 6.69 (s, 1H), 5.51 (q, 1H), 4.04 (s, 3H), 2.27 (s, 3H), 2.23-2.19 (m, 1H), 1.76 (d, 3H), 1.12-1.02 (m, 4H).

Example 181. Synthesis of 228




A-303 **228**

Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(4-fluorophenyl)-1-methyl-1H-pyrazole-5-carboxamide (228)

[000808] To a solution of compound **A-303** (100 mg, 0.43 mmol) in THF (8.0 mL) was added 5-(4-fluorophenyl)-2-methyl-pyrazole-3-carboxylic acid (105 mg, 0.48 mmol) followed by Et_3N (0.18 mL, 1.3 mmol) and T3P (50% in EtOAc, 0.78 mL, 1.3 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (30 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **228** (60 mg, 0.13 mmol, 31% yield) as a solid. Prep. HPLC method: Rt 11.15; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.73 min, 98.4%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 433.1 (M+H), Rt 1.82 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 2.56 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.53 (d, 1H), 7.85-7.81 (m, 3H), 7.77-7.75 (m,

1H), 7.23 (s, 1H), 7.18-7.14 (m, 2H), 5.55 (q, 1H), 4.16 (s, 3H), 2.23-2.16 (m, 1H), 1.79 (d, 3H), 1.11-1.04 (m, 4H).


Example 181. Synthesis of 229



Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-5-carboxamide (229)

[000809] To a stirred solution of compound **A-303** (120 mg, 0.52 mmol) in THF (6.0 mL) was added 5-(difluoromethyl)-2-methyl-pyrazole-3-carboxylic acid (91 mg, 0.52 mmol) followed by Et_3N (0.22 mL, 1.56 mmol) and T3P (50% in EtOAc, 0.93 mL, 1.56 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (30 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **229** (47 mg, 0.12 mmol, 23% yield) as colourless liquid. Prep. HPLC method: Rt 8.38; Column: X-Select (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.94 min, 99.4%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 389.2 (M+H), Rt 2.06 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.55 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm.**^{1}H NMR (400 MHz, CD_3OD):** δ 8.54 (dd, 1H), 8.86 (dd, 1H), 7.76 (dd, 1H), 7.16 (s, 1H), 6.78 (t, 1H), 5.53 (q, 1H), 4.15 (s, 3H), 2.23-2.18 (m, 1H), 1.77 (d, 3H), 1.12-1.05 (m, 4H).

Example 182. Synthesis of 230




A-303 230

Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-(2-(dimethylamino)-2-oxoethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (230)

[000810] To a stirred solution of compound **A-303** (100 mg, 0.43 mmol) in THF (8.0 mL) was added 2-[2-(dimethylamino)-2-oxo-ethyl]-5-(trifluoromethyl)pyrazole-3-carboxylic acid (126 mg, 0.48 mmol) followed by Et_3N (0.18 mL, 1.3 mmol) and T3P (50% in EtOAc, 0.78 mL, 1.3 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (25 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **230** (85 mg, 0.17 mmol, 40% yield) as a solid. Prep. HPLC method: Rt 8.51; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.11 min, 99.5%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 478.1 (M+H), Rt 2.04 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min**. Chiral method:** Rt 1.51 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.53 (d, 1H), 7.92 (s, 1H), 7.86 (d, 1H), 7.35 (s, 1H), 5.61-5.45 (m, 3H), 3.12 (s, 3H), 2.93 (s, 3H), 2.23-2.18 (m, 1H), 1.76 (d, 3H), 1.12-1.04 (m, 4H),

Example 183. Synthesis of 231



Synthesis of ethyl 3-methyl-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylate (A-367) and ethyl 5-methyl-1-(pyridin-3-yl)-1H-pyrazole-3-carboxylate (A-368)

[000811] To a stirred solution of ethyl 3-methyl-1H-pyrazole-5-carboxylate (2.0 g, 12.97 mmol) in DCM (25.0 mL) was added 3-pyridylboronic acid (3.18 g, 25.95 mmol) followed by pyridine (2.09 mL, 25.95 mmol), copper (II) acetate (3.53 g, 19.46 mmol) and molecular sieves. The reaction mixture was stirred for at room temperature for 36 h. The reaction mixture was filtered through sintered funnel and washed with DCM (50 mL). The organic layer was washed with water (2 x 30 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford A-367 (320 mg, 1.3 mmol, 10% yield) and A-368 (150 mg, 0.65 mmol, 5% yield). Prep. HPLC method: Rt 12.0 (**A-368**) and 13.1 (A-367); Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **LCMS**: 232.3 (M+H), Rt 1.68 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.66-8.63 (m, 2H), 7.97-7.94 (m, 1H), 7.59 (dd, 1H), 6.96 (s, 1H), 4.25 (q, 2H), 2.36 (s, 3H), 1.26 (t, 3H). **LCMS**: 232.1 (M+H), Rt 1.49 min; Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min; **^{1}H NMR (400 MHz, CD_3OD):** δ 8.81 (s, 1H), 8.69 (d, 1H), 8.09-8.07 (m, 1H), 7.66 (dd, 1H), 6.81 (s, 1H), 4.39 (q, 2H), 2.41 (s, 3H), 1.40 (t, 3H).


Synthesis of 3-methyl-1-(pyridin-3-yl)-1H-pyrazole-5-carboxylic acid (A-369)

[000812] To a stirred solution of compound **A-367** (50 mg, 0.22 mmol) in methanol (2 mL), THF (2 mL) and water (2 mL) was added lithium hydroxide monohydrate (18 mg, 0.43 mmol). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was concentrated under reduced pressure and treated with ice water (10 mL). The mixture was treated with 1N HCl (1.0 mL) and extracted with ethyl acetate (2 x 30 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated to afford compound **A-369** (38 mg). The crude compound was used for the next step without further purification.

Synthesis of (S)-3-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-(pyridin-3-yl)-1H-pyrazole-5-carboxamide (231)


[000813] To a stirred solution of compound **A-369** (38 mg, 0.19 mmol) in THF (2.0 mL) was added (1S)-1-[3-(2-methyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]propan-1-amine (41 mg, 0.19 mmol) followed by Et_3N (0.08 mL, 0.56 mmol) and T3P (50% in EtOAc, 0.33 mL, 0.56 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (10 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with saturated sodium bicarbonate solution (20 mL), washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **231** (7 mg, 0.02 mmol, 9% yield) as an solid. Prep. HPLC method: Rt 9.69; Column: X-Select C-18 (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.10 min, 98.5%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 404.1 (M+H), Rt 1.11 min, Column: X-Bridge C8 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% TFA in water:ACN (95:5), B: 0.1% TFA in ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 1.53 min, SFC column: Chiralcel OD-H; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.65 (d, 1H), 8.61 (d, 1H), 8.53 (dd, 1H), 7.96-7.92 (m, 2H), 7.88-7.86 (m, 1H), 7.51-7.47 (m, 1H), 6.89 (s, 1H), 5.27 (dd, 1H), 2.65 (s, 3H), 2.39 (s, 3H), 2.25-2.04 (m, 2H), 1.09 (t, 3H).

Example 184. Synthesis of 232



[000814] To a mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) in DCM (10 mL) was added HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1R)-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (120.84 mg, 0.59 mmol), the mixture was stirred at 25 °C for 16 hours. The reaction was quenched with H_2O (10 mL), then extracted with DCM (20 mL x 3). The combined organic phase was washed with brine (30 mL), dried over Na_2SO_4, filtered and concentrated to give the crude product. The crude product was purified from Prep-HPLC ((Boston Prime C18 150×30 mm, 5 μm) A = H_2O (0.05 % NH_4OH) and B = CH_3CN; 17-47 % B over 8 minutes) to give the product as an oil. Analytical SFC (Column: Chiralpak AS-3 150 mm × 4.6 mm I.D., 3 μm Mobile phase: A: CO_2 B:ethanol (0.05% DEA) Gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5min, hold 5% of B for 1.5 min Flow rate: 2.5mL/min Column temp.: 35 °C ABPR: 1500psi) showed 2 peaks at 2.39 min (main peak) and 2.55 min. Then the product was purified by SFC (DAICEL Chiralpak AS-H 250 mm x 30 mm, 5 μm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38°C; 65 mL/min; 20% B; 8.60 min run; 50 injections, Rt of peak 1 = 5.57 min, Rt of Peak 2 = 6.60 min) to give the product of 1-isopropyl-3-methyl-N-[(1R)-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-4-carboxamide (66.75 mg,0.19 mmol, 32 % yield). (Rt = 2.39 min in analytical SFC) as a an oil. **^{1}H NMR** (400MHz, CD_3CN) 400MHz δ = 8.00 (s, 1H), 7.89 (s, 1H), 7.86 (d, 1H), 7.48 - 7.38 (m, 2H), 7.06 (br d, 1H), 5.50 - 5.40 (m, 1H), 4.46 (quin, 1H), 2.44 (s, 3H), 2.39 (s, 3H), 1.69 (d, 3H), 1.48 (d, 6H).**LCMS** R_t = 1.13 min in in 2.0 min chromatography, 10-80AB, MS ESI calcd. for $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.2, found 354.0.

Example 185. Synthesis of 233




A-271 233


[000815] A mixture of 1-isopropyl-3-methyl-pyrazole-4-carboxylic acid (100 mg, 0.59 mmol) , HOBt (160.69 mg, 1.19 mmol), EDCI (227.96 mg, 1.19 mmol), DIPEA (0.33 mL, 2.38 mmol) and (1S)-1-[3-(m-tolyl)-1,2,4-oxadiazol-5-yl]ethanamine (120.84 mg, 0.59 mmol) in DCM (10 mL), the mixture was stirred at 25 °C for 16 hours. The mixture was concentrated to give the residue. The residue was diluted with H_2O (20 mL), and the mixture was extracted with EtOAc (20 mL x 2). The combined organic phase was washed with water (20 mL) and brine (20 mL), dried over Na_2SO_4, filtered and concentrated to give the crude

product. The crude product was purified by prep-HPLC (Boston Prime (150 mm x 30 mm, 5 µm) A = H_2O (0.05% NH_4OH) and B = CH_3CN; 40-60 %B over 9 minutes) to give the crude product. Analytical SFC (Column: Chiralpak AS-3 150 mm × 4.6 mm I.D., 3 µm, Mobile phase: A: CO_2 B:e thanol (0.05% DEA), Gradient: from 5% to 40% of B in 5 min and from 40% to 5% of B in 0.5min, hold 5% of B for 1.5 min, Flow rate: 2.5mL/min, Column temp.: 35°C, ABPR: 1500psi.) showed two peaks at 2.37 min and 2.55 min (main peak). The product was purified by SFC (DAICEL CHIRALPAK AS-H (250 mm x 30 mm, 5 µm); A = CO_2 and B = EtOH (0.1% NH_3H_2O); 38°C; 60 mL/min; 20% B; 8 min run; 10 injections, Rt of peak 1 = 4.8 min, Rt of Peak 2 = 6 min) to give the product (50.77 mg, 0.14 mmol, 24% yield) (Rt = 2.55 min in analytical SFC) as a solid. **^{1}H NMR** (400MHz, CD_3CN) δ_H = 7.96 (s, 1H), 7.86 (s, 1H), 7.83 (br d, 1H), 7.44 - 7.36 (m, 2H), 7.04 (br d, 1H), 5.46 - 5.37 (m, 1H), 4.47 - 4.36 (m, 1H), 2.41 (s, 3H), 2.36 (s, 3H), 1.66 (d, 3H), 1.44 (d, 6H). **LCMS** Rt = 1.20 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{19}H_{24}N_5O_2$ $[M+H]^+$ 354.2, found 354.1.

Example 186. Synthesis of 296 & 234



A-362 296



A-400 234

Synthesis of tert-butyl (R)-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (296)

[000816] To a stirred solution of **A-362** (1.5 g, 7.31 mmol) in 1,4-dioxane (20 mL) was added *N*-Boc -D -alanine (1.52 g, 8.04 mmol), followed by DCC (1.66 g, 8.04 mmol). The reaction was heated at 100 °C for 16 h. The mixture was concentrated to dryness and was diluted with ethyl acetate (100 mL). The organic layer was washed with water (2 x 30 mL) and then with saturated brine solution (1 x 50 mL). The organic layer was dried over magnesium sulphate and concentrated. The crude compound was purified by flash column chromatography eluting 50% EtOAc in hexane. The desired fractions were concentrated to

dryness under reduced pressure to afford 1.5 g of desired compound that was used as it is in next step. 300 mg of the compound was purified by prep HPLC to afford **296**, tert-butyl *N*-[(1R)-1-[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate as a solid (153 mg). **HPLC**: Rt 7.71 min, 99.80%; Column: HPLC-X-Bridge C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% NH_3 in water B: ACN; Flow Rate: 1.2 mL/min. **LCMS**: 359.15 (M+H), Rt 2.03 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **Chiral method:** Rt 3.90 min; SFC column: DIACEL CHIRALPAK-IG (150 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MTBE: IPA(60:40) Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10%B at 10min,hold 10%B till 12Min., Wavelength: 270 nm; Flow: 3 mL/min. **^{1}H NMR (400 MHz, DMSO):** δ_H = 9.02 (d, 1H), 8.3-8.24 (m, 2H), 7.84 (d, 1H), 5.05-5.00 (m, 1H), 1.53 (d, 3H), 1.40 (s, 9H).

Synthesis of (1R)-1-[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (A-400)


[000817] To a stirred solution of compound **296** (1.2 g, 3.35 mmol) in 1,4-dioxane (3 mL) was added 4 M HCl in dioxane (15 mL) at 0 °C. The reaction was slowly brought to room temperature and stirred for 6 h. After completion, the reaction mass was concentrated to dryness to afford compound **A-400** (0.80 g, 2.69 mmol, 80% yield) as a solid. The compound **A-400** was used for the next step without further purification.

Synthesis of 2-methyl-5-(trifluoromethyl)-*N*-[(1R)-1-[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (234)

[000818] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (150 mg, 0.77 mmol) in DCM (10 mL) was added compound **A-400** (273.24 mg, 0.93 mmol). To this reaction was added HATU (323.21 mg, 0.85 mmol), and DIPEA (0.27 mL, 1.55 mmol) and the mixture was stirred at room temperature for 6 h. The reaction was treated with water (10 mL) and extracted with DCM (10 mL). The organic layer was washed with brine solution and dried over anhydrous Na_2SO_4 and concentrated. The crude compound was purified by flash column chromatography. eluting 50% EtOAc in hexane. The desired fractions were concentrated to dryness under reduced pressure to afford **234** as a solid (53 mg, 0.12 mmol, 15% yield). **HPLC**: Rt 9.16 min, 99.8%; Column: X-select CSH C18 (4.6*150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS**: 435 (M+H), Rt 2.15 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm; **Chiral method:** Rt 3.61 min; SFC column: DIACEL CHIRALPAK-IG (150 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MTBE:IPA(60:40) Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10%B at 10min,hold 10%B till 12Min., Wavelength: 270

nm; Flow: 3 mL/min. **^{1}H NMR (400 MHz, DMSO):** δ_H = 9.49 (d, 1H), 9.01 (d, 1H), 8.32-8.24 (m, 2H), 7.45 (s, 1H), 5.52-5.48 (m, 1H), 4.12 (s, 3H), 1.69 (d, 3H).


Example 187. Synthesis of 235



A-364 235

[000819] To a stirred solution 3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxylic acid (90.73 mg, 0.52 mmol) in DCM (5 mL) was added DIPEA (0.27 mL, 1.55 mmol) and HATU (293.83 mg, 0.77 mmol) at 0 °C. To the resulting reaction mixture was added **A-364** (166.98 mg, 0.57 mmol) at 0 °C and stirred at RT for 12 h. The reaction was diluted with water and extracted with DCM (3 x 10 mL). The combined organic layer was dried over sodium sulphate and concentrated to obtain the crude compound. The crude compound was purified by prep HPLC to afford **235** (82 mg, 0.19 mmol, 38% yield) as a solid. **HPLC**: Rt 8.13 min, 99.8%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS**: 417.20 (M+H), Rt 1.85 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO):** δ_H = 9.07 (d, 1H), 9.01 (d, 1H), 8.38 (s, 1H), 8.29-8.26 (m, 2H),7.26 (t, 1H), 5.54-5.45 (m, 1H), 3.94 (s, 3H), 1.65 (d, 3H). **Chiral HPLC:** Rt 3.27 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:20-40% B in 5 min, hold 40 % B till 9 min, 40-20% B in 10 min, hold 20% B till 12 min. Wavelength: 270 nm, Flow: 3 mL/min.

Example 188. Synthesis of 236



A-362 A-402d A-402e 236

Synthesis of tert-butyl (S)-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)carbamate (A-402d)

[000820] To a stirred solution of **A-362** (0.6 g, 2.92 mmol) in 1,4-dioxane (15 mL) was added **A-402c** (0.59 g, 2.92 mmol), DCC (0.6 g, 2.92) mmol and stirred at 100°C for 16 h. The reaction mixture was evaporated under reduced pressure and the residue was diluted with EtOAc (50 ml). The organic material was washed with (2 x 15 mL) water and (1 x 15 mL) saturated brine solution, dried over $MgSO_4$ and evaporated to afford the crude product. The product was purified by flash column chromatography eluting 50% EtOAc in hexane to afford desired **A-402d** (0.60 g, 1.29 mmol, 44% yield) as a solid.


Synthesis of (S)-1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)propan-1-amine (A-402e)

[000821] To a stirred solution of **A-402d** (0.6 g, 1.61 mmol) in 1, 4 –Dioxane (2 mL) was added 4 M HCl in Dioxane (10 mL, 1.61 mmol) at 0 °C and stirred at RT for 6 h. After completion, the reaction mixture was evaporated to dryness to afford **A-402e** (0.48 g, 1.49 mmol, 92% yield) as a solid.

Synthesis of (S)-1-methyl-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1H-pyrazole-5-carboxamide (236)

[000822] To a stirred solution of **A-402e** (0.32 g, 1.03 mmol) and 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (0.2 g, 1.03 mmol) in DCM (15 ml) was added HATU (0.47 g, 1.24 mmol) followed by DIPEA (0.18 mL, 1.03 mmol) and stirred at RT for 6 h. The reaction mixture was quenched with water (10 mL) and diluted with DCM (10 mL). The organic layer was separated, washed with brine solution (1 x 10 mL) and dried over $MgSO_4$ and evaporated to afford the crude product. The crude product was purified by flash column chromatography using 50% EtOAc in hexane as an eluent to afford **236** (58 mg, 0.12 mmol, 12% yield) as a solid. **HPLC**: Rt 7.95 min, 99.7%; Column: X-Bridge C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% NH_3 in water B: ACN; Flow Rate: 1.2 mL/min. **LCMS** : 449.04 (M+H), Rt 2.21 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.43 (d, 1H), 9.02 (d, 1H), 8.28-8.27 (m, 2H), 7.49 (s, 1H), 5.38-5.28 (m, 1H), 4.12 (s, 3H), 2.21 – 2.00 (m, 2H), 1.03 (t, 3H).

Example 189. Synthesis of 237



Synthesis of 4-(trifluoromethyl)pyrimidine-2-carbonitrile (A-403b)

[000823] To a stirred solution of **A-403a** (2 g, 10.96 mmol)) and Zinc cyanide (769.19 mg, 6.57 mmol) in DMF (20 mL) was added $Pd(PPh_3)_4$ (1.3 g, 1.1 mmol) at RT and stirred at 110°C for 16 h. The reaction mixture was quenched using water (50 mL) and diluted with EtOAc (100 mL x 2). The organic layer was separated, dried over Na_2SO_4 and evaporated to afford crude product. The crude product was purified by column chromatography using 100-200 silica and 5-10% EtOAc/Hexane as an eluent to afford **A-403b** (1.8 g, 10.39 mmol, 94% yield).

[000824] Synthesis of (Z)-N'-hydroxy-4-(trifluoromethyl)pyrimidine-2-carboximidamide (A-403c)

[000825] To a stirred solution of **A-403b** (1 g, 5.78 mmol) in ethanol (20 mL) was added hydroxylamine hydrochloride (602.17 mg, 8.67 mmol) and TEA (1.61 mL, 11.55 mmol) at RT and stirred at 70 °C for 16 h. The reaction was quenched with water (100 mL) and diluted with EtOAc (50 mL x 2). The combined organic layer was separated, dried with Na_2SO_4 and evaporated to afford the crude product. The crude product was purified by column chromatography using 100-200 silica and 10-20% EtOAc/Hexane as an eluent to afford **A-403c** (1 g, 4.75 mmol, 82% yield).

Synthesis of tert-butyl (S)-(1-(3-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-403e)

[000826] To a stirred solution of **A-403c** (1 g, 4.85 mmol) and (tert-butoxycarbonyl)-L-alanine (917.92 mg, 4.85 mmol) in 1,4-dioxane (20 mL) was added DCC (1089.31 mg, 5.29 mmol) and stirred at 75 °C for 16 h. The reaction mixture was quenched with water (100 mL) and diluted with EtOAc (50 mL x 2). The combined organic layer was separated, dried over Na_2SO_4 and evaporated to afford the crude product. The crude product was purified by

column chromatography using 100-200 silica and 20-30% EtOAc/Hexane as eluent to afford **A403-e** (1 g, 2.42 mmol, 49% yield).


Synthesis of (S)-1-(3-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A403-f)

[000827] To a stirred solution of **A403-e** (1 g, 2.78 mmol) in 1,4-dioxane (10 mL) was added 4 M hydrochloric acid in dioxane (10 mL, 2.78 mmol) at 0 °C and stirred for 2 h. The reaction mixture was evaporated to afford the crude product. The crude product was then washed with diethyl ether to afford **A-403f** (300 mg, 0.81 mmol, 29% yield) as a solid.

(S)-1-methyl-3-(trifluoromethyl)-N-(1-(3-(4-(trifluoromethyl)pyrimidin-2-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (237)

[000828] To a stirred solution of **A-403f** (0.15 g, 0.51 mmol) and **A-403g** (118.18 mg, 0.61 mmol) in DCM (10 mL) were added HATU (289.37 mg, 0.76 mmol) followed by DIPEA (0.18 mL, 1.01 mmol) at RT and stirred at RT for 2 h. The reaction mixture was diluted with water (100 mL) and DCM (100 mL x 2). The organic layer was separated, dried over Na_2SO_4 and evaporated to afford the crude product. The crude product was purified by column chromatography using 100-200 silica and 30-80% 50% EtOAc in hexane as an eluent to afford **237** (152.86 mg, 0.35 mmol, 69% yield) as a solid. **HPLC**: Rt 8.36 min, 99.7%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 435.95 (M+H), Rt 1.99 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 9.38 (d, 1H), 8.23 (d, 1H), 7.42 (s, 1H), 5.52-5.44 (m, 1H), 4.10 (s, 3H), 1.67 (d, 3H).


Example 190. Synthesis of 238



[000829] To a stirred solution of 3-(difluoromethyl)-1-methyl-1H-pyrazole-5-carboxylic acid (59.77 mg, 0.34 mmol) in DCM (4 mL) was added DIPEA (0.15 mL, 0.85 mmol) and HATU (193.56 mg, 0.51 mmol) at 0 °C. To the resulting reaction mixture **A-364** (100 mg, 0.34 mmol) was added at 0 °C and was stirred at RT for 3 h. After completion, the reaction was diluted with water and extracted with DCM (3 x 10 mL). Combined organic

layer was dried over sodium sulphate and concentrated to obtain the crude product. The crude product was purified by prep HPLC to afford **238** (55 mg, 0.13 mmol, 39% yield) as a solid. **HPLC**: Rt 8.68 min, 99.9%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS**: 417.15 (M+H), Rt 2.11 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO):** δ_H = 9.43 (d, 1H), 9.02 (d, 1H), 8.29 (s, 2H), 7.27 (s, 1H),7.05 (t, 1H), 5.51-5.47 (m, 1H), 4.08 (s, 3H), 1.69 (d, 3H).

Example 191. Synthesis of 239



Synthesis of ethyl 3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-405b)

[000830] To a stirred solution of sodium nitrite (3.52 g, 50.97 mmol) in toluene: water (80:40 mL) was added 2,2,2-trifluoroethanamine hydrochloride (6.91 g, 50.97 mmol) at 0 °C and stirred at 0 °C for 30 min. The resulting reaction mixture was charged with **A-405a** (5 g, 50.97 mmol) and stirred at 100 °C for 6 h. The reaction mixture was quenched with distilled water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic phase was washed with brine solution (150 mL), dried over anhydrous Na_2SO_4, filtered and evaporated under reduced pressure to afford the crude product. The crude product was then purified by column chromatography (100-200 mesh silica; 5-8% EtOAc in hexane as an eluent) to afford **A-405b** (6.1 g, 29.19 mmol, 57% yield) as a solid.

Synthesis of ethyl 2-(2-methoxyethyl)-5-(trifluoromethyl)pyrazole-3-carboxylate (A-405c)

[000831] To a stirred solution of **A-405b** (1 g, 4.8 mmol) in DMF (10 mL) were added potassium carbonate (1.33 g, 9.61 mmol) followed by 1-bromo-2-methoxy-ethane (0.8 g, 5.77 mmol) at 0 °C and stirred at RT for 6 h. The reaction mixture was diluted with EtOAc

(100 mL) and washed with cold water (5 x 25 mL), brine solution (50 mL). The organic layer was separated, dried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to afford the crude product. The crude product was purified by column chromatography (100-200 mesh silica, 10-12% EtOAc in hexane as an eluent) to afford **A-405c** (600 mg, 2.18 mmol, 45% yield) as an oil.


Synthesis of 1-(2-methoxyethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-405d)

[000832] To a stirred solution of **A-405c** (250 mg, 0.94 mmol) in THF (10 mL) was added a solution of Lithium hydroxide (44.98 mg, 1.88 mmol) in water (3 mL) at 0 °C and stirred at RT for 2 h. The reaction mixture was evaporated to afford the crude product. The crude product was diluted with water (5 mL), acidified with 1 N HCl solution to pH 4 and extracted with EtOAc (3 x 15 mL). The combined organic phase was dried and evaporated under reduced pressure to afford **A-405d** (200 mg, 0.83 mmol, 89% yield) as a solid.

Synthesis of (S)-1-(2-methoxyethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (239)

[000833] To a stirred solution of **A-405d** (100 mg, 0.42 mmol) in DCM (10 mL) was added **A-364** (148.47 mg, 0.50 mmol), HATU (239.48 mg, 0.63 mmol) and DIPEA (0.15 mL, 0.84 mmol) at 0 °C and stirred at RT for 3 h. The reaction mixture was diluted with DCM (10 mL) and washed with water (3 x 15 mL) followed by saturated sodium bicarbonate solution (3 x 15 mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography (100-200 mesh silica, 2-4% EtOAc in hexane as an eluent) to afford **239** (35 mg, 0.07 mmol, 17% yield,) as a solid. **HPLC**: Rt 9.67 min, 99.9%; Column: X-select CSH C18 (4.6*150) mm, 5 µm; Mobile phase: 10mM ammonium bicarbonate in water, B: ACN; Flow Rate: 1.0 mL/min. **LCMS**: 479.04 (M+H), Rt 2.10 min, Column: X-select CSH C18 (3.0*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.55 (d, 1H), 9.05 – 8.99 (m, 1H), 8.29 (d, 2H), 7.40 (s, 1H), 5.52-5.46 (m, 1H), 4.81 – 4.64 (m, 2H), 3.68-3.62 (m, 2H), 3.15 (s, 3H), 1.69 (d, 3H). **Chiral method:** Rt 3.12 min, 100 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40% B till 9 min, 40-10% B at 10 min, hold 10 % B till 12 min. Wavelength: 270 nm, Flow: 3 mL/min.

Example 192. Synthesis of 240



Synthesis of 2-(prop-1-en-2-yl)isonicotinonitrile (A-406c)

[000834] To a stirred solution of **A-406a** (5 g, 27.32 mmol) in 1,4 dioxane (50 mL) and water (50 mL) was added **A-406b** (6.89 g, 40.98 mmol) and sodium carbonate (8.69 g, 81.96 mmol) and degassed with argon for 30 minute. The reaction was concentrated and the residue was diluted with water (30 mL) and EtOAc (100 mL) and the organics were washed with saturated brine solution (1 x 30 mL). The organic layer was separated, dried over $MgSO_4$ and concentrated to dryness. The crude was then purified by flash column chromatography eluting 50% EtOAc in hexane. The desired fractions were concentrated to dryness to afford desired product **A-406c** (3 g, 20.241 mmol, 74% yield) as a solid product.

Synthesis of 2-(1-methylcyclopropyl)isonicotinonitrile (A-406d)

[000835] A stirred solution of Diethyl zinc solution 1.0 M in hexanes (41.61 mL, 41.62 mmol) was added to DCM (100 mL) at 0 °C followed by Diiodomethane (3.36 mL, 41.62 mmol) drop wise and the reaction mixture was stirred at the same temperature for 30 min. After that **A-406c** (1.5 g, 10.4 mmol) in DCM (20 mL) was added to the reaction mixture at 0 °C drop wise and stirred at RT for 1.5 h. The reaction was quenched with sat. NH_4Cl at 0 °C and diluted with DCM (100 mL). The organic layer was separated and washed with 1x 30 ml saturated brine solution, dried over $MgSO_4$ and concentrated to obtain the crude product. The crude compound was purified by flash column chromatography eluting 20% EtOAc in hexane to afford desired product **A-406d** (0.40 g, 1.22 mmol, 11% yield) as a solid.

Synthesis of N-hydroxy-2-(1-methylcyclopropyl)isonicotinimidamide (A-406e)

[000836] To a stirred solution of **A-406d** (0.4 g, 2.53 mmol) in ethanol (20 mL) was added Hydroxyl amine hydrochloride (0.31 g, 3.79 mmol) and triethyl amine (0.51 g, 5.06 mmol) and heated to 80 °C for 12 hours. The reaction mixture was concentrated to dryness and the residue was taken up in EtOAc (60 mL). The organic layer was washed with water (2 x 30 mL) then saturated brine solution (1 x 30 mL). The organic layer was then separated,

dried over $MgSO_4$ and concentrated to afford the desired product **A-406e** (0.48 g, 2.04 mmol, 80% yield) as an oil.

Synthesis of tert-butyl (S)-(1-(3-(2-(1-methylcyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (272)

[000837] To a stirred solution of **A-406e** (0.48 g, 2.51 mmol) in 1,4-dioxane (15 mL) was added N-Boc-L-alanine (0.52 g, 2.76 mmol) and DCC (0.57 g, 2.76 mmol) and reaction mixture was stirred at 100 °C for 16 hour. After completion, the reaction was concentrated under reduced pressure and the residue was taken up in EtOAc (30 mL). The organics was washed with water (2 x 10 mL) and saturated brine solution (1 x 10 mL). The organic layer was then separated and dried over $MgSO_4$ before concentration to obtain crude. The crude compound was purified by flash column chromatography eluting 30% EtOAc in hexane to afford the desired compound (0.80 g, 2.28 mmol, 91% yield). From this material was purified 200mg by prep-HPLC to afford (35 mg) of **272**. **HPLC**: Rt 8.00 min, 98.4 %; Column: X-Bridge C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1 NH_3 in water, B: Acetonitrile; Flow Rate: 1.2 mL/min. **LCMS** : 345.15 (M+H), Rt 2.27 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 8.65 (d, 1H), 7.84-7.75 (m, 2H), 7.67 (d, 1H), 5.04-4.96 (m, 1H), 1.55-1.48 (m, 6H), 1.42-1.34 (m, 9H), 1.25-1.20 (m, 2H), 0.92-0.85 (m, 2H). **Chiral method:** Rt 4.91 min, 99.2 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10 % B till 12 min. Wavelength: 290 nm, Flow: 3 mL/min.

Synthesis of (S)-1-(3-(2-(1-methylcyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine hydrochloride (A-406f)


[000838] To a stirred solution of **272** (0.6 g, 1.74 mmol) in 1,4-dioxane (2 mL) was added 4 M hydrochloric acid in dioxane (6 mL) at 0 °C and stirred for 6 h. After completion, the reaction was concentrated to afford the desired product **A-406f** (0.40 g, 1.32 mmol, 76% yield).

Synthesis of (S)-1-methyl-N-(1-(3-(2-(1-methylcyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (240)

[000839] To a stirred solution of **A-406f** (0.22 g, 0.77 mmol) and **1**-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (0.15 g, 0.77 mmol) in DCM (15 mL) was added HATU (0.32 g, 0.85 mmol) followed by DIPEA (0.27 mL, 1.55 mmol) at 0 °C and stirred at RT for 6 h. After completion, the reaction mixture was diluted with water (20 mL) and DCM (30 mL). The organic layer was separated, washed with (1 x 20 mL) saturated

brine solution and dried over $MgSO_4$ and concentrated to obtain the crude product. The crude material was purified by flash column chromatography eluting 30% EtOAc in hexane and concentrated to afford compound **240** (110 mg, 0.25 mmol, 33% yield) as a solid. **HPLC**: Rt 9.20 min, 98.4 %; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 421.20 (M+H), Rt 2.13 min; Column: X-select CSH (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.65 (d, 1H), 7.82 (s, 1H), 7.67 (d, 1H), 7.44 (s, 1H), 5.52-5.44 (m, 1H), 4.12 (s, 3H), 1.68 (d, 3H), 1.52 (s, 3H), 1.22 (d, 2H), 0.92-0.86 (m, 2H). **Chiral method:** Rt 3.71 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-25% B in 5 min, hold 25 % B till 9 min, 25-10% B in 10 min, hold 10% B till 12 min. Wavelength: 290 nm, Flow: 3 mL/min.

Example 193. Synthesis of 241



Synthesis of ethyl 1-benzyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-406g)

[000840] To a stirred solution of **A-405b** (0.5 g, 2.4 mmol) in DMF (10 mL) was added potassium carbonate (663.88 mg, 4.8 mmol) followed by addition of benzyl bromide (0.34 mL, 2.88 mmol) at 0 °C and stirred at RT for 6 h under nitrogen. The reaction mixture was diluted with EtOAc (100 mL) and washed with cold water (20 mL) then brine solution. The organic layer wasdried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to afford **A-406g** (500 mg, 1.55 mmol, 64% yield) as an oil.

Synthesis of 1-benzyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-406h)


[000841] To a stirred solution of **A-406g** (500 mg, 1.68 mmol) in THF (5 mL) was added a solution of Lithium hydroxide.H_2O (140.7 mg, 3.35 mmol) in water (5 mL) at 0 °C

and stirred at RT for 2 h. The reaction mixture was evaporated to afford the crude product. The crude product was acidified with 2 N HCl (5 mL) leading to precipitation. The precipitate formed was filtered and dried under reduced pressure to afford **A-406h** (300 mg, 1.08 mmol, 64% yield) as a solid.

Synthesis of (S)-1-benzyl-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (241)

[000842] To a stirred solution of **A-406h** (200 mg, 0.74 mmol) in DCM (10 mL) was added **A-364** (229.3 mg, 0.890 mmol), HATU (337.7 mg, 0.89 mmol) and DIPEA (0.26 mL, 1.48 mmol) under nitrogen and stirred at RT for 3 h. The reaction mixture was diluted with DCM (25 mL), washed with water (20 mL) and brine. The aqueous layer was again extracted with DCM (25 mL) and washed with brine. The combined organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography using 100-200 mesh silica and 15% EtOAc in hexane as an eluent to afford **241** (35mg, 0.06 mmol, 9% yield) as a solid. **HPLC**: Rt 9.93 min, 95.4 %; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 511.15 (M+H), Rt 2.25 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.57 (d, 1H), 9.02 (d, 1H), 8.30-8.24 (m, 2H), 7.48 (s, 1H), 7.30-7.14 (m, 5H), 5.82-5.72 (m, 2H), 5.51-5.72 (m, 1H), 1.68 (d, 3H). **Chiral method:** Rt 3.61 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) Iso-propyl alcohol, Gradient:10-15% B in 5 min, hold 15%Btill 9 min, 15-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

Example 194. Synthesis of 242



Synthesis of methyl 2-(trifluoromethyl)pyrimidine-4-carboxylate (A-407b)

[000843] To a solution of **A-407a** (1 g, 5.48 mmol) in methanol (15 mL) was added triethyl amine (0.57 mL, 4.11 mmol), 1,1′-ferrocenediyl-bis(diphenylphosphine) (303.72 mg,

0.55 mmol) and palladium (II) acetate (0.06 g, 0.27 mmol) The reaction mixture was heated at 60 °C under carbon monoxide (60 psi) for 16 h. The reaction mixture was filtered through celite and washed with EtOAc (3 x 20 mL). The combined organic phase was dried over anhydrous sodium sulphate, filtered and dried under reduced pressure to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 15-18% EtOAc in hexane as an eluent to afford **A-407b** (700 mg, 3.4 mmol, 62% yield) as a solid.

Synthesis of 2-(trifluoromethyl)pyrimidine-4-carbonitrile (A-407c)

[000844] To a solution of **A-407b** (600 mg, 2.91 mmol) in methanol (5 mL) was added 7 M ammonia in methanol (1.27 mL, 58.22 mmol) and was stirred at 50 °C for 6 h. The reaction mixture was evaporated under reduced pressure to afford a solid which was washed with pentane (20 mL) and dried under reduced pressure to afford the product (500 mg, 2.52 mmol, 87% yield) as a solid. To a solution of oxalyl chloride (166.04 mg, 1.31 mmol) in DMF (15mL) were added pyridine (0.63 mL, 7.85 mmol) and a solution of 2-(trifluoromethyl)pyrimidine-4-carboxamide (500.mg, 2.62 mmol) in DMF (5 mL) at 0°C. The reaction mixture was stirred at the 0 °C for 3 h. and then stirred at RT for 16 h. The reaction mixture was evaporated under reduced pressure to afford the crude product which was diluted with EtOAc (200 mL) and washed with cold water (5 x 30 mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford **A-407c** (400 mg, 2.03 mmol, 78% yield) as a liquid

Synthesis of (Z)-N'-hydroxy-2-(trifluoromethyl)pyrimidine-4-carboximidamide (A-407d)

[000845] To a solution of **A-407c** (400 mg, 2.31 mmol) in ethanol (10 mL) was added hydroxyl amine hydrochloride (284.33 mg, 3.47 mmol) followed by triethyl amine (0.64 mL, 4.62 mmol) and then reaction mixture was stirred at 70 °C for 5 h. The solvent was removed under reduced pressure to afford the crude product which was diluted with water (10 mL) and extracted with EtOAc (3 x 15 mL). The combined organic layer was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford **A-407d** (520 mg, 2.47 mmol, quantitative) as a solid.

Synthesis of (tert-butyl (S)-(1-(3-(2-(trifluoromethyl)pyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-407e)

[000846] To a solution of **A-407d** (515 mg, 2.5 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (520 mg, 2.75 mmol) in 1,4-dioxane (10 mL) was added DCC (566.14 mg, 2.75 mmol) and stirred at 100 °C for 16 h. The reaction mixture was

diluted with EtOAc (50 mL), washed with water (3 x 25 mL), followed by brine (25 mL) and organic layer separated. The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 12-14% of EtOAc in hexane as an eluent to afford **A-407e** (150 mg, 0.38 mmol, 15% yield) as a solid.


Synthesis of (S)-1-(3-(2-(trifluoromethyl)pyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine hydrochloride (A-407f)

[000847] To a solution of **A-407e** (150.41 mg, 0.42 mmol) in 1,4-dioxane (10 mL) was added 4.0 M Hydrogen chloride solution in dioxane (0.74 mL, 20.93 mmol) in a dropwise manner at 0 °C and stirred at RT for 3 h. The solvent was removed under reduced pressure to afford the crude product which was washed with ether (3 x 20 mL) and dried to afford **A-407f** (80 mg, 0.22 mmol, 53% yield) as a solid.

Synthesis of ((S)-1-methyl-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyrimidin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (242)

[000848] To a solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (50.mg, 0.26 mmol), **A-407f** (83.77 mg, 0.28 mmol) and HATU (146.91 mg, 0.39 mmol) in DCM (1 0mL) was added DIPEA (0.13 mL, 0.77 mmol) at 0 °C and stirred at RT for 3 h. The reaction mixture was diluted with DCM (15 mL), washed with water (3 x 15 mL), saturated sodium bicarbonate solution (3 x 15 mL) and brine solution (25 mL) and the organic layer was separated. The organic phase was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 100-200- mesh silica and 25- 30% of EtOAc in hexane as an eluent to afford a solid, which was further purified by prep HPLC purification to afford **242** (10mg, 0.023 mmol, 9% yield) as a solid. **HPLC**: Rt 9.01 min, 98.5%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 10mM Ammonium Bicarbonate in water, B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 434.10 (M-H), Rt 2.02 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.52 (d, 1H), 9.32 (d, 1H), 8.42 (d, 1H), 7.46 (s, 1H), 5.55-5.50 (m, 1H), 4.13 (s, 3H), 1.70 (d, 3H). **Chiral method:** Rt 3.09 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) Iso-propyl-alcohol, Gradient:10-15% B in 5 min, hold 15%Btill 9 min, 15-10% B in 10 min, hold 10% B till 12 min. Wavelength: 270 nm, Flow: 3 mL/min.

Example 195. Synthesis of 243



Synthesis of ethyl 1-(2-(dimethylamino)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-408c)

[000849] To a stirred solution of **A-405b** (0.5 g, 2.4 mmol) in DMF (5 mL) was added potassium carbonate (663.88 mg, 4.8 mmol) followed by the addition of 2-chloro-N,N-dimethyl-ethanamine hydrochloride (415.2 mg, 2.9 mmol) at 0 °C and stirred at RT for 16 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer thus separated treated with brine anddried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to afford the crude product which was purified by combi-flash chromatography in MeOH: DCM as an eluent to afford **A-408c** (300 mg, 1.07 mmol, 45% yield) as an oil.

Synthesis of 1-(2-(dimethylamino)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-408d)


[000850] To a stirred solution of **A-408c** (300 mg, 1.07 mmol) in THF (3 mL) was added a solution of lithium hydroxide.H_2O (90.15 mg, 2.15 mmol) in THF (3 mL) at 0 °C and stirred at RT for 3 h. The reaction mixture was evaporated to afford the crude product. The crude product was acidified with 2 N HCl and was lyophilised to afford **A-408d** (280 mg, 1.06 mmol, 99% yield) as a solid.

(S)-1-(2-(dimethylamino)ethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (243)

[000851] To a stirred solution of **A-408d** (153.46 mg, 0.61 mmol) in DCM (10 mL) were added HATU (290.34 mg, 0.76 mmol) and DIPEA (0.27 mL, 1.53 mmol) at 0 ° C for 10 min and added **A-364** (150.mg, 0.51 mmol), and stirred at RT for 3 h. The reaction mixture

was washed with water (20 mL) and extracted with ethyl acetate (3 times). The combined organic layer obtained was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography and then further purified using prep HPLC to afford **243** (60 mg, 0.12 mmol, 24% yield) as a solid. **HPLC**: Rt 5.91 min, 99.63%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 492.20 (M+H), Rt 1.49 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.59 (d, 1H), 9.02 (d, 1H), 8.31 – 8.25 (m, 2H), 7.37 (s, 1H), 5.51-5.46 (m, 1H), 4.69-4.58 (m, 2H), 2.61 (t, 2H), 2.09 (s, 6H), 1.69 (d, 3H). **Chiral method:** Rt 4.89 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) Iso-propyl-alcohol, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

Example 196. Synthesis of 244




A-364 **244**

[000852] To a stirred solution of 1,3,4-trimethyl-1H-pyrazole-5-carboxylic acid (125.6 mg, 0.81 mmol) in DCM (10 mL) was added **A-364** (0.2 g, 0.68 mmol), HATU (387.12 mg, 1.02 mmol) and DIPEA (0.24 mL, 1.36 mmol) at RT and stirred at RT for 2 h. The reaction mixture was quenched with water (100 mL), diluted with DCM (2 x 100 mL), The combined organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography using 100-200 mesh silica and 30-80% EtOAc in hexane as an eluent to afford **244** (96mg, 0.24 mmol, 36% yield) as a solid. **HPLC**: Rt 8.14 min, 99.9%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 395 (M+H), Rt 1.91 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.08 (d, 1H), 9.03 (d, 1H), 8.30-8.24 (m, 2H), 5.51-5.45 (m, 1H), 3.78 (s, 3H), 2.09-2.06 (m, 6H), 1.67 (d, 3H). **Chiral method:** Rt 4.24 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), -

Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 250 nm, Flow: 3 mL/min.

Example 197. Synthesis of 245



Synthesis of 2-(hydroxymethyl)isonicotinonitrile (A-410b)

[000853] To the stirred solution of **A-410a** (10 g, 96.05 mmol) in methanol (130 mL) was added sulfuric acid (2.57 mL, 48.03 mmol) at RT and stirred for 30 min at refluxe. To the reaction mixture ammonium persulphate (35.07 g, 153.68 mmol) in water (70 mL) was slowly added over 30 min under reflux. The reaction mass was stirred for 4 h at reflux. The reaction mixture was evaporated under reduced pressure to afford a liquid which was treated with saturated solution of sodium bicarbonate to pH 9. The resulting aqueous layer was extracted with DCM (3 x 15 mL). The combined organic layer was separated, dried over sodium sulfate to afford the crude product which was purified using column chromatography, product using 40% of ethyl acetate in hexane as an eluent to afford **A-410b** (3.4 g, 23.06 mmol, 24% yield).

Synthesis of 2-(ethoxymethyl)isonicotinonitrile (A-410c)

[000854] To the stirred solution of **A-410b** (1.g, 7.45 mmol) in THF (15 mL) was added sodium hydride (0.75 g, 18.64 mmol) at 0-5 °C and the reaction mixture was stirred for 30 min at 0-5 °C. To the resulting reaction mixture was added dropwise iodoethane (1.2 mL, 14.91 mmol) and stirred for 3 h at RT. The reaction mixture was diluted with ethylacetate (2 mL) and the reaction quenched with water at 0-5 °C and the aqueous layer extracted with ethyl acetate. The organic layer was dried over sodium sulphate and evaporated to afford **A-410c** (500 mg, 2.59 mmol, 35% yield) as a liquid.

Synthesis of (Z)-2-(ethoxymethyl)-N'-hydroxyisonicotinimidamide (A-410d)

[000855] To a stirred solution of **A-410c** (500.mg, 3.08 mmol) in ethanol (5 mL) was added triethylamine (0.86 mL, 6.17 mmol) and hydroxylamine hydrochloride (321.34 mg, 4.62 mmol) at RT and refluxed for 4 h. The reaction mixture was evaporated to afford the

crude product which was diluted with ethyl acetate andwashed with water. The organic layer separated and the aqueous layer was extracted withethyl acetate (4 x 10 mL). The combined organic layer was dried over sodium sulphate evaporated under reduced pressure to afford **A-410d** (400 mg, 1.97 mmol, 64% yield).

Synthesis of tert-butyl (S)-(1-(3-(2-(ethoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-410e)

[000856] To a stirred solution of **A-410d** (400 mg, 2.05 mmol) in 1,4-dioxane (5 mL) was added *N*,*N*'-Dicyclohexylcarbodiimide (422.76 mg, 2.05 mmol) and N-Boc-L-alanine (387.69 mg, 2.05 mmol) at RT then was refluxed for 4 h. The reaction mixture was filtered and the mother liquor was diluted with ethyl acetate and washed with water (3 x 10 mL). The organic layer was dried over sodium sulphate and evaporated under reduced pressure to obtain the crude product which was purified by column chromatography using 25-30% of ethyl acetate in hexane to afford **A-410e** (370 mg, 1.06 mmol, 52% yield) as a solid.

Synthesis of (S)-1-(3-(2-(ethoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-410f)


[000857] To the stirred solution of **A-410e** (270 mg, 0.77 mmol) in 1,4-dioxane (2mL) was added 4 M HCl in 1,4 dioxane (2 mL) at 0-5 °C and stirred for 2 h at RT. The reaction mixture was evaporated completely under reduced pressure to afford **A-410f** (200 mg, 0.68 mmol, 88% yield) as the hydrochloride salt.

Synthesis of (S)-N-(1-(3-(2-(ethoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (245)

[000858] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (156.36 mg, 0.81 mmol) in DCM (2 mL) was added DIPEA (0.42 mL, 2.42 mmol), HATU (459.44 mg, 1.21 mmol) and **A-410f** (200.mg, 0.8100 mmol) at RT and stirred for 2 h at RT. The reaction mixture was diluted with DCM and washed with water. The organic layer was dried over sodium sulphate and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 30-35% of ethyl acetate in hexane as an eluent to afford **245** (40 mg, 0.09 mmol, 11% yield,) as a solid. **HPLC**: Rt 8.41 min, 99.6%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 425.15 (M+H), Rt 1.95 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.73 (d, 1H), 7.95 (s, 1H), 7.85-7.82 (m, 1H), 7.45 (s, 1H), 5.52-5.44 (m, 1H), 4.63 (s, 2H), 4.13 (s, 3H), 3.60 (q, 2H), 1.68 (d, 3H), 1.20 (t, 3H). **Chiral method:** Rt 3.57 min, 99.5%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5

um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

Example 198. Synthesis of 246



Synthesis of ethyl 1-(pyridin-4-ylmethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-411c)

[000859] To a stirred solution of **A-405b** (500 mg, 2.4 mmol) and 4-(bromomethyl)pyridine, hydrobromide (729.12 mg, 2.88 mmol) at RT in DMF (5 mL) were added K_2CO_3 (274.21 mg, 3.6 mmol) at RT and stirred at RT for 4 h. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic layer was separated, dried over Na_2SO_4 and evaporated to afford crude product which was purified by column chromatography using 30-35% ethyl acetate in hexane as an eluent to afford **A-411c** (250 mg, 0.79 mmol, 32% yield).


Synthesis of 1-(pyridin-4-ylmethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-411d)

[000860] To a stirred solution of **A-411c** (200 mg, 0.67 mmol) in THF (2 mL) was added a solution of lithium hydroxide (84.13 mg, 2.01 mmol) in water (0.2000 mL) at RT and stirred at RT for 2 h. The reaction mixture was evaporated to afford crude product. The crude product was acidified with 6 M HCl. Aqueous layer was lyophilized to afford **A-411d** (180 mg, 0.54 mmol, 81% yield) as a solid.

Synthesis of (S)-1-(pyridin-4-ylmethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (246)

[000861] To a stirred solution of **A-411d** (126.04 mg, 0.46 mmol) and **A-364** (100 mg, 0.39 mmol) in DCM (2 mL) was added HATU (147.26 mg, 0.39 mmol), DIPEA (0.2 mL, 1.16 mmol) and stirred at RT for 2 h. The reaction mixture was diluted with water and DCM. The organic layer was separated, washed with brine solution, dried over Na_2SO_4 and evaporated to afford the crude product. The crude product was purified by column chromatography using silica gel 100-200 mesh and 30-35% EA in Hexane eluent to afford **246** (40 mg, 0.07 mmol, 19% yield) as a solid. **HPLC**: Rt 7.71 min, 98 %; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: Acetonitrile (95:05), B: Acetonitrile; Flow Rate: 1.0 mL/min. **LCMS** : 512.10 (M+H), Rt 1.81 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.60 (d, 1H), 9.01 (d, 1H), 8.48 (d, 2H), 8.28-8.20 (m, 2H), 7.57 (s, 1H), 7.04 (d, 2H), 5.83 (s, 2H), 5.49-5.42 (m, 1H), 1.66 (d, 3H). **Chiral method:** Rt 3.50 min, 99.3 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MEOH + 0.1 % NH3, Gradient:10-40 % B in 5 min, hold 40 % B till 9 min, 40-10 % B at 10 min, hold 10 % B till 12 min, Wavelength: 280 nm, Flow: 3 mL/min.

Example 199. Synthesis of 247 and 248



Synthesis of tert-butyl (S)-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)carbamate (247)

[000862] To a stirred solution of **A-301** (500 mg, 2.82 mmol) in 1, 4 –Dioxane (10 mL) was added (S)-2-((tert-butoxycarbonyl)amino)butanoic acid (630.82 mg, 3.1 mmol) followed by DCC (639.39 mg, 3.1 mmol) and stirred at 70 °C for 5 h. The solvent was removed under reduced pressure and residue was diluted with water (10 mL) and extracted with EtOAc (3 x 15 mL). The combined organic phase was dried and evaporated to afford crude compound, which was purified by column chromatography (15-22% EtOAc in hexane as an eluent) followed by prep HPLC purification to afford **247** (510 mg, 1.47 mmol, 52% yield) as a semi solid. **HPLC**: Rt 9.23 min, 99.8%; Column: X-Select C18 (4.6 x 150) mm, 5 µm; Mobile

phase: A:0.1% Formic acid in water B: Acetonitrile (95:05); Flow Rate: 1.0 mL/min. **LCMS** : 345.20 (M+H), Rt 2.16 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 8.60 (d, 1H), 7.84-7.78 (m, 2H), 7.64 (d, 1H), 4.82–4.80 (m, 1H), 2.32-2.28 (m, 1H), 1.97-1.85 (m, 2H), 1.40 (s, 9H), 1.22-1.21 (m, 2H), 1.22-0.94 (m, 5H). **Chiral method:** Rt 6.21 min, 99.1 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40 % B in 5 min, hold 40 % B till 9 min, 40-10 % B at 10 min, hold 10 % B till 12 min, Wavelength: 292 nm, Flow: 3 mL/min.


Synthesis of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propan-1-amine (A-412f)

[000863] To a stirred solution of **247** (400 mg, 1.16 mmol) in 1, 4 –dioxane (10 ml) was added HCl solution 4.0 M in dioxane (0.41 mL, 11.61 mmol) at 0 °C and the reaction mixture was stirred at RT for 3 h. The solvent was evaporated under reduced pressure, resulting crude product was washed with *n*-pentane (3 x 20 mL) and dried under reduced pressure to afford **A-412f** (390 mg, 0.96 mmol, 82% yield) as a solid.

[000864] Synthesis of (S)-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (248)

[000865] To a stirred solution of **A-412f** (100 mg, 0.36 mmol) and 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (76.05 mg, 0.39 mmol) in DCM (10 ml) were added HATU (203.15 mg, 0.53 mmol) followed by DIPEA (0.12 mL, 0.71 mmol) at 0 °C and stirred at RT for 3 h. The reaction mixture was diluted with DCM (30 mL), washed with water (3 x 25 mL) and sodium bicarbonate solution (3 x 25 mL). The organic phase was separated, dried over anhydrous sodium sulfate, filtered and evaporated to afford the crude product, which was purified by column chromatography (100-200 mesh silica, 8-9.5% of EtOAc in hexane as an eluent) followed by prep HPLC to afford **248** (50 mg, 0.11 mmol, 33% yield) as a solid. **HPLC**: Rt 9.64 min, 99.7%; Column: X-Select C18 (4.6 x 150) mm, 5 µm; Mobile phase: A:10 mM ammonium bicarbonate in water B: Acetonitrile; Flow Rate: 1.0 mL/min. **LCMS** : 421.50 (M+H), Rt 2.14 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.41 (d, 1H), 8.60 (d, 1H), 7.85 (d, 1H), 7.68-7.62 (m, 1H), 7.49 (s, 1H), 5.32 – 5.26 (m, 1H), 4.13 (s, 3H), 2.34-2.22 (m, 1H), 2.21 – 1.97 (m, 2H), 1.06 – 0.92 (m, 7H). **Chiral method:** Rt 3.24 min, 100 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40 % B in 5 min, hold 40 % B till 9 min, 40-10 % B at 10 min, hold 10 % B till 12 min, Wavelength: 292 nm, Flow: 3 mL/min.

Example 200. Synthesis of 249



Synthesis of ethyl 1-(2-oxo-2-(pyrrolidin-1-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-413c)

[000866] To a stirred solution of **A-405b** (500 mg, 2.4 mmol) and 2-chloro-1-pyrrolidin-1-yl-ethanone (390.03 mg, 2.64 mmol) in DMF (5 mL) were added K_2CO_3 (663.88 mg, 4.8 mmol) at 0 °C and stirred at RT for 16 h. The reaction mixture was diluted with ethyl acetate (50 mL) and the organic layer was washed with water (3 x 25 mL) followed by brine (25 mL). The organic phase was dried over anhydrous sodium sulfate, filtered and evaporated to afford crude product which was purified by column chromatography (100-200 mesh silica, 10-1% EtOAc in hexne as an eluent) to afford **A-413c** (480 mg, 1.46 mmol, 61% yield) as a solid.

Synthesis of 1-(2-oxo-2-(pyrrolidin-1-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-413d)


[000867] To a stirred solution of **A-413c** (481.49 mg, 1.51 mmol) in THF (10 mL) was added a solution of lithium hydroxide (54.18 mg, 2.26 mmol) in water (3 mL) at 0°C and stirred at RT for 3 h. The reaction mixture was evaporated to afford the crude product. The crude product was diluted with water (15 mL), acidified with 5 N HCl to pH 4, precipitate collected by filtration and dried to afford **A-413d** (320 mg, 1.09 mmol, 72% yield) as a solid.

Synthesis of (S)-1-(2-oxo-2-(pyrrolidin-1-yl)ethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (249)

[000868] To a stirred solution of **A-413d** (100 mg, 0.34 mmol) and **A-364** (121.41 mg, 0.41 mmol) in DCM (10 mL) were added HATU (195.84 mg, 0.52 mmol) followed by

DIPEA (0.18 mL, 1.03 mmol) at 0°C and stirred at RT for 3 h. The reaction mixture was diluted with DCM (25 mL). The organic layer was separated, washed with sodium bicarbonate solution (3 x 25 mL), brine solution (30 mL) then dried (Na_2SO_4) and concentrated to afford the crude product. The crude product was purified by column chromatography using silica gel 100-200 mesh and 35-40% EA in hexane as an eluent to afford **249** (30 mg, 0.05 mmol, 16% yield) as a solid. **HPLC**: Rt 9.12 min, 99.9 %; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 10mM ammonium bicarbonate in water B: Acetonitrile; Flow Rate: 1.0 mL/min. **LCMS** : 531.90 (M+H), Rt 2.04 min; Column: X-select CSH (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.50 (d, 1H), 9.02 (d, 1H), 8.28-8.24 (m, 2H), 7.49 (s, 1H), 5.52-5.40 (m, 3H), 3.50-3.40 (m, 2H), 3.26-3.18 (m, 2H), 1.90-1.80 (m, 2H), 1.78-1.62 (m, 5H). **Chiral method:** Rt 3.86 min, 100 %; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MEOH + 0.1 % NH3, Gradient:10-40 % B in 5 min, hold 40 % B till 9 min, 40-10 % B at 10 min, hold 10 % B till 12 min, Wavelength: 280 nm, Flow: 3 mL/min.

Example 201. Synthesis of 250



Synthesis of 2-(4-bromopyridin-2-yl)propan-2-ol (A-414b)

[000869] To a stirred solution of **A-414a** (5 g, 23.14 mmol) in THF (50 mL) was added MeMgBr (8.28 g, 69.43 mmol) at 0 °C in dropwise manner and stirred at RT for 2 h. The reaction mixture was diluted withsaturated NH_4Cl solution and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford the crude product which was purified by combi flash using 15% EtOAc /hexane as an eluent to afford **A-414b** (1.9 g, 7.12 mmol, 31% yield) as an oil.

Synthesis of 4-bromo-2-(2-methoxypropan-2-yl)pyridine (A-414c)

[000870] To a stirred solution of **A-414b** (1.9 g, 8.83 mmol) in THF (20 mL), was added NaH (0.42 g, 17.67 mmol) at 0 °C portion-wise and stirred at 0 °C for 20 min. To the resulting reaction mixture was addedMethyl iodide (1.25 g, 8.83 mmol)and stirred at RT for 2 h. The reaction mixture was diluted with water and extracted with EtOAc . The organic

layer was dried over Na_2SO_4, filtered and evaporated to afford **A-414c** (1.6 g, 6.9835 mmol, 79% yield) as an oil.

Synthesis of 2-(2-methoxypropan-2-yl)isonicotinonitrile (A-414d)

[000871] To a stirred solution of **A-414c** (1 g, 4.35 mmol) in 1,4-dioxane (10 mL), was added. $ZnCN_2$ (1.53 g, 13.04 mmol) and the reaction mixture was purged with N_2 gas for 20 min followed by the addition of DPPF (0.24 g, 0.43 mmol), Pd_2dba_3 (398.09 mg, 0.43 mmol). The reaction mixture was stirred at 100 °C for 16 h. The reaction mixture was filtered through a celite bed, diluted with water and extracted with EtOAc . The organic layer was dried over Na_2SO_4, filtered and evaporated to afford **A-414d** which was used for next step without further purification.

Synthesis of (Z)-N'-hydroxy-2-(2-methoxypropan-2-yl)isonicotinimidamide (A-414e)

[000872] To a stirred solution of **A-414d** (700 mg, 3.97 mmol) in ethanol (5 mL) were added $NH_2OH.HCl$ (1.16 g, 11.92 mmol), TEA (0.55 mL, 3.97 mmol) and heated 70 °C for 1 h. The reaction mixture was evaporated to afford the crude product which was diluted in water and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford **A-414e** (600 mg, 2.87 mmol, 72% yield) as an oil.

Synthesis of tert-butyl (S)-(1-(3-(2-(2-methoxypropan-2-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-414f)

[000873] To a solution of **A-414e** (500 mg, 2.39 mmol) in 1,4-dioxane (10 mL) was added (2S)-2-(tert-butoxycarbonylamino)propanoic acid (452.14 mg, 2.39 mmol), DCC (492.26 mg, 2.39 mmol) and heated to 100 °C for 16 h. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic layer was dried with Na_2SO_4, filtered and evaporated to afford the tcrude product which was purified by column chromatography using 100-200 mesh silica and 15% EtOAc/hexane as eluents to afford **A-414f** (320 mg, 0.88 mmol, 37% yield) as an oil.


Synthesis of (S)-1-(3-(2-(2-methoxypropan-2-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-414g)

[000874] To a solution of **A-414f** (300 mg, 0.83 mmol) in DCM (5 mL) was added TFA (0.19 mL, 2.48 mmol) in a dropwise manner and stirred at RT for 4 h. The reaction mixture was diluted with saturated $NaHCO_3$ solution and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford **A-414g** (75 mg, 0.28 mmol, 34% yield) as an oil.

Synthesis of (S)-N-(1-(3-(2-(2-methoxypropan-2-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (250)

[000875] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (53.28 mg, 0.27 mmol) in DCM (3 mL) was added **A-414g** (72.mg, 0.27 mmol), HATU (130.46 mg, 0.34 mmol), DIPEA (0.05 mL, 0.27 mmol) and stirred at RT for 16 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford the crude product which was purified by prep HPLC to afford **250** (25 mg, 0.057 mmol, 21% yield) as an oil. **HPLC**: Rt 8.90 min, 99.67%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 439.15 (M+H), Rt 2.02 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.75 (d, 1H), 8.06 (s, 1H), 7.84-7.80 (m, 1H), 7.45 (s, 1H), 5.52-5.46 (m, 1H), 4.13 (s, 3H), 3.15 (s, 3H), 1.68 (d, 3H), 1.50 (s, 6H). **Chiral method:** Rt 3.06 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

Example 202. Synthesis of 251



Synthesis of ethyl 1-(2-(dimethylamino)-2-oxoethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-415c)

[000876] To a solution of **A-405b** (400.mg, 1.92 mmol) in DMF (5 mL) was added K_2CO_3 (0.33 mL, 3.84 mmol) at 0 °C and stirred for 10 min at 0 °C. To a reaction mixture was added 2-chloro-N,N-dimethylacetamide (0.22 mL, 2.11 mmol) and stirred at RT for 4 h. The reaction mixture was diluted with EtOAc (100 mL) and washed with cold water (5 x 25 mL), organic layer separated. The organic layer was dried over anhydrous sodium sulphate,

filtered and evaporated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 18-22% of EtOAc in hexane as an eluent to afford **A-415c** (355 mg, 1.19 mmol, 62% yield) as an oil.


Synthesis of 2-[2-(dimethylamino)-2-oxo-ethyl]-5-(trifluoromethyl)pyrazole-3-carboxylic acid (A-415d)

[000877] To a solution of **A-415c** (350 mg, 1.19 mmol) in MeCN (10 mL) was added a solution of lithium hydroxide (42.88 mg, 1.79 mmol) in water (3 mL) at 0 °C and stirred at RT for 3 h. The solvent was removed under reduced pressure and the crude product was diluted with water (15 mL) and acidified with 5 N aq. HCl solution to pH 5. The obtained precipitate was filtered, washed with water (10 mL) followed by hexane (10 mL) and dried under reduced pressure to afford **A-415d** (130 mg, 0.49 mmol, 41% yield) as a solid.

Synthesis of (S)-1-(2-(dimethylamino)-2-oxoethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (251)


[000878] To a suspension of **A-415d** (95.mg, 0.36 mmol) and **A-364** hydrochloride salt (126.67 mg, 0.43 mmol) in DCM (10 mL) were added HATU (204.32 mg, 0.54 mmol) followed by DIPEA (0.19 mL, 1.07 mmol) at 0 °C and reaction mixture was stirred at RT for 3 h. The reaction mixture was diluted with DCM (25 mL), washed with water (3 x 20 mL), saturated sodium bicarbonate solution (3 x 20 mL) followed by brine solution (30 mL). The organic phase was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford the crude product, which was purified by column chromatography using 100-200 mesh silica and 25-28% EtOAc in hexane as an eluent to afford **251** (35 mg, 0.069 mmol, 19% yield) as a solid. **HPLC**: Rt 8.86 min, 99.9%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 10mM Ammonium Bicarbonate in water, B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 506.11 (M+H), Rt 2.04 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.52-9.48 (m, 1H), 9.02 (d, 1H), 8.30-8.26 (m, 2H), 7.50 (s, 1H), 5.53 (s, 2H), 5.48-5.45 (m, 1H), 3.00 (s, 3H), 2.78 (s, 3H), 1.66 (d, 3H). **Chiral method:** Rt 4.35 min, 99.7%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 270 nm, Flow: 3 mL/min.

Example 203. Synthesis of 252



[000879] To a stirred solution of 1,3-dimethyl-1H-pyrazole-5-carboxylic acid (57.07 mg, 0.41 mmol) in DCM (2 mL) was added **A-364** (0.1 g, 0.34 mmol), HATU (193.56 mg, 0.51 mmol) and DIPEA (0.18 mL, 1.02 mmol) at RT and stirred at RT for 2 h. The reaction mixture was diluted with DCM and washed with water (3 x 3 mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography using 100-200 mesh silica and 20-25% EtOAc in hexane as an eluent to afford **252** (65 mg, 0.1665 mmol, 49% yield) as a solid. **HPLC**: Rt 8.14 min, 97.43%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 381.50 (M+H), Rt 1.79 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.18 (d, 1H), 9.02 (d, 1H), 8.30-8.26 (m, 2H), 6.75 (s, 1H), 5.48-5.42 (m, 1H), 3.95 (s, 3H), 2.17 (s, 3H), 1.67 (d, 3H). Chiral method: Rt 3.96 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 240 nm, Flow: 3 mL/min.

Example 204. Synthesis of 253



Synthesis of ethyl 1-(pyridin-3-ylmethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylate (A-417b)

[000880] To a stirred solution of **A-405b** (300 mg, 1.44 mmol) and 3-(bromomethyl)pyridine (297.53 mg, 1.73 mmol) in DMF (10 mL) were added K_2CO_3 (398.33 mg, 2.88 mmol) and stirred at RT for 10 h. The reaction mixture was diluted by ethyl acetate (20 mL) and water (20 mL) and the organic layer was separated. The aqueous layer was treated with ethylacetate (20 mL). The combined organic layer was washed with cold water (20 mL), brine, dried over Na_2SO_4 and evaporated to afford the crude product which was purified by column chromatography using silica gel 100-200 mesh and 40% EA in Hexane as an eluent to afford **A-417b** (300 mg, 0.91 mmol, 63% yield) as a solid.

Synthesis of 1-(pyridin-3-ylmethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (A-417c)


[000881] To a stirred solution of **A-417b** (300 mg, 1 mmol) in THF (5 mL) was added lithium hydroxide (48.02 mg, 2.01 mmol) and stirred at 0 °C for 3 h. The solvent was evaporated under reduced pressure and reaction mixture was cooled to 0 °C and acidified using 2 N HCl (5 mL) leading to precipitation. The precipitate was collected by filtration and dried to afford **A-417c** (150 mg, 0.36 mmol, 36% yield).

Synthesis of (S)-1-(pyridin-3-ylmethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (253)

[000882] To a stirred solution of **A-417c** (150 mg, 0.55 mmol) and **A-364** (195.57 mg, 0.66 mmol) in DCM (10 mL) was added HATU (252.37 mg, 0.66 mmol) and DIPEA (0.19 mL, 1.11 mmol) at 0 °C and stirred at RT for 3 h. The reaction mixture was diluted with DCM (25 mL) and water (25 mL) and the organic layer was separated. The aqueous layer was washed with DCM (20 mL). The combined organic layer was washed with brine (30 ml), dried over Na_2SO_4 then evaporated to dryness to afford the crude product which was purified by column chromatography using silica gel 100-200 mesh and 30% EA in Hexane as an eluent to afford **253** (22 mg, 0.043 mmol, 8% yield). **HPLC**: Rt 8.23 min, 99.2%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 512.20 (M+H), Rt 2.00 min, 99.8%; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.60 (d, 1H), 9.01 (d, 1H), 8.45-8.42 (m, 2H), 8.28-8.24 (m, 2H), 7.59-7.56 (m, 1H), 7.50 (s, 1H), 7.33-7.30 (m, 1H), 5.81 (s, 2H), 5.51-5.45 (m, 1H), 1.66 (d, 3H). **Chiral method:** Rt 4.55 min, 99.3%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 %

B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 250 nm, Flow: 3 mL/min.

Example 205. Synthesis of 254



Synthesis of 2-(bromomethyl)isonicotinonitrile (A-418c)

[000883] To the stirred solution of **A-410b** (3 g, 22.36 mmol) in THF (30 mL) was added triphenylphosphine (8.8 g, 33.55 mmol) at RT. Reaction mixture was cooled to 0 °C and carbon-tetrabromide (11.12 g, 33.55 mmol) was added and stirred for 5 h at RT. The reaction mixture was diluted with ethyl acetate and washed with water. The organic layer was dried over sodium sulphate and evaporated under reduced pressure to afford a liquid which was purified by column chromatoghraphy using 30% ethyl acetate in hexane as an eluent to afford **A-418c** (1.27 g, 6 mmol, 27% yield) as a solid.

Synthesis of 2-(cyclopropoxymethyl)isonicotinonitrile (A-418d)

[000884] To a stirred solution of cyclopropanol (0.74 g, 12.69 mmol) in THF (10 mL), maintained 0-5 °C was added sodium hydride (1.01 g, 25.38 mmol) and stirred for 30 min. To the resultant reaction mixture was added **A-418c** (1 g, 5.08 mmol) in THF (10 mL) dropwise. The reaction mixture was stirred for 5 h at RT. The reaction mixture was quenched with ice cold water and extracted with ethyl acetate (2 x 20 mL) and the organic layer was separated. The combined organic layer was dried over sodium sulphate, evaporated to afford the crude product which was purified by column chromatography using 10-15% of ethyl acetate in hexane as an eluent to afford **A-418d** (350 mg, 1.98 mmol, 39% yield).

Synthesis of (Z)-2-(cyclopropoxymethyl)-N'-hydroxyisonicotinimidamide (A-418e)

[000885] To the stirred solution of **A-418d** (500 mg, 2.87 mmol) in ethanol (5 mL) was added triethylamine (0.8 mL, 5.74 mmol) and hydroxylamine hydrochloride (299.18 mg, 4.31 mmol) at RT. The reaction was stirred for 4 h at reflux. The reaction mixture was evaporated to afford a liquid which was diluted with ethyl acetate and washed with water and separated. The aqueous layer was extracted with ethyl acetate (4 x 10 mL). The combined organic layer

was dried over sodium sulphate and evaporated to afford **A-418e** (400 mg, 1.88 mmol, 66% yield).

Synthesis of tert-butyl (S)-(1-(3-(2-(cyclopropoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-418f)

[000886] To the stirred solution of **A-418e** (600 mg, 2.9 mmol) in 1,4-dioxane (6 mL) was added DCC (656.08 mg, 3.18 mmol) and N-Boc-L-alanine (602.61 mg, 3.18 mmol) at RT and stirred for 4 h at reflux. The reaction mixture was filtered and diluted with ethyl acetate, washed with water (3 x 10 mL), dried over sodium sulphate and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 25-30% of ethyl acetated in hexane as an eluent to afford **A-418f** (660 mg, 1.82 mmol, 63% yield).


Synthesis of (S)-1-(3-(2-(cyclopropoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (A-418g)

[000887] To a solution of **A-418f** (500 mg, 1.39 mmol) in 1,4-dioxane (5 mL) was added 4 M HCl/dioxane (5 mL) and stirred for 2 h at RT. The reaction mixture was evaporated to afford **A-418g** (238 mg, 0.79 mmol, 57% yield).

Synthesis of (S)-N-(1-(3-(2-(cyclopropoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (254)

[000888] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (130.82 mg, 0.67 mmol) in DCM (3 mL) was added DIPEA (0.35 mL, 2.02 mmol), HATU (384.39 mg, 1.01 mmol) and **A-418g** (200 mg, 0.67 mmol) and stirred for 2 h at RT. The reaction mixture was diluted with DCM and washed with water (2 x 5mL). Theorganic layer was dried over sodium sulphate and evaporated to afford the crude product which was purified by column chromatography with 20-25% ethyl acetate in hexane as an eluent to afford **254** (60 mg, 0.13 mmol, 20% yield) as a solid. **HPLC**: Rt 8.78 min, 97.7%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 437.25 (M+H), Rt 2.02 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.74 (d, 1H), 7.92 (s, 1H), 7.86-7.84 (m, 1H), 7.45 (s, 1H), 5.51-5.46 (m, 1H), 4.69 (s, 2H), 4.13 (s, 3H), 3.51-3.45 (m, 1H), 1.68 (d, 3H), 0.62-0.58 (m, 2H), 0.52-0.46 (m, 2H). **Chiral method:** Rt 5.14 min, 99.5%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-40% B in 5 min, hold 40 % B till 9 min, 40-10% B in 10 min, hold 10% B till 12 min. Wavelength: 278 nm, Flow: 3 mL/min.

Example 206. Synthesis of 255 and 256



Synthesis of 1-(4-bromopyridin-2-yl)ethan-1-one (A-419b)

[000889] To a stirred solution of **A-419a** (8.g, 43.71 mmol) in THF (50mL) was added MeMgBr (6.25 g, 52.46 mmol) in dropwise manner at 0°C and stirred at RT for 1 h. The reaction mixture was diluted with saturated NH_4Cl solution and extracted with EtOAc. The organic layer dried over Na_2SO_4 filtered and evaporated to afford crude product which was. purified by column chromatography using 100% hexane as an eluent to afford **A-419b** (4 g, 19.4 mmol, 44% yield) as an oil.

Synthesis of 1-(4-bromo-2-pyridyl)ethanol (A-419c)

[000890] To as stirred solution of **A-419b** (2 g, 10 mmol) in methanol (15 mL) was added $NaBH_4$ (758.89 mg, 20 mmol) in a portion-wise manner and stirred at RT for 1 h. The reaction mixture was evaporated to afford the crude product which was diluted with water and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford **A-419c** (2 g, 6.23 mmol, 62% yield) as a semi solid. This material was used in the next step without purification.

Synthesis of 4-bromo-2-(1-methoxyethyl)pyridine (A-419d)

[000891] To a stirred solution of **A-419c** (2 g, 9.9 mmol) in THF (20 mL) was added 60% NaH (0.64 mL, 19.8 mmol) in one portion and was stirred at RT for 30 min. To the reaction mixture was added methyl iodide (2.8 g, 19.8 mmol) and stirred at RT for 1 h. The reaction mixture was diluted with water, extracted with EtOAc and the organic layer separated. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford **A-419d** (2.2 g, 10.2 mmol) as an oil.

Synthesis of 2-(1-methoxyethyl)pyridine-4-carbonitrile (A-419e)

[000892] To a stirred solution of **A-419d** (2 g, 9.26 mmol) in DMF (7mL) was added ZnCN (3.26 g, 27.77 mmol) and the reaction was purged with N_2 for 20 min. To athe reaction mixture was added DPPF (0.51 g, 0.93 mmol), Pd_2dba_3 (0.85 g, 0.93 mmol) and heated at 100°C for 16 h. The reaction was cooled, diluted with water, extracted with EtOAc and the organic layer separated. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 3% EtOAc/hexane as an eluent to afford **A-419e** (1g, 4.62 mmol, 50% yield).

Synthesis of N-hydroxy-2-(1-methoxyethyl)pyridine-4-carboxamidine (A-419f)

[000893] To a stirred solution of **A-419e** (900 mg, 5.55 mmol) in ethanol (10 mL) was added $NH_2OH.HCl$ (3.43 g, 16.65 mmol), TEA (2.32 mL, 16.65 mmol) and heated at 70 °C for 1 h. The reaction mixture was evaporated and crude product was diluted in water, extracted with EtOAc and the organic layer separated. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford **A-419f** (920 mg, 3.9 mmol, 71% yield) as a solid.

Synthesis of tert-butyl N-[(1S)-1-[3-[2-(1-methoxyethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-419g)

[000894] To a stirred solution of **A-419f** (900 mg, 4.61 mmol) in 1,4-dioxane (10 mL) was added (2R)-2-(tert-butoxycarbonylamino)propanoic acid (872.29 mg, 4.61 mmol), DCC (949.7 mg, 4.61 mmol) and stirred at 100 °C for 16 h. The reaction mixture was diluted in water and extracted with ethyl acetate. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford crude product which was purified by column chromatography using 100-200 mesh silica 9% EtOAC/hexane as an eluent to afford **A-419g** (700 mg, 1.99 mmol, 43% yield)

Synthesis of (1S)-1-[3-[2-(1-methoxyethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine (A-419h)

[000895] To a stirred solution of **A-419g** (350 mg, 1 mmol) in DCM (5 mL) was added TFA (0.23 mL, 3.01 mmol) in a dropwise manner and stirred at RT for 3 h. The reaction mixture was diluted with saturated $NaHCO_3$ solution and extracted with ethyl acetate. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford **A-419h** (180 mg, 0.72 mmol, 72% yield) as an oil.

Synthesis of N-((S)-1-(3-(2-((S)-1-methoxyethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (255) and N-((S)-1-(3-(2-((R)-1-methoxyethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-

1H-pyrazole-5-carboxamide (256). Note that the stereochemistry of the ether is randomly assigned

[000896] To a stirred solution of **A-419h** (180 mg, 0.72 mmol) in DCM (5 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (168.87 mg, 0.87 mmol), DIPEA (0.38 mL, 2.17 mmol) andHATU (413.49 mg, 1.09 mmol) and the reaction was stirred at RT for 16 h. The reaction mixture was diluted with water and extracted with DCM and separated. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 30% EtOAc/hexane as an eluent to afford racemic mixture which was purified by chiral chromatography to afford **255** (24 mg, 0.056 mmol, 8% yield) and **256** (20 mg, 0.046 mmol, 6.4% yield).


[000897] The chiral separation was carried out using SFC: Mobile Phase: A) CO_2 B) MeOH+0.1%NH_3 Gradient:10-20% B in 5 min, hold 20-20%B till 4min, 20-40%B at 2min, hold 40-50%B till 2 Min. Column: DIACEL CHIRALPAK-IG (250 x 30 mm, 5 um) Wavelength: 280 nm Flow: 80/min. Note: Stereochemistry has been arbitrarily assigned at one chiral centre.

[000898] **255: HPLC**: Rt 8.30 min, 98.6%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 425.05 (M+H), Rt 1.90 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.75 (d, 1H), 7.93 (s, 1H), 7.86-7.84 (m, 1H), 7.45 (s, 1H), 5.50-5.45 (m, 1H), 4.50 (q, 1H), 4.13 (s, 3H), 3.26 (s, 3H), 1.68 (d, 3H), 1.40 (d, 3H). **Chiral method:** Rt 4.15 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3, Gradient:10-25% B in 5 min, hold 25%Btill 9 min, 25-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

[000899] **256: HPLC**: Rt 8.29 min, 99.2%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 425.05 (M+H), Rt 1.89 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.75 (d, 1H), 7.93 (s, 1H), 7.86-7.84 (m, 1H), 7.45 (s, 1H), 5.54-5.44 (m, 1H), 4.50 (q, 1H), 4.13 (s, 3H), 3.27 (s, 3H), 1.68 (d, 3H), 1.40 (d, 3H). **Chiral method:** Rt 4.62 min, 99.6%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) CO_2 B) MeOH+0.1% NH_3,

Gradient:10-25% B in 5 min, hold 25%Btill 9 min, 25-10% B in 10 min, hold 10% B till 12 min. Wavelength: 280 nm, Flow: 3 mL/min.

Example 207. Synthesis of 257



Synthesis of 2-(2-pyrrolidin-1-ylethyl)-5-(trifluoromethyl)pyrazole-3-carboxylate (A-420c)

[000900] To a stirred solution of **A-405b** (300 mg, 1.44 mmol) in DMF (5 mL) was added 1-(2-chloroethyl)pyrrolidine (385.18 mg, 2.88 mmol), K_2CO_3 (597.49 mg, 4.32 mmol) and stirred at RT for 2 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was dried with Na_2SO_4,filtered and concentrated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 6% EtOAc/hexane as eluent to afford **A-420c** (200 mg, 0.6551 mmol, 45% yield).

Synthesis of 2-(2-pyrrolidin-1-ylethyl)-5-(trifluoromethyl)pyrazole-3-carboxylic acid (A-420d)


[000901] To a stirred solution of **A-420c** (130 mg, 0.43 mmol) in THF (2 mL) and water (2 mL) was added sodium hydroxide solution (51.1 mg of NaOH in 2 mL water). The reaction mixture was stirred at RT for 3 h and concentrated to afford the crude product which was diluted with water and extracted with EtOAc. The aqueous layer was acidified with 2 N HCl then extracted with EtOAc. The combined organic layer was dried over Na_2SO_4, filtered and concentrated to afford **A-420d** (80 mg, 0.29 mmol, 68% yield) as an oil.

Synthesis of (S)-1-(2-(pyrrolidin-1-yl)ethyl)-3-(trifluoromethyl)-N-(1-(3-(2-(trifluoromethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (257)

[000902] To a stirred solution **A-420d** (107.37 mg, 0.39 mmol) and (1S)-1-[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine (100 mg, 0.39 mmol) in DCM (

3mL) was added DIPEA (0.2 mL, 1.16 mmol), HATU (220.89 mg, 0.58 mmol) and stirred at RT for 16 h .The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was dried ($NaSO_4$), filtered and concentrated to afford the crude product which was purified by prep HPLC to afford **257** (10 mg, 0.018 mmol, 5% yield). **HPLC**: Rt 6.37 min, 94.4%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** :518.10 (M+H), Rt 1.53 min, 99.3%; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.59 (d, 1H), 9.02 (d, 1H), 8.30-8.26 (m, 2H), 7.39 (s, 1H), 5.51-5.46 (m, 1H), 4.71-4.63 (m, 2H), 2.85-2.75 (m, 2H), 2.49-2.42 (m, 4H), 1.69 (d, 3H), 1.62-1.56 (m, 4H). **Chiral method:** Rt 7.99 min, 95.6%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM:MeOH (90:10), Isocratic:20% B; Wavelength: 250 nm, Flow: 1.0 mL/min.

Example 208. Synthesis of 258



Synthesis of N'-hydroxy-2-(hydroxymethyl)pyridine-4-carboxamidine (A-421c)

[000903] To a stirred solution of **A-410b** (0.5 g, 3.73 mmol) in ethanol (10 mL) was added hydroxylamine hydrochloride (388.53 mg, 5.59 mmol), TEA (1.04 mL, 7.45 mmol) and stirred for 6 h at 80 °C. The reaction mixture was quenched with water (10 mL) and diluted with EtOAc (100 mL x 2). The organic layer was separated, dried over Na_2SO_4, filtered and evaporated to afford the crude product which was purified by column chromatography using 100-200 silica at 30-80% EtOAc/Hexane as an eluent to afford **A-421c** (0.5 g, 1.49 mmol, 40% yield).

Synthesis of tert-butyl N-[(1S)-1-[3-[2-(hydroxymethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-421e)

[000904] To a stirred solution of **A-421c** (0.5 g, 2.99 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (565.92 mg, 2.99 mmol) in 1,4-dioxane (10 mL) was added DCC (616.14 mg, 2.99 mmol) and the reaction was stirred for 12 h at 100 °C. The reaction mixture was quenched with water (10 mL) and diluted with EtOAc (100 mL x 2).

The combined organic layer was separated, dried over Na_2SO_4, filtered, evaporated under reduced pressure and purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **A-421e** (0.4 g, 1.07 mmol, 36% yield) as a solid.


Synthesis of tert-butyl N-[(1S)-1-[3-[2-(hydroxymethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-421f) hydrochloride salt

[000905] To a stirred solution of **A-421e** (0.4 g, 1.25 mmol) in 1,4-dioxane (5 mL) was added 4 M HCl in 1,4 dioxane (6 mL, 1.25 mmol) and stirred at 0 °C. for 2 h. The reaction mixture was evaporated to afford the crude product which was washed using diethyl ether to afford **A-421f** (0.3 g, 1.05 mmol, 84% yield).

Synthesis of N-[(1S)-1-[3-[2-(hydroxymethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (258)

[000906] To a stirred solution of **A-421f** (0.3 g, 1.17 mmol) and 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (249.55 mg, 1.29 mmol) in DCM (10 mL) was added HATU (666.58 mg, 1.75 mmol) and DIPEA (0.41 mL, 2.34 mmol) and stirred for 2 h at RT. The reaction mixture was quenched with water (100 mL) and DCM (100 mL x 2) was added. The organic layer was separated, dried over Na_2SO_4, then filtered and evaporated to afford the crude product. Purification by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent afforded **258** (15 mg, 0.037 mmol, 3% yield). **HPLC**: Rt 7.20 min, 99.6%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** :397.05 (M+H), Rt 1.70 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.48 (d, 1H), 8.70 (d, 1H), 8.04 (s, 1H), 7.82-7.78 (m, 1H), 7.45 (s, 1H), 5.65-5.55 (m, 1H), 5.51-5.46 (m, 1H), 4.66 (s, 2H), 4.13 (s, 3H), 1.69 (d, 3H). **Chiral method:** Rt 5.38 min, 100%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:50% B; Wavelength: 230 nm, Flow: 1.0 mL/min.

Example 209. Synthesis of 259



Synthesis of 2-(isopropoxymethyl)pyridine-4-carbonitrile (A-422d)

[000907] To a stirred solution of 2-propanol (1.13 g, 18.77 mmol) in THF (20 mL) was added NaH (0.68 g, 28.16 mmol) at 0-5 °C and stirred for 30 min at 0-5 °C. To the resulting reaction was added **A-418c** (1.85 g, 9.39 mmol) in THF (20 mL) in a dropwise manner. The reaction mixture was stirred for 5 h at RT. The reaction mixture was quenched with ice cold water and extracted with ethyl acetate (2 x 20 mL). The combined organic layer was dried over sodium sulphate and concentrated. The crude product was purified by column chromatography, using 10-15% of ethyl acetate in hexane as an eluent to afford **A-422d** (830 mg, 4.39 mmol, 47% yield).

Synthesis of N'-hydroxy-2-(isopropoxymethyl)pyridine-4-carboxamidine (A-422e)

[000908] To a stirred solution of **A-422d** (800 mg, 4.54 mmol) in ethanol (8 mL) was added TEA (1.27 mL, 9.08 mmol) and hydroxylamine hydrochloride (473.2 mg, 6.81 mmol) and stirred for 3 h at reflux. The reaction mixture was concentrated under reduced pressure and diluted with ethyl acetate (20 mL). The organic layer was washed with water (10mL x 2) and the organic layer was evaporated to afford **A-422e** (800 mg, 3.77 mmol, 83% yield).

Synthesis of tert-butyl (S)-(1-(3-(2-(cyclopropoxymethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-422f)

[000909] To a stirred solution of **A-422e** (700 mg, 3.35 mmol) in 1,4-dioxane (7 mL) was added N,N'-dicyclohexylcarbodiimide (DCC) (759.29 mg, 3.68 mmol) and N-Boc-L-alanine (696.29 mg, 3.68 mmol) at RT and the reaction mixture was stirred for 4 h at reflux. The reaction mixture was filtered and diluted with ethyl acetate and washed with water (3 x 10 mL). The organic layer was dried over sodium sulphate and concentrated to afford the crude product which was purified by column chromatography. using 25-30% of ethyl acetated in hexane as eluent to afford **A-422f** (700 mg, 1.84 mmol, 55% yield).

Synthesis of (1S)-1-[3-[2-(isopropoxymethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (A-422g)


[000910] To a stirred solution of **A-422f** (0.7g, 1.93mmol) in 1,4-dioxane (5 mL) was added 4 M HCl/dioxane (5 mL) and stirred for 2 h at RT. The reaction mixture was evaporated under reduced pressure to afford **A-422g** (450 mg, 1.48 mmol, 77% yield) as a hydrochloride salt.

Synthesis of N-[(1S)-1-[3-[2-(isopropoxymethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (259)

[000911] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (116.95 mg, 0.6 mmol) in DCM (2 mL) was added DIPEA (0.09 mL, 0.5 mmol), HATU (286.35mg, 0.7500 mmol) and **A-422g** (150.mg, 0.5 mmol) and stirred for 2 h at RT. The

reaction mixture was diluted with DCM and washed with water (2 x 5 mL). The organic layer was dried over sodium sulphate and concentrated to afford the crude product which was purified by column chromatography using at 30% ethyl acetate in hexane as an eluent to afford **259** (75 mg, 0.16 mmol, 33% yield) as a solid. **HPLC**: Rt 8.73 min, 96.3 %; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** :439.25 (M+H), Rt 2.02 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.72 (d, 1H), 7.96 (s, 1H), 7.84-7.80 (m, 1H), 7.44 (s, 1H), 5.52-5.45 (m, 1H), 4.64 (s, 2H), 4.13 (s, 3H), 3.80-3.72 (m, 1H), 1.68 (d, 3H), 1.18 (d, 6H). **Chiral method:** Rt 8.94 min, 100%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (90:10), Isocratic:20% B; Wavelength: 280 nm, Flow: 1.0 mL/min.

Example 210. Synthesis of 260 and 261



Synthesis of N-[(1S)-1-[3-[2-(bromomethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (A-423a)

[000912] To a stirred solution of **258** (600 mg, 1.51 mmol) in DCM (3 mL) was added CBr_4 (1 g, 3.03 mmol) and PPh_3 (794.18 mg, 3.03 mmol) in portions at 0 °C and stirred at 0 °C for 30 min. The reaction mixture was evaporated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 30%EtOAc:hexane as an eluent to afford **A-423a** (250 mg, 0.45 mmol, 30% yield) as an oil.


Synthesis of (S)-1-methyl-N-(1-(3-(2-(pyrrolidin-1-ylmethyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (261)

[000913] To a stirred solution of **A-423a** (100 mg, 0.22 mmol) in THF (3 mL) was added pyrrolidine (0.04 mL, 0.44 mmol) and DIPEA (0.11 mL, 0.65 mmol) at RT and stirred for 2 h. The reaction mixture was diluted with water and extracted with ethylacetate and the organic layer was dried over Na_2SO_4, filtered and evaporated to afford crude product which was purified by prep HPLC to afford **261** (10mg, 0.02 mmol, 10% yield) as an oil. **HPLC**: Rt 5.90 min, 98.8%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 450.15 (M+H), Rt 1.39 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.69 (d, 1H), 8.16 (s, 1H), 7.97 (s, 1H), 7.81-7.78 (m, 1H), 7.44 (s, 1H), 5.50-5.45 (m, 1H), 4.13 (s, 3H), 3.82 (s, 2H), 1.75-1.65 (m, 6H), 4H merged in solvent peak. **Chiral method:** Rt 5.79 min, 99.6%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) EtOH, Isocratic:15% B; Wavelength: 278 nm, Flow: 1.0 mL/min.

Synthesis of (S)-N-(1-(3-(2-((dimethylamino)methyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (260)

[000914] To a stirred solution **A-423a** (100 mg, 0.2200 mmol) in THF (3mL) was added dimethylamine (0.03 mL, 0.65 mmol) and stirred at RT for 1 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and concentrated to afford the crude product which was purified using prep HPLC to afford **260** (6 mg, 0.014 mmol, 6% yield) as an oil. **HPLC**: Rt 5.76 min, 96.2 %; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** :424.10 (M+H), Rt 1.40 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.69 (d, 1H), 7.99 (s, 1H), 7.82-7.80 (m, 1H), 7.45 (s, 1H), 5.50-5.45 (m, 1H), 4.13 (s, 3H), 3.62 (s, 2H), 2.22 (s, 6H), 1.68 (d, 3H). **Chiral method:** Rt 5.83 min, 99.6%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) EtOH, Isocratic:15% B; Wavelength: 278 nm, Flow: 1.0 mL/min.

Example 211. Synthesis of 262 and 263



Synthesis of 2-(1-methoxyvinyl)isonicotinonitrile (A-424b)

[000915] To a stirred solution of **A-424a** (1.2 g, 6.56 mmol) in ACN (10 mL) was added tributyl(1-methoxyvinyl)stannane (1.8 g, 5.25 mmol), L-Proline (20.83 mg, 0.18 mmol) and sodium hydroxide (7.24 mg, 0.18 mmol) and stirred at 90 °C for 16 h. The reaction mixture was quenched with water (100 mL) and diluted with EtOAc (50 mL). The organic layer was separated, and dried with Na_2SO_4, filtered and evaporated to afford crude reaction product. The crude product was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **A-424b** (1g, 3.12 mmol, 48% yield) as a solid.

Synthesis of 2-(1-methoxycyclopropyl)isonicotinonitrile (A-424c)

[000916] To a solution of diethyl zinc (1.5 g, 12.46 mmol) in DCM was added di-iodo methane (1.01 mL, 12.49 mmol) and stirred for 1 hr 0 °C. To the resulting reaction mixture was added **A-424b** (0.5g, 3.12 mmol) at 0 °C and stirred at RT for 2 h. The reaction mixture was quenched with water and diluted with ethyl acetate. The organic layer was separated and dried over Na_2SO_4 and evaporated to afford the crude product which was purified by column chromatography using 100-200 silica and 10-30% EtOAc/hexane as an eluent to afford **A-424c** (300 mg, 0.86 mmol, 28% yield).

Synthesis of N-hydroxy-2-(1-methoxycyclopropyl)isonicotinimidamide (A-424d)

[000917] To a mixture of **A-424c** (300 mg, 1.72 mmol) and hydroxylamine hydrochloride (143.61 mg, 2.07 mmol) in ethanol (5 mL) was added TEA (0.48 mL, 3.44 mmol) and heated at 80 °C for 16 h. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over sodium sulphate and evaporated to afford the crude product which was further purified by column chromatography using 100-200 silica and 20-50%EtOAc/hexane as an eluent to afford **A-424d** (100 mg, 0.24 mmol, 14% yield).

Synthesis of tert-butyl (S)-(1-(3-(2-(1-methoxycyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (262)

[000918] To a mixture of **A-424d** (0.1 g, 0.48 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (109.6 mg, 0.58 mmol) in 1,4-dioxane (10 mL) was added DCC (198.8 mg, 0.97 mmol) and heated at 100 °C for 16 h. The reaction mixture was diluted with water and extracted with ethyl acetate. The combined organic layer was washed with water and brine, dried over sodium sulphate and evaporated to afford the crude product which was purified by combi-flash chromatography using ethyl acetate: hexane as an eluent, to afford 200 mg crude product which was further purified by prep HPLC to afford **262** as a solid (10 mg, 0.027 mmol, 6% yield). **HPLC**: Rt 8.96 min, 99.8%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water, B: ACN (95:5); Flow Rate: 1 mL/min. **LCMS** : 360.95 (M+H), Rt 2.05 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 8.65 (d, 1H), 7.82-7.79 (m, 2H), 7.67 (d, 1H), 5.05-4.95 (m, 1H), 1.54-1.51 (m, 6H), 1.40-1.36 (m, 9H), 1.24-1.22 (m, 2H), 0.89-0.87 (m, 2H). **Chiral method:** Rt 5.36 min, 100%; SFC column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-hexane + 0.1 % isopropyl amine B) DCM : MeOH (1:1), Isocratic:50% B. Wavelength: 290 nm, Flow: 3 mL/min.

Synthesis of (S)-1-(3-(2-(1-methoxycyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine hydrochloride (A-424f)


[000919] To a stirred solution of **262** (60 mg, 0.17 mmol) in 1,4-dioxane (10 mL) was added 4 M HCl in 1,4-dioxane (10 mL) at 0 °C and stirred at RT for 6 h. The reaction mixture was concentrated to afford **A-424f** (40 mg, 0.14 mmol, 83% yield) as a solid.

Synthesis of (S)-N-(1-(3-(2-(1-methoxycyclopropyl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (263)

[000920] To a stirred solution 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (31.4 g, 0.16 mmol) in DCM (5 mL) was added **A-424f** (31.6 g, 0.16 mmol), HATU (77 mg, 0.2 mmol), DIPEA (0.05 mL, 0.27 mmol) in water (1 mL) and stirred at RT 2 h. The reaction mixture was diluted with water nd DCM. The organic layer was separated, washed with saturated brine solution (20 mL), separated and dried over $MgSO_4$, The solution was evaporated to dryness and the crude product was then purified using prep HPLC to afford **263** (10 mg, 0.023 mmol, 17% yield) as a solid. **HPLC**: Rt 4.71 min, 99 %; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 437.20 (M+H), Rt 2.06 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.61 (d, 1H), 7.87 (s, 1H), 7.69-7.66 (m, 1H), 7.45 (s, 1H), 5.50-5.45 (m, 1H), 4.13 (s, 3H), 1.68 (d, 3H), 1.04-0.97 (m, 4H), 3H merged in solvent peak. **Chiral method:** Rt 6.92 min, 99.66%;

column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% TFA B) DCM: MeOH (50:50), Isocratic:25% B; Wavelength: 254 nm, Flow: 1.0 mL/min.

Example 212. Synthesis of 264



Synthesis of 2-((methylsulfonyl)methyl)isonicotinonitrile (A-425b)

[000921] To a stirred solution of **A-418c** (1 mg, 0.01 mmol) in DMF (10mL) was added sodium methanesulfinate (0.78 mg, 0.01 mmol) and TBAI (0.37mg) at RT and stirred for 7h. The reaction mixture was quenched with ice cold water and extracted with ethyl acetate and the organic layer separated. The organic layer was washed with water (1 x 5 mL), dried over sodium sulphate and evaporated to afford **A-425b** (0.55 mg, 0.0026 mmol, 51% yield).

Synthesis of (Z)-N'-hydroxy-2-((methylsulfonyl)methyl)isonicotinimidamide (A-425c)

[000922] To a stirred solution of **A-425b** (350 mg, 1.78 mmol) in ethanol (4 mL) was added TEA (0.5 mL, 3.57 mmol) and hydroxylamine hydrochloride (185.92 mg, 2.68 mmol) at RT and the reaction mixture was stirred for 3 h at reflux. The reaction mixture was evaporated and the residue diluted with ethyl acetate (20 mL). The organic layer was washed with water (2 x 10 mL), brine, dried over sodium sulphate and evaporated to afford **A-425c** (550 mg, 1.33 mmol, 75% yield) as a solid.

Synthesis of tert-butyl (S)-(1-(3-(6-((methylsulfonyl)methyl)pyridazin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)carbamate (A-425e)

[000923] To a stirred solution of **A-425c** (0.5 g, 1.21 mmol) in 1,4-dioxane (5 mL) was added N,N'-dicyclohexylcarbodiimide (DCC) (0.27 g, 1.33 mmol) and *N*-Boc-L-alanine (0.25 g, 1.33 mmol) at RT and stirred for 12 h at reflux. The reaction mixture was filtered then diluted with ethyl acetate and washed with water (3 x 5 mL). The organic layer was dried over sodium sulphate and evaporated to afford the crude product which was purified by column chromatography using silica gel and 25-30% of ethyl acetated in hexane as an eluent to afford **A-425e** (165 mg, 0.35 mmol, 29% yield) as a solid.


Synthesis of (1R)-1-[3-[2-(methylsulfonylmethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (A-425f)

[000924] To a stirred solution of **A-425e** (150 mg, 0.39 mmol) in 1,4-dioxane (3 mL) was added 4 M HCl in dioxane (3 mL) at RT and stirred for 2h. The reaction mixture was evaporated under reduced pressure to afford **A-425f** (165 mg, 0.37 mmol, 97% yield).

Synthesis of 2-methyl-N-[(1S)-1-[3-[2-(methylsulfonylmethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (264)

[000925] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (95 mg, 0.49 mmol) in DCM (2 mL) was added DIPEA (0.39 mL, 2.25 mmol) and HATU (232.6 mg, 0.6100 mmol) and stirred for 15 min at RT. To the resulting reaction mixture was added **A-425f** (130 mg, 0.41 mmol) and the reaction was stirred for 3 h. The reaction mixture was diluted with DCM. The organic layer was washed with water (3 x 5 mL), dried over sodium sulphate and evaporated to afford a crude product which was purified using column chromatography to afford **264** (25 mg, 0.05 mmol, 13% yield) as a solid. **HPLC**: Rt 7.76 min, 99.8%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 459.85 (M+H), Rt 1.71 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.50-9.47 (m, 1H), 8.81 (d, 1H), 8.10 (s, 1H), 7.96-7.94 (m, 1H), 7.45 (s, 1H), 5.50-5.46 (m, 1H), 4.82 (s, 2H), 4.13 (s, 3H), 3.05 (s, 3H), 1.68 (d, 3H). **Chiral method:** Rt 6.06 min, 99.4%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso propyl amine B) EtOH:MeOH (50:50), Isocratic:25% B; Wavelength: 279 nm, Flow: 1.0 mL/min.

Example 213. Synthesis of 265



Synthesis of tert-butyl N-[(1R)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-426c)

[000926] To a stirred solution of **A-301** (0.5g, 2.82 mmol) in 1,4-dioxane (5 mL) was added N,N'-dicyclohexylcarbodiimide (DCC) (0.64 g, 3.1 mmol) and *N*-Boc-D-alanine (0.59 g, 3.1 mmol) at RT and stirred for 12 h at reflux. The reaction mixture was filtered, diluted with ethyl acetate, washed with water (3 x 5 mL) and separated. The organic layer was dried over sodium sulphate and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using silica and 25-30% of ethyl acetated in hexane as an eluent to afford **A-426c** (0.63 g, 1.79 mmol, 63% yield) as a solid.


Synthesis of (R)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine hydrochloride (A-426d)

[000927] To the stirred solution of **A-426c** (500 mg, 1.51 mmol) in 1,4-dioxane (5 mL) was added 4 M HCl/dioxane (3 mL) at RT and stirred for 2 h. The reaction mixture was evaporated under to afford **A-426d** (350 mg, 0.81 mmol, 53% yield).

Synthesis of N-[(1R)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (265)

[000928] To a stirred solution of 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (174.66 mg, 0.9 mmol) in DCM (2 mL) was added DIPEA (0.39 mL, 2.25 mmol) and HATU (427.66 mg, 1.12 mmol) and stirred for 15 min at RT. To the resulting reaction was added **A-426d** (200 mg, 0.75 mmol) and stirred for 3 h at RT. The reaction was diluted with DCM and the organic layer was washed with water (3 x 5 mL), dried over sodium sulphate and evaporated to afford the crude product which was purified using column chromatography to afford **265** (50 mg, 0.13 mmol, 16% yield) as a solid. **HPLC**: Rt 9.28 min, 99.9%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 10 mM Ammonium Bicarbonate in water; B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 407.06 (M+H), Rt 2.09 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.47 (d, 1H), 8.61 (d, 1H), 7.87 (s, 1H), 7.68-7.66 (m, 1H), 7.45 (s, 1H), 5.50-5.45 (m, 1H), 4.13 (s, 3H), 2.34-2.28 (m, 1H), 1.68 (d, 3H), 1.05-0.98 (m, 4H). **Chiral method:** Rt 9.37 min, 99.7%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% TFA B) Iso propyl Alcohol, Isocratic:10% B; Wavelength: 210-240 nm, Flow: 1.0 mL/min.

Example 214. Synthesis of 266



Synthesis of tert-butyl N-[(1R)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]propyl]carbamate (A-427d)

[000929] To a stirred solution of **A-301** (200 mg, 1.13 mmol) in 1,4-dioxane (4 mL) was added (2R)-2-(tert-butoxycarbonylamino)butanoic acid (275.27 mg, 1.35 mmol) and DCC (279.01 mg, 1.35 mmol) and the reaction mixture was heated at 100 °C for 16 h . The reaction mixture was cooled then diluted with water and extracted with EtOAc. The organic was dried over Na_2SO_4, filtered and evaporated to afford the crude product which was purified by combi flash chromatography using 100-200 mesh silica and 6% EtOAc/hexane as an eluent to afford **A-427d** (210 mg, 0.58 mmol, 51% yield) as an oil.

Synthesis of (1R)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]propan-1-amine hydrochloride (A-427e)


[000930] To a stirred solution of **A-427d** (200 mg, 0.58 mmol) in 1,4-dioxane (2 mL) was added 4 M HCl in 1,4 dioxane (1.12 mL, 2.9mmol) in a dropwise manner. The reaction mixture was stirred at RT for 16 h then the reaction mixture was evaporated to afford **A-427e** (120 mg, 0.38 mmol, 66% yield) as a solid.

Synthesis of N-[(1R)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]propyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (266)

[000931] To a stirred solution of **A-427e** (150 mg, 0.53 mmol) in DCM (5 mL) was added 2-methyl-5-(trifluoromethyl)pyrazole-3-carboxylic acid (103.71 mg, 0.53 mmol), DIPEA (0.19 mL, 1.07 mmol) and HATU (203.15 mg, 0.5300 mmol) and stirred at RT for 16 h. The reaction mixture was diluted with water and extracted with EtOAC. The organic layer was dried over Na_2SO_4 and evaporated to afford the crude product which was purified by column chromatography using 100-200 mesh silica and 20% EtOAc/hexane as an eluent to afford **266** (40 mg, 0.092 mmol, 17% yield) as an oil. **HPLC**: Rt 8.99 min, 96.3%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 421.4 (M+H), Rt 2.09 min,

Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.40 (d, 1H), 8.60 (d, 1H), 7.85 (s, 1H), 7.68-7.62 (m, 1H), 7.49 (s, 1H), 5.34-5.28 (m, 1H), 4.13 (s, 3H), 2.30-2.25 (m, 1H), 2.10-2.02 (m, 2H), 1.10-0.95 (m, 7H). **Chiral method:** Rt 5.25 min, 99.1%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso propyl amine B) DCM:MeOH (50:50), Isocratic:20% B; Wavelength: 295 nm, Flow: 1.0 mL/min.

Example 215. Synthesis of 267, 268, and 269



Synthesis of (S)-N-(1-(3-(2-bromopyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (A-428c)

[000932] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (1.27 g, 6.55 mmol) in DCM (10 mL) was added **A-428a** (2 g, 6.55 mmol), DIPEA (1.14 mL, 6.55 mmol) and HATU (2 .5 g, 6.55 mmol) at RT and stirred for 16 h. The reaction mixture was diluted with EtOAc (2 x 100 mL) and washed with water (100 mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by combi flash column chromatography and 10-15% EtOAc in hexane to afford **A-428c** (0.9 g, 1.88 mmol, 29% yield) as a solid.

Synthesis of (S)-N-(1-(3-(2-cyanopyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (267)

[000933] To a stirred solution of **A-428c** (200 mg, 0.4 mmol) in DMF (3mL) was added $Zn(CN)_2$ (158.22 mg, 1.35 mmol) and purged with N_2 gas for 20 min. To the solution was then added $Pd_2(dba)_3$ (4.12 mg, 0.0045 mmol) and dppf (2.49 mg, 0.0045 mmol). The reaction mixture was stirred at 100 °C for 16 h. The reaction mixture was diluted with EtOAc and washed with water. The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by combi flash column chromatography and 10-15% EtOAc in hexane to afford **267** (18 mg, 0.046 mmol, 10% yield) as a solid. **HPLC**: Rt 8.31 min, 99.7%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 391.8 (M+H), Rt 1.96 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.49 (d, 1H), 8.97 (d, 1H), 8.53 (s, 1H), 8.28-8.24 (m, 1H), 7.45 (s, 1H), 5.52-5.45 (m, 1H), 4.12 (s, 3H), 1.68 (d, 3H). **Chiral method:** Rt 8.72 min, 84.2%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 220 nm, Flow: 1.0 mL/min.

Synthesis of (S)-1-methyl-3-(trifluoromethyl)-N-(1-(3-(2-vinylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (A-428e)

[000934] To a stirred solution of **A-428c** (300 mg, 0.67 mmol) in 1,4-dioxane (4 mL):water (3 mL) was added Na_2CO_3 (214.27 mg, 2.02 mmol) and the reaction mixture was purged with N_2 gas for 20 min. To this solution was added **A-428d** (207.57 mg, 1.35 mmol) and tetrakis(triphenylphosphine)palladium (77.87 mg, 0.07 mmol). The reaction mixture was stirred at 100 °C for 3 h. The reaction mixture was cooled then diluted with DCM and washed with water. The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by flash column chromatography using 100-200 mesh silica and 10-15% EtOAc in hexane to afford **A-428e** (120 mg, 0.24 mmol, 36% yield).

Synthesis of (S)-N-(1-(3-(2-ethylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (268)

[000935] To a stirred solution of **A-428e** (100 mg, 0.25 mmol) in methanol (4 mL) was added 10% Pd/C (27.12 mg, 0.25 mmol). The reaction mixture was stirred at RT under H_2 atmosphere for 30 min. The reaction was filtered through a celite pad and the filtrate was concentrated under reduced pressure. The crude compound was purified by combi flash column chromatography and 20-25% EtOAc in hexane to afford **268** (17 mg, 0.042 mmol, 16% yield) as a solid. **HPLC**: Rt 7.96 min, 97.5%; Column: X-Select CSH C18 (4.6 x 150)

mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 395 (M+H), Rt 1.90 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.68 (d, 1H), 7.80 (s, 1H), 7.75-7.72 (m, 1H), 7.45 (s, 1H), 5.50-5.44 (m, 1H), 4.13 (s, 3H), 2.86 (q, 2H), 1.68 (d, 3H), 1.26 (t, 3H). **Chiral method:** Rt 5.81 min, 100%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 227 nm, Flow: 1.0 mL/min.


Synthesis of (S)-1-methyl-N-(1-(3-(2-(prop-1-en-2-yl)pyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (A-428g):

[000936] To a stirred solution of **A-428c** (302.69 mg, 0.68 mmol) in 1,4-dioxane (4 mL) and water (2 mL) was added Na_2CO_3 (216.19 mg, 2.04 mmol) and the reaction mixture was purged with N_2 gas for 20 min. Then $Pd(PPh_3)_4$ (7.86 mg, 0.01 mmol) and **A-428f** (228.51 mg, 1.36 mmol) were added. The reaction mixture was stirred at 100 °C for 3 h. The reaction mixture was diluted with EtOAc and washed with water. The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by combi flash column chromatography and 10-15% EtOAc in hexane to afford **A-428g** (18 mg, 0.046 mmol, 11% yield) as a solid.

Synthesis of (S)-N-(1-(3-(2-isopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (269)

[000937] To a stirred solution of **A-428g** (144.12 mg, 0.35 mmol) in EtOAc (5mL) was added 10% Pd/C (37.74 mg, 0.35 mmol). The reaction mixture was stirred under hydrogen atmosphere for 30 min . The reaction was filtered through a celite pad and the filtrate was concentrated under reduced pressure. The crude compound was purified by combi flash column chromatography and 20-25% EtOAc in hexane to afford **269** (13 mg, 0.031 mmol, 9% yield) as a solid. **HPLC**: Rt 8.68 min, 95.94%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water:ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 409.4 (M+H), Rt 2.05 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.68 (d, 1H), 7.79 (s, 1H), 7.75-7.72 (m, 1H), 7.45 (s, 1H), 5.50-5.44 (m, 1H), 4.12 (s, 3H), 3.20-3.11 (m, 1H), 1.68 (d, 3H), 1.27 (d, 6H). **Chiral method:** Rt 5.08 min, 100%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 225 nm, Flow: 1.0 mL/min.


Example 216. Synthesis of 270



A-303 270

[000938] To a stirred solution of **A-303** (100.mg, 0.43 mmol) and 1-cyclopentyl-1H-pyrazole-5-carboxylic acid (93.91 mg, 0.52 mmol) in DCM (10 mL) were added HATU (247.69 mg, 0.65 mmol) and DIPEA (0.15 mL, 0.87 mmol) at RT and stirred for 2 h at RT. The reaction mixture was quenched with water (100 mL) diluted with DCM (100 mL x 2) and the organic layer separated. The organic layer was dried over Na_2SO_4, filtered and evaporated. The reaction mixture was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **270** (15 mg, 0.037 mmol, 8% yield). **HPLC**: Rt 8.041 min, 97.7%; Column: X-Select CSH C18 (4.6 x 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 393.10 (M+H), Rt 2.083 min; Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.23 (d, 1H), 8.60 (d, 1H), 7.85 (s, 1H), 7.66-7.64 (m, 1H), 7.52 (d, 1H), 6.90 (d, 1H), 5.58-5.50 (m, 1H), 5.46-5.42 (m, 1H), 2.31-2.24 (m, 1H), 2.05-1.85 (m, 4H), 1.80-1.78 (m, 2H), 1.67 (d, 3H), 1.60-1.57 (m, 2H), 1.03-0.94 (m, 4H). **Chiral method:** Rt 11.51 min, 98.4%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20%B; Wavelength: 292 nm, Flow: 1.0 mL/min.

Example 217. Synthesis of 271



A-430aa A-430bb A-430d

A-430a A-430b A-430c 271

Synthesis of (2S)-2-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoate (A-430bb)

[000939] To a stirred solution of **A-430aa** (1 g, 5.15 mmol) in DCM (2 mL) was added DIPEA (2.69 mL, 15.46 mmol) and HATU (2.94 g, 7.7 3 mmol) and stirred for 10 min. To

resulting reaction mixture was added H-Ala-OtBu HCl (0.94 g, 5.15 mmol) and stirred for 2 h at RT. The reaction mixture was diluted with DCM, washed with water and the organic layer separated. The organic layer was dried over sodium sulphate and evaporated to afford the crude product. The crude product was purified by column chromatography to afford **A-430bb** (1.5 g, 4.2073 mmol, 82% yield) as a solid.

Synthesis of (2S)-2-[[2-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]amino]propanoic acid (A-430d)

[000940] To a stirred solution of **A-430bb** (500 mg, 1.56 mmol) in DCM (10 mL) was added TFA (3 mL, 40.96 mmol) at 0-5 °C in dropwise manner and stirred at RT for 5 h. The reaction mixture was concentrated to afford the crude product as a liquid. The crude product was dissolved in ethyl acetate, washed with water, dried over sodium sulphate and concentrated to afford **A-430d** (400 mg, 1.39 mmol, 89% yield) as an oil.

Synthesis of 2-cyclobutylpyridine-4-carbonitrile (A-430b)

[000941] To a stirred solution of **A-430a** (500 mg, 2.36 mmol) in DMF (4 mL) was added zinc cynide (827.48 mg, 7.07 mmol) and purged with N_2 gas for 10 min. To the resulting reaction mixture was added Pd_2dba_3 (215.95 mg, 0.24 mmol), dppf (130.69 mg, 0.24 mmol) and heated at 100 °C for 16 h. The reaction mixture was diluted with water and extracted with EtOAc and the organic layer separated. The organic layer was dried over Na_2SO_4 , filtered and evaporated to afford the crude product which was purified by combi-flash using 10% EA/hexane as an eluent to afford **A-430b** (90 mg, 0.57 mmol, 24% yield) as an oil.

Synthesis of 2-cyclobutyl-N'-hydroxy-pyridine-4-carboxamidine (A-430c)


[000942] To a stirred solution of **A-430b** (140 mg, 0.88 mmol) in ethanol (3 mL) was added hydroxylamine hydrochloride (184.49 mg, 2.65 mmol), TEA (0.37 mL, 2.65 mmol) and stirred at 100 °C for 16 h. The reaction mixture was diluted with water and extracted using EtOAc and the organic layer separated. The organic layer was dried over Na_2SO_4, filtered and evaporated to afford **A-430c** (160 mg, 0.84 mmol, 95% yield). The crude product was used in the next step without purification.

Synthesis of (S)-N-(1-(3-(2-cyclobutylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (271)

[000943] To a stirred solution of **A-430c** (150 mg, 0.78 mmol) and **A-430d** (208.01 mg, 0.78 mmol) in 1,4-dioxane (4 mL) was added DCC (242.38 mg, 1.18 mmol) and stirred at 100 °C for 16 h. The reaction mixture was diluted with water, extracted with EtOAc, and the organic layer separated. The organic layer was dried over Na_2SO_4 filtered and concentrated to

afford the crude product which was purified by column chromatography using 15% EA/hexane as an eluent to afford **271** (28 mg, 0.064 mmol, 8% yield). **HPLC**: Rt: 8.88 min, 96.5%; Column: X-Select CSH C18 (4.6 x 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 420.9 (M+H), Rt 2.101 min, Column: X-select CSH C18 (3 x 50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ_H = 9.46 (d, 1H), 8.74 (d, 1H), 7.76-7.72 (m, 2H), 7.45 (s, 1H), 5.50-5.43 (m, 1H), 4.13 (s, 3H), 3.80-3.72 (m, 1H), 2.33-2.27 (m, 4H), 2.06-1.96 (m, 1H), 1.88-1.66 (m, 1H), 1.67 (d, 3H). **Chiral method:** Rt 5.574 min, 97.6%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 284 nm, Flow: 1.0 mL/min. Note that under the reaction conditions racemization was observed

Example 218. (S)-*N*-(1-(3-(2-cyclobutylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-5-carboxamide (278) and (R)-*N*-(1-(3-(2-cyclobutylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-5-carboxamide (279). Note that stereochemistry has been randomly assigned




[000944] Racemic **271** (60.0 mg, 0.14 mmol) was separated by chiral HPLC to give the two pure enantiomers. The separation carried out 60 mg of **271** using prep-HPLC SFC giving **278** (Rt 9.046 min, 10 mg, 0.0238 mmol, 20% yield) as a solid and **279** (Rt 9.920 min, 10 mg, 0.0238 mmol, 20% yield) as a solid. Prep-HPLC conditions were: column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) EtOH: MeOH (50:50), Isocratic:10% B; Wavelength: 283 nm, Flow: 1.0 mL/min.

[000945] **278: LCMS** : 421.0 (M+H), Rt 2.087 min, Column: X-select CSH C18 (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.47 (d, 1H), 8.75 (d, 1H), 7.78 (s, 2H), 7.45 (s, 1H), 5.46-5.56 (m, 1H), 4.13 (s, 3H), 3.76-3.84 (m, 1H), 2.20-2.44 (m, 4H), 1.94-2.16 (m, 1H), 1.82-1.92 (m, 1H), 1.68 (d, 3H). **Chiral method:** Rt 9.046 min, 99.6%; column: DIACEL CHIRALPAK-IG (250 mm x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) EtOH: MeOH (50:50), Isocratic:10% B; Wavelength: 283 nm, Flow: 1.0 mL/min.

[000946] **279: HPLC**: Rt 8.892 min, 94.4%; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 421.3 (M+H), Rt 2.092 min, Column: X-select CSH C18 (3*50) mm, 2.5 µm; **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.47 (d, 1H), 8.75 (d, 1H), 7.80 (s, 2H), 7.45 (s, 1H), 5.46-5.56 (m, 1H), 4.13 (s, 3H), 3.68-3.86 (m, 1H), 2.26-2.38 (m, 4H), 1.94-2.19 (m, 1H), 1.82-1.92 (m, 1H), 1.68 (d, 3H). **Chiral method:** Rt 9.920 min, 99%; column: DIACEL CHIRALPAK-IG (250 mm x4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) EtOH: MeOH (50:50), Isocratic:10% B; Wavelength: 283 nm, Flow: 1.0 mL/min.


Example 219. (S)-1-benzyl-N-(1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1H-pyrazole-5-carboxamide (280)



A-303 HCl 280

[000947] To a stirred solution of **A-303** (201.5 mg, 0.76 mmol) and 1-benzyl-1H-pyrazole-5-carboxylic acid (203.67 mg, 0.91 mmol) in DCM (10 mL) was added HATU (430.86 mg, 1.13 mmol) and DIPEA (0.26 mL, 1.51 mmol) at RT. The reaction mixture was stirred at RT for 2 h. The reaction mixture was quenched with water (10 mL) and diluted with DCM (2 x 50mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to give the crude product. The crude compound was purified by column chromatography using 100-200 silica and 30-80% EtOAc/hexane as an eluent to give **280** (42 mg, 0.099 mmol, 13 % yield) as a solid. **HPLC**: Rt 8.358 min, 97.9%; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 415.6 (M+H), Rt 2.042 min, Column: X-select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.31 (d, 1H), 8.60 (d, 1H), 7.84 (s, 1H), 7.54-7.72 (m, 2H), 7.17-7.33 (m, 3H), 7.13 (d, 2H), 7.01 (s, 1H), 5.70-5.84 (m, 2H), 5.40-5.50 (m, 1H), 2.20-2.45 (m, 1H), 1.65 (d, 3H), 0.93-1.05 (m, 4H). **Chiral method:** Rt 5.372 min, 98%; column: DIACEL CHIRALPAK-IA (250 mm x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) EtOH: MeOH (1:1), Isocratic:40% B; Wavelength: 292 nm, Flow: 1.0 mL/min.


Example 220. Synthesis of 273



A-303 273

[000948] To a stirred solution of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (100 mg, 0.43 mmol) in DCM (5.0 mL) at 0 °C was added Et_3N (0.18 mL, 1.3 mmol) followed by benzyl carbonochloridate (111 mg, 0.65 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **273** (35 mg, 0.09 mmol, 21% yield) as a solid. Prep. HPLC method: Rt 12.6; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.37 min, 97.6%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 365.2 (M+H), Rt 2.38 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 5.0 min, SFC column: YMC Amylose-C; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.54 (d, 1H), 7.85 (s, 1H), 7.76-7.71 (m, 1H), 7.40-7.20 (m, 5H), 5.14 (m, 3H), 2.23-2.19 (m, 1H), 1.66 (d, 3H), 1.13-1.05 (m, 4H).


Example 221. Synthesis of 274



A-303 274

[000949] To a stirred solution of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (100 mg, 0.43 mmol) in DCM (5.0 mL) at 0 °C was added Et_3N (0.18 mL, 1.3 mmol) followed by ethyl carbonochloridate (70 mg, 0.65 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **274** (30 mg, 0.09 mmol, 22% yield) as a solid. Prep. HPLC method: Rt 10.5; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.89 min, 98.7%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate: 2.0 mL/min. **LCMS**: 303.1 (M+H), Rt 1.80 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 2.73 min, SFC column: YMC Amylose-C; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, DMSO-d$_6$):** δ 8.60 (d, 1H), 8.09 (d, 1H), 7.85 (s, 1H), 7.65 (d, 1H), 5.08-5.04 (m, 1H), 4.03 (q, 2H), 2.33-2.27 (m, 1H), 1.55 (d, 3H), 1.18 (t, 3H), 1.11-1.01 (m, 4H).

Example 222. Synthesis of 275




A-303 **275**

[000950] To a stirred solution of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (100 mg, 0.43 mmol) in DCM (5.0 mL) at 0 °C was added Et_3N (0.18 mL, 1.3 mmol) followed by isobutyl carbonochloridate (110 mg, 0.81 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **275** (40 mg, 0.11 mmol, 26% yield) as a solid. Prep. HPLC method: Rt 12.1; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile

phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 3.19 min, 96.6%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 331.2 (M+H), Rt 2.36 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 1.42 min, SFC column: YMC Amylose-SA; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 4.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.54 (d, 1H), 7.85 (s, 1H), 7.76-7.75 (m, 1H), 5.13-5.11 (m, 1H), 3.88 (d, 2H), 2.23-2.19 (m, 1H), 1.95-1.93 (m, 1H), 1.66 (d, 3H), 1.13-0.97 (m, 10H).


Example 223. Synthesis of 276



A-303 **276**

[000951] To a stirred solution of **A-432e** (100 mg, 0.43 mmol) in DCM (5.0 mL) at 0 °C was added Et_3N (0.18 mL, 1.3 mmol) followed by cyclopentyl carbonochloridate (0.08 mL, 0.65 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **276** (14 mg, 0.04 mmol, 9% yield) as a solid. Prep. HPLC method: Rt 10.1; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min.**HPLC**: Rt 5.66 min, 99.4%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 10 mM ammonium acetate, B: ACN; Flow Rate: 1.0 mL/min; **LCMS**: 343.2 (M+H), Rt 2.22 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 2.02 min, SFC column: Chiralcel OX-H; mobile phase: 70:30 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.54 (d, 1H), 7.85 (s, 1H), 7.76-7.75 (m, 1H), 5.12-5.07 (m, 2H), 2.23-2.19 (m, 1H), 1.87-1.31 (m, 11H), 1.12-1.06 (m, 4H).


Example 224. Synthesis of 277



A-303 277

[000952] To a stirred solution of (S)-1-(3-(2-cyclopropylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethan-1-amine (100 mg, 0.43 mmol) in DCM (5.0 mL) at 0 °C was added Et_3N (0.18 mL, 1.3 mmol) followed by isopropyl carbonochloridate (80 mg, 0.65 mmol). The reaction mixture was slowly warmed to room temperature and stirred for 4 h. The reaction mixture was treated with saturated $NaHCO_3$ solution (20 mL) and extracted with DCM (2 x 25 mL). The organic layer was washed with brine (20 mL), dried over Na_2SO_4 and concentrated. The crude was purified by preparative HPLC to afford **277** (15 mg, 0.04 mmol, 10% yield) as a solid. Prep. HPLC method: Rt 11.3; Column: X-Bridge (150 x 19 mm), 5.0 µm; Mobile phase: 0.1% TFA in water/acetonitrile; Flow Rate: 15.0 mL/min. **HPLC**: Rt 2.75 min, 94.5%; Column: X-Bridge C8 (50 x 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS**: 317.2 (M+H), Rt 2.15 min, Column: ZORBAX XDB C-18 (50 x 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate: 1.5 mL/min. **Chiral method:** Rt 2.2 min, SFC column: Chiral Pak AD-H; mobile phase: 60:40 (A: B), A = liquid CO_2, B = 0.5% isopropyl amine in methanol; flow rate: 3.0 mL/min; wave length: 210 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.54 (d, 1H), 7.85 (s, 1H), 7.76 (dd, 1H), 5.14-5.11 (m, 1H), 2.23-2.19 (m, 1H), 1.65 (d, 3H), 1.27 (d, 6H), 1.13-1.05 (m, 4H).


Example 225. Synthesis of 297



117 297

[000953] To a solution of 2-methyl-N-[(1S)-1-[3-(3-fluorophenyl)-1,2,4-oxadiazol-5-yl]ethyl]-5-(trifluoromethyl)pyrazole-3-carboxamide (800 mg, 2.09 mmol) in THF (10 mL) was added LiHMDS (2.3 mL, 2.3 mmol) at 0 °C. The mixture was stirred at 0 °C for 30 minutes and at 20 °C for 30 minutes. Then to the mixture was added CH_3I (1.39 mL, 22.33 mmol) at 0 °C. The mixture was stirred at 20 °C for 5 hours. The reaction mixture was quenched with sat. NH_4Cl (20 mL). The layers were separated, the aqueous phase extracted twice with EtOAc (15 mL x 2), the combined organic phases washed with brine (30 mL), and dried over Na_2SO_4 and concentrated to give the crude product. The crude product was purified by flash chromatography on silica gel (EtOAc in PE = 0% to 10% to 15%) to give the product (80 mg, 164.4 μmol, 7% yield) as an oil. **LCMS** R_t = 0.93 min in 1.5 min chromatography, 5-95AB, MS ESI calcd. $C_{17}H_{16}F_4N_5O_2$ $[M+H]^+$ 398.12, found 397.8. **Analytical SFC**: (column (S,S) Whelk-01 100 mm x 4.6 mm I.D., 5 μm, Mobile phase: A: CO_2 B: IPA (0.05% DEA), Gradient: from 5% to 40% of B in 5.5 min, then 5% of B for 1.5 min. Flow rate: 2.5 mL/min Column temp.: 40 °C, ABPR: 100 bar) showed two peaks at 2.82 min (22.5%) and 3.02 min (77.5%). The product was purified by SFC (REGIS (s,s) WHELK-O1 (250 mm x 30 mm, 5 μm); A = CO_2 and B = Neu-IPA ; 38 °C; 60 mL/min; 20% B; 10 min run; 10 injections, Rt of Peak 1 = 6.9 min, Rt of Peak 2 = 8.0 min) to give the product (30.52 mg, 75.8 μmol, 28% yield) (Rt = 3.02 min in analytical SFC) as an oil. **^{1}H NMR** (400MHz DMSO-d_6/D_2O, 80 °C) δ_H = 7.83 (br d, 1H), 7.70 (br d, 1H), 7.65 - 7.54 (m, 1H), 7.39 (dt, 1H), 6.98 (s, 1H), 5.80 - 5.60 (m, 1H), 3.92 (s, 3H), 3.01 (s, 3H), 1.74 (d, 3H). **LCMS** R_t = 1.32 min in 2.0 min chromatography, 10-80AB, MS ESI calcd. $C_{17}H_{16}F_4N_5O_2$ $[M+H]^+$ 398.1, found 398.1.

Example 226. Synthesis of 299




Synthesis of 5-methyl-1-(pyridin-3-yl)-1H-pyrazole-3-carboxylic acid (A-431)

[000954] To a stirred solution of ethyl 5-methyl-1-(pyridin-3-yl)-1H-pyrazole-3-carboxylate (30 mg, 0.13 mmol) in methanol (1.0 mL), THF (1.0 mL) and water (1.0 mL) was added lithium hydroxide monohydrate (11 mg, 0.26 mmol). The reaction mixture was stirred at room temperature for 2 h. The reaction mixture was concentrated under reduced pressure and treated with ice water (10 mL). The mixture was treated with 1N HCl (1.0 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with brine (15 mL), dried over Na_2SO_4 and concentrated to afford **A-431** (22 mg) as a solid. The crude compound was used for the next step without further purification.

Synthesis of (S)-5-methyl-N-(1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propyl)-1-(pyridin-3-yl)-1H-pyrazole-3-carboxamide (299)

[000955] To a stirred solution of **A-431** (22 mg, 0.11 mmol) in THF (3.0 mL) was added (S)-1-(3-(2-methylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)propan-1-amine (23 mg, 0.11 mmol) followed by Et_3N (0.03 mL, 0.21 mmol) and T3P (50% in EtOAc, 0.12 mL, 0.21 mmol). The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was treated with water (10 mL) and extracted with ethyl acetate (2 x 20 mL). The organic layer was washed with saturated sodium bicarbonate solution (10 mL), washed with brine (10 mL), dried over Na_2SO_4 and concentrated. The crude was purified by column chromatography on silica to afford **299** (3.6 mg, 0.009 mmol, 12% yield) as a solid. **HPLC:** Rt 2.27 min; Column: X-Bridge C8 (50 X 4.6) mm, 3.5 µm; Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 2.0 mL/min. **LCMS:** 404.2 (M+H), Rt 1.39 min, Column: ZORBAX XDB C-18 (50 X 4.6 mm), 3.5 µm; Mobile Phase: A: 0.1% HCOOH in water:ACN (95:5), B: ACN; Flow Rate:1.5 mL/min. **Chiral method:** Rt 2.37 min, SFC column: YMC Cellulose-SB; mobile phase: 60:40 (A: B), A = liquid CO2, B = methanol; flow rate: 3.0 mL/min; wavelength: 220 nm. **^{1}H NMR (400 MHz, CD_3OD):** δ 8.87 (d, 1H), 8.68 (dd, 1H), 8.59 (d, 1H), 8.15-8.11 (m, 1H), 7.95 (s, 1H), 7.86 (d, 1H), 7.66 (dd, 1H), 6.80 (s, 1H), 5.43-5.39 (m, 1H), 3.65 (m, 1H), 2.63 (s, 3H), 2.44 (s, 3H), 2.29-2.24 (m, 1H), 2.19-2.13 (m, 1H), 1.12 (t, 3H).

Example 227. Synthesis of 300



Synthesis of tert-butyl N-[[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]methyl]carbamate (A-432)

[000956] To a stirred solution of **A-362** (0.3 g, 1.46 mmol) in 1,4-Dioxane (10 mL) was added *N*-Boc-glycine (0.28 g, 1.61 mmol), DCC (0.33 g, 1.61 mmol) and the reaction mixture was stirred at 100° C for 16 h. The reaction mixture was evaporated under reduced pressure to afford a residue which was diluted using EtOAc (30 mL). The organic layer was washed with water (2 x 10 mL), saturated brine solution (10 mL), dried over $MgSO_4$ and evaporated to dryness to afford the crude product, which was purified by flash column chromatography using 20 % EtOAc in hexane as an eluent. The eluent fractions were evaporated to dryness under reduced pressure to afford 0.38 g of **A-432. Note**: 100 mg **A-432** was further purified by prep HPLC to afford 30 mg of **A-432** as a solid.

Synthesis of [3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]methanamine (A-433)


[000957] To a stirred solution of **A-432** (0.25 g, 0.73 mmol) in 1,4-dioxane (2 mL) was added 4M HCl in dioxane (5 mL) at 0 °C and stirred at RT 6 h. The reaction mixture was evaporated to afford **A-433** (180 mg, 0.5832 mmol, 80 % yield) as a solid.

Synthesis of 2-methyl-5-(trifluoromethyl)-N-[[3-[2-(trifluoromethyl)-4-pyridyl]-1,2,4-oxadiazol-5-yl]methyl]pyrazole-3-carboxamide (300)

[000958] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (0.11 g, 0.57 mmol) in DCM (10 mL) was added **A-433** (0.16 g, 0.57 mmol), HATU (258.57 mg, 0.68 mmol), and DIPEA (0.2 mL, 1.13 mmol) and stirred at RT 6 h. The

reaction mixture was diluted with water (10 mL) and DCM (10 mL) and organic layer separated. The organic layer was washed with saturated brine solution (10 mL), separated, dried over $MgSO_4$ and evaporated to dryness to afford crude product which was then purified by flash column chromatography using 30 % EtOAc in hexane as an eluent to afford the product which was again purified by prep HPLC to afford **300** (45 mg, 0.1025 mmol, 18 % yield) as a solid. **HPLC**: Rt 8.858 min, 95.7%; Column: X-Select CSH C18 (4.6 X 150) mm, 5 µm; Mobile phase: A: 0.1% formic acid in water:ACN (95:05) , B: ACN; Flow Rate: 1.0 mL/min; **LCMS** : 421.15 (M+H), Rt 2.024 min, Column: X-Select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.68 (s, 1H), 9.02 (d, 1H), 8.29-8.27 (m, 2H), 7.43 (s, 1H), 5.0-4.9 (m, 2H), 4.13 (s, 3H).

Example 228. Synthesis N-[[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (301) and Synthesis of N-[[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methyl]-N,2-dimethyl-5-(trifluoromethyl)pyrazole-3-carboxamide (302)



Synthesis of tert-butyl N-[[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methyl]carbamate (A-434)

[000959] To a solution of **A-301** (400 mg, 2.26 mmol) and *N*-Boc-glycine (434.98 mg, 2.48 mmol) in 1,4-dioxane (10 mL) was added N,N'-dicyclohexylcarbodiimide (512.33 mg, 2.48 mmol) at RT. The reaction mixture was heated at 100 °C for 16 h. The reaction mixture was concentrated under reduced pressure to afford the crude product. The crude product was diluted with water and extracted with EtOAc (3 x 25 mL). The combined organic phase was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to

afford the crude product. The residue was further purified by column chromatography (100-200 mesh silica, 22-25 % of EtOAc in hexane as an eluent) to afford **A-434** (350 mg, 1.09 mmol, 48 % yield) as an oil.

Synthesis of [3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methanamine (A-435)

[000960] To a solution of **A-434** (350.mg, 1.11 mmol) in 1,4-dioxane (5 mL) was added hydrogen chloride solution 4.0 M in dioxane (0.78 mL, 22.13 mmol) dropwise at 0 °C. The reaction mixture was allowed stirred at RT for 3 h. The volatile solvent was removed under reduced pressure to afford the crude product. The crude product was washed with hexane (3 x 5 mL) and dried under reduced pressure to afford afford **A-435 (**290 mg, 1.09 mmol, 98 % yield) as a solid.

Synthesis N-[[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (301)


[000961] To a stirred solution of 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (180.mg, 0.93 mmol) and **A-435** (281.2 mg, 1.11 mmol) in DCM (15 mL) was added HATU (528.89 mg, 1.39 mmol) and DIPEA (0.48 mL, 2.78 mmol) at 0 °C and reaction mixture was stirred at RT for 3 h. The reaction mixture was diluted with DCM (25 mL) and washed with water (3 x 20 mL), saturated sodium bicarbonate solution (3 x 20 mL) followed by brine solution (25 mL). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford the crude product, which was purified by column chromatography (22-25 % EtOAc in hexane as an eluent) followed by prep HPLC purification to afford **301** (15 mg, 0.0382 mmol, 4% yield) as a solid. **HPLC**: Rt 8.911 min; Column: X-Select CSH C18 (4.6 X 150) mm, 5 µm; Mobile phase: A: 10 mM ammonium bicarbonate in water, B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 392.9 (M+H), Rt 1.997 min; Column: X-Select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.65 (s, 1H), 8.59 (d, 1H), 7.85 (s, 1H), 7.65-7.64 (m, 1H), 7.42 (s, 1H), 4.86 (s, 1H), 4.13 (s, 3H), 2.3-2.24 (m, 1H), 1.02-0.94 (m, 4H)

Synthesis of N-[[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]methyl]-N,2-dimethyl-5-(trifluoromethyl)pyrazole-3-carboxamide (302)

[000962] To a suspension of sodium hydride (1.83 mg, 0.08 mmol) in THF (1 mL) was added **301** (25 mg, 0.06 mmol) in THF (1 mL) at 0 °C and stirred for 10 min. at 0 °C. To the resulting reaction mixture was added iodomethane (5 mL, 0.08 mmol) and reaction mixture

was stirred at RT for 2 h. Then reaction mixture was quenched with cold water (10 mL) and extracted with EtOAc (3 x 20 mL). The combined organic phase was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford crude product, which was purified by column chromatography (100-200 mesh silica, 15-18 %EtOAc in hexane as an eluent to afford product which was re-purified by prep HPLC purification to afford **302** (25 mg, 0.061 mmol, 96 % yield) as a solid. **HPLC**: Rt 8.353 min; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% formic acid in water:ACN (95:05) , B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 407.20 (M+H), Rt 1.94 min, Column: X-Select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 8.62 (d, 1H), 7.85 (s, 1H), 7.67 (d, 1H), 7.23-7.04 (m, 1H), , 5.12 (s, 2H), 3.97-3.90 (m, 3H), 3.37-3.52 (m, 2H), 3.12 (s, 1H), 2.32-2.25 (m, 1H), 1.0-0.97 (m, 4H)

Example 229. Synthesis of N-[(1S)-1-[3-(6-cyclopropylpyridazin-4-yl)-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (303)



Synthesis of 6-chloropyridazine-4-carbonitrile (A-437)

[000963] A solution of **A-436** (1 g, 5.79 mmol) in 7M ammonia methanol (10 mL) was stirred for 3 h at 80-85 °C. The reaction mixture was cooled to RT and evaporated under reduced pressure to afford a solid. The crude solid was dissolved in $POCl_3$ (3 mL, 32.19 mmol) and stirred for 7 h at reflux temperature. The reaction mixture concentrated under reduced pressure to afford crude product, which was diluted using ethyl acetate and washed with water. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to afford **A-437** (120 mg, 0.8079 mmol, 14 % yield).

Synthesis of 6-cyclopropylpyridazine-4-carbonitrile (A-438)

[000964] To the stirred solution of **A-437** (40.mg, 0.29 mmol) in 1,4-dioxane (2 mL) was added cesium carbonate (140.1 mg, 0.43 mmol) and cyclopropylboronic acid (41.86 mg, 0.49 mmol) at RT. Reaction mixture was purged under argon for 20 min and treated with Pd(dppf)Cl_2 DCM (11.7 mg, 0.01 mmol) at RT and purged with argon for 10 min. The reaction mixture was stirred for 5 h at 100 °C. The reaction mixture was diluted with ethyl acetate and filtered through celite. The organic layer was washed with water (10 mL x 3), concentrated under reduced pressure to afford the crude product which was purified by column chromatography using 20-25 % of ethyl acetate in hexane as an eluent to afford **A-438** (35 mg, 0.228 mmol, 80% yield).


Synthesis of 6-cyclopropyl-N'-hydroxy-pyridazine-4-carboxamidine (A-439)

[000965] To a stirred solution of **A-437** 7 (120 mg, 0.83 mmol) in ethanol (2 mL) was added hydroxylamine hydrochloride (86.17 mg, 1.24 mmol) at RT. The reaction mixture was stirred for 2 h at reflux temperature. The reaction mixture was cooled to RT and evaporated under reduced pressure to afford **A-439** (120 mg, 0.635 mmol, 77 % yield).

Synthesis of N-[1-[3-(6-cyclopropylpyridazin-4-yl)-1,2,4-oxadiazol-5-yl]ethyl]-2-methyl-5-(trifluoromethyl)pyrazole-3-carboxamide (303)

[000966] To a stirred solution of **A-430d** (229.19 mg, 0.86 mmol) in 1,4-dioxane (2 mL) was added **A-439** (140.mg, 0.79 mmol) and DCC (178.03 mg, 0.86 mmol) and stirred for 16 h at reflux temperature. The reaction mixture was filtered and mother liquor was diluted with ethyl acetate and washed with water (3 mL x 2), and separated. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to afford the crude product which was purified using prep-HPLC to afford **303** (45mg, 0.11 mmol, 14 % yield). **HPLC**: Rt 8.209 min; Column: X-Select CSH C18 (4.6 X 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: Acetonitrile (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 408 (M+H), Rt 2.05 min, Column: X-Select CSH C18 (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.49 (d, 1H), 8.05 (d, 1H), (7.45 (s, 1H), 5.57-5.49 (m, 1H), 4.027 (s, 3H), 2.45-2.42 (m, 1H), 1.69 (d, 3H), 1.29-1.26 (m, 1H), 1.17 (d, 4H). **Chiral method:** Rt 5.847 min, 35% + 6.795 (65%); column: PHENOMENE CELLULOSE I (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% TFA B) Isopropyl alcohol IPA, Isocratic:30% B; Wavelength: 254 nm, Flow: 1.0 mL/min. (**NOTE**: Chiral HPLC shows partial racemization)

Example 230. Synthesis of 2-cyclobutyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (304)



Synthesis of methyl 1-cyclobutyl-1H-pyrazole-5-carboxylate (A-442)

[000967] To a stirred solution of **A-440** (2 g, 14.27 mmol) and **A-441** (1.93 mg, 14.27 mmol) in DMF (20 mL) was added K_2CO_3 (3944.63 mg, 28.54 mmol) at RT. The reaction mixture was stirred at 80 °C for 24 h. The reaction mixture was quenched using water (10 mL) and diluted with EtOAc (50 mL x 2). The organic layer was separated, dried over Na_2SO_4, filtered, evaporated under reduced pressure to afford crude product. The crude product was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **A-442**. (0.40 g, 2.039 mmol, 14 % yield) and ethyl 1-cyclobutylpyrazole-3-carboxylate (0.6 g, 3.058 mmol, 21 % yield).


Synthesis of 1-cyclobutyl-1H-pyrazole-5-carboxylic acid (A-443)

[000968] To a stirred solution of **A-442** (0.3 g, 1.54 mmol) in THF (10 mL) and water (5 mL) was added $LiOH.H_2O$ (97.21 mg, 2.32 mmol) at RT. The reaction mixture was for 2 h at RT. The reaction mixture was quenched using water (100 mL) and diluted with EtOAc (50 mL x 2). The organic layer was separated, then the aqueous layer was acidified with 1N HCl leading to precipitation. The precipitate was filtered to afford a solid which was separated and dried under reduced pressure to afford **A-443** (100 mg, 0.5957 mmol, 38 % yield) as a solid.

Synthesis of 2-cyclobutyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (304)

[000969] To a stirred solution of **A-303** (100 mg, 0.37 mmol) and **A-443** (74.76 mg, 0.45 mmol) in DCM (10 mL) was added HATU (213.83 mg, 0.56 mmol) and DIPEA (0.1 3mL, 0.75 mmol) at RT. The reaction mixture was stirred at RT for 2 h. The reaction mixture was quenched with water (10 mL) and diluted with DCM (50 mL x 2). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude product was purified by column chromatography using 100-200 silica and 30-80% EtOAc/hexane as an eluent to afford **304** (15 mg, 0.0388 mmol, 10 % yield) as a solid. **HPLC**: Rt 8.176 min, 95.2%; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 379 (M+H), Rt 2.143 min, Column: X-select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.23 (d, 1H), 8.63 (d, 1H), 7.88 (s, 1H), 7.72 (d, 1H), 7.58 (s, 1H), 6.93 (s, 1H), 5.60-5.56 (m, 1H), 5.45-5.41 (m, 1H), 2.53-2.50 (m, 2H), 2.31-2.29 (m, 3H), 1.78-1.71 (m, 2H), 1.67 (d, 3H), 1.05-1.01 (m, 4H).**Chiral method:** Rt 8.967 min, 98.7%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 292 nm, Flow: 1.0 mL/min.

Example 231. Synthesis of 2-cyclopropyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (305)



Synthesis of 2-cyclopropylpyrazole-3-carboxylic acid (A-445)


[000970] To a stirred solution of **A-444** (0.26 g, 1.44 mmol) in THF (2 mL) and water (0.5 mL) was added lithium hydroxide (0.12 g, 2.89 mmol) at 0 °C and stirred at RT 6 h. The reaction mixture was concentrated under reduced pressure and the residue was acidified with


1N HCl leading to precipitatation which was collected by filtration to afford **A-445** (0.13 g, 0.819 mmol, 57 % yield) as a solid.

Synthesis of 2-cyclopropyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (305)

[000971] To a stirred solution of **A-445** (0.13 g, 0.85 mmol) in DCM (10 mL) was added HATU (0.32 g, 0.85 mmol), **A-303** (0.23 g, 0.850 mmol), and DIPEA (0.3 mL, 1.71 mmol) and stirred at RT for 6 h. The reaction mixture was diluted with DCM (20 mL) and water (10 mL), and the organic layer was separated. The organic layer was washed with water (2 x 10 mL), saturated brine solution (10 mL), dried over $MgSO_4$, and evaporated to dryness to afford the crude product which was purified by flash column chromatography, eluting 30 % EtOAc in hexane to afford **305** (60 mg, 0.1593 mmol, 19 % yield) as an oil. **HPLC**: Rt 7.532 min, 96.8%; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 365.45 (M+H), Rt 1.624 min, Column: X-select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.25 (d, 1H), 8.60 (d, 1H), 7.85 (s, 1H), 7.66 (d, 1H), 7.45 (s, 1H), 6.92 (s, 1H), 5.48-5.45 (m, 1H), 4.43-4.41 (m, 1H), 2.29-2.27 (m, 1H), 1.68 (d, 3H), 1.07-0.92 (m, 8H). **Chiral method:** Rt 12.941 min, 96.7%; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl amine B) DCM: MeOH (1:1), Isocratic:20% B; Wavelength: 292 nm, Flow: 1.0 mL/min.

Example 232. Synthesis of 1-benzyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (306)





Chemical Formula: $C_{23}H_{22}N_6O_2$
Molecular Weight: 414.47

Synthesis of ethyl 1-benzylpyrazole-3-carboxylate (A-447)

[000972] To a stirred solution of **A-440** (5.g, 35.68 mmol) and benzyl chloride (4.14 mL, 35.68 mmol) in ACN (10 mL) was added K_2CO_3 (9.86 g, 71.36 mmol) at RT and stirred for 16 h. The reaction mixture was quenched using water (10 mL) and diluted with EtOAc (50 mL x 2). The organic layer was separated, dried over Na_2SO_4, then filtered. The organic layer was evaporated under reduced pressure. The crude reaction mass was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **A-447** (4.2 g, 16.412 mmol, 46% yield) and ethyl 2-benzylpyrazole-3-carboxylate (1.6 g, 16.115 mmol, 17 % yield).

Synthesis of 1-benzylpyrazole-3-carboxylic acid (A-448)


[000973] To a stirred solution of **A-447** (4.4 g, 19.11 mmol) in THF (20 mL) and water (10 mL) was added $LiOH.H_2O$ (1202.71 mg, 28.66 mmol) at RT and stirred for 2 h. The reaction mixture was quenched using water (100 mL) and diluted with EtOAc (50 mL x 2). The combined organic layer was separated, then the aqueous layer was acidified with 1N HCl to form a precipitate which was collected by filtration. The solid was separated and dried under reduced pressure to afford **A-448** (2 g, 8.902 mmol, 46 % yield).

Synthesis of compound 1-benzyl-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (306)

[000974] To a stirred solution of **A-303** (200 mg, 0.75 mmol) and **A-448** (49.31 mg, 0.31 mmol) in DCM (10 mL) was added HATU (427.66 mg, 1.12 mmol) and DIPEA (0.26 mL, 1.5 mmol) at RT. The reaction mixture was stirred at RT for 2 h. The reaction mixture was quenched with water (10 mL) and diluted with DCM (50 mL). The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated to afford the crude product. The crude compound was purified by column chromatography using 100-200 silica and 30-80% EtOAc/hexane as eluent to afford **306** (10 mg, 0.0234 mmol, 3% yield) as a solid. **HPLC**: Rt 8.433 min; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 415.1 (M+H), Rt 1.966 min, Column: X-select CSH (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 8.95 (d, 1H), 8.59 (d, 1H), 7.97 (d, 1H), 7.84 (s, 1H), 7.64-7.63 (m, 1H), 7.38-7.30 (m, 3H), 7.26-7.24 (m, 2H), 6.73 (d, 1H), 5.45-5.41 (m, 3H), 2.29-2.26 (m, 1H), 1.66 (d, 3H), 1.00-0.96 (m, 4H). **Chiral method:** Rt 7.372 min, 86%; column: YMC

CHIRAL AMYLOSE -SA (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:15% B; Wavelength: 292 nm, Flow: 1.0 mL/min.

Example 233. Synthesis of (S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (307), (S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-5-carboxamide (308) and (S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-(cyclopropylmethyl)-1H-pyrazole-5-carboxamide (309)



Synthesis 2-cyclopentylpyridine-4-carbonitrile (A-449)

[000975] To a stirred solution of isonicotinonitrile (1.5 g, 14.41 mmol) in water (20 mL) and chlorobenzene (20 mL) was added cyclopentanecarboxylic acid (4.93 g, 43.22 mmol), ammonium persulfate (6.58 g, 28.82 mmol), TFA (1.06 mL, 14.41 mmol), silver nitrate (0.24 g, 1.44 mmol). The reaction mixture was stirred at 120 °C for 3 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic layer was dried over Na_2SO_4 filtered and evaporated to afford crude product which was purified by combiflash using 100-

200 mesh silica and 8% ethyl acetate/hexane as an eluent to afford **A-449** (2 g, 11.61 mmol, 80 % yield) as an oil.

Synthesis of 2-cyclopentyl-N-hydroxy-pyridine-4-carboxamidine (A-450)

[000976] To a stirred solution of **A-449** (2 g, 13.87 mmol) was added hydroxylamine hydrochloride (3.83 g, 55.49 mmol) and TEA (5.81 mL, 41.62 mmol) and stirred for 3 h at 70 °C. The reaction mixture was evaporated and diluted using water and EtOAc. The organic layer was separated, dried over Na_2SO_4, filtered and evaporated to afford **A-450** (2.1 g, 11.85 mmol, 85% yield) as an oil.

Synthesis of tert-butyl N-[(1S)-1-[3-(2-cyclopentyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]carbamate (A-451)

[000977] To a stirred solution of **A-450** (1.2 g, 5.85 mmol) and (2S)-2-(tert-butoxycarbonylamino)propanoic acid (1.33 g, 7.02 mmol) in 1,4-Dioxane (8mL) was added DCC (1.45 g, 7.02 mmol) and stirred for 16 h at 100 °C. The reaction was quenched with water (10 mL) and diluted with EtOAc. The organic layer was separated, dried over Na_2SO_4, then filtered and evaporated under reduced pressure to afford crude product. The crude product was purified by column chromatography using 100-200 silica and 22% EtOAc/Hexane as an eluent to afford **A-451** (610 mg, 1.702 mmol, 29% yield) as an oil

Synthesis of (1S)-1-[3-(2-cyclopentyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethanamine hydrochloride (A-452)

[000978] To a stirred solution of **A-451** (600.mg, 1.67 mmol) in 1,4-Dioxane (3 mL) was added 4M HCl in 1,4-Dioxane (2 mL, 14.33 mmol) and stirred at RT for 16 h. The reaction mixture was evaporated under reduced pressure and the crude solid was washed with diethyl ether to afford **A-452** (410 mg, 1.391 mmol, 83 % yield) as a solid.

Synthesis of (S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-(cyclopropylmethyl)-1H-pyrazole-5-carboxamide (309)

[000979] To a stirred solution of **A-452** (70 mg, 0.27 mmol) and 1-(cyclopropylmethyl)-1H-pyrazole-5-carboxylic acid (47.35 mg, 0.28 mmol) in DCM (10 mL) was added HATU (135.44 mg, 0.36 mmol) and DIPEA (0.08 mL, 0.47 mmol) at RT and stirred for 2 h at RT. The reaction was quenched using water (10 mL), diluted with DCM (50 mL), and organic layer was separated. The organic layer was dried over Na_2SO_4, filtered and evaporated under

reduced pressure to afford the crude product which was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **309** (15 mg, 0.0368 mmol, 15 % yield) as a solid. **HPLC**: Rt 8.756 min; Column: X-Select CSH C18 (4.6 X 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 407.03 (M+H), Rt 2.074 min, Column: X-select CSH C18 (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.29 (d, 1H), 8.70 (d, 1H), 7.77 (s, 1H) , 7.71 (d, 1H), 7.53-7.52 (m, 1H), 6.95-6.94 (m, 1H), 5.44-5.41 (m, 1H), 4.32 (d, 2H), 3.29-3.27 (m, 1H), 2.04-1.99 (m, 2H), 1.77-1.71 (m, 4H), 1.68-1.66 (m, 5H), 1.22-1.17 (m, 1H), 0.40-0.34 (m, 2H), 0.30-0.29 (m, 2H). **Chiral method:** Rt 11.397 min, 96.4% min; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20%B; Wavelength: 228 nm, Flow: 1.0 mL/min.


Synthesis of (S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (307)

[000980] To a stirred solution of **A-452** (57.06 mg, 0.19 mmol) and 1-methyl-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (56.36 mg, 0.29 mmol) in DCM (3 mL) was added HATU (110.4 mg, 0.29 mmol) and DIPEA (0.08mL, 0.48 mmol) at RT and stirred for 16 h. The reaction was quenched using water, extracted with DCM, and the organic layer was separated. The organic layer was dried over Na_2SO_4, filtered and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 100-200 silica and 23% EtOAc/Hexane as an eluent to afford **307** (5 mg, 0.0097 mmol, 5 % yield) as a solid. **HPLC**: Rt 9.074 min; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 435.05 (M+H), Rt 2.171 min, Column: X-select CSH C18 (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.47 (d, 1H), 8.70 (d, 1H), 7.78-7.72 (m, 2H), 7.45 (s, 1H), 5.49-5.46 (m, 1H), 4.12 (s, 3H), 2.02-2.00 (m, 2H), 1.77-1.67 (m, 9H). **Chiral method:** Rt 5.287 min, 99.4% min; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20%B; Wavelength: 282 nm, Flow: 1.0 mL/min.

(S)-N-(1-(3-(2-cyclopentylpyridin-4-yl)-1,2,4-oxadiazol-5-yl)ethyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-5-carboxamide (308)

[000981] To a stirred solution of **A-452** (100 mg, 0.34 mmol) and 3-(difluoromethyl)-1-methyl-1H-pyrazole-5-carboxylic acid (71.7 mg, 0.41 mmol) in DCM (10 mL) was added HATU (193.48 mg, 0.51 mmol) and DIPEA (0.12 mL, 0.68 mmol) at RT and stirred for 2 h at RT. The reaction was quenched using water (10 mL), extracted with DCM (50 mL), and the organic layer was separated. The organic layer was dried over Na_2SO_4, filtered and evaporated under reduced pressure to afford the crude product which was purified by column chromatography using 100-200 silica and 30-80% EtOAc/Hexane as an eluent to afford **308** (12.17 mg, 0.029 mmol, 8 % yield) as a solid. **HPLC**: Rt 8.686 min; Column: X-Select CSH C18 (4.6 X 150) mm, 5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN (95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 417.1 (M+H), Rt 2.1 min, Column: X-select CSH C18 (3*50) mm, 2.5 µm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.40 (d, 1H), 8.72 (d, 1H), 7.83 (s, 1H) , 7.78 (d, 1H), 7.26 (s, 1H), 7.19-6.91 (m, 1H), 5.48-5.44 (m, 1H), 4.08 (s, 3H), 3.34-3.30 (m, 1H), 2.04-2.02 (m, 2H), 1.78-1.71 (m, 4H), 1.68-1.66 (m, 5H). **Chiral method:** Rt 7.701 min, 98.4% min; column: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1%Iso-propyl-amine B) DCM: MeOH (1:1), Isocratic:20%B; Wavelength: 227 nm, Flow: 1.0 mL/min.

Example 234. Synthesis of 2-(cyclopropylmethyl)-N-[(1S)-1-[3-(2-cyclopropyl-4-pyridyl)-1,2,4-oxadiazol-5-yl]ethyl]pyrazole-3-carboxamide (310)



A-303 310

[000982] To a stirred solution of **A-303** (100 mg, 0.37 mmol) and 1-(cyclopropylmethyl)-1H-pyrazole-5-carboxylic acid (83.07 mg, 0.45 mmol) in DCM (10 mL) was added HATU (213.83 mg, 0.56 mmol) and DIPEA (0.13 mL, 0.75 mmol) at RT. The reaction mixture was stirred at RT for 2 h. The reaction mixture was quenched with water (10 mL) and diluted with DCM (50 mL x 2). The organic layer was dried over anhydrous sodium sulphate, filtered and concentrated under reduced pressure. The crude compound was purified by column chromatography using 100-200 silica and 30 to 80% EtOAc in hexane as an eluent to afford **310** (20 mg, 0.0525 mmol, 14% yield) as a solid. **HPLC**: Rt 7.97 min; Column: X-Select CSH C18 (4.6 X 150) mm, 3.5 µm; Mobile phase: A: 0.1% Formic acid in water: ACN

(95:05), B: ACN; Flow Rate: 1.0 mL/min. **LCMS** : 379.4 (M+H), Rt 1.801 min, Column: X-select CSH (3*50) mm, 2.5 μm. **^{1}H NMR (400 MHz, DMSO-d6)** δ 9.28 (d, 1H), 8.60 (d, 1H), 7.85 (s, 1H), 7.65 (dd, 1H), 7.53 (d, 1H), 6.95 (d, 1H), 5.44 (quin, 1H), 4.27-4.39 (m, 2H), 2.28-2.46 (m, 1H), 1.68 (d, 3H), 1.10-1.30 (m, 1H), 0.90-1.08 (m, 4H), 0.23-0.47 (m, 4H). **Chiral method:** Rt 6.523 min, 96%: DIACEL CHIRALPAK-IG (250 x 4.6 mm, 5 um), - Mobile Phase: A) n-Hexane+0.1% Iso-propyl-amine B) EtOH: MeOH (50:50), Isocratic:30% B; Wavelength: 293 nm, Flow: 1.0 mL/min.

Example 235. Efficacy of exemplary compounds in the inhibition of KCNT1

KCNT1 - Patch Clamp Assay

[000983] Inhibition of KCNT1 (KNa1.1, Slack) was evaluated using a tetracycline inducible cell line (HEK-TREX). Currents were recorded using the SyncroPatch 384PE automated, patch clamp system. Pulse generation and data collection were performed with PatchController384 V1.3.0 and DataController384 V1.2.1 (Nanion Technologies). The access resistance and apparent membrane capacitance were estimated using built-in protocols. Current were recorded in perforated patch mode (10 μM escin) from a population of cells. The cells were lifted, triturated, and resuspended at 800,000 cells/ml. The cells were allowed to recover in the cell hotel prior to experimentation. Currents were recorded at room temperature. The external solution contained the following (in mM): NaCl 105, NMDG 40, KCl 4, $MgCl_2$ 1, $CaCl_2$ 5 and HEPES 10 (pH = 7.4, Osmolarity ~300 mOsm). The extracellular solution was used as the wash, reference and compound delivery solution. The internal solution contained the following (in mM): NaCl 70, KF 70, KCl 10, EGTA 5, HEPES 5 and Escin 0.01 (pH = 7.2, Osmolarity ~295 mOsm). Escin is made at a 5mM stock in water, aliquoted, and stored at -20°C. The compound plate was created at 2x concentrated in the extracellular solution. The compound was diluted to 1:2 when added to the recording well. The amount of DMSO in the extracellular solution was held constant at the level used for the highest tested concentration. A holding potential of -80 mV with a 100ms step to 0mV was used. Mean current was measured during the step to 0 mV. 100 μM Bepridil was used to completely inhibit KCNT1 current to allow for offline subtraction of non-KCNT1 current. The average mean current from 3 sweeps was calculated and the % inhibition of each compound was calculated. The % Inhibition as a function of the compound concentration was fit with a Hill equation to derive IC_{50}, slope, min and max parameters. If KCNT1 inhibition was less than 50% at the highest tested concentration or if an IC_{50} could not be calculated, then a percent inhibition was reported in place of the IC_{50}.

[000984] Results from this assay are summarized in Table 1 below. In this table, "A" indicates IC_{50} of less than or equal to1 μM; "B" indicates inhibition of between 1 μM to 20 μM; and "C" indicates inhibition of greater than or equal to 20 μM.

Table 1

Patent Compound No.	KCNT1-WT IC_{50} (μM)
192	A
191	B
3	C
4	C
5	A
6	A
7	C
8	B
9	A
10	A
11	B
12	A
13	A
14	B
15	B
16	A
17	B
19	A
20	B
21	A
22	A
23	B
24	A
25	B

26	A
29	A
30	A
31	A
32	B
33	A
34	A
35	B
36	A
37	B
38	A
39	B
40	B
41	B
42	A
43	B
44	C
45	A
46	B
47	A
48	A
49	B
50	B
51	B
52	A
54	C

55	B
56	A
57	B
58	B
59	C
60	A
61	A
62	B
63	A
64	A
65	B
66	A
67	A
68	B
69	A
70	A
71	C
72	A
73	A
74	A
75	A
76	A
77	A
78	B
79	B
80	A

81	A
82	A
83	B
84	A
85	C
86	A
87	B
88	B
89	A
90	B
91	A
92	A
93	B
94	C
95	B
96	A
97	A
98	B
99	B
100	B
101	B
102	A
103	A
104	B
105	A
106	B

107	A
108	A
109	A
110	A
111	A
112	A
113	A
114	C
115	C
116	A
117	A
118	B
119	A
120	A
121	C
122	C
123	C
124	C
125	A
126	A
127	A
128	C
129	C
130	A
131	B
132	A

133	B
134	C
135	B
136	B
137	B
138	A
139	A
140	B
141	B
142	B
143	C
144	B
145	A
146	A
147	B
148	B
149	A
150	A
151	B
152	B
153	A
154	A
155	B
156	C
157	B
160	B

161	B
162	B
163	B
164	B
165	A
166	B
167	B
168	B
169	B
170	A
171	B
172	B
173	B
174	A
175	A
176	B
177	C
178	B
179	B
180	C
181	A
182	C
183	C
184	A
185	B
186	B

187	C
188	B
189	B
193	C
194	A
195	B
196	A
197	A
198	A
199	A
200	B
201	A
202	A
203	A
204	A
205	A
206	A
207	A
208	A
209	A
210	A
211	A
212	A
213	B
214	A
215	B

216	A
217	B
218	A
219	A
220	B
221	A
222	A
223	A
224	A
225	A
226	A
227	A
228	A
229	A
230	A
231	B
232	C
233	A

234	A
235	A
236	A
237	C
238	A
239	A
240	A
241	A
242	B
243	A
244	B
245	B
246	A
248	A
249	A
250	B
251	B
252	B
253	A
254	B
255	A
256	A
257	A
258	B
259	B
260	C

261	C
263	B
264	C
265	A
266	A
267	B
268	A
269	A
270	A
271	A
278	A
279	A
280	A
272	B
247	A
262	B
273	A
274	A
275	A
276	A
277	B
281	B
282	B
283	B
284	B
285	C

286	A
287	A
288	A
289	C
290	A
291	B
292	B
293	C
294	B
295	A
296	B
297	B
299	C
300	A
301	A
302	A
303	A
304	A
305	A
306	A
307	A
308	A
309	A
310	A

Equivalents and Scope

[000985] In the claims articles such as "a," "an," and "the" may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[000986] Furthermore, the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, e.g., in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should it be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements and/or features, certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth *in haec verba* herein. It is also noted that the terms "comprising" and "containing" are intended to be open and permits the inclusion of additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub–range within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

[000987] This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular


embodiment of the invention can be excluded from any claim, for any reason, whether or not related to the existence of prior art.

[000988] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.

****

CLAIMS

1. A pharmaceutical composition comprising a compound of Formula **I**:



(I),

or a pharmaceutically acceptable salt thereof, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be replaced with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-}

$_{6}$haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, -C(O)OR_8, C(O)$N(R_9)_2$, C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$-$N(R_9)_2$, -OC(O)C_{1-6}alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -$N(R_9)_2$, -NR_9-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)$N(R_9)_2$,-NR_{10}C(O)-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N;

and a pharmaceutically acceptable excipient.


2. The pharmaceutical composition of claim 1, wherein one of X, Y, Z, Y', and Z' is N and the other four are CH.

3. The pharmaceutical composition of claim 1, wherein the compound is chosen from: a compound of Formula **I-a**:




(**I-a**),


a compound of Formula **I-b**:


(**I-b**),


a compound of Formula **I-c**:


(**I-c**),


a compound of Formula **I-d**:


(**I-d**),


a compound of Formula **I-e**:


(**I-e**),


a compound of Formula **I-f**:


(**I-f**),


a compound of Formula **I-g**:


(**I-g**),

a compound of Formula **I-h**:



(**I-h**),

a compound of Formula **I-i**:




(**I-i**),

a compound of Formula **I-j**:




(**I-j**),

a compound of Formula **I-k**:




(**I-k**),

a compound of Formula **I-l**:




(**I-l**),

a compound of Formula **I-m**:




(**I-m**),

a compound of Formula **I-n**:




(**I-n**),

a compound of Formula **I-o**:




(**I-o**),

a compound of Formula **I-p**:




(**I-p**),

a compound of Formula **I-q**:




(**I-q**),

a compound of Formula **I-r**:



(**I-r**), or

a compound of Formula **I-s**:



(**I-s**),

or a pharmaceutically acceptable salt thereof, wherein the variables are as defined in claim 1.

4. The pharmaceutical composition of any one of claims 1-3, wherein R_2 is hydrogen or methyl.

5. The pharmaceutical composition of any one of claims 1-4, wherein R_3 is hydrogen, C_{1-6}alkyl, or C_{1-6}alkyl substituted with C_{1-6}alkoxy, -OH, or -C(O)OR_8.

6. The pharmaceutical composition of any one of claims 1-5, wherein R_4 is hydrogen.

7. The pharmaceutical composition of any one of claims 1-4, wherein R_3 and R_4 are taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene.

8. The pharmaceutical composition of claim 7, wherein the C_{3-7}cycloalkylene is selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

9. The pharmaceutical composition of claim 7, wherein the 3-7 membered heterocyclene is selected from the group consisting of oxetanyl, tetrahydrofuranyl, and tetrahydropyranyl.

10. The pharmaceutical composition of any one of claims 1-9, wherein each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-}

$_{6}$alkoxy, C_{1-6}haloalkoxy, C_{3-10}cycloalkyl, O-C_{3-10}cycloalkyl, -OH, -CN, $N(R_9)_2$, and -$C(O)OR_8$.


11. The pharmaceutical composition of any one of claims 1-10, wherein n is 1.

12. The pharmaceutical composition of any one of claims 1-11, wherein R_1 is selected from the group consisting of C_{1-6}alkyl optionally substituted with C_{1-6}alkoxy, $N(R_9)_2$, $C(O)N(R_9)_2$, C_{3-7}cycloalkyl, pyridyl, tetrahydropyranyl, or phenyl, C_{1-6}haloalkyl, C_{3-7}cycloalkyl, phenyl optionally substituted with halogen, and pyridyl optionally substituted with halogen.

13. The pharmaceutical composition of any one of claims 1-12, wherein R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}haloalkyl, C_{3-7}cycloalkyl, and phenyl optionally substituted with halogen.

14. The pharmaceutical composition of any one of claims 1-13, wherein x is 1 or 2.

15. A compound of Formula **II**:



(**II**),

or a pharmaceutically acceptable salt thereof, wherein

one or two selected from X, Y, Z, Y', and Z' are N, and the others are CH, wherein the hydrogen of CH may be replaced with R_5;

R_1 is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy optionally substituted with C_{3-7}cycloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, -$N(R_9)_2$, -C(O)OR_8, -C(O)$N(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -$S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$-$N(R_9)_2$, -OC(O)C_{1-6}alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -$N(R_9)_2$, -NR_9-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)$N(R_9)_2$,-NR_{10}C(O)-R_{11}, -CN, -$S(O)_2$-C_{1-6}alkyl, -$S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -$N(R_9)_2$;


R_8 is selected from the group consisting of hydrogen, C_{1-6}alkyl, and C_{3-10}cycloalkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl; and

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl.

16. A compound of Formula **III**:



(**III**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is C_{1-6}alkyl or C_{3-7}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-7}cycloalkyl is optionally substituted with one or more halogen or C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)N(R_9)$_2$, -C_{1-6}alkylene-CN, -CN, -S(O)$_2$-C_{1-6}alkyl, C_{1-6}alkylene-S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, -OC(O)C_{1-6}alkyl, and -O-C_{1-3}cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, -C(O)OR_8, -C(O)N(R_9)$_2$,-NR_{10}C(O)-R_{11}, -CN, -S(O)$_2$-C_{1-6}alkyl, -S(O)$_2$- N(R_9)$_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached


to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;

wherein the compound is not:




,



, or a pharmaceutically acceptable salt thereof.

17. A compound of Formula **IV**:



(**IV**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is selected from the group consisting of C_{1-6}alkyl, and C_{3-7}cycloalkyl, wherein the C_{1-6}alkyl or C_{3-7}cycloalkyl is optionally substituted with one or more substituents independently selected from halogen and C_{1-6}alkoxy;

R_{12} is C_{1-6}alkyl optionally substituted with one or more halogen or C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}alkoxy, C_{1-6}haloalkyl, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)N(R_9)$_2$, -C_{1-6}alkylene-CN, -CN, -S(O)$_2$-C_{1-6}alkyl, C_{1-

$_6$alkylene-S(O)$_2$-C$_{1-6}$alkyl, -S(O)$_2$- N(R$_9$)$_2$, -OC(O)C$_{1-6}$alkyl, and -O-C$_{3-10}$cycloalkyl optionally substituted with one or more halogen;

R_7 is each independently selected from the group consisting of phenyl, C$_{1-6}$alkoxy, -OH, -O-(C$_{1-6}$alkylene)-phenyl, C$_{3-10}$cycloalkyl, -C(O)OR$_8$, -C(O)N(R$_9$)$_2$,-NR$_{10}$C(O)-R$_{11}$, -CN, -S(O)$_2$-C$_{1-6}$alkyl, -S(O)$_2$- N(R$_9$)$_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C$_{3-10}$cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C$_{1-6}$alkyl, halogen, -OH, C$_{1-6}$alkoxy, and -N(R$_9$)$_2$;

R_8 is hydrogen or C$_{1-6}$alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C$_{1-6}$alkyl, and -(C$_{1-6}$alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents selected from halogen and -OH;


each R_{10} is independently hydrogen or C$_{1-6}$alkyl;

R_{11} is selected from the group consisting of C$_{1-6}$alkyl, C$_{1-6}$alkoxy, and -O-(C$_{1-6}$alkylene)-phenyl; and


n is selected from the group consisting of 0, 1, 2, and 3;

wherein the compound is not:




,


,



, or a pharmaceutically acceptable salt thereof.

18. A compound of Formula **V**:



(**V**),

or a pharmaceutically acceptable salt thereof, wherein

R_1 is phenyl or 3-10 membered heteroaryl, wherein the phenyl or 3-10 membered heteroaryl is optionally substituted with one or more substituents independently selected from halogen and C_{1-6}alkoxy,

R_{12} is selected from the group consisting of C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{3-10}cycloalkyl, 3-10 membered saturated heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from halogen and C_{1-6}alkoxy;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

R_5 is selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkoxy, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, -C_{1-6}alkylene-OH, OH, -C(O)OR_8, -C(O)N$(R_9)_2$, -C_{1-6}alkylene-CN, -CN, -S$(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-S$(O)_2$-C_{1-6}alkyl, -S$(O)_2$- N$(R_9)_2$, -OC(O)C_{1-6}alkyl, and -O-C_{3-10}cycloalkyl;

R_7 is selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, -O-(C_{1-6}alkylene)-phenyl, C_{3-7}cycloalkyl, -C(O)OR_8, -C(O)N$(R_9)_2$,-NR_{10}C(O)-R_{11}, -CN, -S$(O)_z$-C_{1-6}alkyl, -S$(O)_z$- N$(R_9)_2$, 3-7 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N$(R_9)_2$;

R_8 is hydrogen or C_{1-6}alkyl;


each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;


R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

n is selected from the group consisting of 0, 1, 2, and 3;


wherein the compound is not:




,



,












,

















, or a pharmaceutically acceptable salt thereof.








19. The pharmaceutical composition of any one of claims 1-14 or the compound of any one of claims 15-18, wherein the compound is selected from the group consisting of








3	
4	








5	
6	
7	
8	
9	
10	
11	








12	
13	
14	
15	
16	
17	
19	







20	
21	
22	
23	
24	
25	
26	







29	
30	
31	
32	
33	
34	
35	







36	
37	
38	
39	
40	
41	
42	







43	
44	
45	
46	
47	
48	
49	








50	
51	
52	
54	
55	
56	
57	








58	
59	
60	
61	
62	
63	








64	
65	
66	
67	
68	
69	







70	
71	
72	
73	
74	
75	







76	
77	
78	
79	
80	
81	







82	
83	
84	
85	
86	
87	
88	






89	
90	
91	
92	
93	
94	
95	







96	
97	
98	
99	
100	
101	
102	







103	
104	
105	
106	
107	
108	







109	
110	
111	
112	
113	
114	







115	
116	
117	
118	
119	
120	







121	
122	
123	
124	
125	








126	
127	
128	
129	
130	
131	







132	
133	
134	
135	
136	
137	







138	
139	
140	
141	
142	
143	






144	
145	
146	
147	
148	
149	






150	
151	
152	
153	
154	
155	







156	
157	
160	
161	
162	
163	
164	






165	
166	
167	
168	
169	
170	






171	
172	
173	
174	
175	
176	






177	
178	
179	
180	
181	






182	
183	
184	
185	
186	





187	
188	
189	
191	
192	
193	






194	
195	
196	
197	
198	






199	
200	
201	
202	
203	






204	
205	
206	
207	
208	









209	
210	
211	
212	
213	








214	
215	
216	
217	









218	
219	
220	
221	
222	









223	
224	
225	
226	
227	









228	
229	
230	
231	
232	








233	
234	
235	
236	
237	
238	
239	
240	









241	
242	
243	
244	
245	
246	
248	

249	
250	
251	
252	
253	
254	
255	
256	

257	
258	
259	
260	
261	
263	
264	
265	


266	
267	
268	
269	
270	
271	
278	
279	

280	
297	
298	
299	
300	
301	
302	

303	
304	
305	
306	
307	
308	
309	
310	

, or a pharmaceutically acceptable salt thereof.

20. A compound of Formula **VI**:



(**VI**),

or a pharmaceutically acceptable salt thereof, wherein

R_{13} is $C_{1\text{-}6}$alkyl or $C_{3\text{-}10}$cycloalkyl, wherein the $C_{1\text{-}6}$alkyl or $C_{3\text{-}10}$cycloalkyl is optionally substituted with phenyl;

R_{14} is hydrogen;

R_{15} is $C_{1\text{-}6}$alkyl or hydrogen;

R_{16} is $C_{1\text{-}6}$alkyl optionally substituted with one or more halogen, $C_{1\text{-}6}$alkoxy, $C_{3\text{-}10}$ cycloalkyl, or phenyl;


each R_{17} is independently selected from the group consisting of halogen, $C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$haloalkyl, $C_{1\text{-}6}$alkylene-$N(R_{20})_2$, $C_{1\text{-}6}$alkylene-O-$C_{3\text{-}10}$cycloalkyl, $C_{1\text{-}6}$alkoxy optionally substituted with $C_{3\text{-}7}$cycloalkyl, $C_{1\text{-}6}$haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or $C_{1\text{-}6}$alkoxy, 3-10 membered heteroaryl, -$C_{1\text{-}6}$alkylene-OH, $C_{1\text{-}6}$alkylene-$C_{1\text{-}6}$alkoxy, OH, -$N(R_{20})_2$, -$C(O)OR_{19}$, -$C(O)N(R_{20})_2$, -$C_{1\text{-}6}$alkylene-CN, -CN, -$S(O)_2$-$C_{1\text{-}6}$alkyl, $C_{1\text{-}6}$alkylene-$S(O)_2$-$C_{1\text{-}6}$alkyl, -$S(O)_2$- $N(R_{20})_2$, -$OC(O)C_{1\text{-}6}$alkyl, -O-$C_{3\text{-}10}$cycloalkyl optionally substituted with one or more halogen or $C_{1\text{-}6}$alkyl, and $C_{3\text{-}10}$cycloalkyl optionally substituted with one or more substituents selected from halogen, $C_{1\text{-}6}$alkyl, and $C_{1\text{-}6}$alkoxy;

p is selected from the group consisting of 1, 2, and 3;



R_{19} is selected from the group consisting of hydrogen, $C_{1\text{-}6}$alkyl, and $C_{3\text{-}10}$cycloalkyl; and

each R_{20} is independently hydrogen or $C_{1\text{-}6}$alkyl.

21. A pharmaceutical composition comprising a compound of Formula VII



(VII), wherein the compound is a compound of Formula **VII-a** or Formula **VII-b**:

 (**VII-a**),  (**VII-b**),





or a pharmaceutically acceptable salt thereof, wherein

R_{23} is a tert-butyl;

R_{24} is hydrogen;


R_{25} is hydrogen; and

R_{27} is independently selected from the group consisting of halogen and $C_{1\text{-}6}$alkoxy; and a pharmaceutically acceptable excipient.

22. The pharmaceutical composition of claim 21, wherein the compound is selected from the group consisting of  ,  ,  , and  , or a pharmaceutically acceptable salt thereof.

23. A pharmaceutical composition comprising a compound of any one of claims 15-20 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

24. Use of a compound of Formula **I**:

 (**I**),

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a neurological disease or disorder, wherein

X, Y, Z, Y', and Z' are each independently selected from CH and N, wherein the hydrogen of CH may be replaced with R_5, wherein at least 3 selected from X, Y, Z, Y', and Z' are CH;

R_1 is selected from the group consisting of $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein $C_{1\text{-}6}$alkyl, $C_{3\text{-}10}$cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with

one or more substituents each independently selected from the group consisting of halogen, $C(O)N(R_9)_2$, $N(R_9)_2$, C_{3-7}cycloalkyl, phenyl, 3-10 membered heteroaryl, and C_{1-6}alkoxy;

R_{12} is selected from the group consisting of C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl, wherein the C_{1-6}alkyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, or phenyl is optionally substituted with one or more substituents each independently selected from the group consisting of halogen, -OH, -CN, C_{1-6}alkyl, C_{1-6}haloalkyl, and C_{1-6}alkoxy; or

two R_{12} on adjacent carbons can be taken together with the two carbons where R_{12} are attached to form a carbocyclic ring;

x is 0, 1 or 2;

R_2 is hydrogen or C_{1-4}alkyl;

R_3 is selected from the group consisting of hydrogen, C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, and phenyl; and R_4 is selected from C_{1-6}alkyl and hydrogen; or R_3 and R_4 can be taken together with the carbon attached to R_3 and R_4 to form a C_{3-7}cycloalkylene or 3-7 membered heterocyclene; wherein the C_{1-6}alkyl, C_{3-10} cycloalkyl, 3-10 membered heterocyclyl, 3-10 membered heteroaryl, phenyl, C_{3-7}cycloalkylene, or 3-7 membered heterocyclene may be optionally substituted with one or more R_7;

each R_5 is independently selected from the group consisting of halogen, C_{1-6}alkyl, C_{1-6}haloalkyl, C_{1-6}alkylene-$N(R_9)_2$, C_{1-6}alkylene-O-C_{3-10}cycloalkyl, C_{1-6}alkoxy, C_{1-6}alkoxy substituted with C_{3-10}cycloalkyl optionally substituted with one or more halogens, C_{1-6}haloalkoxy, 3-10 membered heterocyclyl optionally substituted with one or more halogens or C_{1-6}alkoxy, 3-10 membered heteroaryl, C_{1-6}alkylene-OH, C_{1-6}alkylene-C_{1-6}alkoxy, OH, $N(R_9)_2$, $-C(O)OR_8$, $C(O)N(R_9)_2$, C_{1-6}alkylene-CN, -CN, $-S(O)_2$-C_{1-6}alkyl, C_{1-6}alkylene-$S(O)_2$-C_{1-6}alkyl, $-S(O)_2$-$N(R_9)_2$, $-OC(O)C_{1-6}$alkyl, -O-C_{3-10}cycloalkyl optionally substituted with one or more halogen or C_{1-6}alkyl, and C_{3-10}cycloalkyl optionally substituted with one or more substituents selected from halogen, C_{1-6}alkyl, and C_{1-6}alkoxy;

n is selected from the group consisting of 0, 1, 2, and 3;

R_7 is each independently selected from the group consisting of phenyl, C_{1-6}alkoxy, -OH, $-N(R_9)_2$, $-NR_9$-SO_2-C_{1-6}alkyl, -O-(C_{1-6}alkylene)-phenyl, C_{3-10}cycloalkyl, $-C(O)OR_8$, $-C(O)N(R_9)_2$,$-NR_{10}C(O)$-R_{11}, -CN, $-S(O)_2$-C_{1-6}alkyl, $-S(O)_2$- $N(R_9)_2$, 3-10 membered heterocyclyl, and 3-10 membered heteroaryl, wherein the phenyl, C_{3-10}cycloalkyl, 3-10 membered heterocyclyl, or 3-10 membered heteroaryl is optionally substituted with one or

more substituents each independently selected from the group consisting of C_{1-6}alkyl, halogen, -OH, C_{1-6}alkoxy, and -N(R_9)$_2$;

R_8 is hydrogen or C_{1-6}alkyl;

each R_9 is independently selected from the group consisting of hydrogen, C_{1-6}alkyl, and -(C_{1-6}alkylene)-OH, or the two R_9 can be taken together with the nitrogen atom attached to the two R_9 to form a heterocycle optionally substituted with one or more substituents each independently selected from halogen and -OH;

each R_{10} is independently hydrogen or C_{1-6}alkyl;

R_{11} is selected from the group consisting of C_{1-6}alkyl, C_{1-6}alkoxy, and -O-(C_{1-6}alkylene)-phenyl; and

when R_3 and R_4 are both hydrogen, at least one selected from X, Y, Z, Y', and Z' is N.

25. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is epilepsy, an epilepsy syndrome, or an encephalopathy.

26. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a genetic or pediatric epilepsy or a genetic or pediatric epilepsy syndrome.

27. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition

associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a cardiac dysfunction.

28. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of epilepsy and other encephalopathies (e.g., epilepsy of infancy with migrating focal seizures (MMFSI, EIMFS), autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE), West syndrome, infantile spasms, epileptic encephalopathy, focal epilepsy, Ohtahara syndrome, developmental and epileptic encephalopathy, Lennox Gastaut syndrome, seizures (e.g., Generalized tonic clonic seizures, Asymmetric Tonic Seizures), leukodystrophy, leukoencephalopathy, intellectual disability, Multifocal Epilepsy, Drug resistant epilepsy, Temporal lobe epilepsy, or cerebellar ataxia).

29. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of cardiac arrhythmia, sudden unexpected death in epilepsy (SUDEP), Brugada syndrome, and myocardial infarction.

30. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from pain and related conditions (e.g. neuropathic pain, acute/chronic pain, migraine).

31. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is a muscle disorder (e.g. myotonia, neuromyotonia, cramp muscle spasms, spasticity).

32. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from itch and pruritis, ataxia and cerebellar ataxias.

33. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from psychiatric disorders (e.g. major depression, anxiety, bipolar disorder, schizophrenia).

34. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder or the disease or condition associated with excessive neuronal excitability and/or a gain-of-function mutation in a gene (e.g., KCNT1) is selected from the group consisting of learning disorders, Fragile X, neuronal plasticity, and autism spectrum disorders.

35. Use of a compound of any one of claims 15-20 or a pharmaceutical composition of any one of claims 1-14 and 21-23, in the manufacture of a medicament for treating a

neurological disease or disorder, a disease or condition associated with excessive neuronal excitability, or a disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1), wherein the neurological disease or disorder, the disease or condition associated with excessive neuronal excitability, or the disease or condition associated with a gain-of-function mutation of a gene (e.g., KCNT1) is selected from the group consisting of epileptic encephalopathy with SCN1A, SCN2A, SCN8A mutations, early infantile epileptic encephalopathy, Dravet syndrome, Dravet syndrome with SCN1A mutation, generalized epilepsy with febrile seizures, intractable childhood epilepsy with generalized tonic-clonic seizures, infantile spasms, benign familial neonatal-infantile seizures, SCN2A epileptic encephalopathy, focal epilepsy with SCN3A mutation, cryptogenic pediatric partial epilepsy with SCN3A mutation, SCN8A epileptic encephalopathy, sudden unexpected death in epilepsy (SUDEP), Rasmussen encephalitis, malignant migrating partial seizures of infancy, autosomal dominant nocturnal frontal lobe epilepsy, KCNQ2 epileptic encephalopathy, and KCNT1 epileptic encephalopathy.