


What is claimed:

1. Use of a ROCK2 inhibitor in the manufacture of a medicament for treating a viral infection in a subject in need thereof.
2. Use of a ROCK2 inhibitor in the manufacture of a medicament for preventing a viral infection in a subject at risk for the viral infection.
3. Use of a ROCK2 inhibitor in the manufacture of a medicament for treating Kawasaki disease, PMIS, or PIMS-TS associated with a viral infection in a subject in need thereof.
4. Use of a ROCK2 inhibitor in the manufacture of a medicament for the treatment or prevention of sequelae resulting from a viral infection in a subject in need thereof.
5. The use of any one of claims 1-4, wherein the ROCK2 inhibitor is a ROCK2 selective inhibitor.
6. The use of any one of claims 1-5, wherein the ROCK2 inhibitor is a ribonucleic acid (RNA).
7. The use of claim 6, wherein the ROCK2 inhibitor is an antisense RNA, a small interfering RNA or a microRNA.
8. The use of any one of claims 1-5, wherein the ROCK2 inhibitor is a compound having the structure of Formula I:



or a pharmaceutically acceptable salt, ester, stereoisomer, polymorph, solvate, N-oxide, or isotopically labeled compound thereof, wherein:

R^1 is selected from the group consisting of $—O—(CH_2)_rCO_2R^{12}$, $—O—(CH_2)_y$, $C(=O)NR^{13}R^{14}$, $—O—(CH_2)_y$, -heteroaryl, $—O—(CH_2)_y$, -cycloalkyl, $—O—C(=O)—(CH_2)_y$, $—NR^{13}R^{14}$, $—O—(CH_2)_z—NR^{13}R^{14}$, $—NH—C(=O)—(CH_2)_y$, $—NR^{13}R^{14}$, $—NH—C(=O)—X—R^{15}$, and $—NH—(CH_2)_y$, $—NR^{13}R^{14}$;

5 R¹² is selected from the group consisting of C₁-C₆ alkyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR¹⁶R¹⁷, —(C₁-C₆ alkyl)-C(=O)NR¹⁶R¹⁷, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted at one or more carbon atoms by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

10 R¹³ and R¹⁴ are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR¹⁶R¹⁷, —(C₁-C₆ alkyl)-C(=O)NR¹⁶R¹⁷, aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₃-C₇ cycloalkyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

15 or R¹³ and R¹⁴ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, C₃-C₇ cycloalkyl, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

X is absent or selected from a -O, NH, and C₁-C₆ alkyl;

20 R¹⁵ is selected from the group consisting of heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

25 or R¹⁵ is selected from —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR¹⁶R¹⁷, —CO₂R¹⁸, —O—(CH₂)_x—CO₂R¹⁸, and —C(=O)NR¹⁶R¹⁷;

30 R¹⁶ and R¹⁷ are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents

independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

or R¹⁶ and R¹⁷ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents

5 independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

R¹⁸ is selected from the group consisting of H, aryl, aralkyl, heteroaryl, C₁-C₆ alkyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR¹⁶R¹⁷, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoroalkyl;

x is selected from 0 to 6;

y is selected from 0 to 6;

z is selected from 2 to 6;

15 each R² is independently selected from the group consisting of lower alkyl, CN, halo, hydroxy, lower alkoxy, amino, and perfluoro lower alkyl;

each R³ is independently selected from the group consisting of lower alkyl, CN, halo, hydroxy, lower alkoxy, amino, and perfluoro lower alkyl;

20 R⁴ is selected from H, —(CH₂)_a—NR⁴³R⁴⁴, —Y—R⁴², —O—(CH₂)_a—CO₂R⁴², —O—(CH₂)_a—C(=O)NR⁴³R⁴⁴, —O—(CH₂)_a-heteroaryl, —O—(CH₂)_a-cycloalkyl, —O—C(=O)—(CH₂)_a—NR⁴³R⁴⁴, —O—(CH₂)_c—N⁴³R⁴⁴, —NH—C(=O)—Y—R⁴⁵, —NH—C(=O)—(CH₂)_a—NR⁴³R⁴⁴;

25 R⁴² is selected from the group consisting of C₁-C₆ alkyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR⁴⁶R⁴⁷, —(C₁-C₆ alkyl)-C(=O)N⁴⁶R⁴⁷, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), each of which may be optionally substituted at one or more carbon atoms by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

5 R⁴³ and R⁴⁴ are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR⁴⁶R⁴⁷, —(C₁-C₆ alkyl)-C(=O)NR⁴⁶R⁴⁷, aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₃-C₇ cycloalkyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

10 or R⁴³ and R⁴⁴ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

Y is absent or selected from O, NH, and C₁-C₆ alkyl;

R⁴⁵ is selected from the group consisting of H, aryl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR⁴⁶R⁴⁷, —CO₂R⁴⁸, —O—(CH₂)₆—CO₂R⁴⁸, and —C(=O)NR⁴⁶R⁴⁷;

15 R⁴⁶ and R⁴⁷ independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

or R⁴⁶ and R⁴⁷ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

25 R⁴⁸ is selected from the group consisting of H, aryl, aralkyl, heteroaryl, C₁-C₆ alkyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR⁴⁶R⁴⁷, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoroalkyl;

a is selected from 0 to 6;

b is selected from 0 to 6;

c is selected from 2 to 6;

5 R^5 is selected from the group consisting of H, C₁-C₆ alkyl, $-(CH_2)_d-C(=O)-$
NR⁵³R⁵⁴, $-C(=O)-(CH_2)_d-NR^{53}R^{54}$, and $-C(=O)-X-R^{55}$;

10 R⁵³ and R⁵⁴ are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, $-(C_1-C_6 \text{ alkyl})-$
NR⁵⁶R⁵⁷, $-(C_1-C_6 \text{ alkyl})-C(=O)NR^{56}R^{57}$, aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may
be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₃-C₇ cycloalkyl, C₁-C₆ alkoxy, hydroxy, amino, cyano
and C₁-C₃ perfluoro alkyl;

15 or R⁵³ and R⁵⁴ may be taken together to form 3-10 membered heterocyclic ring having
up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents
independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, C₃-
C₇ cycloalkyl, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

R⁵⁵ is selected from the group consisting of H, aryl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$,
 $-(C_1-C_6 \text{ alkyl})-NR^{56}R^{57}$, $-CO_2R^{58}$, $-O-(CH_2)_e-CO_2R^{58}$, and $-C(=O)NR^{56}R^{57}$;


20 R⁵⁶ and R⁵⁷ independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-
C₈ alkenyl, C₂-C₈ alkynyl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms,
each of which may be optionally substituted by from 1 to 3 substituents independently
selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, hydroxy, amino, cyano
and C₁-C₃ perfluoro alkyl;

25 or R⁵⁶ and R⁵⁷ may be taken together to form 3-10 membered heterocyclic ring having
up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents
independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo,
hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

- 5 R^{58} is selected from the group consisting of H, aryl, aralkyl, heteroaryl, C_1 - C_6 alkyl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, $-(C_1-C_6 \text{ alkyl})-NR^{56}R^{57}$, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C_1 - C_6 alkoxy, hydroxy, amino, cyano and C_1 - C_3 perfluoroalkyl;
- d is selected from 0 to 6;
- e is selected from 0 to 6;
- R^6 is selected from the group consisting of H, C_1 - C_6 alkyl, $-(CH_2)_r-C(=O)-NR^{63}R^{64}$, $-C(=O)-(CH_2)_r-NR^{63}R^{64}$, and $-C(=O)-X-R^{65}$;
- 10 R^{63} and R^{64} are independently selected from the group consisting of H, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, $-(C_1-C_6 \text{ alkyl})-NR^{66}R^{67}$, $-(C_1-C_6 \text{ alkyl})-C(=O)NR^{66}R^{67}$, aryl, aralkyl, heteroaryl, C_3 - C_7 cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo,
- 15 C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_3 - C_7 cycloalkyl, C_1 - C_6 alkoxy, hydroxy, amino, cyano and C_1 - C_3 perfluoro alkyl;
- or R^{63} and R^{64} may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_1 - C_6 alkoxy, C_3 -
- 20 C_7 cycloalkyl, oxo, hydroxy, amino, cyano and C_1 - C_3 perfluoro alkyl;
- R^{65} is selected from the group consisting of H, aryl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, $-(C_1-C_6 \text{ alkyl})-NR^{66}R^{67}$, $-CO_2R$, $-O-(CH_2)$, $-CO_2R^{68}$, and $-C(=O)NR^{66}R^{67}$;
- R^{66} and R^{67} independently selected from the group consisting of H, C_1 - C_5 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $-(C_1-C_6 \text{ alkyl})-O-(C_1-C_6 \text{ alkyl})$, aryl, aralkyl, heteroaryl,
- 25 C_3 - C_7 cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_1 - C_6 alkoxy, hydroxy, amino, cyano and C_1 - C_3 perfluoro alkyl;


or R⁶⁶ and R⁶⁷ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

- 5 R⁶⁸ is selected from the group consisting of H, aryl, aralkyl, heteroaryl, C₁-C₆ alkyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR⁶⁶R⁶⁷, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoroalkyl;
- 10 r is selected from 0 to 6;
- s is selected from 0 to 6;
- n is selected from 0 to 4;
- m is selected from 0 to 3; and
- p is selected from 0 and 1.
- 15 9. The use of claim 8, wherein the ROCK2 inhibitor is a compound having the structure of Formula II:



or a pharmaceutically acceptable salt, ester, stereoisomer, polymorph, solvate, N-oxide, or isotopically labeled compound thereof, wherein: R¹, R², R³, R⁵, R⁶, m and n are as defined in claim 8.

10. The use of claim 9, wherein the ROCK2 inhibitor is a compound having the structure of Formula III:



5

or a pharmaceutically acceptable salt, ester, stereoisomer, polymorph, solvate, N-oxide, or isotopically labeled compound thereof, wherein:

10 R^{13} and R^{14} are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), —(C₁-C₆ alkyl)-NR¹⁶R¹⁷, —(C₁-C₆ alkyl)-C(=O)NR¹⁶R¹⁷, aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₃-C₇ cycloalkyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

15 or R^{13} and R^{14} may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, C₃-C₇ cycloalkyl, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;


20 R^{16} and R^{17} are independently selected from the group consisting of H, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, —(C₁-C₆ alkyl)-O—(C₁-C₆ alkyl), aryl, aralkyl, heteroaryl, C₃-C₇ cycloalkyl, 3-10 membered heterocyclic ring containing up to 3 heteroatoms, each of which may be optionally substituted by from 1 to 3 substituents

independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;

or R¹⁶ and R¹⁷ may be taken together to form 3-10 membered heterocyclic ring having up to 3 heteroatoms which is optionally substituted by from 1 to 3 substituents independently selected from halo, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₁-C₆ alkoxy, oxo, hydroxy, amino, cyano and C₁-C₃ perfluoro alkyl;


wherein each of R², R³, R⁵, R⁶, m and n are as defined in claim 8.

11. The use of claim 10, wherein the ROCK2 inhibitor is a compound having the structure of Formula IV:



or a pharmaceutically acceptable salt, ester, stereoisomer, polymorph, solvate, N-oxide, or isotopically labeled compound thereof, wherein: R¹³ and R¹⁴ are as defined in claim 10.

12. The use of claim 11, wherein the ROCK2 inhibitor is the compound (2-(3-(4-((1H-indazol-5-yl)amino)quinazolin-2-yl)phenoxy)-N-isopropylacetamide), having the chemical structure of Compound 1:



or a pharmaceutically acceptable salt, ester, stereoisomer, polymorph, solvate, N-oxide, or isotopically labeled compound thereof.

- 5 13. The use according to any one of claims 1 to 12, wherein the viral infection is caused by a coronavirus.
14. The use according to any one of claims 1 to 13, wherein the viral infection is caused by SARS-CoV-1, SARS-CoV-2 or MERS-CoV.
- 10 15. The use according to any one of claims 1 to 14, wherein the viral infection is caused by SARS-CoV-2.
16. The use according to claim 15, wherein the viral infection is caused by a Delta or
- 15 Omicron variant of SARS-CoV-2.
17. The use according to any one of claims 4 to 16, wherein the sequelae resulting from the viral infection include one or more of the group consisting of fatigue, dyspnea, cough, arthralgia, myalgia, headache, chest pain, fever, palpitations, myocardial
- 20 inflammation, ventricular dysfunction, stroke, pulmonary function abnormalities, pulmonary fibrosis, renal dysfunction rash, alopecia, olfactory and/or gustatory dysfunction, sleep dysregulation, cognitive impairment altered, memory impairment, depression, anxiety, changes in mood, and combinations thereof.
- 25 18. The use according to any one of claims 4 to 17, wherein the sequelae is pulmonary fibrosis.
19. The use according to any one of claims 1 to 18, wherein the medicament is to be administered with at least one other therapeutic agent.
- 30 20. The use according to claim 19, wherein the at least one other therapeutic agent is selected from the group consisting of an antiviral agent, a corticosteroid, an anti-inflammatory signal transduction modulator, a β 2-adrenoreceptor agonist bronchodilator, an anticholinergic, a mucolytic agent, hypertonic saline, and
- 35 combinations thereof.

21. The use according to claim 19, wherein the at least one other therapeutic agent comprises remdesivir.